

Research

Open Access

## High mobility group A2 is a target for miRNA-98 in head and neck squamous cell carcinoma

Carla Hebert<sup>1</sup>, Kathleen Norris<sup>1</sup>, Mark A Scheper<sup>2</sup>, Nikolaos Nikitakis<sup>2</sup> and John J Sauk<sup>\*2</sup>

Address: <sup>1</sup>Diagnostic Sciences and Pathology, University of Maryland Baltimore, Baltimore Maryland 21201-1586, USA and <sup>2</sup>Diagnostic Sciences and Pathology, and Greenebaum Cancer Center, University of Maryland Baltimore, Baltimore Maryland 21201-1586, USA

Email: Carla Hebert - [Chebert@umaryland.edu](mailto:Chebert@umaryland.edu); Kathleen Norris - [Knorris@umaryland.edu](mailto:Knorris@umaryland.edu); Mark A Scheper - [Mscheper@umaryland.edu](mailto:Mscheper@umaryland.edu); Nikolaos Nikitakis - [Nnikitakis@umaryland.edu](mailto:Nnikitakis@umaryland.edu); John J Sauk\* - [Jsauk@umaryland.edu](mailto:Jsauk@umaryland.edu)

\* Corresponding author

Published: 14 January 2007

Received: 27 August 2006

*Molecular Cancer* 2007, **6**:5 doi:10.1186/1476-4598-6-5

Accepted: 14 January 2007

This article is available from: <http://www.molecular-cancer.com/content/6/1/5>

© 2007 Hebert et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** HMGA2 expression has been shown to be associated with enhanced selective chemosensitivity towards the topoisomerase (topo) II inhibitor, doxorubicin, in cancer cells. Although the roles of signaling cascades and proteins as regulatory factors in development, neoplasia and adaptation to the environment are becoming well established, evidence for the involvement of regulatory small RNA molecules, such as microRNAs (miRNAs) as important regulators of both transcriptional and posttranscriptional gene silencing is presently mounting.

**Results:** Here we report that HMGA2 expression in head and neck squamous cell carcinoma (HNSCC) cells is regulated in part by miRNA-98 (miR-98). Albeit HMGA2 is associated with enhanced selective chemosensitivity towards topoisomerase (topo) II inhibitor, doxorubicin in HNSCC, the expression of HMGA2 is thwarted by hypoxia. This is accompanied by enhanced expression of miRNA-98 and other miRNAs, which predictably target HMGA2. Moreover, we show that transfection of pre-miR-98<sup>TM</sup> during normoxia diminishes HMGA2 and potentiates resistance to doxorubicin and cisplatin. These findings implicate the role of a miRNA as a key element in modulating tumors in variable microenvironments.

**Conclusion:** These studies validate the observation that HMGA2 plays a prominent role in governing genotoxic responses. However, this may only represent cells growing under normal oxygen tensions. The demonstration that miRNA profiles are altered during hypoxia and repress a genotoxic response indicates that changes in microenvironment in eukaryotes mimic those of lower species and plants, where, for example, abiotic stresses regulate the expression of thousands of genes in plants at both transcriptional and posttranscriptional levels through a number of miRNAs and other small regulatory RNAs.

### Background

High mobility group A2 (HMGA2) protein is a non-histone architectural transcription factor, which is a member

of the HMGA family. This family is constituted by HMGA1a, HMGA1b, HMGA1c, and HMGA2[1,2]. HMGA2 is a chromosomal protein that binds through

their AT binding motifs or AT hooks to the minor groove of AT-rich DNA strands[3]. The two types of HMGA protein, HMGA1 (HMG-I/Y) and HMGA2 (HMGI-C), have similar functions and are coded for by two different genes at chromosomal loci 6p21.3 and 12q15, respectively. HMGA proteins are expressed predominantly during embryogenesis, their expression being absent or greatly reduced in adult tissues [4,5]. However, HMGA proteins have been implicated in the regulation of transcription, differentiation, neoplastic transformation, and integration and expression of viral genomes [6,7]. HMGA2 has been implicated mainly in benign and malignant mesenchymal tumors[8] Nevertheless, increased expression of HMGA2 in oral squamous cell carcinoma has been linked to epithelial-mesenchymal transition, whereby epithelial cells acquire mesenchymal characteristics [8]. Additionally, HMGA2 expression in patients with breast cancer has been shown to be associated with poor prognosis and metastasis [9]. However, the precise role of and the molecular events elicited by HMGA2 in tumorigenesis still need to be defined. Furthermore, HMGA2 expression has been shown to be associated with enhanced selective chemosensitivity towards the topoisomerase (topo) II inhibitor, doxorubicin, in breast cancer HS578T and salivary Pa-4/HMGA2 cells [2]. These studies have suggested that HMGA2 induces a persistent basal H2AX Ser139 phosphorylation and perturbs doxorubicin-elicited DNA damage checkpoint promoting enhanced chemosensitivity towards doxorubicin treatment in HMGA2-expressing cells.

Although proteins have been established as regulatory factors in growth, development and accommodation to alterations in environment, our knowledge of the contribution of regulatory small RNA molecules to such is rapidly evolving [10]. MicroRNAs (miRNAs) are 20–24 nucleotide non-coding RNA molecules that post-transcriptionally control gene expression. They are spawned by RNA polymerase II as large primary miRNA (pri-miRNA) transcripts [11,12]. Some miRNAs are structured in clusters and transcribed as polycistrons [13], while others exist in introns of protein-coding genes [14] and are co-transcribed with host genes. Pri-miRNA transcripts form typical fold-back structures that are identified and processed into ~70nt imperfect stem-loop miRNA precursors (pre-miRNAs) by a microprocessor complex [15,16]. After processing by the microprocessor complex, pre-miRNAs are exported by Ran-GTP dependent transporter exportin-5 [17-19] from the nucleus into cytoplasm, where they are processed by the RNase III enzyme, Dicer, into imperfect dsRNA duplexes that contain both mature and complementary miRNA strands [20-22]. One of the mature miRNA strands forms an effector ribonucleoprotein complex termed RISC (RNA induced silencing complex) which guides the miRNAs to specific mRNAs [23]. RISC

either cleaves or blocks translation of the target mRNA, depending on the degree of sequence complementarity.

Here we report that HMGA2 expression in head and neck squamous cell carcinoma (HNSCC) cells is regulated in part by the expression of miRNA-98 (miR-98). Albeit HMGA2 is associated with enhanced selective chemosensitivity towards topoisomerase (topo) II inhibitor, doxorubicin in HNSCC, the expression of HMGA2 is thwarted by hypoxia. This is accompanied by enhanced expression of miRNA-98 and other miRNAs, which predictably target HMGA2. Moreover, we show that transfection of pre-miR-98™ during normoxia diminishes HMGA2 and potentiates resistance to doxorubicin and cisplatin. These findings implicate the role of a miRNA as a key element in modulating tumors in variable microenvironments.

## Materials and methods

### Cell cultures

All experiments were performed using established cell lines, including primary human oral squamous cell carcinoma (SCC) cell lines from the American Type Culture Collection (SCC-4 and SCC-9), and the UMB-10B cell line (UMB). SCC cell lines were cultured in 1:1 mix of Ham's F12 and Dulbecco's Modified Eagle's Medium (DMEM) with 10% fetal bovine serum, 100 units of penicillin, 100 µg/ml streptomycin and 0.4 g/ml of hydrocortisone (Sigma Chemical Company). Cell lines were subcultured using a disaggregation assay with Trypsin (0.1%) and EDTA (0.01%) in phosphate buffered saline (PBS) pH 7.5. For some experiments cell lines were made hypoxic by placing them in a Heto-Holten Cell house incubator at 1% O<sub>2</sub>, 5% CO<sub>2</sub> and 94% nitrogen for periods up to 1 hr or 5% O<sub>2</sub>, 5% CO<sub>2</sub> and 90% nitrogen for periods up to 8 hr. All agents were administered by syringe in a closed system for hypoxia experiments to avoid reoxygenation effects. The oxygen levels for all systems were monitored using an Instech FO/SYS21OT micro oxygen monitoring system (Instech Labs., Inc.).

### Antibodies

Polyclonal monospecific mouse anti-HMGA2 antibody was obtained from ABNOVA. Monoclonal anti-actin antibody was from Chemicon. HRP-goat-antimouse secondary antibody was obtained from Kirkegaard & Perry Labs. Inc.

### RNA isolation, labeling and miRNA microarray

For these studies RNA was obtained using *mirVana*™ miRNA Isolation Kit (Ambion). The levels and quality of RNA was achieved by the Agilent Bioanalyzer™ and with a spectrophotometer using 230 nm/260 nm ratios. The dendrimer Cy3 and Cy5 dyes were then employed to label RNA sequences. First, poly-A tails were added to the RNA sequences at 3' ends using poly(A) polymerase. A nucle-

otide tag was then added to the poly-A tail in a ligation reaction. In the case of dual-sample experiments, the two sets of RNA sequences were added with tags of two different sequences. The tagged RNA sequences were hybridized to the array with labeling carried out during the second hybridization reaction using dendrimer dyes. The labeled samples were then hybridized over night to a Human\_V7.1C\_051017 miRNA array chip (LC Sciences) using a micro circulation pump. After the hybridization, the chips were subjected to a stringent wash and fluorescence data were collected by using an Axon laser scanner model 4000B (Axon Instruments) and the chips were scanned at a pixel size of 10  $\mu$ M with Cy3 Gain at 460 and the Cy5 Gain at 470 scanning. Data extraction and image process were performed using ArrayPro™ software equipped with a morphological filter (Media Cybernetics).

#### **Selection of plausible miR targets**

To select plausible targets to validate the significance of the detected *miRs* we utilized 1) miRNA viewer [24], which is based on miRanda software for miRNA prediction of potential targets (The default parameters were set at Gao Open Penalty:-8.0; Gap Extend: -2.0; Score Threshold: 50.0; Energy Threshold -20.0 kCal/mol; Scaling Parameter: 2.0) 2) *miRBase Targets* from the Sanger Institute, 3) TargetScanS, and 4) PicTar, which identifies both binding-sites targeted by single microRNA, as well as those that are co-regulated by several microRNAs in a coordinated manner. In that miRNAs may have a multitude of conserved miRNA:UTR pairs, recognizing that hypoxia has been associated with drug resistance, the general principle for these studies was to seek possible target genes that influenced DNA repair, drug action, or resistance.

#### **3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assays for measurement of cell viability**

For these studies, cell lines were seeded into 24-well plates to obtain a confluence of 50–60% on the day of the experiment. The cell lines were treated with various reagents of indicated concentration and medium was changed daily for 3 days. Twenty-four to seventy-two hours after the start of treatment (depending on the cell line), 50  $\mu$ l of 5 mg/mL 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT; Roche) in OptiMEM I (Invitrogen) was added to each well and the plate was incubated at 37°C for an additional 4 hours and 10% SDS in .01 M HCl was added to each well to dissolve the formazan crystals. Absorbance was immediately read at 575 nm in a scanning multiwell spectrophotometer. The results were depicted as percentage of cell viability, reported as the mean  $\pm$  SD of three independent experiments done in triplicate.

#### **Protein lysate preparation and Western blot analysis**

The cell lines were plated in 6-well plates using a density of  $5 \times 10^4$  cells/well and were allowed to grow to 80% confluence. The cell lines were washed twice with cold PBS, lysed in RIPA buffer (50 mM Tris [pH 7.4], 150 mM NaCl, 1% Triton X-100, 1% deoxycholic acid, sodium salt, 0.1% sodium dodecyl sulfate [SDS], 100  $\mu$ g/ml phenylmethylsulfonyl fluoride, 1  $\mu$ g/ml aprotinin, 1 mM dithiothreitol and 1 mM sodium orthovanadate) for 10 min and scraped. The extracts were centrifuged at 40,000 g for 15 min at 4°C. Protein concentrations were measured and equalized using Bio-Rad protein assay (Bio-Rad Laboratories) according to the manufacturer's instructions. Equivalent amounts of protein (50  $\mu$ g) were then separated by SDS-PAGE and then transferred to polyvinylidene difluoride membranes. Equivalent loading was confirmed by staining membranes with Ponceau-S. The membranes were blocked for 1 h in blocking buffer (1 $\times$  Tris-buffered saline, 5% nonfat dry milk, and 0.1% Tween 20), which was subsequently replaced by the primary antibody in blocking buffer, overnight at 4°C. After incubation, the membranes were washed three times in washing buffer (1 $\times$  Tris-buffered saline and 0.1% Tween 20). Primary antibody was detected using horseradish peroxidase-linked goat antimouse (Kirkegaard & Perry Labs. Inc.) or goat antirabbit IgG antibody (Kirkegaard & Perry Labs. Inc.) and visualized with SuperSignal West Pico chemiluminescent substrate. To validate loading of the gels the membranes were washed with PBS/Tween, treated with a stripping buffer (20 mM DTT, 2% SDS, 67.5 mM Tris, pH 6.7) rewash with PBS/Tween, blocked with 10% non-fat dried milk, incubated with anti- $\beta$ -actin antibodies and developed as described above. The bands were scanned and quantified using NIH image software.

#### **Transfection of small interfering RNA**

For transient transfection, cell lines were seeded at 90% confluency, and transfections were carried out by using Lipofectamine 2000 (Invitrogen) according to the instructions of the manufacturer. The oligonucleotides encoding HMGA2 small interfering RNA (siRNA) were 5'-CAGCAATCTGTCGCTAAGGdTdT-3' and 5'-CCITAGCGACAGATTGCTGdTdT-3'. In some studies SCC cell lines grown in D-MEM (GIBCO) supplemented with 10% fetal bovine serum (GIBCO) were transfected with 15 or 5 pmol of Pre-miR-98™ or negative control 1 Precursor miRNAs (Ambion) in 24-well plates using siPort Neo-FX™ (Ambion) according to the manufacturer's protocol. Three days posttransfection, cell lines were analyzed for viability and by Western blot. To study the effects of Anti-miR-98™, cell lines grown as above were transfected in 24-well plates with 30 pmol of Anti-miR-98™ or negative control 1 inhibitors (Ambion) using Lipofectamine 2000. Three days post transfection the cell lines were analyzed by vitality assays, by Northern blot and Western blot.

### Reverse transcription-PCR for HMGA2 and miRNA-98 and Northern blot validation of miRNA-98

RNA was extracted using Trizol reagent (Invitrogen) according to the protocols of the manufacturer followed by DNA-free DNase treatment (Ambion, Austin, TX). Subsequent cDNA synthesis and PCR reactions were carried out using ThermoScript reverse transcription-PCR (RT-PCR) System (Invitrogen). The following primer pairs were used for PCR reaction: HMGA2, 5'-GTGAGCCCTCTCCTAAGAGAC-3' and 5'-CTGCAGTGTCTTCTCCCTTC-3'; glyceraldehyde-3-phosphate dehydrogenase (GAPDH), 5'-GACCACAGTCCATGCCATCAC-3' and 5'-CATACCAGAAATGAGCTTGAC-3'. To validate miRNA by PCR, RNA was extracted with the *mirVana*<sup>™</sup> miRNA Isolation Kit and the *mirVana* qRT-PCR miRNA<sup>™</sup> Detection kit for miRNA-98 and 5S (Ambion). The levels of miRNA-98 were determined using *mirVana*<sup>™</sup> Probe with oligonucleotide positive controls; 5S rRNA and an RNA probe specific for miR-16 miRNA following the manufactures protocols.

### Statistical analysis

Statistical analyses utilized two-tailed Student's test; statistical significance was set as  $p \leq 0.05$ .


### Results and discussion

Poor oxygenation (hypoxia) is present in the majority of human tumors and is associated with poor prognosis due to the protection it affords to radiotherapy and chemotherapy [25]. Here, we sought first to verify the chemoresistance effect of hypoxia to doxorubicin and cisplatin in SCC cell lines, comparing cell lines grown in normoxic and hypoxic atmospheres. These studies revealed all SCC cell lines studied here using MTT cell vitality responded in a dose-dependent manner to doxorubicin. Moreover, when the cell lines were grown under hypoxic conditions, they demonstrated enhanced cell viability (Figure 1). Similar results were also observed for cisplatin (not shown). The mechanisms for chemoresistance to these agents under hypoxia were initially unclear. Existing knowledge dictates that tumors cope with hypoxic microenvironments by bringing forth multiple cellular response pathways that alter gene expression and affect tumor progression by strongly suppressing the rates of mRNA translation during hypoxia [25].

Although the roles of signaling cascades and proteins as regulatory factors in development, neoplasia and adaptation to the environment are becoming well established, the evidence for involvement of regulatory small RNA molecules as important regulators of both transcriptional and posttranscriptional gene silencing is presently mounting [10]. Thus, in the present study we grew SCC cell lines both in a normoxic atmosphere and in an environment that mimics the hypoxia encountered in solid head and

neck tumors and used differential display hybridization to a human miRNA chip, and Real-time RT-PCR and Northern blot validation, to discern miRNA expression profiles. SCC cells that were grown under hypoxic conditions differentially expressed 47 miRNA genes compared to cells grown under normoxic conditions ( $P < 0.04$ ). The following miRNAs were expressed to the greatest extent during hypoxia: hsa-miRNA-572, -214, -563, -637, -628, -191, -210, -498, -373, -98, -148b, -15a, -148-a, -200a, -30b, -429, -7, hsa-let-7e, -7i, and -7g (Table 1). Conversely, the following miRNAs were most diminished in their expression in hypoxic cells: hsa-miRNA-122a, -565, -195, -30e-5p, -374, -19a, -101, -424, -29b, -186, -141, -320, -422b, and -197 (Table 1).

One of the most prevalent changes in expression profiles involved a 4.56 fold increased expression of hsa-miR-98. To validate these findings we grew SCC cell lines in either a normoxic or hypoxic environment as before and assessed the miRNA-98 levels with *mirVana* qRT-PCR miRNA<sup>™</sup> Detection kit. These studies revealed that following hypoxia the relative levels of miRNA-98 increased among the SCC cell lines in an oxygen dependent manner (Figure 2). These changes in miRNA-98 were validated in



**Figure 1**  
**Chemosensitivity towards doxorubicin of SCC cell lines.** Percent cell viability following doxorubicin treatment to assess chemosensitivity towards doxorubicin was measured by MTT assays. SCC cell lines were grown under normoxic (▽ SCC-4, ■ SCC-10B and □ SCC-9) or hypoxic (● SCC-4, ○ SCC-10B, and ▼ SCC-9) conditions and treated with various concentrations of doxorubicin. Significant differences between cell lines grown under hypoxic conditions and normoxic conditions was detected for all three cell lines as determined by two-tailed Student's *t* test ( $p \leq 0.05$ ). The error bars represent the S.E.M. for 3 experiments.

**Table 1: miRNAs differentially expressed following hypoxia in SCC cells.**

miRNAs	Hypoxia Up-regulated	miRNAs	Hypoxia Down-regulated
hsa-miRNA-572	6.16	hsa-miRNA-122a	-13.54
hsa-miRNA-214	5.39	hsa-miRNA-565	- 6.77
hsa-miRNA-563	5.17	hsa-miRNA-195	- 6.36
hsa-miRNA-637	4.96	hsa-miRNA-30e-5p	- 5.93
hsa-miRNA-98	4.56	hsa-miRNA-374	- 4.17
hsa-miRNA-628	4.50	hsa-miRNA-19a	- 3.95
hsa-miRNA-191	4.19	hsa-miRNA-101	- 3.73
hsa-miRNA-210	4.16	hsa-miRNA-424	- 3.68
hsa-miRNA-31	3.97	hsa-miRNA-186	- 3.63
hsa-miRNA-498	2.94	hsa-miRNA-29b	- 3.66
hsa-miRNA-373	2.85	hsa-miRNA-148b	- 3.48
hsa-miRNA-19a	2.46	hsa-miRNA-141	- 3.05
hsa-miRNA-148a	2.30	hsa-miRNA-22	- 2.10
hsa-miRNA-15a	2.28	hsa-miRNA-331	- 2.02
hsa-miRNA-200a	2.20	hsa-miRNA-422b	- 2.01
hsa-miRNA-7	2.20	hsa-miRNA-197	- 2.00
hsa-miRNA-30b	2.19		
hsa-let-7e	2.26		
hsa-let-7g	2.22		
hsa-let-7i	2.12		


SCC cells that were grown under hypoxic conditions differentially expressed 47 miRNA genes compared to cells grown under normoxic conditions ( $P < 0.04$ ). Data represent the fold changes in miRNA expression. hsa-miRNA-98 represents the 5<sup>th</sup> most highly expressed miRNA during hypoxia.

Northern blot analyses using mirVana miR-98<sup>TM</sup> Probe, normalized to 5S rRNA (not shown).


To further appraise the role of miRNA-98 during normoxia and hypoxia SCC cell lines were transfected with the pre-miR-98<sup>TM</sup> or anti-miR-98<sup>TM</sup> (Ambion) using siPORT<sup>TM</sup> NeoFX<sup>TM</sup> Transfection Agent, for 72 hours. The predicted changes in the levels of miRNA-98 were validated by mirVana qRT-PCR<sup>TM</sup> as before (not shown). SCC cell lines transfected with either anti-miR-98<sup>TM</sup> or with pre-miR-98<sup>TM</sup> were then challenged with varying doses of doxorubicin and the surviving fraction of cells as a percentage of control was determined using the MTT assay. Cell lines grown in normoxic conditions revealed greater percent of surviving cells when treated with pre-miR-98<sup>TM</sup> compared to controls and anti-miR-98<sup>TM</sup> transfected cell lines. There was no significant difference between cell lines treated with anti-miR-98<sup>TM</sup> and controls. Cell lines grown under hypoxic conditions and transfected with pre-miR-98<sup>TM</sup> showed a slight increase in surviving cells compared to control cells and anti-miR-98<sup>TM</sup> transfected cell lines. There was no significant difference between control and anti-miR-98<sup>TM</sup> transfected cell lines (Figure 3a). Similar findings were also extended to other double-strand breaks elicited by cisplatin in SCC cells (Figure 3b). These studies revealed that normoxic cell lines treated with pre-miR-98<sup>TM</sup> acquired chemoresistance to doxorubicin and cisplatin.

Next, we sought to assess potential targets for miRNA-98 that might explain the observed chemoresistance to these

drugs during hypoxia or in pre-miR-98<sup>TM</sup> transfected normoxic cell lines. In that miRNA-98 has 487 conserved miRNA:UTR pairs within the human genome by *miRNA viewer*, we selected potential targets that were predicted



**Figure 2**  
**Relative expression of miRNA-98 in normoxia and hypoxia.** The relative levels of miRNA-98 were determined using the *mirVana*<sup>TM</sup> miRNA Isolation Kit and the *mirVana* qRT-PCR miRNA<sup>TM</sup> Detection kit for miR-98 and 5S (Ambion, Austin, TX). Cell lines were grown under normoxic or hypoxic (5% O<sub>2</sub> and 1% O<sub>2</sub>) conditions. The differences between cell lines grown under normoxic condition were significantly different to those grown at 1% O<sub>2</sub> with a  $P \leq 0.05$ . The error bars represent S.E.M. for 3 experiments.



**Figure 3**

**a. Cell viability of doxorubicin treated SCC cell lines following pretreatment with anti-miR-98<sup>TM</sup> and pre-miR-98<sup>TM</sup>.** The SCC-4 cell line was grown under normoxia or hypoxia and transfected with anti-miR-98<sup>TM</sup> or pre-miR-98<sup>TM</sup>. The cell line was then challenged with varying doses of doxorubicin as described in *Methods*. The surviving fraction of cells as a percentage of control was determined using the MTT assay. Cell lines grown in normoxic conditions revealed greater percent of surviving cells when treated with pre-mir-98<sup>TM</sup> (●) compared to controls (▼) and anti-miR-98<sup>TM</sup> (○) transfected cells. ( $p \leq 0.05$ ). There was no significant difference between cell lines treated with anti-miR-98<sup>TM</sup> and controls. Cell lines grown under hypoxic conditions and transfected with pre-miR-98<sup>TM</sup> (▼) showed a slight increase in surviving cells compared to control cell lines (■) and Anti-miR-98<sup>TM</sup> (□) transfected cell lines ( $p \leq 0.05$ ). There was no significant difference between control and anti-miR-98<sup>TM</sup> transfected cell lines. Similar results were observed for SCC-9 and SCC-10B cell lines. The error bars represent S.E.M. for 3 experiments. **b. Cell viability of cisplatin treated cell lines following pretreatment with anti-miR-98<sup>TM</sup> and pre-miR-98<sup>TM</sup>.** The SCC-4 cell line was grown under normoxic or hypoxic conditions and transfected with anti-miR-98<sup>TM</sup> while others were transfected with pre-miR-98<sup>TM</sup>. The cell line was then challenged with varying doses of cisplatin as described in *Methods*. The surviving fraction of cells as a percentage of control was determined using the MTT assay. Cells grown in normoxic conditions revealed greater percent of surviving cells when treated with pre-mir-98<sup>TM</sup> (▼) compared to controls (●) and anti-miR-98<sup>TM</sup> (○) transfected cell lines ( $p \leq 0.05$ ). There was no significant difference between cells treated with anti-miR-98<sup>TM</sup> and controls. Cell lines grown under hypoxic conditions and transfected with pre-miR-98<sup>TM</sup> (■) showed no significant difference in the percent of surviving cells compared to control cells (○) and Anti-miR-98<sup>TM</sup> (□) transfected cell lines. Likewise, there was no significant difference between control and anti-miR-98<sup>TM</sup> transfected cell lines. Similar results were observed for SCC-9 and SCC-10B cell lines. The error bars represents S.E.M. for 3 experiments.

from at least two of the database resources with at least two miRNA target sites. Utilizing *miRNA viewer*, the high mobility group protein HMGI-C (High mobility group AT-hook 2, HMGA2) was one of the highest ranking genes with five potential target sites for *hsa-miRNA-98*. Utilizing

other databases it was found that *PicTar* defined 6 *hsa-miRNA-98* target sites, and *TargetScanS* identified 7 *hsa-miRNA-98* target sites (Figure 4). Notable was that a number of other miRNAs included in the hypoxia profile also were potential regulators for HMGA2. Most prevalent

Position	Predicted Target Region (top) and miRNA98 (bottom)	Seed Match
Position 3-27 of HMGA2 3' UTR	5' ... CCAACG-UUCGAUUU-CUACCUCA...    :   : :         3'  UUGUUAUGUUGAAUGRUGGAGU	8mer
Position 1088-1113 of HMGA2 3' UTR	5' ... ACCUGAAUACCACU--UACCUCA...                   3'  UUGUUAUGUUGAAUGRUGGAGU	7mer-1A
Position 1230-1262 of HMGA2 3' UTR	5' ... GCAAUAUAUCCCAUCUCAAUAUCUCCUCU... :    :              3'  UUGUUAUGU-----UGA---AUGRUGGAGU	7mer-m8
Position 1600-1625 of HMGA2 3' UTR	5' ... AUUACCCUCCAAGUCUGUACCUCAA...               3'  UUGUUAUGUUGAAUGRUGGAGU	7mer-1A
Position 1650-1674 of HMGA2 3' UTR	5' ... UUGAC-UUGCAAAGACCUACCUCC... :    : :         3'  UUGUUAUGUUGAAUGRUGGAGU	7mer-m8
Position 2502-2527 of HMGA2 3' UTR	5' ... CAAUCAAACACAC-UACURCCUCU...                    3'  UUGUUAUGU-UGAAUGRUGGAGU	7mer-m8
Position 2515-2547 of HMGA2 3' UTR	5' ... ACUACUACCUCUUAAGUCCAGUAURCCUCA...                   3'  UUGUUAUGUUGAAU-----GRUGGAGU	7mer-1A

**Figure 4**

**TargetScanS miRNA98 target sites for HMGA2.** In addition to 7mer sites, TargetScanS predicts 8mer sites defined as: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) with a downstream 'A' across from position 1 of the miRNA. In addition to 8mer sites, TargetScanS predicts 7mer sites of two types: 7mer-m8: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) and 7mer-1A: An exact match to positions 2–7 of the mature miRNA (the seed) with a downstream 'A' across from position 1 of the miRNA.

among this later group utilizing *TargetScanS* were hsa-let-7g which possessed 7 target sites for HMGA2 (Figure 5), hsa-let-7e with 7 target sites (Figure 6) and hsa-let-7i with 7 sites (Figure 7). Albeit, the expression of has-let-7g, -7i, and -7i was similar to miRNA-98, being low in normoxia, but increasing following hypoxia (Table 1).

Recognizing that HMGA2 has been shown to perturb doxorubicin-elicited DNA damage checkpoint control and to promote enhanced chemosensitivity towards doxorubicin and cisplatin treatment in HMGA2-expressing breast cancer cells [2], we assessed whether hypoxia altered the expression of HMGA2; SCC cell lines were subjected to hypoxia, harvested and the expression of HMGA2 was determined by Western blot. These studies revealed that hypoxia diminished the levels of HMGA2 protein expression (Figure 8a).

To verify that HMGA2 potentiated genotoxic stress in SCC cell lines we paralleled our studies to those performed in breast cancer cells [2]. In essence, cell lines were transfected with siRNA oligonucleotides for HGMA2 as described above. The oligonucleotides encoding HMGA2 siRNA was compared with an oligonucleotide scramble siRNA as control [2]. These studies revealed that HMGA2 siRNA induced down-regulation of HMGA2 protein expression as assessed by Western blot (Figure 8b lower

panel), and diminished doxorubicin sensitivity by MTT assays (Figure 8c). Recognizing that SCC cell lines became resistant to doxorubicin during periods of hypoxia and following diminished HMGA2 expression after siRNA transfection in normoxic cells we sought to verify that HMGA2 was a feasible target for miRNA-98 in SCC cell lines. To further appraise HMGA2 as a miRNA-98 target, SCC cell lines were transfected with the pre-miR-98<sup>TM</sup> (Ambion) using siPORT<sup>TM</sup> NeoFX<sup>TM</sup> Transfection Agent, for 72 hours. The increased levels of miRNA-98 were validated by miRVana qRT-PCR<sup>TM</sup> as before. These treatments resulted in diminishment in protein levels and mRNA expression of HMGA2 in control the normoxic SCC-4 cell line as well as cell lines exposed to hypoxia (Figure 9a,b); similar results were obtained for SCC-10B and SCC-9 cell lines (not shown). The demonstration that pre-miRNA-98<sup>TM</sup> diminished mRNA expression of HMGA2 was surprising, since the tenet of early studies, using reporter constructs had indicated that complete complementarities between miRNAs and target mRNAs was necessary for cleavage, whereas mismatches result in translational repression. However, the findings reported here are supported by more recent research, which revealed that miRNAs with partial complementarities can induce attenuation in cognate transcript levels [26-29]. In fact, most recently it has been shown that the ability of miRNAs to expedite poly(A) removal does not result from

Position	Predicted Target Region (top) and has-let-7e (bottom)	Seed Match
Position 4-27 of HMGA2 3' UTR	5' ... CAAC-GUUCGAUUU- <b>CU<sup>1</sup>RC<sup>2</sup>CU<sup>3</sup>CA</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	8mer
Position 1089-1113 of HMGA2 3' UTR	5' ... GACCUGAAUA-CCACU <b>RC<sup>1</sup>CU<sup>2</sup>CA</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	7mer-1A
Position 1239-1262 of HMGA2 3' UTR	5' ... CCACUACUCAAU-- <b>AC<sup>1</sup>UC<sup>2</sup>CU<sup>3</sup>CU</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	7mer-m8
Position 1593-1625 of HMGA2 3' UTR	5' ... UUUAAAUAUA <sup>1</sup> CCCU <sup>2</sup> C <sup>3</sup> CAAGUCU <b>RC<sup>4</sup>CU<sup>5</sup>CA</b> A... 3' UGUAUUG-UUGGAGG----- <b>AUGGAGU</b>	7mer-1A
Position 1651-1674 of HMGA2 3' UTR	5' ... UGACU-UGCAAAG <b>AC<sup>1</sup>UC<sup>2</sup>CU<sup>3</sup>CU</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	7mer-m8
Position 2495-2527 of HMGA2 3' UTR	5' ... AUGUUGUACAAUCAAAACA <b>CA<sup>1</sup>CU<sup>2</sup>CC<sup>3</sup>UC<sup>4</sup>U</b> ... 3' UGUAUUGUUUGG----- <b>AGGUGGAGU</b>	7mer-m8
Position 2515-2547 of HMGA2 3' UTR	5' ... ACUACUACCUUAAGUCCCA <b>GU<sup>1</sup>RC<sup>2</sup>CU<sup>3</sup>CA</b> ... 3' UGUAUUGUUGGAGG----- <b>AUGGAGU</b>	7mer-1A

**Figure 5 TargetScanS has-let-7g target sites for HMGA2.** In addition to 7mer sites, TargetScanS predicts 8mer sites defined as: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) with a downstream 'A' across from position 1 of the miRNA. In addition to 8mer sites, TargetScanS predicts 7mer sites of two types: 7mer-m8: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) and 7mer-1A: An exact match to positions 2–7 of the mature miRNA (the seed) with a downstream 'A' across from position 1 of the miRNA.

decreased translation; nor does translational repression by miRNAs require a poly(A) tail, with a 3' histone stem-loop being an effective substitute [30].

Conversely, the use of the anti-miR-98™ provided little information in the present studies. Cell lines grown under normoxia express little miRNA-98 so that the effects of

Position	Predicted Target Region (top) and has-let-7e (bottom)	Seed Match
Position 4-27 of HMGA2 3' UTR	5' ... CAAC-GUUCGAUUU- <b>CU<sup>1</sup>RC<sup>2</sup>CU<sup>3</sup>CA</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	8mer
Position 1089-1113 of HMGA2 3' UTR	5' ... GACCUGAAUA-CCACU <b>RC<sup>1</sup>CU<sup>2</sup>CA</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	7mer-1A
Position 1239-1262 of HMGA2 3' UTR	5' ... CCACUACUCAAU-- <b>AC<sup>1</sup>UC<sup>2</sup>CU<sup>3</sup>CU</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	7mer-m8
Position 1593-1625 of HMGA2 3' UTR	5' ... UUUAAAUAUA <sup>1</sup> CCCU <sup>2</sup> C <sup>3</sup> CAAGUCU <b>RC<sup>4</sup>CU<sup>5</sup>CA</b> A... 3' UGUAUUG-UUGGAGG----- <b>AUGGAGU</b>	7mer-1A
Position 1651-1674 of HMGA2 3' UTR	5' ... UGACU-UGCAAAG <b>AC<sup>1</sup>UC<sup>2</sup>CU<sup>3</sup>CU</b> ... 3' UGUAUUGUUUGGAG <b>GRUGGAGU</b>	7mer-m8
Position 2495-2527 of HMGA2 3' UTR	5' ... AUGUUGUACAAUCAAAACA <b>CA<sup>1</sup>CU<sup>2</sup>CC<sup>3</sup>UC<sup>4</sup>U</b> ... 3' UGUAUUGUUUGG----- <b>AGGUGGAGU</b>	7mer-m8
Position 2515-2547 of HMGA2 3' UTR	5' ... ACUACUACCUUAAGUCCCA <b>GU<sup>1</sup>RC<sup>2</sup>CU<sup>3</sup>CA</b> ... 3' UGUAUUGUUGGAGG----- <b>AUGGAGU</b>	7mer-1A

**Figure 6 TargetScanS has-let-7e target sites for HMGA2.** In addition to 7mer sites, TargetScanS predicts 8mer sites defined as: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) with a downstream 'A' across from position 1 of the miRNA. In addition to 8mer sites, TargetScanS predicts 7mer sites of two types: 7mer-m8: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) and 7mer-1A: An exact match to positions 2–7 of the mature miRNA (the seed) with a downstream 'A' across from position 1 of the miRNA.



Position	Predicted Target Region (top) and has-let-7i (bottom)	Seed Match
Position 1-27 of HMGA2 3' UTR	5' ... GGGCCCAACGUUC GAUUCURCCUC... : 11:1 111            111111  3' UGUCGUGUUUG-----AUGRUGGAGU	8mer
Position 1081-1113 of HMGA2 3' UTR	5' ... UCCAUGAAA GACCUGAAUACCACUURCCUCRA...    11 11            111111  3' UGUCGUGUUUGAUG-----RUGGAGU	7mer-1A
Position 1239-1262 of HMGA2 3' UTR	5' ... CCCACUACUCAAUACURCCUCU... 11 : 11    11111111  3' UGUCGUGUUUGAUGRUGGAGU	7mer-m8
Position 1600-1625 of HMGA2 3' UTR	5' ... UUAACCCUCAAAGUCUG-URCCUCRA... 111: 11:  111111  3' UGUCGUGUUU-GAUGRUGGAGU	7mer-1A
Position 1651-1674 of HMGA2 3' UTR	5' ... GACUUGCAAAGAC--CURCCUC... 111 1:11  111111  3' UGUCGUGUUUGAUGRUGGAGU	7mer-m8
Position 2495-2527 of HMGA2 3' UTR	5' ... AUGUUGUACAAUCAAAACACUACURCCUCU... 1: 1111            11111111  3' UGUCGUGUU-----UGAUGRUGGAGU	7mer-m8
Position 2523-2547 of HMGA2 3' UTR	5' ... UCCAGUA-----URCCUCRU... 111:1            111111  3' UGUCGUGUUUGAUGRUGGAGU	7mer-1A

**Figure 7**

**TargetScanS has-let-7i target sites for HMGA2.** In addition to 7mer sites, TargetScanS predicts 8mer sites defined as: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) with a downstream 'A' across from position 1 of the miRNA. In addition to 8mer sites, TargetScanS predicts 7mer sites of two types: 7mer-m8: An exact match to positions 2–8 of the mature miRNA (the seed + position 8) and 7mer-1A: An exact match to positions 2–7 of the mature miRNA (the seed) with a downstream 'A' across from position 1 of the miRNA.


anti-mir-98<sup>TM</sup> on HMGA2 levels were limited (Figure 9a,b). Transfection with anti-miR-98<sup>TM</sup> had only minor effects on the HMGA2 levels in SCC cell lines grown in hypoxia (Figure 9a,b). This is probably owing to the increased expression during hypoxia of other targeting miRNAs such as *hsa-let-7g*, *hsa-let-7e* and *hsa-let-7i* all of which possess 7 complementary target sites for HMGA2. Inherent in these findings is the recognition that derepression of a mRNA targeted by a miRNAs in an altered micro-environment probably can not be achieved through down-regulation of a single miRNA in that most mRNAs possess multiple complementary target sites with a number of miRNAs.

Until lately, there was only speculation that inhibition by a miRNA to a specific mRNA could be effectively reversed. However, recent studies have shown that CAT-1 mRNA and reporters bearing its 3'UTR can be relieved from miR-122-mediated repression in human Huh7 hepatoma cells when they are subjected to different types of stress. Moreover, that the derepression is accompanied by the release of CAT-1 mRNA from cytoplasmic processing bodies and its entry into polysomes and that the process involves binding of HuR, *trans*-acting AU rich element (ARE) binding protein, to the 3'UTR of CAT-1 mRNA [31]. There is also additional evidence indicating that some miRNA-controlled mRNAs can be relieved from repression by synaptic stimulation [32,33] indicating that miRNA regulation is more pronounced than previously anticipated and that it is able to respond quickly to specific cellular needs.

What is yet to be determined is whether changes in oxygen tension provide a like mechanism for derepression of miRNAs against HMGA2. There is clear evidence that HuR, a major ARE-binding protein is expressed in many different cells and regulates the stability, subcellular shuttling, and/or translation of ARE-mRNAs (reviewed [34]). Moreover, the level of cytoplasmic HuR increases during cell proliferation, differentiation, and in response to stress stimuli [35-38] decreasing with senescence [39]. Also, the high level of cytoplasmic HuR in cancer cells is in keeping with the ability of HuR to increase the stability and translation of ARE-mRNAs that regulate cell proliferation and neovascularization by stabilizing VEGF following hypoxia. HuR has also been shown to shuttle between nucleus and cytoplasm during hypoxia and to stabilize HIF-1 $\alpha$  and EGFR mRNAs [34]. Consequently, we searched the AERD database [40] but were not able to demonstrate any of the five clusters containing adenylate uridylylate (AU)-rich elements in their 3'-untranslated region (3'-UTR) that would afford HMGA2 stabilization or derepression by HuR. Furthermore, since HuR shuttles from the nucleus to the cytoplasm during hypoxia it would appear that HMGA2 would be overwhelmed by HuR, making it unlikely that HMGA mRNA is stabilized like other ARE-mRNAs.


## Conclusion

These studies validate the observation that HMGA2 plays a prominent role in governing genotoxic responses. However, this may only represent cells growing under normal

**Figure 8**

**a. Effect of hypoxia and normoxia on the protein expression of HMGA2 by Western blot.** SCC-4, SCC-9 and SCC-10B cell lines were grown under normoxic and hypoxic conditions (1%  $\text{O}_2$ ), harvested and the levels of HMGA2 assayed by Western blot, as described in *Methods*. **b. Effect of HMGA2 small interfering RNA on cell viability following doxorubicin treatment in SCC cell lines.** The SCC-4 cell line grown under normoxic conditions was transfected with oligonucleotide encoding HMGA2 small interfering RNA or a scramble siRNA sequence. The dose dependent effects of doxorubicin on cell viability was measured by MTT assays. The differences between control cell line and cell lines treated with HMGA2 siRNA was statistically significant ( $p \leq 0.01$ ) as determined by Student's *t* test. The panel below depicts the effects of HMGA2 siRNA and the scramble counterpart on HMGA2 in Western blot. Similar results were obtained for SCC-9 and SCC-10B cell lines. The error bars represent the S.E.M. for 3 experiments.

oxygen tensions. The demonstration that miRNA profiles are altered during hypoxia and repress a genotoxic response indicates that changes in microenvironment in eukaryotes mimic lower species and plants, where, for example, abiotic stresses regulate the expression of thousands of genes in plants at both transcriptional and post-transcriptional levels through a number of miRNAs and other small regulatory RNAs[41]. Yet to be established is

**Figure 9**

**a. The effect of pre-miR-98<sup>TM</sup> and anti-miR-98<sup>TM</sup> on the protein levels of HMGA2 during normoxia and hypoxia.** The SCC-4 cell line was transfected with pre-miR-98<sup>TM</sup> or Anti-miR-98<sup>TM</sup> and grown under hypoxic or normoxic conditions. Similar results were obtained for SCC-10B and SCC-9 cell lines (not shown). **b. The effect of pre-miR-98<sup>TM</sup> and anti-miR-98<sup>TM</sup> on the relative expression of HMGA2 during normoxia and hypoxia.** The mRNA expression of HMGA2 was significantly diminished both during normoxia and hypoxia following transfection with pre-miR-98<sup>TM</sup> (\*\* $p \leq 0.01$ ). Anti-miR98<sup>TM</sup> transfection had a significant effect on enhancing HMGA2 expression when cells lines were grown during normoxia (\* $p \leq 0.05$ ) but were ineffective under hypoxic conditions. The mRNA expression of the control levels of each condition was set as 1. The error bars represents S.E.M. for 3 experiments.

to what extent miRNA expression is regulated by defined signaling cascades and transcription factors.

#### Authors' contributions

CH performed Western blots, Gene transfection

KN maintained Cell Cultures and performed Drug assays

MS performed miRNA target analysis

NN carried out data analysis and was co-writer

JJ initiated research design performed data analysis and was a co-writer

## Acknowledgements

The research was supported in part by a grant from U.P.H.S./N.I.D.C.R./N.I.H. 5R01DE012606-07

## References

- Reeves R: **Molecular biology of HMGA proteins: hubs of nuclear function.** *Gene* 2001, **277**:63-81.
- Boo LM, Lin HH, Chung V, Zhou B, Louie SG, O'Reilly MA, Yen Y, Ann DK: **High mobility group A2 potentiates genotoxic stress in part through the modulation of basal and DNA damage-dependent phosphatidylinositol 3-kinase-related protein kinase activation.** *Cancer Res* 2005, **65**:6622-6630.
- Reeves R, Beckerbauer L: **HMGI/Y proteins: flexible regulators of transcription and chromatin structure.** *Biochim Biophys Acta* 2001, **1519**:13-29.
- Zhou X, Benson KF, Przybys K, Liu J, Hou Y, Cherath L, Chada K: **Genomic structure and expression of the murine Hmgi-c gene.** *Nucleic Acids Res* 1996, **24**:4071-4077.
- Hirring-Folz U, Wilda M, Rippe V, Bullerdiek J, Hameister H: **The expression pattern of the Hmgi-c gene during development.** *Genes Chromosomes Cancer* 1998, **23**:350-357.
- Sarhadi VK, Wikman H, Salmenkivi K, Kuosma E, Sioris T, Salo J, Karjalainen A, Knuutila S, Anttila S: **Increased expression of high mobility group A proteins in lung cancer.** *J Pathol* 2006, **209**:206-212.
- Sgarra R, Rustighi A, Tessari MA, Di Bernardo J, Altamura S, Fusco A, Manfioletti G, Giancotti V: **Nuclear phosphoproteins HMGA and their relationship with chromatin structure and cancer.** *FEBS Lett* 2004, **574**:1-8.
- Rogalla P, Drechsler K, Frey G, Hennig Y, Helmke B, Bonk U, Bullerdiek J: **HMGI-C expression patterns in human tissues. Implications for the genesis of frequent mesenchymal tumors.** *Am J Pathol* 1996, **149**:775-779.
- Wikman H, Kettunen E, Seppanen JK, Karjalainen A, Hollmen J, Anttila S, Knuutila S: **Identification of differentially expressed genes in pulmonary adenocarcinoma by using cDNA array.** *Oncogene* 2002, **21**:5804-5813.
- Carrington JC, Ambros V: **Role of microRNAs in plant and animal development.** *Science* 2003, **301**:336-338.
- Cai X, Hagedorn CH, Cullen BR: **Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs.** *Rna* 2004, **10**:1957-1966.
- Lee Y, Kim M, Han J, Yeom KH, Lee S, Baek SH, Kim VN: **MicroRNA genes are transcribed by RNA polymerase II.** *Embo J* 2004, **23**:4051-4060.
- Lee Y, Jeon K, Lee JT, Kim S, Kim VN: **MicroRNA maturation: stepwise processing and subcellular localization.** *Embo J* 2002, **21**:4663-4670.
- Rodriguez A, Griffiths-Jones S, Ashurst JL, Bradley A: **Identification of mammalian microRNA host genes and transcription units.** *Genome Res* 2004, **14**:1902-1910.
- Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ: **Processing of primary microRNAs by the Microprocessor complex.** *Nature* 2004, **432**:231-235.
- Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R: **The Microprocessor complex mediates the genesis of microRNAs.** *Nature* 2004, **432**:235-240.
- Bohnsack MT, Czaplinski K, Gorlich D: **Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs.** *Rna* 2004, **10**:185-191.
- Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U: **Nuclear export of microRNA precursors.** *Science* 2004, **303**:95-98.
- Yi R, Qin Y, Macara IG, Cullen BR: **Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs.** *Genes Dev* 2003, **17**:3011-3016.
- Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G, Mello CC: **Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control C. elegans developmental timing.** *Cell* 2001, **106**:23-34.
- Hutvagner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, Zamore PD: **A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA.** *Science* 2001, **293**:834-838.
- Ketting RF, Fischer SE, Bernstein E, Sijen T, Hannon GJ, Plasterk RH: **Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in C. elegans.** *Genes Dev* 2001, **15**:2654-2659.
- Bhattacharya A, Toth K, Mazurchuk R, Sperry JA, Slocum HK, Pendyala L, Azrak R, Cao S, Durrani FA, Rustum YM: **Lack of microvessels in well-differentiated regions of human head and neck squamous cell carcinoma A253 associated with functional magnetic resonance imaging detectable hypoxia, limited drug delivery, and resistance to irinotecan therapy.** *Clin Cancer Res* 2004, **10**:8005-8017.
- Website: [<http://cbio.mskcc.org/cgi-bin/mirnaviewer/mirnaviewer.pl>].
- Koumenis C, Wouters BG: **"Translating" Tumor Hypoxia: Unfolded Protein Response (UPR)-Dependent and UPR-Independent Pathways.** *Mol Cancer Res* 2006.
- Bagga S, Bracht J, Hunter S, Massirer K, Holtz J, Eachus R, Pasquinelli AE: **Regulation by let-7 and lin-4 miRNAs results in target mRNA degradation.** *Cell* 2005, **122**:553-563.
- Lim LP, Lau NC, Garrett-Engel P, Grimson A, Schelter JM, Castle J, Bartel DP, Linsley PS, Johnson JM: **Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs.** *Nature* 2005, **433**:769-773.
- Yu Z, Raabe T, Hecht NB: **MicroRNA Mirn 122a reduces expression of the posttranscriptionally regulated germ cell transition protein 2 (Tnp2) messenger RNA (mRNA) by mRNA cleavage.** *Biol Reprod* 2005, **73**:427-433.
- Martin SE, Caplen NJ: **Mismatched siRNAs downregulate mRNAs as a function of target site location.** *FEBS Lett* 2006, **580**:3694-3698.
- Wu L, Fan J, Belasco JG: **MicroRNAs direct rapid deadenylation of mRNA.** *Proc Natl Acad Sci U S A* 2006, **103**:4034-4039.
- Bhattacharyya SN, Habermacher R, Martiny-Bar C, Closs EI, Filipowicz W: **Relief of microRNA-mediated translational repression in human cells subjected to stress.** *Cell* 2006, **125**:1111-1124.
- Ashraf SI, McLoon AL, Sclarsic SM, Kunes S: **Synaptic protein synthesis associated with memory is regulated by the RISC pathway in Drosophila.** *Cell* 2006, **124**:191-205.
- Schratt GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, Kiebler M, Greenberg ME: **Corrigendum: A brain-specific microRNA regulates dendritic spine development.** *Nature* 2006, **441**:902.
- Sheflin LG, Zou AP, Spaulding SW: **Androgens regulate the binding of endogenous HuR to the AU-rich 3'UTRs of HIF-1 alpha and EGF mRNA.** *Biochem Biophys Res Commun* 2004, **322**:644-651.
- Ming XF, Stoecklin G, Lu M, Looser R, Moroni C: **Parallel and independent regulation of interleukin-3 mRNA turnover by phosphatidylinositol 3-kinase and p38 mitogen-activated protein kinase.** *Mol Cell Biol* 2001, **21**:5778-5789.
- Wang W, Caldwell MC, Lin S, Furneaux H, Gorospe M: **HuR regulates cyclin A and cyclin B1 mRNA stability during cell proliferation.** *Embo J* 2000, **19**:2340-2350.
- Wang W, Yang X, Lopez de Silanes I, Carling D, Gorospe M: **Increased AMP:ATP ratio and AMP-activated protein kinase activity during cellular senescence linked to reduced HuR function.** *J Biol Chem* 2003, **278**:27016-27023.
- Atasoy U, Watson J, Patel D, Keene JD: **ELAV protein HuA (HuR) can redistribute between nucleus and cytoplasm and is upregulated during serum stimulation and T cell activation.** *J Cell Sci* 1998, **111 (Pt 21)**:3145-3156.
- Briata P, Ilengo C, Corte G, Moroni C, Rosenfeld MG, Chen CY, Gherzi R: **The Wnt/beta-catenin->Pitx2 pathway controls the turnover of Pitx2 and other unstable mRNAs.** *Mol Cell* 2003, **12**:1201-1211.
- Bakheet T, Williams BR, Khabar KS: **ARED 3.0: the large and diverse AU-rich transcriptome.** *Nucleic Acids Res* 2006, **34**:D111-4.
- Sunkar R, Zhu JK: **Novel and stress-regulated microRNAs and other small RNAs from Arabidopsis.** *Plant Cell* 2004, **16**:2001-2019.