

Editorial



OPEN ACCESS

Received: Aug 10, 2019

Accepted: Aug 13, 2019

Correspondence to

Young-Hoon Jeong, MD, PhD

Department of Internal Medicine,
Gyeongsang National University School
of Medicine and Cardiovascular Center,
Gyeongsang National University Changwon
Hospital, 11 Samjeongja-ro, Seongsan-gu,
Changwon 51472, Korea.
E-mail: goodoctor@naver.com

Copyright © 2019. The Korean Society of
Cardiology

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License ([https://
creativecommons.org/licenses/by-nc/4.0](https://creativecommons.org/licenses/by-nc/4.0))
which permits unrestricted noncommercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Young-Hoon Jeong 
<https://orcid.org/0000-0003-0403-3726>

Conflict of Interest

Dr. Jeong has received honoraria for lectures
from AstraZeneca, Sanofi-Aventis, Daiichi
Sankyo, Otsuka, Han-mi Pharmaceuticals
and Yuhan Pharmaceuticals; and research
grants or support from AstraZeneca, Han-mi
Pharmaceuticals, Yuhan Pharmaceuticals, and
Haemonetics.

Dr. Gurbel reports serving as a consultant
fees/receiving honoraria from Daiichi Sankyo,
Bayer, AstraZeneca, Merck, Boehringer, New
Haven Pharmaceuticals, Janssen, and CSL and
receiving grants from the National Institutes
of Health, Daiichi Sankyo, CSL, AstraZeneca,
Harvard Clinical Research Institute,

Temporal Variability of Platelet Reactivity Phenotype: Another Barrier to Personalized Antiplatelet Strategy Guided by Platelet Function Testing

Young-Hoon Jeong , MD, PhD¹, Udaya S. Tantry, PhD², and Paul A. Gurbel, MD³

¹Department of Internal Medicine, Gyeongsang National University School of Medicine and Cardiovascular Center, Gyeongsang National University Changwon Hospital, Changwon, Korea

²Sinai Center for Thrombosis Research, Sinai Hospital of Baltimore, Baltimore, MD, USA

³Inova Center for Thrombosis Research and Drug Development, Inova Heart and Vascular Institute, Fairfax, VA, USA

► See the article “Temporal Variability of Platelet Reactivity in Patients Treated with Clopidogrel or Ticagrelor” in volume 49 on page 1052.

Dual antiplatelet therapy (DAPT) with aspirin and P2Y₁₂ receptor inhibitor is the standard treatment in patients with acute coronary syndrome (ACS) or undergoing percutaneous coronary intervention (PCI).¹⁾ The availability of different P2Y₁₂ receptor inhibitors (clopidogrel, prasugrel and ticagrelor) with different levels of antiplatelet potency has enabled physicians to adopt individualized antiplatelet treatment. In addition, the development of point-of-care platelet function testing (PFT) to measure platelet reactivity to adenosine diphosphate can be fit for use in clinical practice. Like the therapeutic range of international normalized ratio (between 2 and 3) during warfarin treatment, consensus and clinical studies have suggested the therapeutic range of platelet reactivity during P2Y₁₂ inhibitor by applying uniform cutoff values of high- and low-platelet reactivity (HPR and LPR) (Table 1).¹⁾ In the near future, a tailored DAPT approach guided by PFT (escalation or de-escalation) may be an important way to precision medicine.

Available PFTs can be classified as point-of-care assays (VerifyNow, Multiplate, thromboelastography) and laboratory-based methods (light transmission aggregometry, vasodilator-stimulated phosphoprotein).¹⁾ The point-of-care assays would be preferred in terms of practical aspect, but selection of assays needs to depend on their availability and local site experience. Although simple application with consensus-defined cutoffs of HPR/LPR can be a practical guide for tailored antiplatelet treatment, confirmative clinical evidences are required to strengthen the concept of a therapeutic window in PCI-treated patients. The cutoffs for HPR/LPR can be changed according to the clinical setting (stable

Table 1. Consensus cutoffs to determine HPR and LPR¹⁾

Device	HPR cutoff	LPR cutoff
VerifyNow assay (PRU)	>208	<85
Multiplate analyzer (U)	>46	<19
VASP-P assay (PRI, %)	>50	<16
TEG platelet mapping assay (mm)	>47	<31

HPR = high platelet reactivity; LPR = low platelet reactivity; PRI = platelet reactivity index; PRU = P2Y₁₂ reaction units; TEG = thromboelastography; VASP = vasodilator-stimulated phosphoprotein.

Haemonetics, New Haven Pharmaceuticals, Duke Clinical Research Institute, Sinnowa, and Coramed. Dr. Gurbel has patents in the field of platelet function testing.

The other author reports no conflicts of interest.

Author Contributions

Writing - original draft: Jeong YH; Writing - review & editing: Tantry US, Gurbel PA.

The contents of the report are the author's own views and do not necessarily reflect the views of the *Korean Circulation Journal*.

angina vs. ACS), the phase of the disease entity (early vs. late phase), the complexity of PCI (simple vs. complex PCI) and the risk for bleeding complications. Of importance, how to adjust antiplatelet regimen in high-risk patients out of the therapeutic window may affect the consequent clinical outcomes remains to be verified. In addition, the optimal timing of PFT to predict post-PCI events remains a topic of debate. Although single measurements of on-treatment platelet reactivity have been conducted in most of previous clinical studies, its clinical usefulness may be questionable.¹⁾ Patients' phenotype of HPR and LPR may change over time since influencing factors are subject to change over time.²⁻⁵⁾

In this edition of the *Korean Circulation Journal*, Yun et al.⁵⁾ have published their study to evaluate the variability of platelet reactivity over time in ACS patients treated with PCI (n=198) during clopidogrel or ticagrelor administration. Platelet reactivity was measured using multiple electrode platelet aggregometry (Multiplate analyzer; Roche Diagnostic GmbH, Mannheim, Germany), in which HPR was defined as ≥ 47 U, and LPR was defined as ≤ 18 U.¹⁾ Platelet reactivity in the clopidogrel group increased over time (38.2 ± 21.7 U at 48 hours vs. 44.7 ± 25.5 U at 6 months; $p=0.018$), whereas this value in the ticagrelor group was not significantly changed (21.4 ± 12.6 U at 48 hours vs. 22.8 ± 13.8 U at 6 months; $p=0.392$). Between 48 hours and 6 months, 43% of patients changed their response status (HPR or normal response) in the clopidogrel group, and 13% in the ticagrelor group ($p<0.001$). They concluded that ticagrelor treatment resulted in less temporal variability of platelet reactivity than clopidogrel treatment in terms of HPR.

Although this study covers the important issue regarding tailored strategy of antiplatelet treatment, there are several important concerns to be pointed. Because the investigators did not evaluate the degree of drug adherence and change of concomitant medication over time, there would be the risks of unrecognized noncompliance and change in drug-drug interaction. In addition, they did not check time interval between last-dose administrations and blood sampling for measurement. The ONSET/OFFSET (Onset and Offset of Antiplatelet Effects Comparing Ticagrelor, Clopidogrel, and Placebo With Aspirin) study showed that platelet reactivity during P2Y₁₂ inhibitor administration can be variable over 8 hours after the last-dose administration, especially reversible-binding ticagrelor.⁶⁾ More importantly, the levels of platelet reactivity during the acute (at 48 hours) vs. chronic phase (at 6 months) can be different. In addition, the activity of disease entity (stable angina vs. ACS) and administered dose of study drug (loading vs. maintenance) could influence the level of platelet reactivity at the measuring point. Between 1 and 6 months, platelet reactivity on clopidogrel was not significantly changed (41.4 ± 22.3 U vs. 44.7 ± 25.5 U; $p=0.256$), and this value on ticagrelor did not differ significantly (20.0 ± 12.2 U vs. 22.8 ± 13.8 U; $p=0.071$) in the present study. Previous studies also showed the similar level of on-clopidogrel platelet reactivity and low prevalence of change in the response status (6.5–15.7%) during the chronic phase.²⁻⁴⁾

Potent P2Y₁₂ inhibitors inevitably increased the risk of major bleeding in ACS patients, compared with clopidogrel.^{7,8)} Meanwhile, the antiplatelet effect on ticagrelor vs. prasugrel treatment appears greater in ACS patients.⁹⁾ Platelet reactivity at 30-day post-PCI in patients on ticagrelor was lower than those on prasugrel following acute myocardial infarction (21.1 ± 26.1 vs. 67.3 ± 62.5 P2Y₁₂ reaction units measured by VerifyNow; $p<0.001$). In the present study, ticagrelor treatment almost abolished the risk of HPR, but more than 50% of the patients had LPR phenotype (≤ 18 U).⁵⁾ Absolute increase of non-CABG TIMI major bleeding (+0.6%) seemed similar between ticagrelor and prasugrel, compared with




Figure 1. Conceptual therapeutic window of platelet reactivity during irreversible and reversible P2Y₁₂ receptor inhibitors.⁶⁾

clopidogrel.⁷⁾⁸⁾ Compared with the short plasma exposure of thienopyridine (prasugrel and clopidogrel) active metabolites (up to 6 hours), ticagrelor has significant 24-hour systemic exposure of a direct active compound (reversible-binding property)⁶⁾; this unique profile of ticagrelor may be characterized by its fast and strong platelet inhibition with a wider therapeutic window (**Figure 1**). The dedicated clinical studies to indicate the different cutoffs of LPR during thienopyridine or ticagrelor treatment are needed to guide the tailored strategy to target the therapeutic window of platelet reactivity.

Specific considerations may be warranted for East Asian patients, who have a different benefit/risk ratio compared with the Caucasian population.¹⁾¹⁰⁾ During antithrombotic treatment, East Asian patients have shown a lower risk of atherothrombotic event and a higher tendency of bleeding; this phenomenon was first denominated in 2011 by Jeong and coinvestigators who developed the concept of “East Asian Paradox”, suggesting a different therapeutic targeting of antiplatelet effect in East Asian patients.¹⁰⁾ In East Asian population with ACS, reduced-dose ticagrelor and prasugrel may provide more acceptable clinical efficacy and safety comparing with standard-dose treatment, but there are no large-scale prospective studies to confirm the clinical benefit of this antiplatelet strategy.

The experience has accumulated over the past decade on studies of PFT and its use has been proposed as optional methods to aid clinical decision making in the selective scenarios.¹⁾ Although the results of proof-of-concept studies made a guided choice of antiplatelet regimen attractive, the robustness of the evidence and limitations of “therapeutic window” concept based on PFT (e.g., temporal variability of platelet reactivity over time) still may not allow routine recommendation of PFT use in clinical practice. Nevertheless, escalation strategy may be required when thrombotic risk outweighs bleeding risk (e.g., clopidogrel treatment after complex PCI), whereas de-escalation strategy may be desired when bleeding risk outweighs thrombotic risk (e.g., chronic ticagrelor treatment in East Asian patients with ACS). The results of ongoing clinical trials will further fine-tune the personalized strategy of P2Y₁₂ inhibitor in high-risk patients undergoing PCI.

REFERENCES

1. Sibbing D, Aradi D, Alexopoulos D, et al. Updated expert consensus statement on platelet function and genetic testing for guiding P2Y₁₂ receptor inhibitor treatment in percutaneous coronary intervention. *JACC Cardiovasc Interv* 2019;12:1521-37.
[CROSSREF](#)
2. Jaitner J, Stegherr J, Morath T, et al. Stability of the high on-treatment platelet reactivity phenotype over time in clopidogrel-treated patients. *Thromb Haemost* 2011;105:107-12.
[PUBMED](#) | [CROSSREF](#)
3. Campo G, Parrinello G, Ferraresi P, et al. Prospective evaluation of on-clopidogrel platelet reactivity over time in patients treated with percutaneous coronary intervention relationship with gene polymorphisms and clinical outcome. *J Am Coll Cardiol* 2011;57:2474-83.
[PUBMED](#) | [CROSSREF](#)
4. Hochholzer W, Ruff CT, Mesa RA, et al. Variability of individual platelet reactivity over time in patients treated with clopidogrel: insights from the ELEVATE-TIMI 56 trial. *J Am Coll Cardiol* 2014;64:361-8.
[PUBMED](#) | [CROSSREF](#)
5. Yun KH, Cho JY, Rhee SJ, Oh SK. Temporal variability of platelet reactivity in patients treated with clopidogrel or ticagrelor. *Korean Circ J* 2019;49:1052-61.
[PUBMED](#) | [CROSSREF](#)
6. Gurbel PA, Bliden KP, Butler K, et al. Randomized double-blind assessment of the ONSET and OFFSET of the antiplatelet effects of ticagrelor versus clopidogrel in patients with stable coronary artery disease: the ONSET/OFFSET study. *Circulation* 2009;120:2577-85.
[PUBMED](#) | [CROSSREF](#)
7. Wallentin L, Becker RC, Budaj A, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2009;361:1045-57.
[PUBMED](#) | [CROSSREF](#)
8. Wiviott SD, Braunwald E, McCabe CH, et al. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2007;357:2001-15.
[PUBMED](#) | [CROSSREF](#)
9. Alexopoulos D, Xanthopoulou I, Mavronasiou E, et al. Randomized assessment of ticagrelor versus prasugrel antiplatelet effects in patients with diabetes. *Diabetes Care* 2013;36:2211-6.
[PUBMED](#) | [CROSSREF](#)
10. Huo Y, Jeong YH, Gong Y, et al. 2018 update of expert consensus statement on antiplatelet therapy in East Asian patients with ACS or undergoing PCI. *Sci Bull (Beijing)* 2019;64:166-79.
[CROSSREF](#)