

# Polyphasic taxonomy of *Aspergillus* section *Fumigati* and its teleomorph *Neosartorya*

R.A. Samson<sup>1\*</sup>, S. Hong<sup>2</sup>, S.W. Peterson<sup>3</sup>, J.C. Frisvad<sup>4</sup> and J. Varga<sup>1,5</sup>

<sup>1</sup>CBS Fungal Biodiversity Centre, Uppsalalaan 8, NL-3584 CT Utrecht, The Netherlands; <sup>2</sup>Korean Agricultural Culture Collection, NIAB, Suwon, 441-707, Korea; <sup>3</sup>Microbial Genomics and Bioprocessing Research Unit, National Center for Agricultural Utilization Research, 1815 N. University Street, Peoria, IL 61604, U.S.A.; <sup>4</sup>BioCentrum-DTU, Building 221, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark; <sup>5</sup>University of Szeged, Faculty of Science and Informatics, Department of Microbiology, P.O. Box 533, H-6701 Szeged, Hungary

\*Correspondence: Robert A. Samson, [r.samson@cbs.knaw.nl](mailto:r.samson@cbs.knaw.nl)

**Abstract:** The taxonomy of *Aspergillus* section *Fumigati* with its teleomorph genus *Neosartorya* is revised. The species concept is based on phenotypic (morphology and extrolite profiles) and molecular ( $\beta$ -tubulin and calmodulin gene sequences) characters in a polyphasic approach. Four new taxa are proposed: *N. australensis* *N. ferenczii*, *N. papuaensis* and *N. warcupii*. All newly described and accepted species are illustrated. The section consists of 33 taxa: 10 strictly anamorphic *Aspergillus* species and 23 *Neosartorya* species. Four other *Neosartorya* species described previously were not available for this monograph, and consequently are relegated to the category of doubtful species.

**Taxonomic novelties:** *Neosartorya australensis*, *N. ferenczii*, *N. papuaensis*, *N. warcupii*.

**Key words:** *Aspergillus* section *Fumigati*, extrolite profiles, *Neosartorya*, phylogenetics, polyphasic taxonomy.

## INTRODUCTION

*Aspergillus* section *Fumigati* includes species characterised by uniseriate aspergilli, columnar conidial heads in shades of green and flask shaped vesicles (Raper & Fennell 1965). Teleomorphic species belonging to the “*Aspergillus fischeri* series” of the *A. fumigatus* group (Raper & Fennell 1965) were placed in the genus *Neosartorya* (family Trichocomaceae) by Malloch & Cain (1972). Section *Fumigati* includes more than 20 *Neosartorya* species and 10 anamorphic species (Pitt *et al.* 2000; Samson 2000; Horie *et al.* 2003; Hong *et al.* 2005, 2006, 2007).

*Aspergillus fumigatus* Fresenius is an ubiquitous filamentous fungus in the environment, and also an important human pathogen (Raper & Fennell 1965). Several *Neosartorya* species have been described as causal agents of human diseases including invasive aspergillosis, osteomyelitis, endocarditis and mycotic keratitis (Coriglione *et al.* 1990; Summerbell *et al.* 1992; Padhye *et al.* 1994; Lonial *et al.* 1997; Jarv *et al.* 2004; Balajee *et al.* 2005, 2006). All of the *Neosartorya* species produce heat-resistant ascospores that are frequently encountered in different food products (Gomez *et al.* 1994; Samson 1989; Tournas 1994). The several mycotoxins produced by these species may cause serious health hazard (Fujimoto *et al.* 1993; Frisvad & Samson 1990; Larsen *et al.* 2007). Some species also have valuable properties for mankind; e.g. *N. fischeri* strains produce fiscalins which effectively inhibit the binding of substance P to the human neurokinin receptor (Wong *et al.* 1993), while *A. fumigatus* strains produce pyripyropenes, potent inhibitors of acyl-CoA:cholesterol acyltransferase (Tomoda *et al.* 1994), the immunosuppressant restrictocins (Müllbacher & Eichner 1984), ribotoxins (Lin *et al.* 1995) and fumagillin that has amebicidal activity (McCowen *et al.* 1951). *Neosartorya spinosa*

can be used for the complete enzymatic recovery of ferulic acid from corn residues (Shin *et al.* 2006).

Here we present an overview of the species belonging to *Aspergillus* section *Fumigati* based on analysis of macro- and micromorphology, extrolite profiles and  $\beta$ -tubulin, calmodulin, ITS and actin gene sequences of the isolates. We also describe four new homothallic *Neosartorya* species found in soil samples in Australia and Papua New Guinea using this polyphasic approach and list synonymies.

## MATERIALS AND METHODS

### Source of microorganisms

The fungi examined included type strains or representatives of all species available for examination in *Aspergillus* section *Fumigati*. Some atypical isolates collected in Australia and Papua New Guinea were also examined to clarify their taxonomic status (Table 1).

### Morphology and physiology

The strains (Table 1) were grown for 7 d as 3-point inoculations on Czapek agar, Czapek yeast autolysate agar (CYA), oat meal agar (OA) and malt extract agar (MEA) plates at 25 °C, and on CYA at 37 °C. For *Neosartorya* species Hay infusion agar and SNA agar have also been used for inducing the anamorphs (medium compositions in Samson *et al.* 2004). In some species e.g. *N. tatenoi* the anamorph could only be produced when growing the cultures at 30 or 37 °C on MEA + 40 % sucrose.

**Table 1.** *Aspergillus* section *Fumigati* isolates used in this study.

Species	Isolate No.*	Source
<i>A. brevipes</i>	CBS 118.53 <sup>T</sup>	Soil, Australia
<i>A. duricaulis</i>	CBS 481.65 <sup>T</sup>	Soil, Buenos Aires, Argentina
<i>A. fumigati</i> affinis	IBT12703 <sup>T</sup>	Soil, U.S.A.
<i>A. fumigatus</i>	CBS 133.61 <sup>T</sup> = NRRL 163	Chicken lung, U.S.A.
<i>A. fumisynnematus</i>	IFM 42277 <sup>T</sup>	Soil, Venezuela
<i>A. lentulus</i>	CBS 117887 <sup>T</sup> = NRRL 35552 = KACC 41940	Man, U.S.A.
<i>A. novofumigatus</i>	IBT 16806 <sup>T</sup>	Soil, Ecuador
<i>A. unilateralis</i>	CBS 126.56 <sup>T</sup>	Rhizosphere, Australia
<i>A. viridinutans</i>	CBS 127.56 <sup>T</sup>	Rabbit dung, Australia
<i>A. turcosus</i>	KACC 42090 = IBT 27920	Air conditioner, Inchen, Korea
	KACC 42091 <sup>T</sup> = IBT 27921	Air conditioner, Seoul, Korea
	KACC 41955 = CBS 117265 = IBT 3016	Car air conditioner, Seoul, Korea
<i>N. assulata</i>	KACC 41691 <sup>T</sup>	Tomato soil, Buyeo, Korea
<i>N. aurata</i>	CBS 466.65 <sup>T</sup>	Jungle soil, Brunei
<i>N. aureola</i>	CBS 105.55 <sup>T</sup>	Soil, Tafo, Ghana
<i>N. australensis</i> sp. nov.	CBS 112.55 <sup>T</sup> = NRRL 2392 = IBT 3021	Garden soil, Adelaide, Australia
<i>N. coreana</i>	KACC 41659 <sup>T</sup> = NRRL 35590 = CBS 121594	Tomato soil, Buyeo, Korea
<i>N. denticulata</i>	CBS 652.73 <sup>T</sup> = KACC 41183	Soil under <i>Elaeis guineensis</i> , Suriname
	CBS 290.74 = KACC 41175	<i>Acer pseudoplatanus</i> , Netherlands
<i>N. fennelliae</i>	CBS 598.74 <sup>T</sup>	Eye ball of <i>Oryctolagus cuniculus</i> , U.S.A.
	CBS 599.74	Eye ball of <i>Oryctolagus cuniculus</i> , U.S.A.
<i>N. ferenczii</i> sp. nov.	CBS 121594 <sup>T</sup> = IBT 27813 = NRRL 4179	Soil, Australia
<i>N. fischeri</i>	CBS 544.65 <sup>T</sup> = NRRL 181	Canned apples
<i>N. galapagensis</i>	CBS 117522 <sup>T</sup> = IBT 16756 = KACC 41935	Soil, Ecuador
	CBS 117521 = IBT 16763 = KACC 41936	Soil, Ecuador
<i>N. glabra</i>	CBS 111.55 <sup>T</sup>	Rubber scrub from old tire, Iowa, U.S.A.
<i>N. hiratsukae</i>	CBS 294.93 <sup>T</sup>	Aloe juice, Tokyo, Japan
<i>N. laciniosa</i>	KACC 41657 <sup>T</sup> = NRRL 35589 = CBS 117721	Tomato soil, Buyeo, Korea
<i>N. multiplicata</i>	CBS 646.95 <sup>T</sup> = 'BT 17517	Soil, Mouli, Taiwan
<i>N. nishimurae</i>	IFM 54133 = IBT 29024	Forest soil, Kenya
<i>N. nishimurae</i>	CBS 116047	Cardboard, Netherlands
<i>N. papuensis</i> sp. nov.	CBS 841.96 <sup>T</sup> = IBT 27801	Bark of <i>Podocarpus</i> sp. (Podocarpaceae), bark, Myola, Owen Stanley Range, Northern Province, Papua New Guinea
<i>N. pseudofischeri</i>	NRRL 20748 <sup>T</sup> = CBS 208.92	Human vertebrate, U.S.A.
<i>N. quadricincta</i>	CBS 135.52 <sup>T</sup> = NRRL 2154	Cardboard, York, U.K.
	CBS 107078	Soil, Korea
	CBS 100942	Fruit juice, Netherlands
	CBS 253.94	Canned oolong tea beverage, Japan (type strain of <i>N. primulina</i> )
<i>N. spathulata</i>	CBS 408.89 <sup>T</sup>	Soil under <i>Alocasia macrorrhiza</i> , Taiwan
<i>N. spinosa</i>	CBS 483.65 <sup>T</sup>	Soil, Nicaragua
<i>N. stramenia</i>	CBS 498.65 <sup>T</sup>	Soil from maple-ash-elm forest, Wisconsin, U.S.A.
<i>N. tatenoi</i>	CBS 407.93 <sup>T</sup>	Soil of sugarcane, Timbauba, Brazil
	CBS 101754	Fruit, Yunnan, China (type strain of <i>N. delicata</i> )
<i>N. udagawae</i>	CBS 114217 <sup>T</sup>	Soil, Brazil
	CBS 114218	Soil, Brazil
<i>N. warcupii</i> sp. nov.	NRRL 35723 <sup>T</sup>	Arid soil, Finder"s Range, Australia

\* CBS = Centraalbureau voor Schimmelcultures, Utrecht, the Netherlands; IBT = Institute for Biotechnology, Lyngby, Technical University of Denmark; IFM = Institute for Food Microbiology (at present, the Research Center for Pathogenic Fungi and Microbial Toxicoses, Chiba University), Chiba, Japan; KACC = Korean Agricultural Culture Collection, Suwon, Korea; NRRL = Agricultural Research Service Culture Collection, Peoria, Illinois, U.S.A.; T = type strain.

## Analysis for extrolites

Extrolites were analysed using the HPLC-diode array detection method of Frisvad & Thrane (1987, 1993) as modified by Smedsgaard (1997). Extrolites were analyzed from cultures grown on CYA, OA and YES agar using three agar plugs (Smedsgaard 1997).

## Isolation and analysis of nucleic acids

Isolates used for the molecular studies were grown on 2 mL of malt peptone broth [10 % (v/v) malt extract (Brix 10) and 0.1 % (w/v) bacto peptone (Difco)], in 15 mL tubes. The cultures were incubated at 25 °C for 7 d. DNA was extracted from the cells using the Masterpure™ yeast DNA purification kit (Epicentre Biotechnol.) following the instructions of the manufacturer. Fragments containing the ITS region were amplified using primers ITS1 and ITS4 as described (White *et al.* 1990). Amplification of partial  $\beta$ -tubulin gene was performed using the primers Bt2a and Bt2b and methods of Glass & Donaldson (1995). Amplifications of the partial calmodulin and actin genes were as described (Hong *et al.* 2005, 2007). Sequencing reactions were performed with the Big Dye Terminator Cycle Sequencing Ready Reaction Kit and carried out for both strands. All the sequencing reactions were purified by gel filtration through Sephadex G-50 (Amersham Pharmacia Biotech, Piscataway, NJ) equilibrated in double-distilled water and analyzed on the ABI PRISM 310 Genetic Analyzer (Applied Biosystems). The complementary sequences were corrected with the MT Navigator software (Applied Biosystems). Unique ITS,  $\beta$ -tubulin, actin and calmodulin sequences were deposited in GenBank (<http://www.ncbi.nlm.nih.gov>) with accession numbers DQ534140, DQ534141 and EU20279–EU220287.

## Data analysis

Sequence alignments were performed using CLUSTAL-X (Thompson *et al.* 1997) and improved manually. The neighbour-joining (NJ) method was used for the phylogenetic analysis. For NJ analysis, the data were first analysed using the Tamura–Nei distance calculation with gamma-distributed substitution rates (Tamura & Nei 1993), which were then used to construct the NJ tree with MEGA v. 3.1 (Kumar *et al.* 2004). A bootstrap analysis was performed with 1 000 replications to determine the support for each clade.

PAUP v. 4.0 b10 software was used for parsimony analysis (Swofford 2002). Alignment gaps were treated as a fifth character state and all characters were unordered and of equal weight. Maximum parsimony analysis was performed for all data sets using the heuristic search option with random addition order (100 reps) and tree bisection-reconnection (TBR) branch-swapping algorithm. Branches of zero length were collapsed and all multiple, equally parsimonious trees were saved. The robustness of the trees obtained was evaluated by 1 000 bootstrap replications (Hillis & Bull 1993). Sequences from an *A. clavatus* isolate were used as outgroups in these experiments.

## RESULTS AND DISCUSSION

### Phylogenetic analysis

We examined the phylogenetic relatedness of species belonging to *Aspergillus* section *Fumigati* using sequence analysis of partial  $\beta$ -tubulin, calmodulin and actin genes including sequences of all known species. ITS sequences were determined from the new species and the species most closely related to them in the  $\beta$ -tubulin tree. The partial  $\beta$ -tubulin gene alignment included 453 characters. Among the polymorphic sites, 102 were found to be phylogenetically informative. The Neighbour-joining tree based on partial  $\beta$ -tubulin genes sequences is shown in Fig. 1. The topology of the tree is the same as one of the 419 maximum parsimony trees constructed by the PAUP programME (length: 465 steps, consistency index: 0.6710, retention index: 0.6467). The calmodulin data set included 549 characters with 85 parsimony informative characters. The Neighbour-joining tree shown in Fig. 2 has the same topology as one of the 9 maximum parsimony trees (tree length: 323, consistency index: 0.7585, retention index: 0.6422). The actin data set included 390 characters with 104 parsimony informative characters. The Neighbour joining tree shown in Fig. 3 has the same topology as one of the 312 maximum parsimony trees (tree length: 397, consistency index: 0.6675, retention index: 0.7130). The ITS data set included 501 characters with 26 parsimony informative characters. The Neighbour joining tree shown in Fig. 4 has the same topology as one of the 57 maximum parsimony trees (tree length: 77, consistency index: 0.7532, retention index: 0.7765).

The four *Neosartorya* isolates representing new species were found to be different from all known species of *Aspergillus* section *Fumigati* based on either their  $\beta$ -tubulin, calmodulin or actin gene sequences. However, one of them (NRRL 4179) had identical ITS sequences with *N. denticulata* (Fig. 4). This isolate was found to be closely related to a clade including *N. fennelliae* and *N. denticulata* on all other trees.

Possible synonymies of some species described previously have also been examined during this study. Based on multilocus sequence analyses Hong *et al.* (2007) discussed the synonymy of *N. botucatensis*, *N. paulistensis* and *N. takaki* with *N. spinosa* (Raper & Fennell) Kozak. (1972). *N. spinosa* and the synonyms have roughly circular arrangements of projections on the ascospore convex walls. *N. spinosa* produces echinulate ascospores with spines ranging from < 0.5  $\mu$ m up to 5(–7)  $\mu$ m long with verruculose and small triangular projections or sometimes with circularly arranged projections.

*N. otanii* Takada, Y. Horie & Abliz (2001) was described on the basis of its rapid growth on Czapek and malt extract agars, lenticular ascospores with two widely separated equatorial crests, tuberculate or lobate-reticulate convex surface, and globose to broadly ellipsoidal conidia with a microtuberculate wall. The morphology of *N. otanii* resembles *N. fennelliae*, although Takada *et al.* (2001) reported small differences of the ascospore ornamentation, which was not confirmed in our SEM studies. The  $\beta$ -tubulin gene sequences of *N. otanii* (GenBank accession numbers AB201363 and AB201362) were identical with *N. fennelliae* (KACC 42228) (Fig. 5A). These *N. fennelliae* isolates produced ascospores after mating with the *N. fennelliae* type strains (data not shown). *N. otanii* is probably synonymous with *N. fennelliae*, but mating experiments with *N. fennelliae* and *N. otanii* are needed for its confirmation.




Fig. 1. Neighbour-joining tree based on  $\beta$ -tubulin sequence data of *Aspergillus* section *Fumigati*. Numbers above branches are bootstrap values. Only values above 70 % are indicated.




Fig. 2. Neighbour-joining tree based on calmodulin sequence data of *Aspergillus* section *Fumigati*. Numbers above branches are bootstrap values. Only values above 70 % are indicated.




Fig. 3. Neighbour-joining tree based on actin sequence data of *Aspergillus* section *Fumigati*. Numbers above branches are bootstrap values. Only values above 70 % are indicated.





Fig. 4. Neighbour-joining tree based on ITS sequence data of selected species of *Aspergillus* section *Fumigati*. Numbers above branches are bootstrap values. Only values above 70 % are indicated.



**Fig. 5 A.** Neighbour-joining tree based on  $\beta$ -tubulin sequences showing the relationship of *N. otanii* and *N. fennelliae*. **B.** Neighbour-joining trees based on  $\beta$ -tubulin, calmodulin and actin sequence data of *Neosartorya* spp. showing the relationship of *N. primulina*, *N. quadricincta*, *N. tatenoi* and *N. delicata*.

These experiments could not be carried out because the ex type cultures of *N. otanii* were not available.

*Neosartorya primulina* Udagawa, Toyaz. & Tsub. (1993) was characterised by its restricted growth on Czapek agar, chalky-buff ascospores with a very irregular ornamentation composed of several narrow crests and verrucose hemispheres. The ascospore ornamentation and anamorph morphology resembles those of *N. quadricincta*. Furthermore, the ex type culture (CBS 253.94) of *N. primulina* showed nearly identical sequences with strains of *N. quadricincta* for  $\beta$ -tubulin, calmodulin and actin genes (Fig. 5B). *N. primulina* is reduced to synonymy with *N. quadricincta*.

*Neosartorya delicata* H.Z. Kong (1997) was described based on its ellipsoid or nearly clavate vesicles, and ascospores with

conspicuous spines, joining one spine to another by fairly prominent ridges and reticulate ornamentation, the ridges spreading to the equatorial crests. This species has identical ascospore morphology with *N. tatenoi* (Fig. 36), and both taxa were clustered into a clade in three gene trees (99.6 % in  $\beta$ -tubulin, 98.5 % in calmodulin and 97.3 % in actin gene sequences) (Fig. 5B). Therefore, we consider *N. delicata* as a synonym of *N. tatenoi*.

*Neosartorya nishimurae* (Takada *et al.* 2001), *N. indohii*, *N. tsurutae* (Horie *et al.* 2003), *N. takakii* (Horie *et al.* 2001) and *N. sublevispora* (Someya *et al.* 1999) ex-type cultures were not available for this monograph of *Aspergillus* section *Fumigati*, and because we could not study them, they are listed as doubtful species.

**Table 2.** Extrolites produced by species assigned to *Aspergillus* section *Fumigati*.

Species	Extrolites produced
<i>Aspergillus brevipes</i>	roquefortine C, meleagrins-like
<i>Aspergillus duricaulis</i>	pseurotin A, fumagillin, asperpentyn, duricaulic acid and asperdurin, phthalides, chromanols, cyclopaldic acid, 3-O-methylcyclopolic acid
<i>Aspergillus fumigati</i>	auranthine, cycloechinuline, fumigaclavines, helvolic acid, neosartorin, palitantin, pyripropenes A, E, O & S, tryptoquivalone
<i>Aspergillus fumigatus</i>	fumagillin, fumitoxins, fumigaclavines A & C, fumitremorgins, gliotoxin, trypacidin, pseurotins, helvolic acid, pyripropens, methyl-sulochrin, verruculogen, fumiquinazolines
<i>Aspergillus fumisynnematus</i>	neosartorin, pyripropens, fumimycin
<i>Aspergillus lentulus</i>	cyclopiazonic acid, pyripropenes A, E & O, terrein, auranthine, neosartorin
<i>Aspergillus novofumigatus</i>	aszonalenin, cycloechinuline, fiscalins, helvolic acid, neosartorin, palitantin, terrein, territrem B
<i>Aspergillus turcosus</i>	kotanins and several unique but not yet elucidated secondary metabolites
<i>Aspergillus unilateralis</i>	mycophenolic acid, other unique secondary metabolites
<i>Aspergillus viridutans</i>	viriditoxin, 13-O-methylviriditin, phomalgin A, variotin, viriditin, wasabidienone B0, B1, viriditin, 4-acetyl-6,8-dihydroxy-5-methyl-2-benzopyran-1-1 A
<i>Neosartorya assulata</i>	indole alkaloids and apolar metabolites
<i>Neosartorya aurata</i>	helvolic acid, yellow unidentified compounds
<i>Neosartorya aureola</i>	fumagillin, tryptoquivalone, tryptoquivalone, pseurotin A and viriditoxin (FRR 2269 also produces helvolic acid)
<i>Neosartorya australensis</i>	wortmannin-like, aszonalenin-like
<i>Neosartorya coreana</i>	aszonalenins
<i>Neosartorya denticulata</i>	gliotoxin, viriditoxin
<i>Neosartorya fennelliae</i>	asperfuran, aszonalenin, fumigaclavine, viridicatumtoxin
<i>Neosartorya ferenczii</i>	asperfuran, aszonalenin, fumigaclavine, viridicatumtoxin, gliotoxin-like, fumigatins, aszonalenin-like
<i>Neosartorya fischeri</i>	terrein, fumitremorgins A & C, tryptoquivalone A, trypacidin, TR-2, verruculogen, sarcin, aszonalenins, fischerin, neosartorin, fiscalins, helvolic acid
<i>Neosartorya galapagensis</i>	gregatins
<i>Neosartorya glabra</i>	asperpentyn, avenaciolide, wortmannin-like compound
<i>Neosartorya hiratsukae</i>	avenaciolide
<i>Neosartorya laciniata</i>	aszonalenins, tryptoquivalone, tryptoquivalone
<i>Neosartorya multiplicata</i>	helvolic acid
<i>Neosartorya papuensis</i>	wortmannin-like
<i>Neosartorya pseudofischeri</i>	asperfuran, cytochalasin-like compound, fiscalin-like compound, pyripropens, gliotoxin
<i>Neosartorya quadricincta</i>	quinolactacin, aszonalenins
<i>Neosartorya spinosa</i>	aszonalenins, 2-pyrovoylaminobenzamide, pseurotin
<i>Neosartorya spathulata</i>	xanthocillins, aszonalenins
<i>Neosartorya stramenia</i>	quinolactacin, avenaciolide
<i>Neosartorya tatenoi</i>	aszonalenins
<i>Neosartorya udagawae</i>	fumigatin, fumagillin, tryptoquivalone, tryptoquivalone
<i>Neosartorya warcupii</i>	wortmannin-like, aszonalenin-like, chromanols-like, tryptoquivalone-like and tryptoquivalone-like

## Morphology and extrolite production

The atypical *N. glabra* isolate NRRL 4179 (Raper & Fennell 1965) produced asperfuran, aszonalenin, fumigaclavine, viridicatumtoxin, and fumigatins, extrolites common in *N. fennelliae*, but none of the extrolites produced by *N. glabra*. However, in contrast with the heterothallic *N. fennelliae*, this isolate is homothallic. It is closely related to *N. denticulata* based on phylogenetic analysis of sequence data, although their ascospore ornamentations are strikingly different (Figs. 21, 23). Ascospore ornamentation of NRRL 4179 is similar to that of the heterothallic *N. fennelliae* (Fig. 22) with equatorial crests much narrower, while *N. denticulata* has denticulate ascospores without equatorial crests. Isolate NRRL 4179 exhibited 72 % nuclear DNA relatedness to *N. fennelliae* and only 60 % relatedness to *N. glabra* isolates (Peterson 1992). This isolate also yielded different mtDNA and *Sma*I-digested repetitive DNA patterns from those of all the other *Neosartorya* strains examined (Rinyu *et al.* 2000). Hybridisation experiments were also carried out with *Neurospora crassa* mating type genes (the *A* idiomorph with about 6 kb flanking sequences, or the *a* idiomorph flanked by about 2 kb genomic DNA on either side) to the *Eco*RI digested DNA of several teleomorphic and asexual *Aspergillus* strains. Hybridisation to a 1.9 kb band was observed for both mating-type strains of *N. fennelliae* and isolate NRRL 4179 (Rinyu *et al.* 2000). Based on these observations, isolate NRRL 4179 seems to be closely related to *N. fennelliae* strains. These results are in agreement with those found using carbon source utilisation tests and isoenzyme analysis of these strains (Varga *et al.* 1997).

Strain NRRL 35723 was isolated from soil in Australia, and produced compounds structurally related to wortmannin, aszonalenin, chromanols, tryptoquivalins and tryptoquivalons. This isolate was markedly different from all other known *Neosartorya* species in secreting a bluish pigment after 7 d incubation on MEA and CYA plates. The microtuberculate ascospore ornamentation of this isolate is similar to those of *N. laciniosa*, *N. glabra* and *N. galapagensis* (Hong *et al.* 2007). However, it grew more slowly on

CYA than these species, and phylogenetic data also indicate that this isolate represents a new species.

CBS 112.55 was isolated from garden soil in Adelaide, Australia, and produced compounds similar to wortmannin and aszonalenin and some unique metabolites, while CBS 841.96 was isolated from *Podocarpus* bark in Papua New Guinea, and produced a compound related to wortmannins and some unique compounds the structures of which have not yet been elucidated (Table 2). The ascospore ornamentations of these isolates were microtuberculate, similarly to those of *N. glabra* and *N. galapagensis*. However, both isolates produced cream-coloured colonies on CYA in contrast with *N. glabra* which produces greyish green colonies. In phylogenetic analysis they were unrelated to any other *Neosartorya* species, justifying their treatment as new species. We propose four new homothallic and monotypic *Neosartorya* species; *N. ferenczii* (NRRL 4179), *N. warcupii* (NRRL 35723), *N. australensis* (CBS 112.55) and *N. papuensis* (CBS 841.96).

## Identification

Traditionally the identification of members of section *Fumigati* were done using the colony patterns and the morphology of the conidiogenous structures, conidia, ascomata and ascospores. Ascospore ornamentation has been studied by Scanning electron microscopy, but our studies have shown that different species have similar ascospore shape and surface structure. Several species such *A. fumigatus*, *A. novofumigatus*, *fumigatiaffinis*, *A. fumisynnematus* and *A. lentulus* show strong morphological resemblance and in the light microscope these species can be difficult to be separated. The anamorphs of *Neosartorya udagawae* and *N. fennelliae* also show a similar morphology. Therefore we recommend that for a correct species identification, sequence analysis should be carried out. Our experience with sequencing the calmodin and  $\beta$ -tubulin gene revealed good species delimitation and recognition. All sequences of the ex type cultures of section *Fumigati* are available from specialised databases and also from GenBank.

---



---

## List of accepted species belonging to *Aspergillus* section *Fumigati*

The list of known species of *Neosartorya* and anamorphic species from the section *Fumigati* (Horie *et al.* 2003; Hong *et al.* 2005, 2006, 2007) is still expanding. With the species proposed here, there are now 23 *Neosartorya* species (including four new taxa) and 10 *Aspergillus* species in this group, 33 species in total and they are illustrated below.

### Strict anamorphic species:

*Aspergillus brevipes* Smith

*Aspergillus duricaulis* Raper & Fennell

*Aspergillus fumigatiaffinis* Hong, Frisvad & Samson

*Aspergillus fumigatus* Fresenius

= *A. anomalus* Pidoplichko & Kirilenko

= *A. fumigatus* var. *acolumnaris* Rai *et al.*

= *A. fumigatus* var. *ellipticus* Raper & Fennell

= *A. fumigatus* mut. *helvola* Rai *et al.*

= *A. phialiseptus* Kwon-Chung

= *A. neoellipticus* Kozakiewicz

= *Aspergillus arvii* Aho, Horie, Nishimura & Miyaji

*Aspergillus fumisynnematus* Horie, Miyaji, Nishimura, Taguchi & Udagawa

*Aspergillus lentulus* Balajee & Marr



*Aspergillus novofumigatus* Hong, Frisvad & Samson  
*Aspergillus turcosus* Hong, Frisvad & Samson  
*Aspergillus unilateralis* Thrower  
 = *A. brevipes* var. *unilateralis* (Thrower) Kozakiewicz  
*Aspergillus viridinutans* Ducker & Thrower  
 = *A. fumigatus* var. *sclerotiorum* Rai, Agarwal & Tewari

#### Teleomorph species:

*Neosartorya assulata* Hong, Frisvad & Samson [anamorph: *A. assulatus* Hong, Frisvad & Samson]  
*Neosartorya aurata* (Warcup) Malloch & Cain [anamorph: *A. igneus* Kozakiewicz]  
*Neosartorya aureola* (Fennell & Raper) Malloch & Cain [anamorph: *A. aureoluteus* Samson & Gams]  
*Neosartorya australensis* Samson, Hong & Varga, **sp. nov.**  
*Neosartorya coreana* Hong, Frisvad & Samson [anamorph: *A. coreanus* Hong, Frisvad & Samson]  
*Neosartorya denticulata* Samson, Hong & Frisvad [anamorph: *A. denticulatus* Samson, Hong & Frisvad]  
*Neosartorya fennelliae* Kwon-Chung & Kim [anamorph: *A. fennelliae* Kwon-Chung & Kim]  
 = *Neosartorya otanii* Takada, Horie & Abliz [anamorph: *A. otanii* Takada, Horie & Abliz]  
*Neosartorya ferenczii* Varga & Samson, spec. nov.  
*Neosartorya fischeri* (Wehmer) Malloch & Cain [anamorph: *A. fischeranus* Kozakiewicz]  
*Neosartorya galapagensis* Frisvad, Hong & Samson [anamorph: *A. galapagensis* Frisvad, Hong & Samson]  
*Neosartorya glabra* (Fennell & Raper) Kozakiewicz [anamorph: *A. neoglaber* Kozakiewicz]  
*Neosartorya hiratsukae* Udagawa, Tsubouchi & Horie [anamorph: *A. hiratsukae* Udagawa, Tsubouchi & Horie]  
*Neosartorya laciniosa* Hong, Frisvad & Samson [anamorph: *A. lacinosus* Hong, Frisvad & Samson]  
*Neosartorya multiplicata* Yaguchi, Someya & Udagawa [anamorph: *A. multiplicatus* Yaguchi, Someya & Udagawa]  
*Neosartorya papuensis* Samson, Hong & Varga, **sp. nov.**  
*Neosartorya pseudofischeri* Peterson [anamorph: *A. thermomutatus* (Paden) Peterson]  
*Neosartorya quadricincta* (Yuill) Malloch & Cain [anamorph: *A. quadricingens* Kozakiewicz]  
 = *Neosartorya primulina* Udagawa, Toyazaki & Tsubouchi [anamorph: *A. primulinus* Udagawa, Toyazaki & Tsubouchi]  
*Neosartorya spinosa* (Raper & Fennell) Kozakiewicz [anamorph: *A. spinosus* Kozakiewicz]  
 = *Aspergillus fischeri* var. *spinus* Raper & Fennell 1965 (basionym)  
 = *Sartorya fumigata* var. *verrucosa* Udagawa & Kawasaki  
 = *Neosartorya botucatensis* Horie, Miyaji & Nishimura [anamorph: *A. botucatensis* Horie, Miyaji & Nishimura]  
 = *Neosartorya paulistensis* Horie, Miyaji & Nishimura [anamorph: *A. paulistensis* Horie, Miyaji & Nishimura]  
 ? = *Neosartorya takakii* Horie, Abliz & Fukushima [anamorph: *A. takakii* Horie, Abliz & Fukushima]  
*Neosartorya spathulata* Takada & Udagawa [anamorph: *A. spathulatus* Takada & Udagawa]  
*Neosartorya stramenia* (Novak & Raper) Malloch & Cain [anamorph: *A. paleaceus* Samson & Gams]  
*Neosartorya tatenoi* Horie, Miyaji, Yokoyama, Udagawa & Campos-Takagi [anamorph: *A. tatenoi* Horie, Miyaji, Yokoyama, Udagawa & Campos-Takagi]  
 = *Neosartorya delicata* Kong [anamorph: *A. delicatus* Kong]  
*Neosartorya udagawae* Horie, Miyaji & Nishimura [anamorph: *A. udagawae* Horie, Miyaji & Nishimura]  
*Neosartorya warcupii* Peterson, Varga & Samson, **sp. nov.**

#### Doubtful species:

*Neosartorya sublevispora* Someya, Yaguchi & Udagawa [anamorph: *A. sublevisporus* Someya, Yaguchi & Udagawa]  
*Neosartorya indohii* Horie [anamorph: *A. indohii* Horie]  
*Neosartorya tsurutae* Horie [anamorph: *A. tsurutae* Horie]  
*Neosartorya nishimurae* Takada, Horie & Abliz [anamorph: *A. nishimurae* Takada, Horie & Abliz]

***Aspergillus brevipes*** Smith, Trans. Br. mycol. Soc. 35: 241. 1952. Fig. 6.

**Type:** CBS 467.91, from soil, New South Wales, Australia

**Other no. of the type:** ATCC 16899; CBS 118.53; IFO 5821; IMI 16034; IMI 51494; NRRL 2439; WB 4772 = IBT 22571; WB 4078 = IBT 22572

#### Description

Colony diam (7 d): CYA25: 12–15 mm; MEA25: 30–34 mm; YES25: 23–25 mm; OA25: 28–33 mm; CYA37: 16–19 mm; CREA: weak growth, no acid production

Colony colour: purple red

Conidiation: abundant

Reverse colour (CZA): dull yellow turning to reddish brown

Colony texture: velutinous

Conidial head: short columnar

Stipe: 15–50 (–100)  $\mu\text{m}$ , occasionally septate, heavy walled

Vesicle diam, shape: 10–18  $\mu\text{m}$ , pear shaped

Conidium size, shape, surface texture: 2.8–3.5  $\mu\text{m}$ , globose, spinulose

**Cultures examined:** CBS 467.91; WB 4772; WB 4078; CBS 118.523 = IBT 3051, all from the same original source

**Diagnostic features:** short heavy walled stipes, finely spinulose conidia, purple red colony colour, coloured vesicles and phialides and dark blue conidia; characterised by its vesicles borne at an angle to the stipe, as in *A. viridinutans* and *A. duricaulis*

**Similar species:** *A. duricaulis*

**Distribution:** Australia

**Ecology and habitats:** soil

**Extrolites:** Roquefortine C, cf. meleagrins, red metabolite (not structure elucidated)

**Pathogenicity:** not reported

**Note:** previous reports on viriditoxin production of *A. brevipes* (Weisleder & Lillehoj 1971; Cole & Cox 1981) were based on studies of a mixed culture of *A. brevipes* and *A. viridinutans* (Peterson SW, pers. comm.)

***Aspergillus duricaulis*** Raper & Fennell, The genus *Aspergillus*, 249. 1965. Fig. 7.

**Type:** CBS 481.65, from soil, Buenos Aires, Argentina

**Other no. of the type:** ATCC 16900; IMI 172282; JCM 01735; IBT 23177; NRRL 4021; VKM F-3572; WB 4021

#### Description

Colony diam (7 d): CYA25: 21–25 mm; MEA25: 20–22 mm; YES25: 40–44 mm; OA25: 40–44 mm; CYA37: 21–25 mm; CREA: poor growth, no acid production

Colony colour: lily green to slate olive

Conidiation: heavy in central areas

Reverse colour (CZA): colourless to pinkish drab

Colony texture: velutinous

Conidial head: loosely columnar

Stipe: 5–50  $\times$  3.5–5.5  $\mu\text{m}$ , smooth thick walled

Vesicle diam, shape: 7–14  $\mu\text{m}$ , flask shaped

Conidium size, shape, surface texture: (2.8–)3–3.3(–3.3)  $\mu\text{m}$ , globose, echinulate

**Cultures examined:** IMI 172282 = IBT 23177; CBS 481.65

**Diagnostic features:** echinulate conidia and weakly coloured reverse on CYA distinguish it from other anamorphic species

**Similar species:** *A. brevipes*

**Distribution:** Argentina

**Ecology and habitats:** soil

**Extrolites:** pseurotin A, fumagillin (found here), asperpentyn (Muhlenfeld & Achenbach 1988), duricaulic acid and asperdurin (Achenbach *et al.* 1985a), phthalides and chromanols (Achenbach *et al.* 1982a, 1985b), cyclopaldic acid and 3-O-methylcyclopolic acid (Brillinger *et al.* 1978; Achenbach *et al.* 1982b)

**Pathogenicity:** not reported

***Aspergillus fumigatiaffinis*** Hong, Frisvad & Samson, Mycologia 97: 1326. 2005. Fig. 8.

**Type:** CBS 117186, from soil, Socorro County, Sevilleta National Wildlife Refuge, New Mexico, U.S.A.

**Other no. of the type:** KACC 41148; IBT 12703

#### Description

Colony diam (7 d): CYA25: 46–49 mm; MEA25: 53–60 mm; YES25: 67–74; CYA37: 65–70; CREA: weak growth, good acid production

Colony colour: white, with center dull green

Conidiation: limited

Reverse colour (CZA): yellowish to greyish orange

Colony texture: floccose

Conidial head: short columnar

Stipe: 6–8  $\mu\text{m}$  in diam.

Vesicle diam, shape: 18–24  $\mu\text{m}$ , globose-subglobose

Conidium size, shape, surface texture: 2–3  $\mu\text{m}$ , globose-subglobose, smooth

**Diagnostic features:** has comparatively small (sub)globose vesicles (16–24  $\mu\text{m}$ ); able to grow at 10 °C, and unable to grow at 50 °C

**Similar species:** *A. fumigatus*, *A. lentulus*, *A. novofumigatus*, *A. fumigatiaffinis*

**Distribution:** U.S.A., Spain

**Ecology and habitats:** kangaroo rat, soil, human

**Extrolites:** auranthine, cycloechinoline, fumigaclavines, helvolic acid, neosartorin, palitantin, pyripropenes A, E, O & S, tryptoquivaline, tryptoquivalone

**Pathogenicity:** pathogenic to humans (Alcazar-Fuoli *et al.* 2007)

**Note:** exhibits high MICs to amphotericin B and several triazoles (Alcazar-Fuoli *et al.* 2007)



Fig. 6. *Aspergillus brevipes*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C–I. Conidiophores. J. Conidia. Scale bars = 10 μm.




Fig. 7. *Aspergillus duricaulis*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C–H. Conidiophores. I. Conidia. Scale bars = 10 µm.




Fig. 8. *Aspergillus fumigatiaffinis*. A–C. Colonies 7 d 25 °C. A. CYA. B. MEA 25 °C. C. MEA 37 °C. D–H. Conidiophores. I. Conidia. Scale bars = 10 µm.

***Aspergillus fumigatus*** Fresenius, Beitr. Mykol. 81: 18. 1863. Fig. 9.

- = *Aspergillus fumigatus* var. *acolumnaris* Rai, Agarwal & Tewari (1971)
- = *Aspergillus fumigatus* var. *albus* Rai, Tewari & Agarwal (1974)
- = *Aspergillus fumigatus* var. *cellulosae* Sartory, Sartory & Mey. (1935)
- = *Aspergillus fumigatus* var. *coeruleus* Malchevsk. (1939)
- = *Aspergillus fumigatus* var. *ellipticus* Raper & Fennell (1965)
- = *Aspergillus fumigatus* var. *fulviruber* Rai, Tewari & Agarwal (1974)
- = *Aspergillus fumigatus* var. *fumigatus* Fresen. (1863)
- = *Aspergillus fumigatus* var. *griseibrunneus* var. Rai & Singh (1974)
- = *Aspergillus fumigatus* var. *helvolus* Yuill (1937)
- = *Aspergillus fumigatus* var. *lunzinense* Svilv. (1941)
- = *Aspergillus fumigatus* var. *minus* Sartory (1919)
- = *Aspergillus neoellipticus* Kozak. (1989)
- = *Aspergillus phialoseptus* Kwon-Chung (1975)
- = *Aspergillus bronchialis* Blumentritt (1901)
- = *Aspergillus septatus* Sartory & Sartory (1943)
- = *Aspergillus arvii* Aho, Horie, Nishimura & Miyaji (1994)

**Type:** IMI 016152, from chicken lung, Connecticut, U.S.A.

**Other no. of the type:** Thom 118; QM 1981; WB 163; CBS 133.61; NRRL 163; ATCC 1022; LSHB Ac71; NCTC 982; KACC 41143

#### Description

Colony diam (7 d): CYA25: 21–67 mm; MEA25: 25–69 mm; YES25: 48–74 mm; OA25: 34–62 mm, CYA37: 60–75 mm, CREA: poor growth, no or very weak acid production

Colour: greyish turquoise or dark turquoise to dark green to dull green

Conidiation: abundant, rarely less abundant

Reverse colour (CYA): creamy, yellow to orange

Colony texture: velutinous, st. floccose (define the abbreviation st.)

Conidial head: columnar

Stipe: 50–350 × 3.5–10 µm

Vesicle diam, shape: 10–26 µm, pyriform to subclavate, sometimes subglobose, but rarely globose

Conidia length, shape, surface texture: 2–3.5(–6) µm, globose to ellipsoidal, smooth to finely rough

**Cultures examined:** ATCC 32722, AF71, AF 293, AF294, CBS 112389, CBS 487.65, CBS 133.61, CBS 545.65, CBS 457.75, CBS 542.75, CBS 113.26, CBS 110.46, CBS 120.53, CBS 132.54, CBS 123.59, CBS 158.71, CBS 180.76, CBS 143.89, CBS 148.89, CBS 488.90, CBS 287.95, CBS 100076, CBS 109032, CBS 386.75, CBS 286.95, CEA10, IMI 376380, NRRL 1979

**Diagnostic features:** Rapid growing velutinous colonies, abundant and fast conidiation, thick stipe (ca. 6–10 µm), large pyriform to semi-clavate vesicle is representative morphological features of the species. However, the characteristics are various according to strains, and some strains have exceptional characteristics. The species grows at 50 °C, no growth at 10 °C.

**Similar species:** *A. fumigati*affinis, *A. fumisynnematus*, *A. lentulus*, *A. novofumigatus*, *A. viridinutans*.

**Distribution:** Worldwide distribution, cosmopolitan fungus (Pringle *et al.* 2005)

**Ecology and habitats:** soil, human

**Extrolites:** fumagillin, fumitoxins, fumigaclavines A & C, fumitremorgins, fumiquinazolines, gliotoxin, helvolic acid, pseurotins, pyripyropens, methyl-sulochrin, trypacidin, verruculogen

**Pathogenicity:** pathogenic to humans (Raper & Fennell 1965; Marr *et al.* 2002)

**Note:** no growth at 10 °C, growth at 50 °C; some isolates carry dsRNA mycoviruses (Anderson *et al.* 1996)

***Aspergillus fumisynnematus*** Horie, Miyaji, Nishimura, Taguchi *et al.* Udagawa, Trans. Mycol. Soc. Japan: 34: 3–7. 1993. Fig. 10.

**Type:** IFM 42277, from soil, Sabaneta, Coro City, Falcon State, Venezuela

#### Description

Colony diam (7 d): CYA25: 44–48 mm; MEA25: 56–60 mm; YES25: 35–39 mm; OA25: 42–46; CYA37: 57–61 mm, CREA: poor growth and no acid production

Colony colour: greenish grey

Conidiation: limited

Reverse colour (CZA): orange white to orange grey

Colony texture: floccose

Conidial head: short columnar

Stipe: 210 × 6–8.5(–10) µm

Vesicle diam, shape: 16–20(–25) µm, hemispherical

Conidium size, shape, surface texture: 2.8–3.2 × 2.4–2.8 µm, broadly ellipsoidal, verruculose

**Cultures examined:** IFM 42277

**Diagnostic features:** production of synnemata on MEA with age (1.4–2.3 mm in height, 30–40 µm in diam.)

**Similar species:** *A. fumigatus*, *A. lentulus*, *A. novofumigatus*, *A. fumigati*affinis

**Distribution:** Brazil, Venezuela, Spain

**Ecology and habitats:** soil, human

**Extrolites:** neosartorin, pyripyropens (found here), fumimycin (Kwon *et al.* 2007)

**Pathogenicity:** pathogenic to humans (Alcazar-Fuoli *et al.* 2007; Yaguchi *et al.* 2007)

**Note:** growth at 10 °C, no growth at 50 °C

***Aspergillus lentulus*** Balajee & Marr, Eukaryot. Cell 4: 631.2005. Fig. 11.

**Type:** FH5, from clinical specimens of patients hospitalised at the Fred Hutchinson Cancer Research Center, U.S.A.

**Other no. of the type:** KACC 41940, NRRL 35552; IBT 27201

#### Description

Colony diam: CYA25: (19–)25–56 mm, MEA25: (30)40–70 mm; YES25: 42–80 mm; OA25: 44–59 mm; CYA37: 54–70 mm, CREA: weak growth, no acid production

Colour: white with interspersed grey green conidia

Conidiation: usually poor, but abundant in some isolates

Reverse colour (CYA): pale yellow to grey orange, greyish brown

Colony texture: floccose

Conidial head: short columnar

Stipe: 20–500 × 4–7 µm, smooth, sometimes sinuous and constricted neck

Vesicle diam, shape: (6–)10–25 µm, globose to pyriform, usually subglobose




Fig. 9. *Aspergillus fumigatus*. A–C. Colonies 7 d 25 °C. A. CYA. B. MEA. C. CYA 37 °C. after 3 d. D–I. Conidiophores. D–F. *A. fumigatus*. G–H. *A. fumigatus* var. *ellipticus*. I. Atypical conidiophore of CBS 133.61. J. Conidia of *A. fumigatus* var. *ellipticus*. K. Conidia of *A. fumigatus*. Scale bars = 10 µm.




Fig. 10. *Aspergillus fumisynnematus*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C–H. Conidiophores. I. Conidia. Scale bars = 10 µm.






Fig. 11. *Aspergillus lentulus*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C. Macroscopic view of the columnar conidial heads. D–I. Conidiophores. J. Conidia. Scale bars = 10 µm.

Conidia length, shape, surface texture: 2–3.2 µm, globose to broadly ellipsoidal, smooth to finely roughened

**Cultures examined:** KACC 41391 = CBS 116886, KACC 41392, KACC 41393, KACC 41681, KACC 41682, KACC 41642, KACC 41394, KACC 41395, KACC 41939 = FH7 = IBT 27209, KACC 41941 = FH4 = IBT 27210, KACC 41942 = FH220 = IBT 27202, KACC 41940 = FH5 = IBT 27201 = NRRL 35552

**Diagnostic features:** slow and poor conidiation, floccose colony texture, short columnar conidial heads, thin stipe (<7µm), globose vesicle; growth at 10 °C and no growth at 50 °C

**Similar species:** *A. fumigatiaffinis*, *A. fumigatus*, *A. fumisynnematus*, *A. novofumigatus*, *A. viridinutans*

**Distribution:** Korea, U.S.A., Japan, Australia, Netherlands, Spain etc. It is assumed that the species is distributed worldwide.

**Ecology and habitats:** soil, human, dolphin

**Extrolites:** cyclopiazonic acid, pyripyropenes A, E & O, terrein, auranthine, neosartorin

**Pathogenicity:** pathogenic to humans (Balajee *et al.* 2005b; Alhambra *et al.* 2006; Alcazar-Fuoli *et al.* 2007; Yaguchi *et al.* 2007; Lau *et al.* 2007)

**Note:** exhibits high MICs to amphotericin B and several triazoles (Balajee *et al.* 2004, 2005b)

***Aspergillus novofumigatus*** Hong, Frisvad & Samson, *Mycologia* 97: 1326. 2005. Fig. 12.

**Type:** CBS 117520, from soil, Galapagos Islands, Ecuador

**Other no. of the type:** IBT 16806

#### Description

Colony diam (7 d): CYA25: 33–48 mm; MEA25: 48–60 mm; YES25: 44–55 mm; OA25: 54–67 mm; CYA37: 49–52 mm; CREA: weak growth, no acid production

Colony colour: deep green to grey green

Conidiation: in central areas

Reverse colour (CZA): greyish orange to yellowish orange

Colony texture: velutinous

Conidial head: short columnar

Stipe: 50–500 × 4–7 µm in diam

Vesicle diam, shape: (13–)15–30 µm subglobose to flask shaped

Conidium size, shape, surface texture: 2.5–3 µm, ellipsoidal, smooth

**Cultures examined:** CBS 117520 = IBT 16806, CBS 117519 = IBT 16755

**Diagnostic features:** has nearly flask-shaped and comparatively large vesicles (15–30 mm); growth at 10 °C, no growth at 50 °C

**Similar species:** *A. fumigatus*, *A. lentulus*, *A. fumisynnematus*, *A. fumigatiaffinis*

**Distribution:** Galapagos Islands, Ecuador

**Ecology and habitats:** soil

**Extrolites:** aszonalenin, cycloechinuline, fiscalins, helvolic acid, neosartorin, palitantin, terrein, territrem B

**Pathogenicity:** not reported

***Aspergillus turcosus*** Hong, Frisvad & Samson, Antonie van Leeuwenhoek (in press). Fig. 13.

**Type:** KACC 42091, from air conditioner, Seoul, South Korea

**Other no. of the type:** IBT 27921

#### Description

Colony diam: CYA25: 32–41 mm; MEA25: 42–53 mm; YES25: 48–52 mm; OA25: 46–52 mm; CYA37: 48–56; CREA poor growth, no acid production

Colony colour: grey-turquoise to grey-green

Conidiation: abundant

Reverse colour (CZA): yellowish orange to greyish orange

Colony texture: velutinous

Conidial head: short columnar

Stipe: 80–100 × 4–7 µm

Vesicle diam, shape: 15–25 µm, flask shaped to globose

Conidium size, shape, surface texture: 2.5–3.5 µm, subglobose, smooth

**Cultures examined:** KACC 42091 = IBT 27921, KACC 42090 = IBT 27920, KACC 41955 = IBT 3016

**Diagnostic features:** Velutinous colony, grey-turquoise (green) colony colour and yellowish orange reverse on MEA and CYA, phialides cover distal two-thirds of the vesicle and growth at both 10 and 50 °C

**Similar species:** -

**Ecology and habitats:** air conditioner

**Distribution:** South Korea

**Extrolites:** Kotanins and several unique compounds but not yet elucidated secondary metabolites

**Pathogenicity:** not reported

***Aspergillus unilateralis*** Thrower, *Austral. J. Bot.* 2: 355. 1954. Fig. 14.

= *A. brevipes* var. *unilateralis* (Thrower) Kozakiewicz

**Type:** CBS 126.56, from rhizosphere of *Hibbertia fasciculata* and *Epacris impressa*, Australia

**Other no. of the type:** ATCC 16902; IFO 8136; IMI 062876; NRRL 577, QM 8163; WB 4366; WB 4779; IBT 3210

#### Description

Colony diam: CZA25: 30 mm; MEA25: 60–70 mm in 14 d, CRWEA: poor growth, no acid production

Colony colour: slate olive

Conidiation: limited

Reverse colour (CZA): nearly black

Colony texture: thin, brittle, folded in central area

Conidial head: diminutive, with few divergent spore chains

Stipe: 5–30 × 1.2–2.2 µm

Vesicle diam, shape: 4–8.5 µm, irregularly globose

Conidium size, shape, surface texture: 2.5–3.5 µm, globose, coarsely echinulate

**Cultures examined:** CBS 126.56; CBS 283.66 = IBT 3211




Fig. 12. *Aspergillus novofumigatus*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C–H. Conidiophores. I. Conidia. Scale bars = 10 μm.




Fig. 13. *Aspergillus turcosus*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C–I. Conidiophores. J. Conidia. Scale bars = 10 µm.




Fig. 14. *Aspergillus unilateralis*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C. Macroscopic view of the conidial heads. D–I. Conidiophores. J. Conidia. Scale bars = 10 μm.

**Diagnostic features:** phialides clustered on one side of the vesicle, echinulate conidia, slow growth rate and dark reverse on CYA

**Similar species:** -

**Distribution:** Australia

**Ecology and habitats:** soil

**Extrolites:** mycophenolic acid, other unique secondary metabolites

**Pathogenicity:** not reported

***Aspergillus viridinutans*** Ducker & Thrower, Austral. J. Bot. 2: 355. 1954. Fig. 15.

= *A. fumigatus* var. *sclerotiorum* J.N. Rai, S.C. Agarwal & J.P. Tewari

**Type:** CBS 127.56, from dung of rabbit, Frankston, Victoria, Australia

**Other no. of the type:** ATCC 16901; IMI 062875; IMI 062875ii; NRRL 4365; WB 4081; WB 4782; WB 4365

#### Description

Colony diam (7 d): CYA25: 20–40 mm; MEA25: 11–15 mm; YES25: 24–28 mm; OA25: 29–31 mm; CYA 37: 25–28 mm; CREA: poor growth, no acid production

Colony colour: Niagara green

Conidiation: limited on CZA, abundant on MEA

Reverse colour: colourless (CZA), yellowish green to light brownish olive (MEA)

Colony texture: centre raised, velutinous on MEA

Conidial head: columnar

Stipe: 20–35 × 3.3–4.4 μm

Vesicle diam, shape: 7.5–12 μm, flask shaped to subglobose

Conidium size, shape, surface texture: 2–2.8 μm, globose, delicately roughened

**Cultures examined:** CBS 127.56

**Diagnostic features:** “nodding” conidial heads, Niagara green colony colour

**Similar species:** none

**Ecology and habitats:** soil, dung, human

**Distribution:** Australia, Sri Lanka, Zambia, Russia (Varga *et al.* 2000b)

**Extrolites:** viriditoxin, 13-O-methylviriditin, phomaligin A, variotin, viriditin, wasabidienone B0, B1, viriditin A (Omolo *et al.* 2000), 4-acetyl-6,8-dihydroxy-5-methyl-2-benzopyran-1-1 A (Aldridge *et al.* 1966)

**Pathogenicity:** pathogenic to humans (Katz *et al.* 2005, Yaguchi *et al.* 2007, Alcazar-Fuoli *et al.* 2007)

**Notes:** this is a highly variable species; further taxonomic studies needed to clarify the taxonomic position of the isolates assigned to it (Varga *et al.* 2000a, b); exhibits high MICs to some azoles (Alcazar-Fuoli *et al.* 2007)




Fig. 15. *Aspergillus viridinutans*. A–B. Colonies 7 d 25 °C. A. CYA. B. MEA. C–I. Conidiophores. J. Conidia. Scale bars = 10 µm.

***Neosartorya assulata*** Hong, Frisvad & Samson [anamorph: *A. assulatus* Hong, Frisvad & Samson], Antonie van Leeuwenhoek (in press). Fig. 16.

**Type:** KACC 41691, from Tomato field soil, Buyeo, Korea

**Other no. of the type:** IBT 27911

**Morphological characteristics**

Colony diam (7 d): CYA25: (19–)37–41 mm; MEA25: 47–58 mm; YES25: 28–31 mm; OA25: 36–40; CYA37: 32–68 mm

Colony colour: white

Conidiation: abundant

Reverse colour (CYA): yellowish white to pale yellow

Colony texture: radially sulcate

Conidial head: short columnar

Stipe: 3–7.5 µm wide

Vesicle diam, shape: 10–18 µm, subclavate

Conidium size, shape, surface texture: 2–3 µm, subglobose to ovoid, smooth

Homothallic

Cleistothecia: 120–250 µm, white to yellowish

Ascospores: 5–6 µm, lenticular, with two well-separated equatorial crests and convex surface decorated with several large, round flaps

**Cultures examined:** KACC 41691 = IBT 27911, IBT 27910

**Diagnostic features:** well developed long and round flaps on convex surface of ascospore with two distinct equatorial crests; grow on MEA and CZA much slower than *N. pseudofischeri*

**Similar species:** *N. pseudofischeri*

**Distribution:** Korea

**Ecology and habitats:** soil

**Extrolites:** some indole alkaloids and some apolar metabolites

**Pathogenicity:** not reported

***Neosartorya aurata*** (Warcup) Malloch & Cain [anamorph: *A. igneus* Kozakiewicz], Raper & Fennell 1965. Fig. 17.

**Type:** CBS 466.65, from jungle soil, Berakas, Muama, Brunei

**Other no. of the type:** ATCC 16894; IFO 8783; IMI 075886; IMI 075886ii; NRRL 4378; QM 7860; WB 4378; IBT 3028

**Morphological characteristics**

Colony diam (7 d): CYA25: 13–15 mm; MEA25: 30–42 mm; YES25: 17–29 mm; OA25: 31–35 mm; CYA37: 13–16 mm, CREA: weak growth and no acid production

Colony colour (MEA): orange to ochraceous orange

Conidiation: sparse

Reverse colour (CZA): orange to dull brown

Colony texture: velutinous

Conidial head: loosely columnar

Stipe: 60–120 × 2–4 µm

Vesicle diam, shape: 10–16 µm, flask shaped

Conidium size, shape, surface texture: 2.5–3 µm, globose, punctate

Homothallic

Cleistothecia: 50–150 µm, orange, surrounded by a loose tangle of

encrusted orange hyphae

Ascospores: 6–6.5 × 4.5–5 µm, lenticular, with two narrow equatorial crests and convex walls finely reticulate

**Cultures examined:** CBS 466.65; WB 4379; IFO 9817

**Diagnostic features:** bright orange colour of the colony on MEA, restricted growth on CZA

**Similar species:** *N. stramenia*

**Distribution:** Brunei

**Ecology and habitats:** soil

**Extrolites:** helvolic acid, yellow unidentified compounds

**Pathogenicity:** not reported

***Neosartorya aureola*** (Fennell & Raper) Malloch & Cain [anamorph: *A. aureoluteus* Samson & Gams], Mycologia 47: 71–75. 1955. Fig. 18.

**Type:** CBS 105.55, from soil, Tafo, Ghana

**Other no. of the type:** ATCC 16896; IFO 8105; IMI 061451; IMI 061451ii; MUCL 13579; NRRL 2244; QM 1906; WB 2244; IBT 3027

**Morphological characteristics**

Colony diam (7 d): CYA25: 64–80 mm; MEA25: 77–90 mm; YES25: 70–75 mm; OA25: 55–59 mm; CYA37: 75–80 mm, CREA: poor growth, no acid production

Colony colour (CZA): apricot to light cadmium yellow

Conidiation: sparse

Reverse colour (CZA): yellow ochre to ochraceous

Colony texture: radially furrowed at center, slightly zonate

Conidial head: loosely columnar

Stipe: 50 × 2.5–4.5 µm

Vesicle diam, shape: 6–9 µm, clavate to flask shaped

Conidium size, shape, surface texture: 3–3.3 µm, globose to subglobose, delicately echinulate

Homothallic

Cleistothecia: 175–500 µm, pale lemon yellow, surrounded by loose wefts of dark golden yellow hyphae

Ascospores: 6–7 × 4.4–5 µm, lenticular, with two prominent equatorial crests and with convex surfaces conspicuously echinulate

**Cultures examined:** CBS 105.55; WB 2391

**Diagnostic features:** yellow to golden pigmentation of hyphae surrounding the cleistothecia

**Similar species:** *N. udagawae*, *A. viridinutans*

**Distribution:** Suriname, Ghana, Liberia, Fiji

**Ecology and habitats:** soil, canned passionfruit

**Extrolites:** fumagillin, tryptoquivaline, tryptoquivalone, pseurotin A and viriditoxin (FRR 2269 also produces helvolic acid)

**Pathogenicity:** not reported






Fig. 16. *Neosartorya assulata*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 5  $\mu$ m.





Fig. 17. *Neosartorya aurata*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.



**Fig. 18.** *Neosartorya aureola*. A–B. Colonies 14 d 25 °C. A. MEA. B. OA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 μm, except D = 30 μm, E = 15 μm, I = 1 μm.

***Neosartorya australensis*** Samson, Hong & Varga, **sp. nov.** (Fig. 19) – MycoBank MB492203.

Homothallica; cleistothecia superficialia, luteoalba vel dilute lutea, globosa vel subglobosa, 150–380 µm diam, in hyphis hyalinis vel luteoalbis laxo obtectis. Asci octospori, globosi vel subglobosi, 12–14 µm diam, evanescentes. Ascospores 4.5–7.5 µm diam, cristis angustis, aequatoriis binis, pagina convexa sublaevigata. Mycelium ex hyphis hyalinis, ramosis, septatis, laeviparietinis constans. Capitula conidialia curta, columnaria. Conidiophora ex hyphis aeriis exorientata, uniseriata, stipitibus 8–14 µm; vesiculae ampulliformes, 12–30 µm diam; phialides 7.5–9 × 2–3 µm, dimidium supernum vesiculae obtegentes. Conidia subglobosa vel ellipsoidea, laevia, 3.5–5 µm diam. Coloniae in agar MEA in 7 diebus et 25 °C celeriter crescentes, 40–45 mm diam, albae, capitulis conidialibus paucis. Coloniae in agar CYA in 7 diebus et 25 °C 30–35 mm diam, cremeoalbae, centro ab hyphis aeralibus laxo obtecto; capitula conidialia pauca; colonia reversa luteoalba vel luteobrunnea.

Holotype of *Neosartorya australensis*, here designated as CBS 112.55<sup>T</sup> (dried culture), isolated from garden soil, Adelaide, Australia.

Homothallic, cleistothecia superficial, yellowish white to pale yellow, globose to subglobose, 150–380 µm in diam., surrounded by a loose covering of hyaline to yellowish white hyphae. Asci 8-spored, globose to subglobose 12–14 µm, evanescent at maturity. Ascospores lens-shaped, 4.5–7.5 µm, with two equatorial crests, convex surfaces smooth to microtuberculate. Mycelium composed of hyaline, branched, septate, smooth-walled hyphae. Conidial heads short, columnar. Conidiophores arising from aerial hyphae often curling, uniseriate, stipes 12–30 µm; vesicles flask-shaped, 8–14 µm in diam.; phialides 7.5–9 × 2–3 µm, covering the upper half of vesicle. Conidia subglobose to ellipsoidal, smooth, 2.0–3.2 µm. Colonies on MEA growing rapidly, 40–45 mm in 7 d at 25 °C, white. Conidial heads produced few in number. Colonies on CYA, 30–35 mm in 7 d at 25 °C, creamy white, loosely overgrown by aerial hyphae in center. Conidial heads few in number. Reverse yellowish white to pale yellow.

**Etymology:** isolated from soil in Australia

**Extrolites:** wortmannin-like, aszonalenin-like

**Distinguishing features:** conidiophores often curled

**Other no. of the type:** IMI 061450; NRRL 2392; IBT 3021; WB 2392; Warcup SA14

**Diagnostic features:** smooth or microtuberculate 4.5–7.5 µm ascospores

**Similar species:** *N. glabra*

**Distribution:** Australia

**Ecology and habitats:** soil

**Pathogenicity:** not reported

***Neosartorya coreana*** Hong, Frisvad & Samson [anamorph: *A. coreanus* Hong, Frisvad & Samson], *Int. J. Syst. Evol. Microbiol.* 56: 477. 2006. Fig 20.

**Type:** CBS 117059, from tomato field soil, Buyeo, Korea

**Other no. of the type:** KACC 41659 = NRRL 35590 = IBT 24945

**Morphological characteristics**

Colony diam (7 d): CYA25: 41–62 mm; MEA25: 57–66 mm; YES25:

50–74 mm; OA25: 54–58 mm; CYA37: 70–74 mm, CREA: poor growth, no acid production

Colony colour: white to yellowish white

Conidiation: sparse

Reverse colour (CYA): pale to light orange

Colony texture: radially sulcate

Conidial head: columnar

Stipe: 3–4 µm wide

Vesicle diam, shape: 8–13(–15) µm, subclavate

Conidium size, shape, surface texture: 2.5–3.5 µm, subglobose to broadly elliptical, smooth

Homothallic

Cleistothecia: 200–300 µm, white to light yellow

Ascospores: 4–5 µm, with two well-separated but often bent equatorial crests up to 2 µm, convex surface reticulate

**Cultures examined:** CBS 117059

**Diagnostic features:** rugose to weak reticulate ascospores with two often bent crests, but without the equatorial rings of small projections

**Similar species:** *N. spinosa*, *N. lacinosus*

**Distribution:** South Korea, Australia

**Ecology and habitats:** soil, strawberry

**Extrolites:** aszonalenins

**Pathogenicity:** not reported in humans (although isolated from the air sacks of an ostrich: Katz *et al.* 2005)

***Neosartorya denticulata*** Samson, Hong & Frisvad [anamorph: *A. denticulatus* Samson, Hong & Frisvad], Antonie van Leeuwenhoek (in press). Fig. 21.

**Type:** CBS 652.73, from Soil under *Elaeis guineensis*, Suriname

**Other no. of the type:** KACC 41183

**Morphological characteristics**

Colony diam (7 d): CYA25: 22–24 mm; MEA25: 35–40 mm; CYA37: 35–38 mm; CREA: poor growth, no acid production

Colony colour: white

Conidiation: only on the marginal area

Reverse colour (CYA): yellowish white to pale yellow

Colony texture: loosely overgrown by aerial hyphae in the centre, sulcate in marginal areas

Conidial head: short columnar

Stipe: 3–4.5 µm wide

Vesicle diam, shape: 7–12 µm, spatulate

Conidium size, shape, surface texture: 2–3 µm, subglobose to broadly elliptical, smooth

Homothallic

Cleistothecia: 140–230 µm, yellowish white to pale yellow

Ascospores: 4–5 µm, denticulate with a prominent equatorial furrow

**Cultures examined:** CBS 652.73

**Diagnostic features:** denticulate ascospore surface and lacking equatorial crests make this a distinctive species

**Similar species:** *N. fennelliae*, *N. ferenczii*




Fig. 19. *Neosartorya australiensis*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 μm, except D = 30 μm, E = 15 μm, I = 1 μm.




Fig. 20. *Neosartorya coreana*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.




Fig. 21. *Neosartorya denticulata*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 5  $\mu$ m.

**Distribution:** Netherlands, Suriname

**Ecology and habitats:** soil, sycamore

**Extrolites:** gliotoxin, viriditoxin

**Pathogenicity:** not reported

***Neosartorya fennelliae*** Kwon-Chung & Kim [anamorph: *A. fennelliae* Kwon-Chung & Kim], *Mycologia* 66: 628. 1974. Fig. 22.

**Type:** CBS 598.74 & CBS 599.74, from eye ball of *Oryctolagus cuniculus*, U.S.A.

**Other no. of the type:** ATCC 24325 & ATCC 24326, NRRL 5534 & NRRL 5535

#### Morphological characteristics

Colony diam (7 d): CYA25: 25–30 mm; MEA25: 44–48 mm; YES25: 30–34 mm; OA25: 34–38 mm; CYA37: 50–58 mm; CREA: poor growth and no acid production

Colony colour: grey

Conidiation: abundant

Reverse colour (CZA): white

Colony texture: velutinous

Conidial head: short columnar

Stipe: 150–250 × 4–6 µm

Vesicle diam, shape: 10–17 µm, flask-shaped

Conidium size, shape, surface texture: 2.2–2.5(–2.8) µm, globose to subglobose to ellipsoid, smooth or finely roughened

Heterothallic

Cleistothecia: 150–450 µm, white

Ascospores: 5.5–7.7 × 3.2–5 µm, with two equatorial crests, convex surfaces delicately roughened

**Cultures examined:** CBS 598.74, CBS 599.74

**Diagnostic features:** heterothallic

**Similar species:** *N. denticulata*, *N. ferenczii*

**Distribution:** U.S.A., Japan, South Korea

**Ecology and habitats:** soil, mirne sludge, rabbit

**Extrolites** asperfuran, aszonalenin, fumigaclavine, viridicatumtoxin

**Pathogenicity:** not reported in humans

**Note:** no growth at 47 °C

***Neosartorya ferenczii*** Varga & Samson, **sp. nov.** (Fig. 23) – MycoBank MB504847.

Homothallica; cleistothecia superficialia, luteoalba vel dilute lutea, globosa vel subglobosa, 180–350 µm diam, in hyphis hyalinis vel luteoalbis laxe obtectis. Asci octospori, globosi vel subglobosi, 12–16 µm diam, evanescentes. Ascosporae 3.5–5.5 µm diam, cristis angustis, aequatoriis binis, pagina convexa sublaevigata. Mycelium ex hyphis hyalinis, ramosis, septatis, laeviparietinis constans. Capitula conidialia curta, columnaria. Conidiophora ex hyphis aeriis exorientia, uniseriata, stipitibus 100–150 × 4–5 µm; vesiculae ampulliformes, 10–14 µm diam; phialides 7.5–9 × 2–3 µm, dimidium supernum vesiculae obtegentes. Conidia globosa vel subglobosa, laevia, 2–2.5 µm diam. Coloniae in agar MEA in 7 diebus et 25 °C celeriter crescentes, 35–40 mm diam, albae, capitulis conidialibus paucis. Coloniae in agar CYA in 7 diebus et 25 °C 20–30 mm diam, cremeoalbae, centro ab hyphis

aerialibus laxe obtecto; capitulis conidialibus paucis; colonia reversa luteoalba vel pallide lutea.

Holotype of *Neosartorya ferenczii*, here designated as CBS 121594<sup>T</sup> (dried culture), isolated from soil in Australia.

Homothallic, cleistothecia superficial, yellowish white to pale yellow, globose to subglobose, 180–350 µm in diam., surrounded by a loose covering of hyaline to yellowish white hyphae. Asci 8-spored, globose to subglobose 12–16 µm, evanescent at maturity. Ascospores lens shaped, 3.5 × 5.5 µm, with two narrow equatorial crests, convex surface nearly smooth, microtuberculate. Mycelium composed of hyaline, branched, septate, smooth-walled hyphae. Conidial heads short, columnar. Conidiophores arising from aerial hyphae, uniseriate, stipes 100–150 × 4–5 µm; vesicles subclavate, 8–14 µm in diam; phialides 7.5–9 × 2–3 µm, covering the upper half of vesicle. Conidia globose to subglobose, smooth, 2–2.5 µm. Colonies on MEA growing rapidly, 35–40 mm in 7 d at 25 °C, white. Conidial heads produced few in number. Colonies on CYA, 20–30 mm in 7 d at 25 °C, creamish white, loosely overgrown by aerial hyphae in center. Conidial heads few in number. Reverse yellowish white to pale yellow (12A23) (Kornerup & Wanscher 1978).

**Etymology:** named after Prof. Lajos Ferenczy, eminent mycologist.

**Extrolites:** asperfuran, aszonalenin, fumigaclavine, viridicatumtoxin, gliotoxin-like, fumigatins and aszonalenin-like

**Type:** CBS 121594, from soil, Australia

**Other no. of the type:** IBT 27813, NRRL 4179; Warcup SA57

**Diagnostic features:** ascospore ornamentation similar to that of *N. fennelliae*, but with equatorial crests much narrower, and markedly different from those of *N. denticulata*

**Similar species:** *N. fennelliae*, *N. denticulata*

**Distribution:** Australia

**Ecology and habitats:** soil

**Extrolites:** asperfuran, aszonalenin, fumigaclavine, viridicatumtoxin, gliotoxin-like, fumigatins, and aszonalenin-like

**Pathogenicity:** not reported

***Neosartorya fischeri*** (Wehmer) Malloch & Cain [anamorph: *A. fischeranus* Kozakiewicz], *Can. J. Bot.* 50: 2621. 1973. Fig. 24.

= *Aspergillus fischeri* Wehmer, *Centr. Bakteriolog. Parasitenk. Abt. II* 18: 390. 1907.

= *Sartorya fumigata* Vuill., *Compt. rendu Acad. Sci. Paris* 184: 136. 1927.

**Type:** CBS 544.65, from canned apples, Wehmer

**Other no. of the type:** ATCC 1020; DSM 3700; IMI 211391; NRRL 181; QM 1983; Thom 4651.2, WB 181; IBT 3018

#### Morphological characteristics

Colony diam (7 d): CYA25: 45–68 mm; MEA25: 66–80 mm; YES25: 70–80 mm; OA25: 58–80 mm; CYA37: 65–84 mm; CREA: poor growth and no acid production

Colony colour (CZA): white to pale yellow to buff

Conidiation: sparse

Reverse colour (CZA): colourless to flesh coloured






Fig. 22. *Neosartorya fennelliae*. A–B. Colonies 14 d 25 °C. A. MEA. B–C. Crossing of mating types on MEA. D–E. Ascospores. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.




Fig. 23. *Neosartorya ferencii*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascogonia. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.




Fig. 24. *Neosartorya fischeri*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C. Macroscopic view of the columnar conidial heads. D–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.

Colony texture: velutinous  
 Conidial head: columnar  
 Stipe: 300–500 × 4–7 µm  
 Vesicle diam, shape: 12–18 µm, flask shaped  
 Conidium size, shape, surface texture: 2–2.5 µm, globose to subglobose, microtuberculate  
 Homothallic  
 Cleistothecia: up to 400 µm, light cream, borne singly or in small clusters within a loose hyphal envelope  
 Ascospores: 7–8 × 3–4 µm, convex surfaces bearing anastomosing ridges (reticulate)

**Cultures examined:** CBS 544.65; WB 4075; CBS 317.89; CBS 584.90; CBS 118441; NRRL 181; NRRL 4075; NRRL 4161; NRRL 4585

**Diagnostic features:** reticulate ascospore ornamentation

**Similar species:** *N. tatenoi*

**Distribution:** worldwide

**Ecology and habitats:** Soil, (milled) rice, cotton, potatoes, groundnuts, leather, paper products, canned products, human

**Extrolites:** terrein, fumitremorgins A & C, tryptoquivaline A, trypacidin, TR-2, verruculogen, sarcin, aszonalenins, fischerin, neosartorin, fiscalins, helvolic acid

**Pathogenicity:** pathogenic to animals and humans (Coriglione *et al.* 1990; Lonial *et al.* 1997; Mellado *et al.* 2006; Chim *et al.* 1998; Gori *et al.* 1998)

***Neosartorya galapagensis*** Frisvad, Hong & Samson [anamorph: *A. galapagensis* Frisvad, Hong & Samson], Antonie van Leeuwenhoek (in press). Fig. 25.

**Type:** CBS 117522, from soil, Galapagos Islands, Ecuador

**Other no. of the type:** KACC 41935 = IBT 16756

**Morphological characteristics**

Colony diam (7 d): CYA25: 25–40 mm; MEA25: 26–35 mm; YES25: 39–44 mm; OA25: 34–41 mm; CYA37: 44–65 mm; CREA poor growth and no acid production

Colony colour: white

Conidiation: sparse

Reverse colour (CYA): golden yellow

Colony texture: strongly funiculose

Conidial head: columnar

Stipe: 2–4 µm wide

Vesicle diam, shape: 4–11 µm, (sub)clavate

Conidium size, shape, surface texture: 2.3–3 µm, globose to subglobose, smooth

Homothallic

Cleistothecia: 90–220 µm, yellowish white, surrounded by a loose covering of aerial hyphae

Ascospores: 5 µm, with two distinct equatorial crests 1–2 µm wide, convex surface of ascospores microtuberculate

**Cultures examined:** CBS 117522 = IBT 16756; CBS 117521 = IBT 16763

**Diagnostic features:** colonies funiculose, the *Aspergillus* anamorph arises from bundles of aerial hyphae, ascospores with

two wide conspicuous equatorial crests and with microtuberculate convex surface

**Similar species:** *N. glabra*, *N. australensis*

**Distribution:** Galapagos Islands (Ecuador)

**Ecology and habitats:** soil

**Extrolites:** gregatins

**Pathogenicity:** not reported

***Neosartorya glabra*** (Fennell & Raper) Kozakiewicz [anamorph: *A. neoglaber* Kozakiewicz], Mycol. Pap. 161: 56. 1989. Fig. 26.

**Type:** CBS 111.55, from rubber scrub of an old tire, Iowa, U.S.A.

**Other no. of the type:** ATCC 16909; IFO 8789; IMI 061447; IMI 061447ii; NRRL 2163; QM 1903; WB 2163

**Morphological characteristics**

Colony diam (7 d): CYA25: 24–43 mm; MEA25: 49–66 mm; YES25: 45–54 mm; OA25: 55–76 mm; CYA37: 30–80 mm; CREA: poor growth and no acid production

Colony colour (CZA): white to pale yellow to buff

Conidiation: sparse

Reverse colour (CZA): colourless to light pink

Colony texture: velutinous

Conidial head: columnar

Stipe: 300–500 × 4–7 µm

Vesicle diam, shape: 10–18 µm, flask shaped

Conidium size, shape, surface texture: 2.5–3.5 µm, globose to subglobose, microtuberculate

Homothallic

Cleistothecia: 100–500 µm, yellowish white

Ascospores: 6.5–7.5 × 4.5–5 µm, lenticular, with two equatorial crests of 1–1.5 µm, convex surfaces finely roughened

**Cultures examined:** CBS 111.55; IMI 144207; IMI 102073; CBS 165.63

**Diagnostic features:** has smaller and whiter cleistothecia and relatively straight equatorial crests and smoother walled convex surfaces compared to *N. laciniosa*, *N. coreana* and *N. spinosa*; *N. glabra* grows somewhat slower than the other species and grows well at comparatively low temperatures; can be distinguished from *N. papuensis* and *N. australensis* using sequence data or extrolite profiles


**Similar species:** *N. papuensis*, *N. australensis*

**Distribution:** U.S.A., Morocco, Denmark, Australia, Netherlands, South Korea

**Ecology and habitats:** soil, foods, indoor

**Extrolites:** asperpentyn, avenaciolide, wortmannin-like compound

**Pathogenicity:** not reported



**Fig. 25.** *Neosartorya galapagensis*. A–B. Colonies 14 d 25 °C. A. CYA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 5  $\mu$ m.




Fig. 26. *Neosartorya glabra*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.



Fig. 27. *Neosartorya hiratsukae*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.

***Neosartorya hiratsukae*** Udagawa, Tsubouchi & Horie [anamorph: *A. hiratsukae* Udagawa, Tsubouchi & Horie], Trans. Mycol. Soc. Japan 32: 23. 1991. Fig. 27.

**Type:** NHL 3008, from pasteurised aloe juice, Tokyo, Japan

**Other no. of the type:** CBS 294.93; NRRL 20819

#### Morphological characteristics

Colony diam (7 d): CZA25: 14–15 mm; CYA25: 12–14 mm; MEA25: 26–39 mm; YES25: 42–45 mm; OA25: 42–45 mm; CYA37: 27–30 mm; CREA: rather poor growth and no acid production

Colony colour: greyish green

Conidiation: moderate

Reverse colour (CZA): light brown

Colony texture: velutinous

Conidial head: short columnar

Stipe: 120–380 × 5–7 µm

Vesicle diam, shape: 15–24 µm, flask-shaped

Conidium size, shape, surface texture: 2–2.5 µm, globose to subglobose, smooth or delicately roughened

Homothallic

Cleistothecia: 130–220 µm, light cream coloured

Ascospores: 4.5–5 µm, lenticular, with two closely appressed equatorial crests, convex surfaces finely reticulate

**Cultures examined:** CBS 294.93; IFM 50770 = IBT 27913

**Diagnostic features:** restricted growth on CZA, small cleistothecia, finely reticulate ascospores

**Similar species:** *N. fischeri*, *N. tatenoi*

**Distribution:** Japan, Brazil, South Korea

**Ecology and habitats:** soil, fruit juice, indoor air, human

**Extrolites:** avenaciolide

**Pathogenicity:** pathogenic to humans (Guarro *et al.* 2002; Mellado *et al.* 2006; Alcazar-Fuoli *et al.* 2007)

**Note:** no growth above 48 °C; some isolates carry dsRNA mycoviruses which are efficiently transmitted both through ascospores and conidia to the progeny (Varga *et al.* 1998)

***Neosartorya laciniosa*** Hong, Frisvad & Samson [anamorph: *A. lacinosus* Hong, Frisvad & Samson], Int. J. Syst. Evol. Microbiol. 56: 477. 2006. Fig. 28.

**Type:** CBS 117721, from tomato field soil, Buyeo, Korea

**Other no. of the type:** NRRL 35589 = KACC 41657

#### Morphological characteristics

Colony diam (7 d): CYA25: 38–58 mm; MEA25: 53–67 mm; YES25: 60–78 mm; OA25: 52–59 mm; CYA37: 70–80 mm; CREA: poor growth and no acid production

Colony colour: white to pale yellow

Conidiation: sparse

Reverse colour (CYA): greyish orange to yellowish orange

Colony texture: sulcate, granular

Conidial head: columnar

Stipe: 3–4 µm wide

Vesicle diam, shape: 10–14 µm, subclavate

Conidium size, shape, surface texture: 2.5–3.5 µm, globose to

subglobose, smooth

Homothallic

Cleistothecia: 300–400 µm, white to light yellow

Ascospores: 4–5 µm, broadly lenticular, with two distinct straight equatorial crests which are up to 2 µm

**Cultures examined:** CBS 117721; IBT 6660; KACC 41648; CBS 117719 = KACC 41652; KACC 41644

**Diagnostic features:** cleistothecia surrounded by a loose covering of hyaline to yellowish white, 2–4 µm wide hyphae; microtuberculate ascospores with two bent crests and two distinct equatorial rings of small projections

**Similar species:** *N. spinosa*, *N. coreana*

**Distribution:** South Korea, U.S.A., Pakistan, Netherlands, Suriname, Dominican Republic, Kenya

**Ecology and habitats:** soil

**Extrolites:** aszonalenins, tryptoquivaline, tryptoquivalone

**Pathogenicity:** not reported

***Neosartorya multiplicata*** Yaguchi, Someya & Udagawa [anamorph: *A. multiplicatus* Yaguchi, Someya & Udagawa], Mycoscience 35: 309. 1994. Fig. 29.

**Type:** PF 1154, from soil, Taiwan

**Other no. of the type:** CBS 646.95, IBT 17517

#### Morphological characteristics

Colony diam (7 d): CYA25: 24–36 mm; MEA25: 35–50 mm; YES25: 38–42 mm; OA28–43 mm; CYA37: 41–80 mm; CREA: poor growth and no acid production

Colony colour: white

Conidiation: sparse

Reverse colour (CYA): greyish yellow to olivaceous buff

Colony texture: floccose

Conidial head: loosely columnar

Stipe: 20–160 × 2.5–4 µm

Vesicle diam, shape: 4–8 µm, flask-shaped to irregular

Conidium size, shape, surface texture: 2.5–4 µm, globose to subglobose, smooth

Homothallic

Cleistothecia: 100–300 µm, cream coloured

Ascospores: 4–5 µm, with a shallow furrow but without distinct equatorial crests, ornamented on surfaces by several linear ridges presenting ribbed or somewhat reticulate pattern

**Cultures examined:** CBS 646.95

**Diagnostic features:** can be distinguished from other species of *Neosartorya* by its almost globose ascospores, which have ribbed ornamentation with several linear ridges, and by the reduced production of its conidial heads on common media

**Similar species:** none


**Distribution:** Taiwan

**Ecology and habitats:** soil

**Extrolites:** helvolic acid

**Pathogenicity:** not reported





**Fig. 28.** *Neosartorya laciniosa*. A–B. Colonies 14 d 25 °C. A. MAA. B. CYA. C–E. Ascogonia. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.




Fig. 29. *Neosartorya multiplicata*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C. Macroscopic view of the columnar conidial heads D–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.




Fig. 30. *Neosartorya papuensis*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.

***Neosartorya papuensis*** Samson, Hong & Varga, sp. nov. (Fig. 30) – MycoBank MB505571.

Homothallica; cleistothecia superficialia, luteoalba vel dilute lutea, globosa vel subglobosa, 200–350 µm diam, in hyphis hyalinis vel luteoalbis laxo obtectis. Asci octospori, globosi vel subglobosi, 14–20 µm diam, evanescentes. Ascospores 5.5–7.5 µm diam, cristis angustis, aequatoris binis, pagina convexa sublaevigata. Mycelium ex hyphis hyalinis, ramosis, septatis, laeviparietinis constans. Capitula conidialia curta, columnaria. Conidiophora ex hyphis aeriis exorientia, uniseriata, stipitibus 80–120 × 4–5 µm; vesiculae ampulliformes, 10–14 µm diam; phialides 7.5–9 × 2–3 µm, dimidium supernum vesiculae obtegentes. Conidia globosa vel subglobosa, laevia, 2–3 µm diam. Coloniae in agar MEA in 7 diebus et 25 °C celeriter crescentes, 35–40 mm diam, albae, capitulis conidialibus paucis. Coloniae in agar CYA in 7 diebus et 25 °C 20–30 mm diam, cremeoalbae, centro ab hyphis aeralibus laxo obtecto; capitula conidialia pauca; colonia reversa luteoalba vel pallide lutea.

Holotype of *Neosartorya papuensis*, here designated as CBS 841.96<sup>T</sup> (dried culture), isolated from *Podocarpus* (Podocarpaceae), bark, Myola, Owen Stanley Range, Northern Province, Papua New Guinea.

Homothallic, cleistothecia superficial, yellowish white to pale yellow, globose to subglobose, 200–350 µm in diam., surrounded by a loose covering of hyaline to yellowish white hyphae. Asci 8-spored, globose to subglobose 14–20 µm, evanescent at maturity. Ascospores 5.5–7.5 µm, with two equatorial crests, convex surface smooth microtuberculate. Mycelium composed of hyaline, branched, septate, smooth-walled hyphae. Conidial heads short, columnar. Conidiophores arising from aerial hyphae, uniseriate, stipes 100–150 × 4–5 µm; vesicles flask-shaped, 10–14 µm in diam.; phialides 7.5–9 × 2–3 µm, covering the upper half of vesicle. Conidia globose to subglobose, smooth, 2–3 µm. Colonies on MEA growing rapidly, 35–40 mm in 7 d at 25 °C, white. Conidial heads few in number. Colonies on CYA, 30–35 mm in 7 d at 25 °C, producing sectors, creamy white, loosely overgrown by aerial hyphae in center. Conidial heads few in number. Reverse yellowish white to pale yellow (12A23) (Kornerup and Wanscher 1978).

**Etymology:** isolated in Papua New Guinea

**Extrolites:** wortmannin-like

**Distinguishing features:** smooth microtuberculate 5.5–7.5 µm, ascospores

**Other no. of the type:** IBT 27801

**Cultures examined:** CBS 841.96

**Similar species:** *N. galapagensis*, *N. glabra*, *N. australensis*

**Distribution:** Papua New Guinea

**Pathogenicity:** not reported

***Neosartorya pseudofischeri*** Peterson [anamorph: *A. thermomutatus* (Paden) Peterson], Mycol. Res. 86: 547. 1992. Fig. 31.

**Type:** NRRL 20748, from human vertebrae, Atlanta, Georgia, U.S.A.

**Other no. of the type:** CBS 208.92

**Holotype:** 404.67, moldy cardboard, Victoria, British Columbia, Canada

#### Morphological characteristics

Colony diam (7 d): CYA25: 60–70 mm; MEA25: 90 mm in 7 d

Colony colour: white to pale creamish

Conidiation: sparse

Reverse colour (CZA): clear or faintly yellowish

Colony texture: velutinous

Conidial head: loosely columnar

Stipe: 200–300 × 4–7 µm

Vesicle diam, shape: 10–17 µm, subglobose

Conidium size, shape, surface texture: 3–4 µm, globose to subglobose, smooth

Homothallic

Cleistothecia: 150–300 µm, white

Ascospores: 4.5–6 µm, subglobose, with two equatorial crests of 1 µm wide, convex surfaces with raised flaps resembling triangular projections

**Cultures examined:** CBS 208.92, CBS 404.67

**Diagnostic features:** distinctly ornamented ascospores

**Similar species:** -

**Distribution:** U.S.A., Canada, Netherlands, South Korea, Spain, Denmark, Estonia

**Ecology and habitats:** soil, indoor, human

**Extrolites:** asperfuran, cytochalasin-like compound, fiscalin-like compound, pyripyropens, gliotoxin

**Pathogenicity:** pathogenic to humans (Padhye *et al.* 1994; Matsumoto *et al.* 2002; Jarv *et al.* 2004; Balajee *et al.* 2005a; Alcazar-Fuoli *et al.* 2007; Lau *et al.* 2007) and animals (Barrs *et al.* 2007)

***Neosartorya quadricincta*** (J.L. Yuill) Malloch & Cain [anamorph: *A. quadricingens* Kozakiewicz], Can. J. Bot. 50: 2621. 1973. Fig. 32.

= *Neosartorya primulina* Udagawa, Toyaz. & Tsub. [anamorph: *A. primulinus* Udagawa, Toyaz. & Tsub.]

**Type:** CBS 135.52, from cardboard, York, U.K.

**Other no. of the type:** ATCC 16897; IMI 048583; IMI 048583ii; NRRL 2154; QM 6874; WB 2154

#### Morphological characteristics

Colony diam (7 d): CYA25: 26–42 mm; MEA25: 52–59 mm; YES25: 36–59 mm; OA25: 47–55 mm; CYA37: 50–58 mm; CREA: poor growth and no acid production

Colony colour (CZA): white to light tan

Conidiation: sparse

Reverse colour (CZA): colourless to flesh coloured

Colony texture: floccose

Conidial head: loosely columnar


Stipe: 400–500 × 2–7 µm

Vesicle diam, shape: 10–20 µm, flask shaped

Conidium size, shape, surface texture: 2–3 µm, elliptical to globose, microtuberculate

Homothallic

Cleistothecia: up to 300 µm, buff to light tan



**Fig. 31.** *Neosartorya pseudofischeri*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.



Fig. 32. *Neosartorya quadricincta*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 μm, except D = 30 μm, E = 15 μm, I = 1 μm.

Ascospores: 4–5 µm, with two prominent equatorial crests, each duplicated by a some-what less prominent band, reticulate

**Cultures examined:** CBS 135.52; WB 2221; WB 4175; CBS 100942

**Diagnostic features:** presence of 4 equatorial crests on ascospores, reticulate ascospore ornamentation

**Similar species:** -

**Distribution:** Suriname, South Korea, U.K., Netherlands, Australia

**Ecology and habitats:** Soil, pectin, cardboard, fruit juice, mango pulp

**Extrolites:** quinolactacin, aszonalenins

**Pathogenicity:** not reported

**Note:** some isolates carry dsRNA mycoviruses (Varga *et al.* 1998)

***Neosartorya spathulata*** Takada & Udagawa [anamorph: *A. spathulatus* Takada & Udagawa], Mycotaxon 24: 395. 1985. Fig. 33.

**Type:** CBS 408.89 & CBS 409.89, from cultivated soil under *Alocasia macrorrhiza*, Taiwan

**Other no. of the type:** IMI 308593 & IMI 308593; NHL 2948, NHL 2949; NRRL 20549 & NRRL 20550

#### Morphological characteristics

Colony diam (7 d): CZA25: 33–38 mm, MEA25: 80 mm; OA25: 40–46 mm

Colony colour: greyish green

Conidiation: abundant

Reverse colour (CZA): uncoloured

Colony texture: velutinous

Conidial head: loosely columnar

Stipe: 500–1500 × 11–18(–25) µm and 60–250 × 4–10 µm

Vesicle diam, shape: 25–52 µm and 8–15 µm, flask-shaped

Conidium size, shape, surface texture: 3–5.5 × 2–4.5 µm, ellipsoidal, smooth

Heterothallic

Cleistothecia: 100–260 µm, pale yellow to light yellow

Ascospores: 3.5–4 µm, lenticular, with two equatorial crests, convex surfaces nearly smooth

**Cultures examined:** CBS 408.89 & CBS 409.89

**Diagnostic features:** yellowish cleistothecia, ascospores with large equatorial crests and smooth surface, two types of conidial heads (diminutive??)

**Similar species:** -

**Distribution:** Taiwan

**Ecology and habitats:** soil

**Extrolites:** xanthocillins, aszonalenins

**Pathogenicity:** not reported

***Neosartorya spinosa*** (Raper & Fennell) Kozakiewicz [anamorph: *A. spinosus* Kozakiewicz], Mycol. Pap. 161: 58. 1989. Fig. 34.

= *Aspergillus fischeri* var. *spinus* Raper & Fennell 1965 (basionym)

= *Sartorya fumigata* var. *verrucosa* Udagawa & Kawasaki

= *Neosartorya botucatensis* Y. Horie, Miyaji & Nishim. [anamorph: *A. botucatensis* Y. Horie, Miyaji & Nishim.]

= *Neosartorya paulistensis* Y. Horie, Miyaji & Nishim. [anamorph: *A. paulistensis* Y. Horie, Miyaji & Nishim.]

? = *Neosartorya takakii* Horie, Abliz & K. Fukush. [anamorph: *A. takakii* Horie, Abliz & K. Fukush.]

**Type:** CBS 483.65, from soil, Nicaragua

**Other no. of the type:** ATCC 16898; IFO 8782; IMI 211390; NRRL 5034; WB 5034; IBT 3022

#### Morphological characteristics

Colony diam (7 d): CYA25: 41–70 mm; MEA25: 55–75 mm; YES25: 55–80 mm; OA25: 56–64 mm; CYA37: 67–85 mm; CREA: poor growth and no acid production

Colony colour (CZA): white to pale yellow to buff

Conidiation: sparse

Reverse colour (CZA): colourless to light pink

Colony texture: velutinous

Conidial head: columnar

Stipe: 300–500 × 4–7 µm

Vesicle diam, shape: 12–18 µm, flask shaped

Conidium size, shape, surface texture: 2–2.5 µm, globose to subglobose, microtuberculate

Homothallic

Cleistothecia: 200–300 µm, cartridge buff

Ascospores: 4.5 µm, with two widely separated equatorial crests, with convex surfaces bearing spinelike projections

**Cultures examined:** CBS 483.65

**Diagnostic features:** have echinulate ascospores with spines ranging from <0.5 µm up to 7 µm long, or with verruculose and small triangular, sometimes circularly arranged, projections

**Similar species:** *N. coreana*, *N. laciniosa*

**Distribution:** Nicaragua, Kenya, Denmark, Dominican Republic, U.S.A., Belgium, Sudan, Japan, India, Pakistan, South Korea

**Ecology and habitats:** Soil, fruit juice, human

**Extrolites:** aszonalenins, 2-pyrovoylaminobenzamide, pseurotin

**Pathogenicity:** pathogenic to humans (Summerbell *et al.* 1992; Mellado *et al.* 2006; Gerber *et al.* 1973)

***Neosartorya stramenia*** (R.O. Novak & Raper) Malloch & Cain [anamorph: *A. paleaceus* Samson & Gams], Can. J. Bot. 50: 2622. 1972. Fig. 35.

**Type:** CBS 498.65, soil from maple-ash-elm forest, Wisconsin, U.S.A.

**Other no. of the type:** ATCC 16895; IFO 9611; IMI 172293; WB 4652

#### Morphological characteristics

Colony diam (7 d): CYA25: 10–40; MEA25: 40–59 mm; YES25: 58–62 mm; OA: 56–60 mm; CYA37: 45–49 mm; CREA: poor growth and no acid production




Fig. 33. *Neosartorya spathulata*. A–B. Colonies 14 d 25 °C. A. MEA. B–C. Crossing of mating types on MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.






Fig. 34. *Neosartorya spinosa*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.

Colony colour (CZA): mustard-yellow  
 Conidiation: sparse  
 Reverse colour (CZA): yellow-orange  
 Colony texture: granulose  
 Conidial head: loosely columnar  
 Stipe: 80–140 × 3.5–5.5 µm, heavy walled, septate, coloured in terminal areas  
 Vesicle diam, shape: 10–12 µm, flask shaped to globose  
 Conidium size, shape, surface texture: 2.5–3 µm, globose, microverrucose  
 Homothallic  
 Cleistothecia: 50–175 µm, cartridge buff  
 Ascospores: 4.5–5.5 µm, with two widely separated flexuous equatorial crests, convex surfaces finely echinulate

**Cultures examined:** CBS 498.65; IFO 31358

**Diagnostic features:** faster growth rate and pronounced echinulate ascospore ornamentation distinguishes this species from *N. aurata*

**Similar species:** *N. aurata*

**Distribution:** U.S.A., Argentina

**Ecology and habitats:** Soil, salt grass (*Distichlis scoparia*)

**Extrolites:** quinolactacin, avenaciolide

**Pathogenicity:** not reported

***Neosartorya tatenoi*** Horie, Miyaji, Yokoyama, Udagawa & Campos-Takagi [anamorph: *A. tatenoi* Y. Horie, M. Miyaji, K. Yokoy., Udagawa & Campos-Takagi], Trans. Mycol. Soc. Japan 33: 395. 1992. Fig. 36.

= *Neosartorya delicata* H.Z. Kong [anamorph: *A. delicatus* H.Z. Kong]

**Type:** CBM FA 0022, from soil, Brazil

**Other no. of the type:** CBS 407.93; IBT 21589

#### **Morphological characteristics**

Colony diam (7 d): CYA25: 35–39 mm; MEA25: 31–39 mm; YES25: 57–74 mm; OA25: 50–55 mm; CYA37: 72–78 mm; CREA: poor growth and no acid production  
 Colony colour: pale yellow to yellowish white  
 Conidiation: sparse  
 Reverse colour (CZA): orange white to pale orange  
 Colony texture: velutinous to floccose  
 Conidial head: short columnar  
 Stipe: 270 × 4–7.5 µm  
 Vesicle diam, shape: 10–20 µm, hemispherical to flask-shaped  
 Conidium size, shape, surface texture: 2–3(–3.5) µm, globose to ovoid, smooth  
 Homothallic  
 Cleistothecia: 140–360 × 140–310 µm, hyaline to pale yellowish brown  
 Ascospores: 5–5.5 µm, lenticular, with two equatorial crests, convex surfaces with distinctly and narrowly reticulate ridges

**Cultures examined:** CBS 407.93; NRRL 4584

**Diagnostic features:** distinct narrowly reticulate ascospore ornamentation

**Similar species:** *N. fischeri*, *N. multiplicata*

**Distribution:** Brazil, Dominican Republic

**Ecology and habitats:** soil

**Extrolites:** aszonalenins

**Pathogenicity:** not reported

***Neosartorya udagawae*** Horie, Miyaji & Nishim. [anamorph: *A. udagawae* Horie, Miyaji & Nishim.], Mycoscience 36: 199. 1995. Fig. 37.

**Type:** CBM FA-0703 & CBM FA-0702, from soil, Brazil

**Other no. of the type:** CBS 114217 & CBS 114218

#### **Morphological characteristics**

Colony diam (7 d): CYA25: 33–36 mm; MEA25: 63–68 mm; YES25: 64–68 mm; OA25: 51–55 mm; CYA37: 61–65 mm; CREA: poor growth and no acid production  
 Colony colour (CZA): dull green  
 Conidiation: abundant  
 Reverse colour (CZA): light orange to greyish orange  
 Colony texture: velutinous  
 Conidial head: columnar  
 Stipe: up to 530 × 4–6 µm  
 Vesicle diam, shape: 12–15 µm, hemispherical to flask shaped  
 Conidium size, shape, surface texture: 2.6–3.2 × 2.4–2.6 µm, subglobose to broadly ellipsoidal, smooth  
 Heterothallic  
 Cleistothecia: 310–620 × 280–530 µm, yellowish white to light yellow, surrounded by a loose covering of hyaline to pale yellowish brown hyphae  
 Ascospores: 5–5.5 × 4–5 µm, broadly lenticular, with two equatorial or often irregular crests, convex surfaces tuberculate

**Cultures examined:** CBS 114217, CBS 114218

**Diagnostic features:** heterothallic species, with characteristic tuberculate ascospore ornamentation

**Similar species:** *N. aureola*, *A. viridinutans*

**Distribution:** Brazil, U.S.A., Spain, Japan

**Ecology and habitats:** Soil, human

**Extrolites:** fumigatin, fumagillin, tryptoquivaline, tryptoquivalone

**Pathogenicity:** pathogenic to humans (Balajee *et al.* 2006; Moragues *et al.* 2006)




Fig. 35. *Neosartorya stramenia*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 µm, except D = 30 µm, E = 15 µm, I = 1 µm.




Fig. 36. *Neosartorya tatenoi*. A–B. Colonies 14 d 25 °C. A. MEA. B. OA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.




Fig. 37. *Neosartorya udagawae*. A–B. Colonies 14 d 25 °C. A. MEA. B. Crossing of mating types on MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10  $\mu$ m, except D = 30  $\mu$ m, E = 15  $\mu$ m, I = 1  $\mu$ m.




Fig. 38. *Neosartorya warcupii*. A–B. Colonies 14 d 25 °C. A. OA. B. MEA. C–E. Ascomata. F–G. Asci and ascospores. H. Ascospores. I. SEM of ascospores. J–L. Conidiophores. M. Conidia. Scale bars = 10 μm, except D = 30 μm, E = 15 μm, I = 1 μm.

***Neosartorya warcupii* Peterson, Varga & Samson, sp. nov.**  
(Fig. 38) – MycoBank MB505572.

Homothallica; cleistothecia superficialia, alba vel dilute lutea, globosa vel subglobosa, 200–350 µm diam, in hyphis hyalinis vel luteoalbis laxe obtectis. Asci octospori, globosi vel subglobosi, 4.5–7 µm diam, evanescentes. Ascospores 5.5–7 µm diam, cristis angustis, aequatoris binis, pagina convexa sublaevigata. Mycelium ex hyphis hyalinis, ramosis, septatis, laeviparietinis constans. Capitula conidialia curta, columnaria. Conidiophora ex hyphis aeriis exorientia, uniseriata, stiptibus 100–150 × 4–5 µm; vesiculae ampulliformes, 10–25 µm diam; phialides 7.5–9 × 2–3 µm, dimidium supernum vesiculae obtegentes. Conidia subglobosa vel ellipsoidea, laevia, 1.8–1.5 µm diam. Coloniae in agar MEA in 7 diebus et 25 °C celeriter crescentes, 35–40 mm diam, albae, capitulis conidialibus paucis. Coloniae in agar CYA in 7 diebus et 25 °C 20–30 mm diam, cremeoalbae, centro ab hyphis aerialibus laxe obtecto; capitula conidialia pauca; colonia reversa luteobrunnea vel atrobrunnea.

Holotype of *Neosartorya warcupii*, here designated as NRRL 35723<sup>T</sup> (dried culture), isolated from soil, Finder's Range, Australia.

Homothallic, cleistothecia superficial, yellowish white to pale yellow, globose to subglobose, 180–350 µm in diam., surrounded by a loose covering of hyaline to yellowish white hyphae. Asci 8-spored, globose to subglobose 10–16 µm, evanescent at maturity. Ascospores lens shaped 4.5–7 µm, with two prominent equatorial crests, convex surface smooth to microtuberculate. Mycelium composed of hyaline, branched, septate, smooth-walled hyphae. Conidial heads short, columnar. Conidiophores arising from aerial hyphae, uniseriate, stipes 100–150 × 4–6 µm; vesicles subclavate to subglobose, 12–18 µm in diam; phialides 7.5–9 × 2–3 µm, covering the upper half of vesicle. Conidia globose to subglobose, smooth, 1.8–2.5 µm. Colonies on MEA growing rapidly, 35–40 mm in 7 d at 25 °C. Colonies on CYA, 18–22 mm in 7 d at 25 °C, creamish white, sectors frequently produced. Conidial heads few in number. Reverse bluish in colour.

**Etymology:** named after Prof. J. H. Warcup, eminent mycologist, who isolated this culture.

**Extrolites:** wortmannin-like, aszonalenin-like, chromanol-like, tryptoquivaline-like and tryptoquivalone-like

**Distinguishing features:** secretes a blue pigment to the medium in 7–10 d; relatively slow growth on CYA at 25 °C

**Distribution:** Australia

**Ecology and habitats:** soil

**Pathogenicity:** not reported

## REFERENCES

- Achenbach H, Muhlenfeld A, Brillinger GU (1985b). Metabolites from microorganisms. 30. Phthalides and chromanols from *Aspergillus duricaulis*. *Liebigs Annalen der Chemie* **1985**: 1596–1638.
- Achenbach H, Muhlenfeld A, Kohl W, Brillinger GU (1985a). Metabolic products of microorganisms. 31. Duricaulic acid, a new natural product of the phthalimidine type from *Aspergillus duricaulis*. *Zeitschrift für Naturforschung Section B* **40**: 1219–1225.
- Achenbach H, Muhlenfeld A, Weber B, Brillinger GU (1982a). Highly substituted chromanols from cultures of *Aspergillus duricaulis*. *Tetrahedron Letters* **23**: 4659–4660.
- Achenbach H, Muhlenfeld A, Weber B, Kohl W, Brillinger GU (1982b). Investigations on metabolites of microorganisms. 27. Cyclopaldic acid and 3-O-methyl cyclopaldic acid, 2 antibioticly active substances from *Aspergillus duricaulis*. *Zeitschrift für Naturforschung Section B* **1982**: 1091–1097.
- Alcazar-Fuoli L, Mellado E, Alastruey-Izquierdo A, Cuenca-Estrella M, Rodriguez-Tudela JL (2007). Antifungal susceptibility pattern of sequences based identified isolates of *Aspergillus* section *Fumigati*. *Antimicrobial Agents and Chemotherapy* (in press)
- Aldridge DC, Grove JF, Truner WB (1966). 4-acetyl-6,8-dihydroxy-5-methyl-2-benzopyran-1-1 A metabolite of *Aspergillus viridinutans*. *Journal of the Chemical Society C* **1966**: 126–129.
- Alhambra A, Moreno JM, Dolores Moragues M, Brena S, Quindós G, Pontón J, del Palacio A (2006). Aislamiento de *Aspergillus lentulus* en un enfermo crítico con EPOC y aspergilosis invasora. *VIII Congreso Nacional Micología, Barcelona, Spain*, D14.
- Anderson MJ, Gull K, Denning DW (1996). Molecular typing by random amplification of polymorphic DNA and M13 Southern hybridization of related paired isolates of *Aspergillus fumigatus*. *Journal of Clinical Microbiology* **34**: 87–93.
- Balajee SA, Gribskov J, Brandt M, Ito J, Fothergill A, Marr KA (2005a). Mistaken identity: *Neosartorya pseudofischeri* and its anamorph masquerading as *Aspergillus fumigatus*. *Journal of Clinical Microbiology* **43**: 5996–5999.
- Balajee SA, Gribskov JL, Hanley E, Nickle D, Marr KA (2005b). *Aspergillus lentulus* sp. nov., a new sibling species of *A. fumigatus*. *Eukaryotic Cell* **4**: 625–632.
- Balajee SA, Nickle D, Varga J, Marr KA (2006). Molecular studies reveal frequent misidentification of *Aspergillus fumigatus* by morphotyping. *Eukaryotic Cell* **5**: 1705–1712.
- Balajee SA, Weaver M, Imhof A, Gribskov J, Marr KA (2004). *Aspergillus fumigatus* variant with decreased susceptibility to multiple antifungals. *Antimicrobial Agents and Chemotherapy* **48**: 1197–1203.
- Barrs VR, Beatty JA, Lingard AE, Malik R, Krockenberger MB, Martin P, O'Brien C, Angles JM, Dowden M, Halliday C (2007). Feline sino-orbital aspergillosis: an emerging clinical syndrome? *Australian Veterinary Journal* **85**: N23.
- Brillinger GU, Beberle W, Weber B, Achenbach H (1978). Metabolic products of microorganisms. 167. Cyclopaldic acid from *Aspergillus duricaulis*. 1. Production, isolation and biological properties. *Archives of Microbiology* **116**: 245–252.
- Chim CS, Ho PL, Yuen KY (1998). Simultaneous *Aspergillus fischeri* and herpes simplex pneumonia in a patient with multiple myeloma. *Scandinavian Journal of Infectious Diseases* **30**: 190–191.
- Cole RJ, Cox RH (1981). *Handbook of toxic fungal metabolites*. New York: Academic Press.
- Coriglione G, Stella G, Gafa L, Spata G, Oliveri S, Padhye AA, Ajello L (1990). *Neosartorya fischeri* var. *fischeri* (Wehmer) Malloch and Cain 1972 (anamorph: *Aspergillus fischerianus* Samson and Gams 1985) as a cause of mycotic keratitis. *European Journal of Epidemiology* **6**: 382–385.
- Filttenborg O, Frisvad JC, Svendsen JA (1983). Simple screening method for molds producing intracellular mycotoxins in pure culture. *Applied and Environmental Microbiology* **45**: 581–585.
- Frisvad JC, Samson RA (1990). Chemotaxonomy and morphology of *Aspergillus fumigatus* and related taxa. In: *Modern concepts in Penicillium and Aspergillus classification*. Samson RA, Pitt JI, eds. New York: Plenum Press: 201–208.
- Frisvad JC, Thrane U (1987). Standardized high performance liquid chromatography of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection). *Journal of Chromatography A* **404**: 195–214.
- Frisvad JC, Thrane U (1993). Liquid chromatography of mycotoxins. *Journal of Chromatography Library* **54**: 253–372.
- Fujimoto H, Ikeda M, Yamamoto K, Yamazaki M (1993). Structure of fischerin, a new toxic metabolite from an ascomycete, *Neosartorya fischeri* var. *fischeri*. *Journal of Natural Products* **56**: 1268–1275.
- Geiser DM, Frisvad JC, Taylor JW (1998). Evolutionary relationships in *Aspergillus* section *Fumigati* inferred from partial -tubulin and hydrophobin DNA sequences. *Mycologia* **90**: 831–845.
- Gerber J, Chomicki J, Brandsberg JW, Jones R, Hammerman KJ (1973). Pulmonary aspergillosis caused by *Aspergillus fischeri* var. *spinus*: report of a case and value of serologic studies. *American Journal of Clinical Pathology* **60**: 861–866.
- Glass NL, Donaldson GC (1995). Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* **61**: 1323–1330.
- Gomez MM, Pflug IJ, Busta FF (1994). Resistance of *Neosartorya fischeri* to wet and dry heat. *Journal of Pharmacological Science and Technology* **48**: 16–23.
- Gori S, Pellegrini G, Filippini F, Capanna SD, Biancofiore G, Mosca F, Lofaro A (1998). Pulmonary aspergillosis caused by *Neosartorya fischeri* (*Aspergillus fischerians*) in a liver transplant recipient. *Journal de Mycologie Medicale* **8**: 105–107.
- Guarro J, Kallas EG, Godoy P, Karenina A, Gene J, Stchigel A, Colombo AL (2002). Cerebral aspergillosis caused by *Neosartorya hiratsukae*, Brazil. *Emerging Infectious Diseases* **8**: 989–991.
- Hillis DM, Bull JJ (1993). An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.

- Hong SB, Cho HS, Shin HD, Frisvad JC, Samson RA (2006). Novel *Neosartorya* species isolated from soil in Korea. *International Journal of Systematic and Evolutionary Microbiology* **56**: 477–486.
- Hong SB, Go SJ, Shin HD, Frisvad JC, Samson RA (2005). Polyphasic taxonomy of *Aspergillus fumigatus* and related species. *Mycologia* **97**: 1316–1329.
- Hong SB, Shin HD, Hong JB, Frisvad JC, Nielsen PV, Varga J, Samson RA (2007). New taxa of *Neosartorya* and *Aspergillus* in *Aspergillus* section *Fumigati*. *Antonie van Leeuwenhoek* (in press)
- Horie Y, Abliz P, Fukushima K, Okada K, Campos Takaki GM (2003). Two new species of *Neosartorya* from Amazonian soil, Brazil. *Mycoscience* **44**: 397–402.
- Horie Y, Abliz P, Fukushima K, Okada K, Gusmao NB (2001). *Neosartorya takakii*, a new species from soil in Brazil. *Mycoscience* **42**: 91–95.
- Horie Y, Miyaji M, Nishimura K, Franco MF, Coelho KIR (1995b). New and interesting species of *Neosartorya* from Brazilian soil. *Mycoscience* **36**: 199–204.
- Horie Y, Miyaji M, Nishimura K, Franco MF, Labuki KRC (1995a). Two new species of *Neosartorya* from Brazilian soil. *Mycoscience* **36**: 159–165.
- Horie Y, Miyaji M, Nishimura K, Taguchi H, Udagawa S (1993). *Aspergillus fumisynnematus*, a new species from Venezuelan soil. *Transactions of the Mycological Society of Japan* **34**: 3–7.
- Horie Y, Miyaji M, Yokoyama K, Udagawa S, Campos-Takaki GM (1992). *Neosartorya tatenoi*, a new species from Brazilian soil. *Transactions of the Mycological Society of Japan* **33**: 395–399.
- Jarv H, Lehtmaa J, Summerbell RC, Hoekstra ES, Samson RA, Naaber P (2004). Isolation of *Neosartorya pseudofischeri* from blood: first hint of pulmonary aspergillosis. *Journal of Clinical Microbiology* **42**: 925–928.
- Katz ME, Dougall AM, Weeks K, Cheetham BF (2005). Multiple genetically distinct groups revealed among clinical isolates identified as atypical *Aspergillus fumigatus*. *Journal of Clinical Microbiology* **43**: 551–555.
- Katz ME, Mcloon M, Burrows S, Cheetham BF (1998). Extreme DNA sequence variation in isolates of *Aspergillus fumigatus*. *FEMS Immunology and Medical Microbiology* **20**: 283–288.
- Kornerup A, Wanscher JH (1978). *Methuen handbook of colour*, 3rd ed. Eyre Methuen, London.
- Kumar S, Tamura K, Nei M (2004). MEGA3: Integrated Software for Molecular Evolutionary Genetics Analysis and Sequence Alignment. *Briefings in Bioinformatics* **5**: 150–163.
- Kwon YJ, Sohn MJ, Zheng CJ, Kim WG (2007). Fumimycin: a peptide deformylase inhibitor with an unusual skeleton produced by *Aspergillus fumisynnematus*. *Organic Letters* **9**: 2449–2451.
- Kwon-Chung KJ, Kim SJ (1974). A second heterothallic *Aspergillus*. *Mycologia* **66**: 628–638.
- Larsen TO, Smedsgaard J, Nielsen KF, Hansen MA, Samson RA, Frisvad JC (2007). Production of mycotoxins by *Aspergillus lentulus* and other medically important and closely related species in section *Fumigati*. *Medical Mycology* **45**: 225–232.
- Lau A, Chen S, Sorrell T, Carter D, Malik R, Martin P, Halliday C (2007). Development and clinical application of a panfungal PCR assay to detect and identify fungal DNA in tissue specimens. *Journal of Clinical Microbiology* **45**: 380–385.
- Lin A, Huang K, Hwu L, Tzean SS (1994). Production of type II ribotoxins by *Aspergillus* species and related fungi in Thailand. *Toxicon* **33**: 105–110.
- Lonial S, Williams L, Carrum G, Ostrowski M, McCarthy PJr (1997). *Neosartorya fischeri*: an invasive fungal pathogen in an allogeneic bone marrow transplant patient. *Bone Marrow Transplantation* **19**: 753–755.
- Malloch, D. and Cain, R. F. 1972. The Trichocomataceae: Ascomycetes with *Aspergillus*, *Paecilomyces*, and *Penicillium* imperfect states. *Canadian Journal of Botany* **50**: 2613–2628.
- Marr KA, Patterson T, Denning D (2002). Aspergillosis. Pathogenesis, clinical manifestations, and therapy. *Infectious Disease Clinics of North America* **16**: 875–894.
- Matsumoto N, Shiraga H, Takahashi K, Kikuchi K, Ito K (2002). Successful treatment of *Aspergillus* peritonitis in a peritoneal dialysis patient. *Pediatric Nephrology* **17**: 243–245.
- McCowen MC, Callender M, Lawlis JF Jr (1951). Fumagillin (H-3), a new antibiotic with amebicidal properties. *Science* **113**: 202–203.
- Mellado E, Alcazar-Fuoli L, Garcia-Effro G, Alastruey-Izquierdo NA, Cuenca-Estrella M, Rodríguez-Tudela JL (2006). New resistance mechanisms to azole drugs in *Aspergillus fumigatus* and emergence of antifungal drugs-resistant *A. fumigatus* atypical strains. *Medical Mycology* **44**: S367–S371.
- Moragues MD, Brena S, Miranda I, Laporta R, Muñoz M, Pontón J, del Palacio A (2006). Colonización pulmonar por *Neosartorya udagawae* en un paciente trasplantado de pulmón. *VIII Congreso Nacional Micología, Barcelona, Spain*, D15.
- Muhlenfeld A, Achenbach H (1988). Asperopentyn, a novel acetylenic cyclohexene epoxide from *Aspergillus duricaulis*. *Phytochemistry* **27**: 3853–3855.
- Müllbacher A, Eichner RD (1984). Immunosuppression *in vitro* by a metabolite of a human pathogenic fungus. *Proceedings of the National Academy of Sciences U.S.A.* **81**: 3835–3837.
- Omolo JO, Anke H, Chhabra S, Sterner O (2000). New variotin analogues from *Aspergillus viridinutans*. *Journal of Natural Products* **63**: 975–977.
- Padhye AA, Godfrey JH, Chandler FW, Peterson SW (1994). Osteomyelitis caused by *Neosartorya pseudofischeri*. *Journal of Clinical Microbiology* **32**: 2832–2836.
- Peterson SW (1992). *Neosartorya pseudofischeri* sp. nov. and its relationship to other species in *Aspergillus* section *Fumigati*. *Mycological Research* **96**: 547–554.
- Pringle A, Baker DM, Platt JL, Wares JP, Latge JP, Taylor JW (2005). Cryptic speciation in the cosmopolitan and clonal human pathogenic fungus *Aspergillus fumigatus*. *Evolution* **59**: 1886–1899.
- Raper KB, Fennell DI (1965). *The genus Aspergillus*. Baltimore: Williams & Wilkins.
- Rinyu E, Varga J, Kozakiewicz Z, Ferenczy L (2000). Phenotypic and genotypic variability within *Aspergillus* section *Fumigati*. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification*. (Samson RA, Pitt JI, eds). Amsterdam: Harwood Academic Publishers: 483–492.
- Rydholm C, Szakács G, Lutzoni F (2006). Low genetic variation and no detectable population structure in *Aspergillus fumigatus* compared to closely related *Neosartorya* species. *Eukaryotic Cell* **5**: 650–657.
- Samson RA (1989). Filamentous fungi in food and feed. *Journal of Applied Bacteriology* (Suppl) **67**: 27S–35S.
- Samson RA, Hoekstra ES, Frisvad JC (eds) (2004). *Introduction to food and airborne fungi*. 7th ed. Utrecht: Centraalbureau voor Schimmelcultures.
- Samson RA, Nielsen PV, Frisvad JC (1990). The genus *Neosartorya*: Differentiation by scanning electron microscopy and mycotoxin profiles. In: *Modern concepts in Penicillium and Aspergillus classification*. (Samson RA, Pitt JI, eds). New York: Plenum Press: 455–467.
- Shin HD, McClendon S, Le T, Taylor F, Chen RR (2006). A complete enzymatic recovery of ferulic acid from corn residues with extracellular enzymes from *Neosartorya spinosa* NRRL 185. *Biotechnology and Bioengineering* **95**: 1108–1115.
- Smedsgaard J (1997). Micro-scale extraction procedure for standardized screening of fungal metabolite production in cultures. *Journal of Chromatography A* **760**: 264–270.
- Someya A, Yaguchi T, Udagawa S (1999). *Neosartorya sublevispora*, a new species of soil-borne Eurotiales. *Mycoscience* **40**: 405–409.
- Summerbell RC, de Repentigny L, Chartrand C, St Germain G (1992). Graft-related endocarditis caused by *Neosartorya fischeri* var. *spinosa*. *Journal of Clinical Microbiology* **30**: 1580–1582.
- Swofford T (2000). *PAUP: Phylogenetic analysis using parsimony*. v. 4.0. Sunderland: Sinauer Associates.
- Takada M, Horie Y, Abliz P (2001). Two new heterothallic *Neosartorya* from African soil. *Mycoscience* **42**: 361–367.
- Takada M, Udagawa S (1985). A new species of heterothallic *Neosartorya*. *Mycotaxon* **24**: 395–402.
- Tamura K, Nei M (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **10**: 512–526.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997). The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* **25**: 4876–4882.
- Tomoda H, Hishida H, Kim YK, Obata R, Sunazaka T, Omura S, Bordner J, Guadliana M, Dormer PG, Smith AB (1994). Relative and absolute stereochemistry of pyrropropene A, a potent bioavailable inhibitor of acyl-CoA:cholesterol acyltransferase (ACAT). *Journal of the American Chemical Society* **116**: 12097–12098.
- Tournas V (1994). Heat-resistant fungi of importance to the food and beverage industry. *Critical Reviews in Microbiology* **20**: 243–263.
- Udagawa S, Toyazaki N, Tsubouchi H (1993). *Neosartorya primulina*, a new species of food-borne ascomycetes. *Mycotaxon* **47**: 359–366.
- Udagawa S, Tsubouchi H, Horie Y (1991). *Neosartorya hiratsukae*, a new species of food-borne Ascomycetes. *Transactions of the Mycological Society of Japan* **32**: 23–29.
- Varga J, Rinyu E, Kevei É, Tóth B, Kozakiewicz Z (1998). Double-stranded RNA mycoviruses in species of *Aspergillus* sections *Circumdati* and *Fumigati*. *Canadian Journal of Microbiology* **44**: 569–574.
- Varga J, Rinyu E, Kiss I, Botos B, Kozakiewicz Z (1997). Carbon source utilization and isoenzyme analysis as taxonomic aids for toxigenic *Neosartorya* species and their relatives. *Acta Microbiologica et Immunologica Hungarica* **44**: 1–11.
- Varga J, Tóth B, Rigó K, Debets F, Kozakiewicz Z (2000). Genetic variability within the *Aspergillus viridinutans* species. *Folia Microbiologica* **45**: 423–428.
- Varga J, Vida Z, Tóth B, Debets F, Horie Y (2000). Phylogenetic analysis of newly described *Neosartorya* species. *Antonie van Leeuwenhoek* **77**: 235–239.
- Wang L, Yokoyama K, Miyaji M, Nishimura K (2000). Mitochondrial cytochrome b gene analysis of *Aspergillus fumigatus* and related species. *Journal of Clinical Microbiology* **38**: 1352–1358.



- Weisleder D, Lillehoj EB (1971). Structure of viriditoxin, a toxic metabolite of *Aspergillus viridinutans*. *Tetrahedron Letters* **48**: 4705–4706.
- White TJ, Bruns T, Lee S, Taylor J (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: A guide to methods and applications*. Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. New York: Academic Press: 315–322.
- Wong SM, Musza LL, Kydd GC, Kullnig R, Gillum AM, Cooper R (1993). Fiscalins: new substance P inhibitors produced by the fungus *Neosartorya fischeri*. Taxonomy, fermentation, structures, and biological properties. *Journal of Antibiotics* **46**: 545–553.
- Yaguchi T, Horie Y, Tanaka R, Matsuzawa T, Ito J, Nishimura K (2007). Molecular phylogenetics of multiple genes on *Aspergillus* section *Fumigati* isolated from clinical specimens in Japan. *Japanese Journal of Medical Mycology* **48**: 37–46.
- Yaguchi T, Someya A, Udagawa S (1994). A new species of *Neosartorya* from Taiwan soil. *Mycoscience* **35**: 309–313.