

Fcγ receptor IIIb polymorphism and use of glucocorticoids at baseline are associated with infusion reactions to infliximab in patients with rheumatoid arthritis

Ayumi Okuyama,¹ Hayato Nagasawa,¹ Katsuya Suzuki,¹ Hideto Kameda,^{1,2} Harumi Kondo,² Kouichi Amano,¹ Tsutomu Takeuchi^{1,2}

► Additional data (supplementary figure) are published online only. To view these files please visit the journal online <http://ard.bmj.com>

¹Rheumatology/Clinical Immunology, Saitama Medical Center, Saitama Medical University, Saitama, Japan
²Department of Internal Medicine, School of Medicine, Keio University, Tokyo, Japan

Correspondence to

Professor Tsutomu Takeuchi, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan; tsutake@z5.keio.jp

Accepted 29 August 2010
Published Online First
27 October 2010

ABSTRACT

Objective Infusion reaction is a major adverse event in patients with rheumatoid arthritis (RA) treated with infliximab. The possible factors including Fcγ receptor (FcγR) polymorphism associated with the development of infusion reactions in patients with RA receiving infliximab were prospectively examined.

Methods 96 patients with RA were enrolled and scheduled to receive infliximab at a dose of 3 mg/kg at weeks 0, 2 and 6 and every 8 weeks thereafter. Genetic polymorphisms for FcγR were examined in *FCGR3A* 176F/V and *FCGR3B* NA1/2 alleles by allele-specific PCR analysis.

Results An infusion reaction was observed in 17 patients (18%) during 52 weeks of treatment with infliximab. The *FCGR3B* NA1/NA1 genotype was found in 75% of the patients with infusion reactions and in only 37% of those without ($p=0.01$), whereas the *FCGR3A* 176F/V genotype was equally distributed in the patients with or without infusion reactions. Glucocorticoids were used in 53% of the patients who developed an infusion reaction and in 80% of those without an infusion reaction ($p=0.02$). A multivariable logistic regression model showed that the *FCGR3B* NA1/NA1 genotype and use of glucocorticoids at baseline could be used as independent predictive factors for infusion reactions (OR 6.1 (95% CI 1.9 to 24.3) and OR 0.26 (95% CI 0.08 to 0.84), respectively). The presence of anti-infliximab antibody during infliximab treatment was also associated with infusion reactions.

Conclusion *FCGR3B* NA1/NA1 genotype, use of glucocorticoids and the presence of anti-infliximab antibody accounted for nearly all patients with RA who developed infusion reactions.

INTRODUCTION

Biological agents targeting a specific molecule provide an effective means for therapeutic management of rheumatoid arthritis (RA) due to their specificity and powerful functional capabilities, which has resulted in a paradigm shift in the treatment strategy of this disease.¹⁻⁴ Despite their effectiveness, several adverse drug reactions associated with the use of biological agents have been identified, such as opportunistic infections and the development of hypersensitivity/infusion reactions. For example, approximately 10-23% of patients with RA experience infusion reactions (including fever, malaise, headaches, erythema and urticaria) to infliximab,

a chimeric monoclonal IgG1 antibody against tumour necrosis factor α (TNF α), in combination with methotrexate (MTX).⁵⁻⁹ Although much less frequent, severe infusion reactions such as anaphylactic/anaphylactoid symptoms are also observed in patients with RA.⁸ Despite the fact that most of these reactions are only mild or moderate in severity, they may lead to discontinuation of treatment, which is of particular concern and highly relevant to daily clinical practice.⁹ It is tempting to speculate that biological agents may induce these reactions in a portion of susceptible patients by causing hypersensitivity to the given biological agents,¹⁰ in part through the immunogenicity of the agents or by direct effects on cellular functions through the Fc portion of the products.¹¹

The Fc portion of IgG-based biological agents can bind to Fcγ receptors (FcγR) for clearing the agents and even affecting cellular functions including phagocytosis, antibody-dependent cellular cytotoxicity and neutrophil activation.¹² FcγR consist of three major families which are encoded by eight genes: *FCGR1A*, *FCGR1B*, *FCGR1C*, *FCGR2A*, *FCGR2B*, *FCGR2C*, *FCGR3A* and *FCGR3B*. Functional allelic polymorphisms leading to distinct effector capabilities have been identified in the receptors FcγRIIa, FcγRIIb, FcγRIIIa and FcγRIIIb.¹²

Since the *FCGR2A* 131 H/R allele strongly influences the ability of FcγRIIa to bind human IgG2 but has only limited effects on IgG1 and IgG3 binding,¹² we focused on the *FCGR3A* 158F/V and *FCGR3B* NA1/NA2 alleles and prospectively examined the possible association of these *FCGR* alleles, development of antibody to infliximab and clinical parameters with development of infusion reactions to infliximab in patients with RA.

METHODS

Patients and treatment

Consecutive patients with RA who fulfilled the 1987 revised criteria of the American College of Rheumatology for the classification of RA¹⁵ and satisfied the Japanese guidelines for the use of anti-TNF biological agents¹⁴ were invited to participate in the study. Ninety-six patients showing incomplete response to MTX were enrolled after obtaining their written informed consent. The 2008 Declaration of Helsinki and the 2008 Ethical Guidelines for Clinical Research by the Japanese Ministry of Health, Labour and Welfare were



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://ard.bmj.com/info/unlocked.dtl>

strictly followed in this study. The patients were scheduled to receive infliximab at a dose of 3 mg/kg at weeks 0, 2, 6 and subsequently every 8 weeks added to MTX at the Saitama Medical Center between September 2003 and March 2008. The management of infusion was performed using a previously reported protocol¹⁵ approved by the University Institutional Review Board. We did not perform any premedication including histamine antagonists, paracetamol and additional glucocorticoids until the first infusion reaction developed in individual patients. Blood samples were taken for genotyping before initiation of infliximab, and anti-infliximab antibody (AIA) was measured at each infusion visit during the 52-week study.

Assessment of efficacy and safety

Patients were followed up longitudinally with examinations at baseline and at each regular infusion visit and emergency visit during the 52-week study. To monitor disease activity and disability, the 28-joint Disease Activity Score based on C reactive protein (DAS28-CRP) and serum levels of CRP, matrix metalloproteinase-3 (MMP-3) and Health Assessment Questionnaire-Disease Index (HAQ-DI) were determined. The attending physicians recorded any adverse drug reactions at baseline, at each regular infusion visit and emergency visit during the 52-week study period.

Measurement of AIA

AIA was measured using an ELISA kit (Immunodiagnostik, Bensheim, Germany). Briefly, serum samples were diluted with phosphate buffered saline (PBS) and added to 96-well plastic plates coated with the F(ab)₂ portion of infliximab to avoid interference with rheumatoid factor (RF). After incubating overnight at 4°C, the wells were washed with PBS and horseradish peroxidase-labelled infliximab was added. After incubation for 60 min, the plates were extensively washed with PBS, followed by addition of substrates into each well. The OD₄₅₀-OD₆₂₀ was recorded using an ELISA reader.

Determination of FcγR polymorphisms

Heparinised venous blood was collected from patients and chromosomal DNA was isolated using phenol-chloroform extraction. Genetic polymorphisms for FcγR were examined in FCGR3A 176F/V and FCGR3B NA1/2 alleles by allele-specific PCR analysis, as previously described.^{16 17} PCR products were separated on 3% agarose gels and visualised under ultraviolet light using a photoimager.

Statistical analysis

Baseline variables of patients with RA were analysed for association with development of infusion reactions using a χ² test for categorical variables and Student t test for continuous variables. Univariate logistic regression analysis was used to screen for potential predictor variables, and a stepwise selection process was used to generate a multivariate model for potential predictors of infusion reactions. All statistical analyses were performed on a Mac OS X platform (Sun Microsystems, Palo Alto, California, USA) using JMP Version 8.0.2.

RESULTS

Clinical characteristics of the patients with RA who developed infusion reactions

As shown in table 1, the development of infusion reactions, concomitant medications and the continuation of infliximab varied in individual patients. The observed infusion reactions were typically mild or moderate and included chills, fever, erythema and urticaria; however, one patient developed a severe anaphylactoid reaction and hypotension. Prior to the study, this patient had been enrolled in a clinical trial for infliximab, and the severe infusion reaction developed in this study after the second infusion, which represented a 3-year interval since he last received infliximab in the clinical trial. The most common manifestation in the patients was skin eruption (n=10), followed by fever (n=3), nausea/vomiting (n=3) and headache (n=2). Although these manifestations were not considered severe, nearly all of

Table 1 Demographic and clinical characteristics of patients with infusion reactions

No	Gender	Age (year)	Duration (month)	DAS28 at baseline	CRP (mg/dl)	RF (IU/ml)	ANA (×)	MMP-3 (ng/ml)	HAQ-DI	Clinical manifestation	Time of infusion reaction (weeks)	MTX (mg/week)	PRED (mg/day)	Adherence to infliximab
1	F	51	60	6.9	7.8	12	160	346	2.6	Chills, fever, dyspnoea	38	13	0	Discontinue
2	F	52	9	6.6	6.9	444	0	188	2.1	Hot flushes, headache, subfever	1	8	10	Continue
3	M	69	624	5.7	3.9	191	80	180	1.8	Anaphylactoid reaction, hypotension	2 (after 3-year interval)	6	5	Discontinue
4	F	40	50	4.8	2.9	864	80	200	1.3	Urticaria	14	12.5	0	Discontinue
5	F	50	97	3.5	0.3	23	80	121	0.3	Urticaria	30	10.5	5	Discontinue
6	F	68	48	4.5	3.2	42	80	95	0.13	Erythema	14	8	0	Discontinue
7	M	65	240	4.8	2.8	1080	320	85	2.6	Hot flushes, headache, nausea	14	10	3	Discontinue
8	F	52	255	5.7	3.3	845	160	191	1.3	Erythema	30	8	0	Discontinue
9	F	31	29	4.8	0.1	17	1280	98	0.5	Nausea	44	8	0	Discontinue
10	F	60	210	4.2	1.8	250	80	359	0.6	Urticaria	14	8	0	Discontinue
11	F	52	168	3.5	0.3	15	1280	121	0.3	Urticaria	2	6	0	Discontinue
12	F	57	360	6.6	6.4	98	1280	360	1.5	Erythema	14	6	4	Discontinue
13	F	33	21	6.6	5.2	122	80	612	1.1	Erythema	14	10	4	Continue
14	F	37	9	7.5	5.6	<5	0	280	1.9	Fever	14	6	10	Discontinue
15	F	37	60	4.3	1.4	446	80	ND	0.8	Erythema	30	8	0	Continue
16	F	63	4	7.4	11.0	1060	80	412	2.5	Nausea, vomiting	14	8	7.5	Discontinue
17	F	64	134	6.0	0.8	<5	1280	253	1.9	Erythema	22	6	0	Discontinue

ANA, antinuclear antibody; CRP, C reactive protein; DAS28, 28-joint Disease Activity Score; HAQ-DI, Health Assessment Questionnaire-Disease Index; MMP, matrix metalloproteinase; MTX, methotrexate; ND, not done; PRED, prednisolone; RF, rheumatoid factor.

the patients (15 of 17) discontinued infliximab due to the development of these infusion reactions.

Demographics and clinical characteristics of the patients

As shown in table 2, the mean age of the patients was 54 years and 83% were women. The mean disease duration was 8 years, RF positivity was 90%, the mean DAS28 score was 5.2 and the mean serum HAQ-DI level was 1.5, suggesting that the patients with RA enrolled in this study were established, active and disabled.

There were no significant differences in the demographics and clinical characteristics between patients who developed infusion reactions and those who did not, with the exception of the concomitant use of glucocorticoids. Glucocorticoids were used in 53% of patients with infusion reactions compared with 80% of patients without infusion reactions. In contrast, no significant differences in the demographics and clinical characteristics between patients who continued infliximab and those who discontinued its use were demonstrated in this study.

FcγR polymorphism

Allele distributions in the patients with RA for the *FCGR3A* 176F/V and *FCGR3B* NA1/NA2 polymorphisms are summarised in table 3. In this cohort of patients, the distribution of the *FCGR3A* high-affinity genotype V/V was 6% whereas the V/F and F/F genotypes were observed with a frequency of 48% and 46%, respectively. The occurrence of the V/V genotype was slightly lower than in the healthy Japanese population (8.6%).¹⁸ The V/V genotype was enriched in patients with RA with infusion reactions (12%) over those without such reactions (5%), but the difference was not significant. The distributions of the genotypes for the *FCGR3A* 176F/V allele between patients with and without adherence to infliximab were comparable.

The high-affinity genotype of *FCGR3B* (NA1/NA1) was the most prevalent genotype in the patients with RA (44%), while the NA1/NA2 and NA2/NA2 genotypes were found in 33% and 23% of patients, respectively. The *FCGR3B* NA1/NA1 genotype was found in 70% of patients with infusion reactions but was present in only 37% of patients without infusion reactions, indicating that this genotype is associated with the development of infusion reactions. In contrast, the low-affinity genotypes NA1/NA2 and NA2/NA2 were only observed in 18% and

6%, respectively, of patients who developed infusion reactions, which is much lower than the 37% and 27%, respectively, of those without reactions. Analyses confirmed that the distribution of the *FCGR3B* genotypes between patients with and without infusion reactions was significantly different ($p=0.01$). On the other hand, such differences in distribution of the *FCGR3B* genotypes were not observed between those with and without adherence to infliximab.

The presence of AIA during each visit was found to be positive in 19% of the patients. The development of AIA was significantly higher in patients with infusion reactions than in those without (65% and 9%, respectively, $p<0.001$), whereas no difference was observed in patients with or without adherence to infliximab (15% and 22%, respectively, $p=0.40$).

Potential predictive variables for infusion reactions

The impact of the presence or absence of risk factors identified on the development of infusion reactions is summarised and compared in figure 1. In order to create a multivariate model of potential predictors of infusion reactions, we first screened a series of baseline variables which included age, gender, duration of RA, DAS score, levels of CRP, MMP-3, HAQ-DI, RF and anti-nuclear antibody, doses of MTX and prednisolone equivalent, glucocorticoid use and *FCGR3A* V/V+V/F and *FCGR3B* NA1/NA1 genotypes by univariate logistic regression. As shown in table 4, the *FCGR3B* NA1/NA1 genotype and use of glucocorticoids were significant predictive variables, consistent with the above analyses. Even by stepwise selection, these variables were identified as significant factors associated with infusion reactions (data not shown).

We next subjected these two variables to multivariate logistic regression analyses which allowed for differences in patients with or without infusion reactions to be adjusted. From these analyses, the *FCGR3B* NA1/NA1 genotype and use of glucocorticoids were finally identified as significant independent variables associated with the development of infusion reactions. The ORs of the *FCGR3B* NA1/NA1 genotype and the use of glucocorticoids were 6.1 (95% CI 1.9 to 24.3, $p=0.002$) and 0.26 (95% CI 0.08 to 0.84, $p=0.025$), respectively (table 5).

Finally, we examined the potential additive effect of the three identified factors on the development of infusion reactions. Patients with RA with only one factor had a rate of infusion

Table 2 Patient characteristics and infliximab-related outcome

Subject	Total (n=96)	Infusion reaction during 52 weeks			Adherence to infliximab during 52 weeks		
		Yes (n=17)	No (n=79)	p Value	Continued (n=46)	Discontinued (n=50)	p Value
Age (years)	54±13	52±12	55±13	0.82	56±12	53±14	0.16
Female (%)	83	88	81	0.55	89	78	0.14
Disease duration (years)	8±9	11±14	8±8	0.18	8±7	9±10	0.57
RF positivity (%)	90	88	90	0.84	91	88	0.60
RF (IU/ml)	213±321	324±394	189±300	0.10	190±319	145±324	0.79
ANA positivity (%)	32	41	30	0.39	30	34	0.71
ANA titre (×)	273±616	377±150	251±69	0.20	341±813	211±346	0.16
Stage I+II (%)	44	47	43	0.46	33	43	0.05
Class 1+2 (%)	65	82	62	0.18	65	64	0.53
DAS28	5.2±1.3	5.4±1.6	5.2±1.3	0.69	5.1±1.5	5.4±1.2	0.22
CRP (mg/dl)	4.0±3.3	4.1±3.0	3.9±3.4	0.87	3.5±3.4	4.3±3.2	0.26
MMP-3 (ng/ml)	293±287	146±38	301±309	0.36	340±374	255±189	0.17
HAQ-DI	1.5±0.8	1.4±0.9	1.5±0.7	0.76	1.5±0.8	1.5±0.7	0.96
MTX dose (mg/week)	8±3	8±2	8±3	0.35	8±4	9±3	0.95
Use of glucocorticoids (%)	75	53	80	0.02*	74	76	0.81
Glucocorticoid dose (mg/day)	4±3	3±4	5±3	0.92	5±4	4±3	0.19

ANA, antinuclear antibody; CRP, C reactive protein; DAS28, 28-joint Disease Activity Score; HAQ-DI, Health Assessment Questionnaire-Disease Index; MMP, matrix metalloproteinase; MTX, methotrexate; RF, rheumatoid factor.

Table 3 Association of Fcγ receptor polymorphism and anti-infliximab antibody (AIA) with infliximab-related outcome

Subject	Total (n=96)	Infusion reaction during 52 weeks		p Value	Adherence to infliximab during 52 weeks		p Value
		Yes (n=17)	No (n=79)		Continued (n=46)	Discontinued (n=50)	
<i>FCGR3A</i> 176F/V							
F/F	44 (46)	7 (41)	37 (47)	0.58	22 (48)	22 (44)	0.53
F/V	46 (48)	8 (47)	38 (48)		20 (44)	26 (52)	
V/V	6 (6)	2 (12)	4 (5)		4 (9)	2 (4)	
<i>FCGR3B</i> NA1/NA2							
NA1/NA1	42 (44)	13 (77)	29 (37)	0.01*	16 (35)	26 (52)	0.23
NA1/NA2	32 (33)	3 (18)	29 (37)		18 (39)	14 (28)	
NA2/NA2	22 (23)	1 (6)	21 (27)		12 (26)	10 (20)	
AIA							
(+)	18 (19)	11 (65)	7 (9)	<0.01*	7 (15)	11 (22)	0.40
(-)	78 (81)	6 (35)	72 (91)		39 (85)	39 (78)	

Values are numbers (%) unless otherwise indicated.




Figure 1 Percentage of patients with rheumatoid arthritis who developed an infusion reaction during the 52-week study in those with or without anti-infliximab antibody, those with or without *FCGR3B* NA1/NA1 genotype and those with or without concomitant use of glucocorticoids.

Table 4 Univariate logistic regression analysis for potential predictors of infusion reaction

Variables	Estimates	SE	OR	p Value
Age (years)	-0.02	0.02	0.98	0.37
Gender (female vs male)	0.24	0.40	1.62	0.55
Duration (years)	0.04	0.03	1.04	0.12
DAS28	0.08	0.21	1.08	0.69
CRP (mg/dl)	0.01	0.08	1.01	0.87
MMP-3 (per 10 ng/ml)	-0.01	0.01	0.99	0.55
HAQ-DI	-0.12	0.39	0.89	0.75
RF (per 10 IU/ml)	0.01	0.00	1.00	0.14
ANA (×)	0.00	0.00	1.00	0.46
MTX (mg/week)	0.03	0.09	1.03	0.77
MTX duration (months)	0.00	0.01	1.00	0.72
Use of glucocorticoids	-0.63	0.28	0.53	0.03*
Dose of prednisolone (mg/day)	-0.13	0.09	0.88	0.14
<i>FCGR3A</i> V/V+V/F vs F/F	0.11	0.27	1.12	0.67
<i>FCGR3B</i> NA1/1 vs NA1/2+2/2	0.86	0.31	2.37	<0.01*

ANA, antinuclear antibody; CRP, C reactive protein; DAS28, 28-joint Disease Activity Score; HAQ-DI, Health Assessment Questionnaire-Disease Index; MMP, matrix metalloproteinase; MTX, methotrexate; RF, rheumatoid factor.

reactions of 11%, whereas those with two or three factors had rates of 53% and 100%, respectively (see figure in online supplement), demonstrating that these three factors additively account for the infusion reactions to infliximab observed in the 96 patients with RA examined in this study.

DISCUSSION

The results of this study show that the high-affinity *FCGR3B* NA1/NA1 genotype and the absence of glucocorticoid use at baseline are the main independent predictive factors for the development of infusion reactions to infliximab in Japanese patients with RA. This study is the first to demonstrate a role for the *FCGR3B* NA1/NA1 genotype in infusion reactions to infliximab.

Mild to moderate and severe infusion reactions were observed in 18% and 1%, respectively, of the Japanese patients with RA treated with infliximab during the 52-week course of this study. This result is consistent with previous studies showing that the incidence of infusion reactions to infliximab was approximately 10–23% of patients with RA per year.^{5-7 8 15} The most frequent manifestations of the infusion reactions observed in this study were consistent with those reported for infusion reactions to infliximab in clinical practice settings in both the USA⁵ and Japan.⁸ A significant proportion of these patients subsequently discontinued infusion, implying that infusion reactions may account, as well as lack of efficacy, for the low retention rate of infliximab observed in the first year of treatment,^{9 19-21} while others do not.^{22 23} The inconsistency of the retention rate for infliximab among studies may result from the variable management strategies used to reduce infusion reactions, which include premedication,⁵ gradual increases in infusion speed,⁷ adjustment of infusion intervals²⁴ and dose increments,²⁵ among others.

It has been shown using radiolabelled infliximab in patients with RA that the formation of infliximab and anti-infliximab complexes were found in non-responders, one of which showed an anaphylactoid reaction, resulting in higher liver/spleen uptake and rapid clearance of infliximab.¹¹ It was recently observed that the IgE class of AIA is associated with the development of severe infusion reactions,²⁶ whereas true IgE-mediated hypersensitivity is not related to acute infusion reactions in most cases.²⁷ This possibility in one patient who developed an anaphylactoid reaction in this cohort remains to be addressed.

Table 5 Multivariate model of predictors of infusion reaction by logistic regression

	Estimates	OR	95% CI	p Value
<i>FCGR3B</i> NA1/1: NA2/2+NA1/2	0.90	6.1	1.9 to 24.3	<0.01
Glucocorticoid use	-0.68	0.26	0.08 to 0.84	0.03

We found that 19% of the patients with RA developed AIA during the 52-week study period. The reported prevalence of AIA in patients with RA varies from 12% to 44% and appears to be inversely proportional to the serum levels of infliximab and therapeutic response.²⁸ The dose of MTX did not significantly affect the development of AIA in our study (data not shown). In this study, infusion reactions frequently occurred between 14 and 30 weeks after initiating infliximab. Since the infusion interval was increased to 8 weeks after induction at 0, 2 and 6 weeks, the serum trough levels of infliximab probably lowered at the infusions with 8-week intervals, supporting the above observation in this study. However, the precise mechanisms related to AIA and the development of infusion reactions are not fully understood and require further study.

Numerous studies have analysed the possible association between *FCGR2A* and *FCGR3A* polymorphisms and the efficacy of biological agents against TNF α in patients with RA.^{29–31} However, the association between these polymorphisms and the adverse effects of anti-TNF α has not been fully explored. In this regard, the *FCGR3B* NA2 allele has been shown to be associated with urinary tract infections in patients with RA treated with MTX or etanercept.³² Fc γ RIIIb is expressed exclusively on neutrophils, eosinophils¹² and basophils.³³ The isoform containing the *FCGR3B* NA1 allele produces larger phagocytic, oxidative burst and degranulation responses than the *FCGR3B* NA2 allele.³² Thus, the *FCGR3B* NA1/NA1 genotype with high affinity to Ig present on the surface of neutrophils, eosinophils and basophils may account for the higher incidence of infusion reactions to infliximab. In addition, optimal ligand concentrations leading to formation of immune complexes may allow binding to Fc γ RIIIb and subsequent activation of cells.³⁴ Recently, variation in the copy number of *FCGR3B* has been shown to be associated with susceptibility to systemic autoimmunity.³⁵ Copy number variation of *FCGR3B* may play a role in infusion reactions and warrants further examination.

One may speculate that glucocorticoids may interfere with the binding, activation and effector function of immune cells, thereby reducing the severity and frequency of infusion reactions. Some of these effects may be shared with MTX. In this study, the concomitant use of oral glucocorticoids was significantly associated with a reduced risk of developing an infusion reaction, consistent with recent reports.^{9–36} Therefore, not only the moderate- to high-dose glucocorticoids given as a premedication, but also the low-dose daily glucocorticoids may be a potent inhibitor of infusion reactions. Fc γ RIIIb is a phosphatidyl inositol-linked cell surface protein and thus lacks any self-kinase activity. Instead, Fc γ RIIa is coupled with Fc γ RIIIb to transduce signals. As one of the functional polymorphisms in Fc γ RIIa is the 131H allele which confers receptor affinity to IgG2 subclass, it is unlikely that this receptor plays a direct role in binding to the IgG1 monoclonal antibody infliximab. Nevertheless, the possibility that other polymorphisms of *FCGR2A* associated with receptor function contribute to infusion reactions should be examined.

There are a number of limitations that warrant mention. First, the number of patients examined in this study was insufficient

to demonstrate an association between *FCGR3A* polymorphism and infusion reactions. Second, in addition to the small number of patients, other aspects of the study design also imposed limitations. For example, the study was designed to monitor infusion reactions during 52 weeks, meaning that infusion reactions appearing after 52 weeks could not be addressed. Third, this was an open-label study which may have affected the incidence and severity of infusion reactions and the response and retention rates. However, the incidence and the types of infusion reactions in this study were comparable to those of previous reports, as were the other results related to the efficacy and retention rate of infliximab.^{37–38} Also, AIA is typically developed during the treatment period and this information would not be available before starting infliximab; it is apparent that the presence of AIA is limited in its use as a predictive marker. When the two risk factors excluding AIA are used as predictive variables, they were still associated with approximately 40% of infusion reactions observed in this study (data not shown).

In summary, we have shown that the *FCGR3B* NA1/NA1 genotype and the absence of glucocorticoid usage are predictive factors of infusion reactions in patients with RA. Premedication for infliximab may therefore not be necessary for all patients but only for those with the *FCGR3B* NA1/NA1 genotype without daily glucocorticoids.

Acknowledgements The authors thank all medical staff of Saitama Medical University and Keio University.

Funding Part of this study was supported by a Research Grant-In-Aid for Scientific Research by the Ministry of Health, Labour and Welfare of Japan.

Competing interests None of authors are industry employees. TT has received consultant fees from Mitsubishi-Tanabe Pharma, Bristol-Myers-Squibb and Novartis, and lecture fees from Mitsubishi-Tanabe Pharma, Takeda Pharmaceutical, Abbott, Eisai Pharma and Chugai Pharma. HK received honoraria from Mitsubishi-Tanabe Pharma, Centocor, Wyeth Japan, Takeda Pharmaceutical, Abbott, Eisai Pharma and Chugai Pharma.

Patient consent Obtained.

Ethics approval This study was conducted with the approval of the internal review board of Saitama Medical University.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Scheinecker C, Redlich K, Smolen JS. Cytokines as therapeutic targets: advances and limitations. *Immunity* 2008;**28**:440–4.
- Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum* 2008;**59**:762–84.
- van der Kooij SM, le Cessie S, Goekoop-Ruiterman YP, et al. Clinical and radiological efficacy of initial vs delayed treatment with infliximab plus methotrexate in patients with early rheumatoid arthritis. *Ann Rheum Dis* 2009;**68**:1153–8.
- Nam JL, Winthrop KL, van Vollenhoven RF, et al. Current evidence for the management of rheumatoid arthritis with biological disease-modifying antirheumatic drugs: a systematic literature review informing the EULAR recommendations for the management of RA. *Ann Rheum Dis* 2010;**69**:976–86.
- Wasserman MJ, Weber DA, Guthrie JA, et al. Infusion-related reactions to infliximab in patients with rheumatoid arthritis in a clinical practice setting: relationship to dose, antihistamine pretreatment, and infusion number. *J Rheumatol* 2004;**31**:1912–17.
- Kapetanovic MC, Larsson L, Truedsson L, et al. Predictors of infusion reactions during infliximab treatment in patients with arthritis. *Arthritis Res Ther* 2006;**8**:R131.
- Lequerré T, Vittecoq O, Klemmer N, et al. Management of infusion reactions to infliximab in patients with rheumatoid arthritis or spondyloarthritis: experience from an immunotherapy unit of rheumatology. *J Rheumatol* 2006;**33**:1307–14.
- Takeuchi T, Tatsuki Y, Nogami Y, et al. Postmarketing surveillance of the safety profile of infliximab in 5000 Japanese patients with rheumatoid arthritis. *Ann Rheum Dis* 2008;**67**:189–94.
- Du Pan SM, Dehler S, Ciurea A, et al. Comparison of drug retention rates and causes of drug discontinuation between anti-tumor necrosis factor agents in rheumatoid arthritis. *Arthritis Rheum* 2009;**61**:560–8.
- Radstake TR, Svenson M, Eijsbouts AM, et al. Formation of antibodies against infliximab and adalimumab strongly correlates with functional drug levels and clinical responses in rheumatoid arthritis. *Ann Rheum Dis* 2009;**68**:1739–45.

11. **van der Laken CJ**, Voskuyl AE, Roos JC, *et al.* Imaging and serum analysis of immune complex formation of radiolabelled infliximab and anti-infliximab in responders and non-responders to therapy for rheumatoid arthritis. *Ann Rheum Dis* 2007;**66**:253–6.
12. **Hogarth PM**. Fc receptors are major mediators of antibody based inflammation in autoimmunity. *Curr Opin Immunol* 2002;**14**:798–802.
13. **Arnett FC**, Edworthy SM, Bloch DA, *et al.* The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;**31**:315–24.
14. **Koike R**, Takeuchi T, Eguchi K, *et al.* Update on the Japanese guidelines for the use of infliximab and etanercept in rheumatoid arthritis. *Mod Rheumatol* 2007;**17**:451–8.
15. **Cheifetz A**, Smedley M, Martin S, *et al.* The incidence and management of infusion reactions to infliximab: a large center experience. *Am J Gastroenterol* 2003;**98**:1315–24.
16. **Edberg JC**, Langefeld CD, Wu J, *et al.* Genetic linkage and association of Fcγ₃ receptor IIIA (CD16A) on chromosome 1q23 with human systemic lupus erythematosus. *Arthritis Rheum* 2002;**46**:2132–40.
17. **Magnusson V**, Johanneson B, Lima G, *et al.* Both risk alleles for Fcγ₃RIIA and Fcγ₃RIIA are susceptibility factors for SLE: a unifying hypothesis. *Genes Immun* 2004;**5**:130–7.
18. **Kyogoku C**, Tsuchiya N, Matsuta K, *et al.* Studies on the association of Fc gamma receptor IIA, IIB, IIIA and IIIB polymorphisms with rheumatoid arthritis in the Japanese: evidence for a genetic interaction between *HLA-DRB1* and *FCGR3A*. *Genes Immun* 2002;**3**:488–93.
19. **Hyrich KL**, Lunt M, Watson KD, *et al.* Outcomes after switching from one anti-tumor necrosis factor alpha agent to a second anti-tumor necrosis factor alpha agent in patients with rheumatoid arthritis: results from a large UK national cohort study. *Arthritis Rheum* 2007;**56**:13–20.
20. **Buch MH**, Bingham SJ, Bryer D, *et al.* Long-term infliximab treatment in rheumatoid arthritis: subsequent outcome of initial responders. *Rheumatology* 2007;**46**:1153–6.
21. **Kievit W**, Adang EM, Franssen J, *et al.* The effectiveness and medication costs of three anti-tumour necrosis factor alpha agents in the treatment of rheumatoid arthritis from prospective clinical practice data. *Ann Rheum Dis* 2008;**67**:1229–34.
22. **Geborek P**, Crnkic M, Petersson IF, *et al.* Etanercept, infliximab, and leflunomide in established rheumatoid arthritis: clinical experience using a structured follow up programme in southern Sweden. *Ann Rheum Dis* 2002;**61**:793–8.
23. **Zink A**, Listing J, Kary S, *et al.* Treatment continuation in patients receiving biological agents or conventional DMARD therapy. *Ann Rheum Dis* 2005;**64**:1274–9.
24. **Haraoui B**, Cameron L, Ouellet M, *et al.* Anti-infliximab antibodies in patients with rheumatoid arthritis who require higher doses of infliximab to achieve or maintain a clinical response. *J Rheumatol* 2006;**33**:31–6.
25. **Finckh A**, Simard JF, Gabay C, *et al.* Evidence for differential acquired drug resistance to anti-tumour necrosis factor agents in rheumatoid arthritis. *Ann Rheum Dis* 2006;**65**:746–52.
26. **Vultaggio A**, Matucci A, Nencini F, *et al.* Anti-infliximab IgE and non-IgE antibodies and induction of infusion-related severe anaphylactic reactions. *Allergy* 2010;**65**:657–61.
27. **Cheifetz A**, Mayer L. Monoclonal antibodies, immunogenicity, and associated infusion reactions. *Mt Sinai J Med* 2005;**72**:250–6.
28. **Emi Aikawa N**, de Carvalho JF, Artur Almeida Silva C, *et al.* Immunogenicity of anti-TNF-alpha agents in autoimmune diseases. *Clin Rev Allergy Immunol* 2010;**38**:82–9.
29. **Tutuncu Z**, Kavanaugh A, Zvaifler N, *et al.* Fcγ₃ receptor type IIIA polymorphisms influence treatment outcomes in patients with inflammatory arthritis treated with tumor necrosis factor alpha-blocking agents. *Arthritis Rheum* 2005;**52**:2693–6.
30. **Kastbom A**, Bratt J, Ernestam S, *et al.* Fcγ₃ receptor type IIIA genotype and response to tumor necrosis factor alpha-blocking agents in patients with rheumatoid arthritis. *Arthritis Rheum* 2007;**56**:448–52.
31. **Tsukahara S**, Ikari K, Sato E, *et al.* A polymorphism in the gene encoding the Fcγ₃ IIIA receptor is a possible genetic marker to predict the primary response to infliximab in Japanese patients with rheumatoid arthritis. *Ann Rheum Dis* 2008;**67**:1791–2.
32. **Hughes LB**, Criswell LA, Beasley TM, *et al.* Genetic risk factors for infection in patients with early rheumatoid arthritis. *Genes Immun* 2004;**5**:641–7.
33. **Meknache N**, Jönsson F, Laurent J, *et al.* Human basophils express the glycosylphosphatidylinositol-anchored low-affinity IgG receptor Fcγ₃RIIIB (CD16B). *J Immunol* 2009;**182**:2542–50.
34. **Arora T**, Padaki R, Liu L, *et al.* Differences in binding and effector functions between classes of TNF antagonists. *Cytokine* 2009;**45**:124–31.
35. **Fanciulli M**, Norsworthy PJ, Petretto E, *et al.* *FCGR3B* copy number variation is associated with susceptibility to systemic, but not organ-specific, autoimmunity. *Nat Genet* 2007;**39**:721–3.
36. **Augustsson J**, Eksborg S, Ernestam S, *et al.* Low-dose glucocorticoid therapy decreases risk for treatment-limiting infusion reaction to infliximab in patients with rheumatoid arthritis. *Ann Rheum Dis* 2007;**66**:1462–6.
37. **Tanaka Y**, Takeuchi T, Inoue E, *et al.* Retrospective clinical study on the notable efficacy and related factors of infliximab therapy in a rheumatoid arthritis management group in Japan: one-year clinical outcomes (RECONFIRM-2). *Mod Rheumatol* 2008;**18**:146–52.
38. **Nagasawa H**, Kameda H, Sekiguchi N, *et al.* Normalisation of physical function by infliximab in patients with RA: factors associated with normal physical function. *Clin Exp Rheumatol* 2010;**28**:365–72.