

POSTER PRESENTATION

Open Access

# P04.83. What factors influence the use of integrative medicine (IM) modalities by infectious disease (ID) physicians?

K Shere-Wolfe<sup>1\*</sup>, J Tilburt<sup>2</sup>, C D'Adamo<sup>1</sup>, B Berman<sup>1</sup>, M Chesney<sup>3</sup>

From International Research Congress on Integrative Medicine and Health 2012  
Portland, Oregon, USA. 15-18 May 2012

## Purpose

The purpose was to assess factors that may influence the use of IM modalities by ID physicians in their practice.

## Methods

In a 2010 national survey of 1000 practicing ID physicians, participants were asked to report the extent (major, minor or not at all) to which the following considerations played a role in their recommendation/referral of IM modalities: (1) Knowledge of how and when to use them; (2) Amount of clinical research showing clear benefit; (3) Insurance; (4) Cost; (5) Reliable referral base; (6) Concern for professional reputation; (7) Fear of judgment from colleagues; (8) Insufficient regulatory oversight of supplements; and (9) Potential drug interactions with botanicals/supplements.

**Table 1**

Factor	Number of respondents	Major role (%)	Minor or Not at all (%)
Drug Interactions	293	82	18
Research	294	80	20
Knowledge	294	72	28
Insurance	292	24	76
Cost	293	39	61
Referral base	288	39	61
Professional reputation	293	14	86
Fear of judgement	293	4	96
Regulation oversight	294	69	31

## Results

A total of 311 (31%) ID physicians responded to the survey. The mean age was 49 and 64% of respondents were male. Their responses to the questions are listed below.

## Conclusion

For ID physicians, factors that were considered a major influence on the use of IM modalities included: potential drug interactions, clinical research, knowledge of IM modalities, and regulatory oversight. Factors that played a minor/no role in the use of IM modalities included fear of judgment and concern for professional reputation.

## Author details

<sup>1</sup>University of Maryland, Baltimore, USA. <sup>2</sup>Mayo Clinic, Rochester, USA.

<sup>3</sup>University of California, San Francisco, USA.

Published: 12 June 2012

doi:10.1186/1472-6882-12-S1-P353

**Cite this article as:** Shere-Wolfe et al.: P04.83. What factors influence the use of integrative medicine (IM) modalities by infectious disease (ID) physicians? *BMC Complementary and Alternative Medicine* 2012 12(Suppl 1): P353.

<sup>1</sup>University of Maryland, Baltimore, USA

Full list of author information is available at the end of the article