

## Brief Communication



# Thapsigargin Increases IL-2 Production in T Cells at Nanomolar Concentrations

Ki-Hyang Kim, Sang-Hyun Kim, Ho-Hyun Jung, Jun-Hyeok Moon, Seong-Un Jeong, Kyeongae Yu, Chong-Kil Lee<sup>\*</sup>

College of Pharmacy, Chungbuk National University, Cheongju 28644, Korea



**Received:** May 2, 2018  
**Revised:** Jul 27, 2018  
**Accepted:** Jul 30, 2018

### \*Correspondence to

**Chong-Kil Lee**

College of Pharmacy, Chungbuk National University, 1 Chungdae-ro, Seowon-gu, Cheongju 28644, Korea.  
E-mail: cklee@chungbuk.ac.kr

**Copyright** © 2018. The Korean Association of Immunologists

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Conflict of Interest

The authors declare no potential conflicts of interest.

### Abbreviations

AIDS, acquired immunodeficiency syndrome; BAPTA/AM, 1,2-bis(2-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid acetoxymethyl ester; Con A, concanavalin A; DC, dendritic cell; DG, diacylglycerol; ER, endoplasmic reticulum; IFN- $\gamma$ , Interferon-gamma; IP<sub>3</sub>, inositol 1,4,5-triphosphate; LPS, lipopolysaccharide; MLR, mixed lymphocyte reaction; PKC, protein kinase C; PMA, phorbol myristate acetate; TGN, thapsigargin

## ABSTRACT

Thapsigargin (TGN) is a potent and selective inhibitor of sarco-endoplasmic Ca<sup>2+</sup>-ATPase, leading to rapid elevation of cytoplasmic Ca<sup>2+</sup> concentration. Previous reports have shown that TGN increases the production of various cytokines from macrophages and dendritic cells. Here, we examine the effects of TGN on murine T cells. Nanomolar concentrations of TGN are a significant inducer of IL-2 production with full activity at 50 nM. Micromolar concentrations of TGN, however, are inhibitory to IL-2 production and T cell proliferation. The IL-2 production-inducing activity of TGN is much more prominent when T cells are primed with concanavalin A or anti-CD3 mAb, and is due to the increase of cytoplasmic Ca<sup>2+</sup> concentration. TGN at 50 nM does not affect interferon-gamma or IL-4 production from T cells. Thus, the present study shows that low nanomolar concentrations of TGN could be useful in potentiating IL-2 production from antigen-primed T cells.

**Keywords:** Thapsigargin; Cytoplasmic Ca<sup>2+</sup>; T cells; IL-2; Phorbol myristate acetate

## INTRODUCTION

Intracellular Ca<sup>2+</sup> plays a vital second messenger in the activation of lymphocytes (1). For instance, stimulation of T cell receptors induces hydrolysis of phosphatidylinositol 4,5-bisphosphate to produce the second messengers, inositol 1,4,5-triphosphate (IP<sub>3</sub>) and diacylglycerol (DG). IP<sub>3</sub> induces an increase in intracellular Ca<sup>2+</sup> concentration. IP<sub>3</sub> binds to its receptor on the surface of the endoplasmic reticulum (ER), and triggers the opening of the Ca<sup>2+</sup> channel, resulting in the release of Ca<sup>2+</sup> into the cytoplasm (2). DG induces activation of protein kinase C (PKC). The combined action of IP<sub>3</sub> and DG leads to the activation and proliferation of T cells. Ca<sup>2+</sup> signaling is essential not only for T cell activation, but also for immune tolerance to self-antigens (2). Pharmacological agents such as A23187, which increase intracellular Ca<sup>2+</sup> concentrations, are mitogenic for lymphocytes (3) and synergize with direct PKC activators, such as phorbol esters, in inducing activation and proliferation of lymphocytes (4). The combined use of ionomycin and phorbol myristate acetate (PMA) is an effective method to produce a variety of cytokines (5).

**Author Contributions**

Conceptualization: Lee CK; Data curation: Kim KH; Formal analysis: Kim KH; Investigation: Kim KH, Kim SH, Jung HH, Yu K; Validation: Moon JH, Jeong SU; Writing - review & editing: Lee CK.

Thapsigargin (TGN), a sesquiterpene lactone isolated from a plant *Thapsia garganica*, increases intracellular  $\text{Ca}^{2+}$  concentration without the involvement of  $\text{IP}_3$  (6,7). TGN elevates the cytosolic  $\text{Ca}^{2+}$  concentration from intracellular stores by inhibiting ER  $\text{Ca}^{2+}$ -ATPase. Much like ionomycin, TGN induces the activation of several intracellular signaling pathways, resulting in the production of a variety of cytokines. TGN elevates the production of IL-6 from mouse peritoneal macrophages (8), interferon-gamma (IFN- $\gamma$ ), IL-1 $\beta$ , and IL-6 from lipopolysaccharide (LPS)-stimulated rat peritoneal macrophages (9), tumor necrosis factor- $\alpha$  and IL-6 from LPS-activated macrophages (10), IFN- $\gamma$  from murine macrophages and human peripheral blood mononuclear cells (11), and IL-6 from bone marrow-derived dendritic cells (DCs) (12). TGN has also been reported to inhibit the production of nitric oxide, IL-10, and RANTES in macrophages stimulated with LPS, or LPS plus IFN- $\gamma$  (13).

The effects of TGN on the activation of normal T cells are not clearly defined yet. Here, we examine the effects of TGN on T cells. Our results show that TGN at low nanomolar concentrations is a significant inducer of IL-2 production, but is inhibitory to IL-2 production and T cell proliferation in micromolar concentrations. Our results also show that the IL-2 production-inducing activity of TGN is much more prominent when T cells are primed with concanavalin A (Con A) or anti-CD3 mAb.

## MATERIALS AND METHODS

### Reagents

1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid acetoxyethyl ester (BAPTA/AM), phorbol 12-myristate 13-acetate, and TGN were purchased from Sigma-Aldrich (St. Louis, MO, USA). Anti-mouse CD3 $\epsilon$  mAb was purchased from BD Biosciences (San Diego, CA, USA).

### Generation of DCs from bone marrow cells

DCs were generated as described previously (14). Briefly, the bone marrow cells obtained from femurs of C57BL/6 mice were cultured in 6-well plates ( $5 \times 10^6$  cells/well) in a culture medium with 40 ng/ml granulocyte macrophage-colony stimulating factor and 40 ng/ml IL-4 (both from Creagene, Seongnam, Korea). After 3 days, the non-adherent cells were removed by gently shaking the plate and then replacing the medium. On day 4, the non-adherent cells were again removed by the same method. On day 6, half the culture medium was replaced with fresh medium. The DCs were harvested by gentle pipetting on day 7.

### Preparation of T cells

Total T cells were purified from the spleens of BALB/c mice by adding the spleen cells to a nylon wool column and incubating for 1 h to remove adherent cells. CD4 $^+$  T cells were isolated from the adherent cell-depleted spleen cell population using the CD4 $^+$  isolation kit (Milteny Biotec, Bergisch Gladbach, Germany).

### Mixed lymphocyte reaction (MLR)

To assay the effects of TGN on lymphocyte proliferation in the MLR, CD4 $^+$  T cells isolated from BALB/c mice were seeded in 96-well round bottomed plates ( $2 \times 10^5$  cells/well), and were stimulated with DCs ( $1 \times 10^5$  cells/well) generated from bone marrow cells of C57BL/6 mice for 72 h in the presence of various concentrations of TGN. DNA synthesis was measured by incorporating [ $^3\text{H}$ ]-thymidine, which was added before the final 18 h of culture.

### XTT assay

The DOBW cells were cultured for 48 h in the presence of TGN. To measure cell viability, 50  $\mu$ l of a solution containing 1 mg/ml XTT [2,3-bis(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5-carboxanilide] (Sigma-Aldrich) and 0.383 mg/ml phenazine methosulfate (PMS, Fluka, Buchs, Switzerland) diluted in PBS were added to each well and incubated for 4 h at 37°C. Absorbance was measured at 460 nm.

### Cytokine measurements

The amounts of cytokines in the cell culture supernatants were measured by using ELISA kits (BD Biosciences) according to the manufacturer's instructions.

### Statistical analysis


The statistical significance of the difference between values of the control and treatment groups in the various assays was assessed using Student's *t*-test.

## RESULTS AND DISCUSSION

The T cell proliferation-inducing activity of TGN was examined in an allogeneic MLR. In this experiment, the highest concentration of TGN was 3  $\mu$ M, and the lower TGN concentrations were the ones diluted in 3-fold. As shown in **Fig. 1A**, TGN produced a 'bell-shaped' dose-responsive curve, with full response at approximately 100 nM (**Fig. 1A**). High concentrations of TGN (3  $\mu$ M), however, significantly reduced alloantigen-primed T cell proliferation. TGN also increased T cell proliferation in mouse splenocyte cultures that were primed with Con A (**Fig. 1B**). The maximum level of T cell proliferation occurred at approximately 100 nM. Con A-primed T cell proliferation was also significantly reduced by TGN when its concentration was 3  $\mu$ M. The cytotoxic activity of TGN was examined using DOBW cells, which are cells from an ovalbumin-specific CD4 T cell hybridoma (15), using an XTT assay. TGN was cytotoxic to DOBW cells at 1  $\mu$ M or above (**Fig. 1C**).

The T cell proliferation-inducing activity of TGN is due to the increased production of IL-2 from TGN-stimulated T cells (**Fig. 2**). The effects of TGN on IL-2 production was first examined in mouse splenocyte cultures primed with or without 1  $\mu$ g/ml Con A (**Fig. 2A**). TGN alone increased IL-2 production significantly, with maximum response at 50 nM. The amounts of IL-2 produced from 25 nM TGN-stimulated lymphocytes were almost the same as that of 1  $\mu$ g/ml Con A-stimulated lymphocytes. It is noteworthy that the IL-2 production-inducing activity of TGN reaches its maximum level at 50 nM, and diminishes at higher concentration of TGN (200 nM). This phenomenon may explain why TGN exerts a bell-shaped dose-response curve in **Fig. 1A and B**. This phenomenon may also explain the reason for the earlier observation which showed that TGN alone did not induce IL-2 production from lymphocytes (16). In that study, TGN was used at 2  $\mu$ M, which is cytotoxic and inhibitory to IL-2 production according to our present study. TGN profoundly increased IL-2 production when used simultaneously with Con A (**Fig. 2A**). In fact, the IL-2 production-inducing activity of low nanomolar concentrations of TGN was previously noted in lymphocytes isolated from acquired immunodeficiency syndrome (AIDS) patients (17).

The IL-2 production-inducing activity of TGN was also examined using T cells purified from mouse spleens. TGN alone increased IL-2 production from purified T cells significantly, with maximum response at 50 nM (**Fig. 2B**). Furthermore, TGN increased IL-2 production profoundly when used along with anti-CD3 mAb (**Fig. 2B**). We then examined the




**Figure 1.** Effects of TGN on T cell proliferation. (A) Indicated amounts of TGN was added to allogeneic MLRs, which is composed of DCs ( $1 \times 10^5$  cells/well) generated from bone marrow cells of C57BL/6 mice and CD4<sup>+</sup> T cells ( $2 \times 10^5$  cells/well) isolated from spleens of BALB/c mice. DNA synthesis was measured by [<sup>3</sup>H]-thymidine incorporation for the final 18 h of 72 h culture period. The highest concentration of TGN was 3  $\mu$ M, and the lower TGN concentrations were the ones diluted in 3-fold. The dotted lines indicate the amounts of [<sup>3</sup>H]-thymidine incorporation in the absence of TGN. (B) Indicated amounts of TGN were added to cultures of splenocytes isolated from C57BL/6 mouse spleens along with Con A (1  $\mu$ g/ml). After 48 h, DNA synthesis was measured by [<sup>3</sup>H]-thymidine incorporation for the final 18 h of the 48 h culture period. (C) The same concentrations of TGN as Fig. 1A were added to cultures of CD4<sup>+</sup> DOBW cells. After 48 h, cell viability was measured using an XTT assay. Each data point represents the mean  $\pm$  standard deviation of values obtained from 3 individual experiments. OD, optical density. \* $p < 0.05$ , \*\* $p < 0.01$  compared with untreated control.

combination effects of various concentrations of TGN with 20 ng/ml PMA, which mimics the action of DG in PKC activation. TGN together with PMA induced IL-2 production potently in T cells purified from mouse spleens (Fig. 2C).

We added various concentrations of BAPTA/AM, an intracellular Ca<sup>2+</sup> chelator, to cultures of T cells stimulated with PMA (20 ng/ml) and TGN (25 nM), to confirm that the increase of intracellular Ca<sup>2+</sup> is the main mechanism of TGN's activity on IL-2 production. BAPTA/AM blocked IL-2 production dose-dependently, with complete nullification at approximately 20  $\mu$ M (Fig. 2D), confirming that TGN's activity on IL-2 production is mostly due to the increase of intracellular Ca<sup>2+</sup>.


Subsequently, we examined whether TGN could also increase any cytokine production other than IL-2 in antigen-primed T cells. In this experiment, the total T cells purified from mouse spleens were cultured in plates pre-coated with anti-CD3 mAb (1  $\mu$ g/ml) in the presence or absence of TGN. TGN (25 nM) alone or in combination with anti-CD3 mAb did not induce IFN- $\gamma$  or IL-4 production significantly (Fig. 3).

Thapsigargin Increases IL-2 Production



**Figure 2.** Effects of TGN on IL-2 production from T cells. (A) Indicated amounts of TGN and 1 μg/ml Con A were added simultaneously to cultures of splenocytes (5×10<sup>5</sup> cells/well) isolated from C57BL/6 mouse spleens. After 48 h, the culture supernatants were harvested and measured for IL-2 content. (B) CD4<sup>+</sup> T cells (2×10<sup>5</sup> cells/well) isolated from spleens of C57BL/6 mice were cultured in the presence or absence of TGN in 96-well culture plates that were pre-coated with anti-CD3 mAb (1 μg/ml) overnight. After 48 h, the culture supernatants were harvested and measured for IL-2 content. (C) Indicated amounts of TGN and 20 ng/ml PMA were added simultaneously to cultures of CD4<sup>+</sup> T cells (2×10<sup>5</sup> cells/well) isolated from spleens of C57BL/6 mice. After 48 h, the culture supernatants were harvested and measured for IL-2 content. (D) Indicated amounts of BAPTA/AM and 50 nM TGN were added to CD4<sup>+</sup> T cell cultures which were performed in 96-well plates that were pre-coated with anti-CD3 mAb (0.1 μg/well) overnight. After 48 h, the culture supernatants were harvested and measured for IL-2 content. Each data point represents the mean±standard deviation of values obtained from 3 individual experiments. \*p<0.05, \*\*p<0.01 compared with untreated control; \*p<0.05, \*\*\*p<0.01 compared with matched group.

TGN does not activate PKC; instead it discharges intracellular Ca<sup>2+</sup> stores by specific inhibition of the ER Ca<sup>2+</sup>-ATPase resulting in rapid increase of intracellular Ca<sup>2+</sup> concentration (7). Due to this activity, TGN induces acute responses in a large number of cell types (18,19). Thus, TGN did not receive much attention as a pharmacological agent that modulates immune responses. TGN, however, came to attention in recent years since the development of TGN analogue-based prodrug for the treatment of solid tumors (20). TGN has been known to increase cytokine production in macrophages and DCs (9-12,21). The effects of TGN on cytokine production by T cells are not clearly defined yet. TGN has been reported to have no effect on IL-2 production in Jurkat T cells (16), while there are reports showing that TGN increases IL-2 production from lymphocytes isolated from AIDS patients (17). This discrepancy might be due to the difference in the concentration of TGN used, as well



**Figure 3.** Effects of TGN on IFN- $\gamma$  and IL-4 production from T cells. CD4<sup>+</sup> T cells isolated from C57BL/6 mouse spleens were cultured in the presence or absence of TGN in 96-well culture plates that were pre-coated with anti-CD3 mAb (0.1  $\mu$ g/well) overnight. After 48 h, the culture supernatants were harvested and measured for IL-2 content. Each data point represents the mean  $\pm$  standard deviation of values obtained from 3 individual experiments.

as difference in the cell type used. The present study shows that TGN exerts a ‘bell-shaped’ dose-response curve in inducing T cell proliferation in allogeneic MLR and Con A-stimulated lymphocyte cultures. It also shows that TGN alone is a significant inducer of IL-2 production only in low nanomolar concentrations with full activity at approximately 50 nM. In addition, this study shows that the IL-2 production-inducing activity of TGN is much more potent when T cells are primed with Con A or anti-CD3. Thus, the present study sheds a new light on the potentiation of antigen-primed T cells using low nanomolar concentrations of TGN.

## ACKNOWLEDGEMENTS

This study was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (No. MRC, 2017R1A5A2015541), and the research grant of the Chungbuk National University in 2015.

## REFERENCES

- Oh-hora M, Rao A. Calcium signaling in lymphocytes. *Curr Opin Immunol* 2008;20:250-258. [PUBMED](#) | [CROSSREF](#)
- Fracchia KM, Pai CY, Walsh CM. Modulation of T cell metabolism and function through calcium signaling. *Front Immunol* 2013;4:324. [PUBMED](#) | [CROSSREF](#)
- Åkerman KE, Andersson LC. Direct mitogenic effect of ionophore A23187 on isolated human T helper lymphocytes. *Eur J Immunol* 1984;14:286-288. [PUBMED](#) | [CROSSREF](#)
- Truneh A, Albert F, Golstein P, Schmitt-Verhulst AM. Early steps of lymphocyte activation bypassed by synergy between calcium ionophores and phorbol ester. *Nature* 1985;313:318-320. [PUBMED](#) | [CROSSREF](#)
- Ai W, Li H, Song N, Li L, Chen H. Optimal method to stimulate cytokine production and its use in immunotoxicity assessment. *Int J Environ Res Public Health* 2013;10:3834-3842. [PUBMED](#) | [CROSSREF](#)

6. Gouy H, Cefai D, Christensen SB, Debré P, Bismuth G.  $Ca^{2+}$  influx in human T lymphocytes is induced independently of inositol phosphate production by mobilization of intracellular  $Ca^{2+}$  stores. A study with the  $Ca^{2+}$  endoplasmic reticulum-ATPase inhibitor thapsigargin. *Eur J Immunol* 1990;20:2269-2275.  
[PUBMED](#) | [CROSSREF](#)
7. Thastrup O, Cullen PJ, Drøbak BK, Hanley MR, Dawson AP. Thapsigargin, a tumor promoter, discharges intracellular  $Ca^{2+}$  stores by specific inhibition of the endoplasmic reticulum  $Ca^{2+}$ -ATPase. *Proc Natl Acad Sci U S A* 1990;87:2466-2470.  
[PUBMED](#) | [CROSSREF](#)
8. Bost KL, Mason MJ. Thapsigargin and cyclopiazonic acid initiate rapid and dramatic increases of IL-6 mRNA expression and IL-6 secretion in murine peritoneal macrophages. *J Immunol* 1995;155:285-296.  
[PUBMED](#)
9. Harmatha J, Buděšínský M, Vokáč K, Kostecká P, Kmoníčková E, Zídek Z. Trilobolide and related sesquiterpene lactones from *Laser trilobum* possessing immunobiological properties. *Fitoterapia* 2013;89:157-166.  
[PUBMED](#) | [CROSSREF](#)
10. Chen BC, Hsieh SL, Lin WW. Involvement of protein kinases in the potentiation of lipopolysaccharide-induced inflammatory mediator formation by thapsigargin in peritoneal macrophages. *J Leukoc Biol* 2001;69:280-288.  
[PUBMED](#)
11. Kmoníčková E, Melkusová P, Harmatha J, Vokáč K, Farghali H, Zídek Z. Inhibitor of sarco-endoplasmic reticulum  $Ca^{2+}$ -ATPase thapsigargin stimulates production of nitric oxide and secretion of interferon-gamma. *Eur J Pharmacol* 2008;588:85-92.  
[PUBMED](#) | [CROSSREF](#)
12. Peters LR, Raghavan M. Endoplasmic reticulum calcium depletion impacts chaperone secretion, innate immunity, and phagocytic uptake of cells. *J Immunol* 2011;187:919-931.  
[PUBMED](#) | [CROSSREF](#)
13. Kmoníčková E, Canová NK, Farghali H, Holý A, Zídek Z. Modulator of intracellular  $Ca^{2+}$ , thapsigargin, interferes with in vitro secretion of cytokines and nitric oxide. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2005;149:321-324.  
[PUBMED](#) | [CROSSREF](#)
14. Lee YH, Im SA, Kim JW, Lee CK. Vanilloid receptor 1 agonists, capsaicin and resiniferatoxin, enhance MHC class I-restricted viral antigen presentation in virus-infected dendritic cells. *Immune Netw* 2016;16:233-241.  
[PUBMED](#) | [CROSSREF](#)
15. Harding CV, Song R. Phagocytic processing of exogenous particulate antigens by macrophages for presentation by class I MHC molecules. *J Immunol* 1994;153:4925-4933.  
[PUBMED](#)
16. Sei Y, Reich H. Thapsigargin induces IL-2 receptor alpha-chain in human peripheral and Jurkat T cells via a protein kinase C-independent mechanism. *Immunol Lett* 1995;45:75-80.  
[PUBMED](#) | [CROSSREF](#)
17. Scharff O, Foder B, Thastrup O, Hofmann B, Møller J, Ryder LP, Jacobsen KD, Langhoff E, Dickmeiss E, Christensen SB, et al. Effect of thapsigargin on cytoplasmic  $Ca^{2+}$  and proliferation of human lymphocytes in relation to AIDS. *Biochim Biophys Acta* 1988;972:257-264.  
[PUBMED](#)
18. Quynh Doan NT, Christensen SB. Thapsigargin, origin, chemistry, structure-activity relationships and prodrug development. *Curr Pharm Des* 2015;21:5501-5517.  
[PUBMED](#) | [CROSSREF](#)
19. Cubillos-Ruiz JR, Bettigole SE, Glimcher LH. Tumorigenic and immunosuppressive effects of endoplasmic reticulum stress in cancer. *Cell* 2017;168:692-706.  
[PUBMED](#) | [CROSSREF](#)
20. Mahalingam D, Wilding G, Denmeade S, Sarantopoulos J, Cosgrove D, Cetnar J, Azad N, Bruce J, Kurman M, Allgood VE, et al. Mipsagargin, a novel thapsigargin-based PSMA-activated prodrug: results of a first-in-man phase I clinical trial in patients with refractory, advanced or metastatic solid tumours. *Br J Cancer* 2016;114:986-994.  
[PUBMED](#) | [CROSSREF](#)
21. Chen B, Stiff P, Sloan G, Kash J, Manjunath R, Pathasarathy M, Oldenburg D, Foreman KE, Nickoloff BJ. Replicative response, immunophenotype, and functional activity of monocyte-derived versus CD34(+)-derived dendritic cells following exposure to various expansion and maturational stimuli. *Clin Immunol* 2001;98:280-292.  
[PUBMED](#) | [CROSSREF](#)