

Common antigens between hydatid cyst and cancers

Shima Daneshpour, Mehran Bahadoran¹, Seyed Hossein Hejazi¹, Abas Ali Eskandarian¹, Mehdi Mahmoudzadeh², Hossein Yousofi Darani¹

Department of Parasitology and Mycology, ¹Department of Parasitology, Faculty of Medicine, ²Department of Internal Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: Different research groups reported a negative correlation between cancers and parasitological infections. As an example, the prevalence of a hydatid cyst among patients with cancer was significantly lower than its prevalence among normal population. Tn antigens exist both in cancer and hydatid cyst. This common antigen may be involved in the effect of parasite on cancer growth. So in this work, common antigens between hydatid cyst and cancers have been investigated.

Materials and Methods: Different hydatid cyst antigens including hydatid fluid, laminated and germinal layer antigens, and excretory secretory antigens of protoscolices were run in SDS PAGE and transferred to NCP paper. In western immunoblotting, those antigens were probed with sera of patients with different cancer and also sera of non-cancer patients. Also, cross reaction among excretory secretory products of cancer cells and antisera raised against different hydatid cyst antigen was investigated.

Results: In western immunoblotting, antisera raised against laminated and germinal layers of hydatid cyst reacted with excretory secretory products of cancer cells. Also, a reaction was detected between hydatid cyst antigens and sera of patients with some cancers.

Conclusion: Results of this work emphasize existence of common antigens between hydatid cyst and cancers. More investigation about these common antigens is recommended.

Key Words: Cancer, common antigen, hydatid cyst

Address for correspondence:

Prof. Hossein Yousofi Darani, Infectious Diseases and Tropical Medicine Research Center, Isfahan University of Medical Sciences, Isfahan, Iran.

E-mail: yousofidarani@gmail.com

Received: 24.04.2013, Accepted: 28.08.2013

INTRODUCTION

An adverse relationship between some parasitic infections and cancer in human population has been

reported by different research groups.^[1-3] Anticancer activity of some parasites such as *Trypanosoma cruzi*,^[4-10] *Toxoplasma gondii*,^[11-18] *Toxocara canis*,^[19] *Acanthamoeba castellanii*,^[20] and *Plasmodium Yoelii*^[21] have been shown in experimental animals. *In vitro* investigations also revealed that some parasites such as *Trypanosoma cruzi*,^[9,22] hydatid cyst protoscolices,^[23] and *Toxoplasma gondii*^[11,24] show anticancer activities. Moreover, it has been shown that cancer-associated mucin-type O-glycan compositions are made by parasites.^[25,26]

Unusual O-glycans such as the α -N-cetyl galactosamine-O-serine/threonine (Tn), sialyl-Tn or Thomsen-

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.175242

Copyright: © 2016 Daneshpour. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: Daneshpour S, Bahadoran M, Hejazi SH, Eskandarian AA, Mahmoudzadeh M, Darani HY. Common antigens between hydatid cyst and cancers. Adv Biomed Res 2016;5:9.

Friedenreich (TF) antigens presented in cancer cells have important roles in metastasis, cell adhesion, and invasion.^[27] Parasite *O*-glycans and mucin-like molecules seem to have important functions in the interaction of helminthes and their hosts.^[28] It has been shown that mucin-type *O*-glycan structures of cancers are expressed by helminthes. As an example, the adult worm of *Schistosoma mansoni* and its schistosomula express Tn antigen.^[26] In other studies, expression of Tn antigen in adult and larval stages of *Echinococcus granulosus* was shown.^[25] Another carbohydrate antigen, called Tk, is expressed at the surface of human colorectal carcinomas. This antigen was found in *Taenia crassiceps*, *Mesocestoides voga*, and *Taenia hydatigena*.^[29] More carbohydrate antigens such as sialyl-Tn antigens and TF antigens have also been recognized as common antigens of cancers and parasites.^[25,30-32] We considered these unusual *O*-glycan antigens as common antigens of cancers and parasites. These antigens received great attention because antibody response to them yields protection against certain cancers.^[33,34] Moreover, these antigens may be involved in anticancer activities of some parasites.^[35] So in this work, immunological cross reaction between cancers and hydatid cyst, larval stage of *Echinococcus granulosus*, has been investigated.

MATERIALS AND METHODS

In this experimental study, *Echinococcus granulosus* hydatid cysts were collected from sheep or cattle from a slaughter house in Isfahan, Iran. Hydatid fluid was aspirated, collected, and examined for the presence of protoscolices. The fluids were then centrifuged at 2000×g, for 2 min, and the supernatant was concentrated using the lyophilizer apparatus and kept at -20 as hydatid fluid antigen. The sediment which was compact protoscolices washed with isotonic saline, sonicated, and kept at -20 as crude protoscolice antigen. Also, culture medium was added to compact alive protoscolices, and following 48 hours incubation was centrifuged at 2000×g, for 2 min and the supernatant was used as excretory secretory molecules of protoscolices (ES antigen). Laminated and germinal layers were removed from the cyst, homogenized, and then sonicated and kept at -20 as laminated and germinal layers antigen.

Polyspecific rabbit antisera were raised against four hydatid cyst antigens by injecting 4 rabbits fortnightly with 1 ml of emulsions containing equal volume of each antigen and ferund's adjuvant (complete for the first and incomplete for the boosters). Polyacrylamide gel electrophoresis with sodium dodecyl sulfate (SDS-PAGE) was performed as described by Laemmli^[36]

under non-reducing condition using the BIO RAD apparatus. Western immunoblotting was performed as described by Towbin *et al.*^[36] using the BIO RAD apparatus. Pooled sera for each cancer was prepared by mixing 0.5 ml of 10 individual patients sera. For preparing cancer cell ES antigens, Baby Hamster Kidney (BHK) cell line was purchased from Pasture institute cell bank in Tehran, Iran. 6 million cells were incubated in 5 ml culture medium for 48 hours. Then they were centrifuged at 2000×g for 5 min and the supernatant was kept as excretory secretory antigen of cancer cells.

RESULTS

Different hydatid cyst antigens including: hydatid fluid, crude antigen of protoscolices, excretory secretory antigen, laminated and germinal layers antigens were run in SDS-PAGE and were transferred to NCP papers. Each antigen probed with pooled sera (10 sera for each cancer) of patients with (i) breast cancer, (ii) colon cancer, (iii) blood cancer, (iv) normal sera. Breast cancer sera reacted with a band in hydatid fluid antigen about 40-50 KDa. This reaction was not clear for other cancers and was not observed with normal sera.

Also, hydatid cyst fluid antigen was probed with different cancer sera in western immunoblotting. Results of this experiment revealed that breast cancer sera reacted with hydatid cyst antigen [Figure 1].

In another experiment, excretory secretory products of cancer cells were probed with different rabbit antisera in western immunoblotting. Antiserum raised against laminated and germinal layer of hydatid cyst reacted strongly with excretory secretory products of cancer cells [Figure 2].




Figure 1: Western immunoblotting of hydatid cyst fluid antigen probed with (from left) normal sera, sera of patients with breast cancer and a patient serum with hydatid cyst




Figure 2: Western immunoblotting of cancer cells ES antigens probed with the sera of rabbits raised against hydatid cyst Laminated and germinal layers antigen (both lanes)

DISCUSSION

In this investigation, it has been shown that rabbit antisera raised against hydatid cyst antigens specially laminated and germinal layers antigen reacted with excretory secretory products of cancer cells. Also, patient's sera with cancers especially breast cancer reacted with antigens of hydatid cyst.

Mucin-type *O*-glycans have important roles in cancer metastasis.^[27] These molecules have also important functions in the interaction of helminthes and their hosts.^[28] Nyame *et al.* showed that mucin-type *O*-glycan structures of cancers are expressed by the adult worm of *Schistosoma mansoni*.^[26] These molecules have also been shown in another parasites such as *Echinococcus granulosus*, *Taenia crassiceps*, *Mesocostoides voga*, and *Taenia hydatigena*.^[29] According to these investigations, mucin-type *O*-glycans are expressed in both cancer cells and helminthic parasites. These findings are consistent with our results and emphasize existence of common antigens both in cancers and helminthes. Tn antigens which exist both in cancers and parasites are glycosylated molecules so the cross reaction between cancer patient's sera and hydatid cyst antigen may be related to carbohydrate branches of the glycoproteins.

Immunological cross reaction of hydatid cyst antigens and cancer patient's sera has been reported in previous investigations.^[37,38] In a study, only 6.3 of 270 cancer patients sera reacted with hydatid fluid antigens in ELISA. However, the cross reaction was not detectable in western immunoblotting with the 8 kDa band.^[38] The smallest 8 kDa subunit of antigen B is a peptide band while we assumed that immunological cross reaction between hydatid cyst

antigens and cancer patient's sera may be mainly related to carbohydrate branches of glycoproteins. In the present investigation, we did not get antibody reaction against 8kDa band in western blotting, instead the reaction was against 40-50 kDa bands. Moreover, the majority of sera of patients with breast cancer reacted with these bands in western blotting.

In conclusion, in this study for the first time we showed that a mutual cross reaction exists between antigens of hydatid cyst and excretory secretory products of cancer cells. More investigation is needed to confirm that this cross reaction is related to carbohydrate molecules.

ACKNOWLEDGMENT

This work was supported by a grant from Isfahan University of Medical Sciences

REFERENCES

1. Akgül H, Tez M, Ünal AE, Keskek M, Sayek I, Özçelik T. Echinococcus against cancer: Why not? Cancer 2003;98:1999-2000.
2. Plumelle Y, Gonin C, Edouard A, Bucher BJ, Thomas L, Brebion A, *et al.* Effect of Strongyloides stercoralis infection and eosinophilia on age at onset and prognosis of adult T-cell leukemia. Am J Clin Pathol 1997;107:81-7.
3. Yousofi Darani H, Yousefi M. Parasites and cancers: Parasite antigens as possible targets for cancer immunotherapy. Future Oncol 2012;8:7.
4. Batmonkh Z, Kallinikova VD, Pakhorukova LV, Kravtsov EG, Karpenko LP, Dalin MV. In vivo anticancer activity of lysates from trypanosoma cruzi of different genetic groups. Bull Exp Biol Med 2006;142:470-3.
5. Kallinikova VD, Batmonkh T, Kosobokova EN, Pakhorukova LV, Ogloblina TA, Kravtsov EG, *et al.* Antibodies against Trypanosoma cruzi in intact mice and their oncoprotective effect. Med Parazitol (Mosk) 2008;1:11-5.
6. Kallinikova VD, Borisova EN, Pakhorukova LV, Ogloblina TA, Batmonkh Ts, Kravtsov EG, *et al.* Immunization against Trypanosoma cruzi and tumor growth in mice. Med Parazitol (Mosk) 2006;4:9-12.
7. Kallinikova VD, Matekin PV, Ogloblina TA, Leïkina MI, Kononenko AF, Sokolova NM, *et al.* Anticancer properties of flagellate protozoan Trypanosoma cruzi Chagas, 1909. Izv Akad Nauk Ser Biol 2001;3:299-311.
8. Kinnamon KE, Poon BT, Hanson WL, Waits VB. Activity of anticancer compounds against Trypanosoma cruzi-infected mice. Am J Trop Med Hyg 1998;58:804-6.
9. Sheklakova LA, Kallinikova VD, Karpenko LP. Genetic heterogeneity of Trypanosoma cruzi and its direct anticancer effect in cultured human tumor cells. Bull Exp Biol Med 2003;135:89-92.
10. Zenina AV, Kravtsov EG, Tsetsegsaikhan B, Yashina NV, Dalin MV, Karpenko LP, *et al.* The study of immunological component in antitumor effect of Trypanosoma cruzi. Bull Exp Biol Med 2008;145:352-4.
11. Choo JD, Lee JS, Kang JS, Lee HS, Yeom JY, Lee YH. Inhibitory effects of toxoplasma antigen on proliferation and invasion of human glioma cells. J Korean Neurosurg Soc 2005;37:129-36.
12. Hibbs JB Jr, Lambert LH, Remington JS. Resistance to murine tumors conferred by chronic infection with intracellular protozoa, Toxoplasma gondii and Besnoitia jellisoni. J Infect Dis 1971;124:587-92.
13. Kim JO, Jung SS, Kim SY, Kim TY, Shin DW, Lee JH, *et al.* Inhibition of Lewis lung carcinoma growth by Toxoplasma gondii through induction of Th1 immune responses and inhibition of angiogenesis. J Korean Med Sci 2007;22:S38-46.

14. Miyahara K, Honda U, Tose S, Sakurai H, Igarashi I, Saito A, *et al.* Therapeutic effects of Toxoplasma lysate antigen on 20-methylcholanthrene-induced BALB/c mouse tumors. *J Vet Med Sci* 1992;54:7-12.
15. Miyahara K, Yokoo N, Sakurai H, Igarashi I, Sakata Y, Yoshida Y, *et al.* Antitumor activity of Toxoplasma lysate antigen against methylcholanthrene-induced tumor-bearing rats. *J Vet Med Sci* 1992;54:221-8.
16. Pyo KH, Jung BK, Chai JY, Shin EH. Suppressed CD31 Expression in Sarcoma-180 Tumors after Injection with Toxoplasma gondii Lysate Antigen in BALB/c Mice. *Korean J Parasitol* 2010;48:171-4.
17. Suzuki Y, Kobayashi A. Antitumor effect of intralesional injection with formalin-fixed Toxoplasma gondii organisms on Lewis lung carcinoma in Toxoplasma-infected mice. *Cancer Lett* 1985;25:247-54.
18. Suzuki Y, Muto M, Kobayashi A. Antitumor effect of formalin-fixed Toxoplasma gondii organisms on EL4 lymphoma in Toxoplasma-infected mice. *J Biol Response Mod* 1986;5:288-93.
19. Darani HY, Shirzad H, Mansoori F, Zabardast N, Mahmoodzadeh M. Effects of Toxoplasma gondii and Toxocara canis antigens on WEHI-164 fibrosarcoma growth in a mouse model. *Korean J Parasitol* 2009;47:175-7.
20. Pidherney MS, Alizadeh H, Stewart GL, McCulley JP, Niederkorn JY. In vitro and in vivo tumoricidal properties of a pathogenic/free-living amoeba. *Cancer Lett* 1993;72:91-8.
21. Chen L, He Z, Qin L, Li Q, Shi X, Zhao S, *et al.* Antitumor effect of malaria parasite infection in a murine lewis lung cancer model through induction of innate and adaptive immunity. *PLoS One* 2011;6:e24407.
22. Atayde VD, Jasiulionis MG, Cortez M, Yoshida N. A recombinant protein based on Trypanosoma cruzi surface molecule gp82 induces apoptotic cell death in melanoma cells. *Melanoma Res* 2008;18:172-83.
23. Yousofi Darani H, Soozangar N, Khorami S, Taji F, Yousofi M, Shirzad H. Hydatid cyst protoscolices induce cell death in WEHI-164 fibrosarcomacells and inhibit the proliferation of baby hamster kidney fibroblasts in vitro. *J Parasitol Res* 2012;2012:304183.
24. Shirzad H, Khorami S, Soozangar N, Yousefi M, Darani HY. Toxoplasma gondii but not Leishmania major or Trichomonas vaginalis decreases cell proliferation and increases cell death on fibrosarcoma cancer cells in culture medium. *World J Vaccine* 2012;2:105-8.
25. Alvarez Errico D, Medeiros A, Míguez M, Casaravilla C, Malgor R, Carmona C, *et al.* O-glycosylation in Echinococcus granulosus: Identification and characterization of the carcinoma-associated Tn antigen. *Exp Parasitol* 2001;98:100-9.
26. Nyame K, Cummings R and Damian R. Schistosoma mansoni synthesizes glycoproteins containing terminal O-linked N-acetylglucosamine residues. *J Biol Chem* 1987;262:7990-5.
27. Baldus SE, Engelmann K, Hanisch FG. MUC1 and the MUCs: A family of human mucins with impact in cancer biology. *Crit Rev Clin Lab Sci* 2004;41:189-231.
28. Thomas PG, Harn Jr DA. Immune biasing by helminth glycans. *Cell Microbiol* 2004;6:13-22.
29. Ubillos L, Medeiros A, Cancela M, Casaravilla C, Saldaña J, Domínguez L, *et al.* Characterization of the carcinoma-associated Tn antigen in helminth parasites. *Exp Parasitol* 2007;116:129-36.
30. Casaravilla C, Freire T, Malgor R, Medeiros A, Osinaga E, Carmona C. Mucin-type O-glycosylation in helminth parasites from major taxonomic groups: Evidence for widespread distribution of the Tn antigen (GalNAc-Ser/Thr) and identification of UDP-GalNAc: Polypeptide N-acetylgalactosaminyltransferase activity. *J Parasitol* 2003;89:709-14.
31. Freire T, Casaravilla C, Carmona C, Osinaga E. Mucin-type O-glycosylation in Fasciola hepatica: Characterisation of carcinoma-associated Tn and sialyl-Tn antigens and evaluation of UDP-GalNAc: Polypeptide N-acetylgalactosaminyltransferase activity. *Int J Parasitol* 2003;33:47-56.
32. Thors C, Jansson B, Helin H, Linder E. Thomsen-Friedenreich oncofetal antigen in Schistosoma mansoni: Localization and immunogenicity in experimental mouse infection. *Parasitology* 2006;132:73-81.
33. Livingston PO, Zhang S, Lloyd KO. Carbohydrate vaccines that induce antibodies against cancer. 1. Rationale. *Cancer Immunol Immunother* 1997;45:1-9.
34. Slovin SF, Keding SJ, Ragupathi G. Carbohydrate vaccines as immunotherapy for cancer. *Immunol Cell Boils* 2005;83:418-28.
35. Darani HY, Yousefi M. Parasites and cancers: Parasite antigens as possible targets for cancer immunotherapy. *Future Oncol* 2012;8:1529-35.
36. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-5.
37. Dar FK, Buhidma MA, Kidwai SA. Hydatid false positive serological test results in malignancy. *Br Med J (Clin Res Ed)* 1984;288:1197-7.
38. Pfister M, Gottstein B, Cerny T, Cerny A. Immunodiagnosis of echinococcosis in cancer patients. *Clin Microbiol Infect* 2008;5:693-7.

Source of Support: Isfahan University of Medical Sciences, Isfahan, Iran.
Conflicts of Interest: None declared.