

Carcinoma of an unknown primary: are EGF receptor, Her-2/neu, and c-Kit tyrosine kinases potential targets for therapy?

C Massard^{*1}, J-J Voigt², A Laplanche³, S Culine⁴, A Lortholary⁵, R Bugat⁶, C Theodore¹, F Priou⁷, M-C Kaminsky⁸, T Lesimple⁹, X Pivot¹⁰, B Coudert¹¹, J-Y Douillard¹², Y Merrouche¹³ and K Fizazi¹

¹Department of Medicine, Institut Gustave Roussy, 39 rue Camille Desmoulins, Villejuif 94805, France; ²Department of Pathology, Centre Claudius Regaud, Toulouse, France; ³Department of Statistics, Institut Gustave Roussy, Villejuif, France; ⁴Department of Medicine, Centre Val d'Aurelle, Montpellier, France; ⁵Department of Medicine, Centre Paul Papin, Anger, France; ⁶Department of Medicine, Institut Claudius Regaud, Toulouse, France; ⁷Department of Medicine, Centre Hospitalier La Roche/Yon, La Roche/Yon, France; ⁸Department of Medicine Centre Alexis Vautrin, Nancy, France; ⁹Department of Medicine, Centre Eugène Marquis, Rennes, France; ¹⁰Department of Medicine, Centre Hospitalier Universitaire, Besançon, France; ¹¹Department of Medicine, Centre G.F. Leclerc, Dijon, France; ¹²Department of Medicine, Centre René Gauducheau, Nantes, France; ¹³Department of Medicine, Institut de Cancérologie de la Loire, Saint-Etienne, France

Carcinomas of an unknown primary site (CUP) are heterogeneous tumours with a median survival of only 8 months. Tyrosine kinase inhibitors are promising new drugs. The aim of this study was to determine the expression of EGF-receptor, Her-2/neu, and c-Kit tyrosine kinases in CUP. Paraffin-embedded specimens were obtained from 54 patients with a CUP who were included in the GEFCAPI 01 randomised phase II trial. Immunohistochemistry was performed using the Dako autostainer with antibodies directed against HER-2/neu protein, EGFR protein, and c-Kit protein (CD117). EGFR expression was found in 36 out of 54 samples (66%). In contrast, Her-2/neu overexpression and c-Kit positivity were only detected in 4 and 10% of patients, respectively. No significant association was found between the expression of the tyrosine kinase receptors and prognosis. EGFR expression was significantly associated with response to cisplatin-based chemotherapy: the response rates were 50 and 22% in patients with EGFR-positive tumours and EGFR-negative tumours, respectively ($P < 0.05$). This study shows that EGFR is frequently expressed in CUP. This finding may prompt clinical trials investigating EGFR inhibitors in this setting. In contrast, c-Kit expression and Her-2/neu overexpression occur infrequently in CUP. EGFR expression was correlated to tumour chemosensitivity.

British Journal of Cancer (2007) 97, 857–861. doi:10.1038/sj.bjc.6603942 www.bjcancer.com

Published online 18 September 2007

© 2007 Cancer Research UK

Keywords: carcinoma of an unknown primitive; tyrosine kinase receptor; EGFR; c-Kit; Her-2/neu overexpression

Carcinoma of an unknown primary site (CUP) rank among the 10 most frequent cancers worldwide and its prognosis is notoriously poor with median survival rates attaining 8 months (Pavlidis and Fizazi, 2005). Carcinoma of an unknown primary site are heterogeneous tumours whose origin is unidentifiable at the time of the diagnosis but they share the unique clinical feature of metastatic disease: a slow local development and a high metastatic potential. Despite advances in tumour imaging and pathology, patients with CUP still account for about 5% of all cancer patients. The primary site remains frequently unknown but necroptic studies indicate that the primary sites are most often the pancreas, lung, gut, and kidney.

The treatment of patients presenting with a CUP remains a daily challenge for physicians. Although systemic chemotherapy is usually recommended, the optimal regimen remains to be determined. To improve the poor outcome of patients, several recent studies have focused on the introduction of new cytotoxic agents such as gemcitabine (GC), irinotecan (IC), and taxanes which exhibit a broad spectrum of clinical activity (Greco *et al*,

2002; Culine *et al*, 2003). However, no randomised trial has provided a clear evidence of a survival benefit for CUP patients (Fizazi, 2006).

In addition, the biology of CUP is still very poorly understood (Fizazi, 2006). Growth factor receptors with tyrosine kinase activities such as EGFR, HER-2/neu, and c-Kit have recently emerged as promising targets for novel therapeutic agents, especially in some poorly chemosensitive neoplasms (Krause and Van Etten, 2005). This prompted us to study receptor tyrosine kinases, which could be potential targets for novel therapies in patients with CUP.

We therefore studied tumour specimens from 54 patients with CUP enrolled in one of a large prospective randomised phase II trial (Culine *et al*, 2003) to determine EGFR, Her-2/neu, and c-Kit protein expression.

MATERIALS AND METHODS

Patients

Tissues were obtained from the original biopsy specimens of 54 patients who were enrolled in a randomised phase II trial (GEFCAPI 01) conducted by the French Study Group of

*Correspondence: Dr C Massard; E-mail: christophe.massard@igr.fr
Received 22 March 2007; revised 13 July 2007; accepted 20 July 2007;
published online 18 September 2007

Carcinomas of an unknown primary from August 1999 to November 2000 (Culine *et al*, 2003). The study was approved by the Ethics Committee in Montpellier. All patients signed a written informed consent before participation. Patients were randomly assigned to one of the two treatment regimens at the time of study entry and could receive chemotherapy combinations including cisplatin with GC or IC. Patients were excluded if they had any of the following features: a CUP subset requiring specific treatment (i.e. women with adenocarcinoma with axillary lymph node involvement alone, women with papillary serous carcinoma of the peritoneum, patients with squamous carcinoma exclusively involving cervical or inguinal lymph nodes, carcinomas with neuroendocrine features, and patients with carcinoma in a potentially resectable site) and those with symptomatic brain metastasis. To exclude other malignancies (lymphoma, melanoma, sarcoma, and neuroendocrine carcinoma), immunoperoxidase staining with antibodies against leukocyte common antigen, cytokeratin, neuroendocrine markers (chromogranin and synaptophysin), and melanoma markers (S-100 protein and homatropine methylbromide-45) was recommended in poorly differentiated carcinoma. All specimens were centrally reviewed by a single pathologist (JJV). Well-differentiated adenocarcinoma was the most common histologic type. The dominant visceral sites of disease were the bone, lung, and liver, whereas the mediastinum was the most common site of lymph node involvement. The results of this clinical trial have already been published (Culine *et al*, 2003). The study demonstrated the activity of both regimens in CUP patients. Response to chemotherapy was evaluated by CT scan after every two cycle of chemotherapy and independent radiologic review was carried out. Moreover, a simple prognostic model was established with Performans Status and LDH serum in CUP patients (Culine *et al*, 2002).

Immunohistochemistry

Formalin-fixed, paraffin-embedded sections were immunostained for EGFR, HER-2/neu, and c-Kit. Five-micrometer sections were cut, deparaffinised, and rehydrated. There is no SOP (Standardized Operation Procedures) for tissue collection. The slides were processed using an autostainer (Dako, Carpinteria, CA, USA) with a streptavidin-biotin complex and diaminobenzidine as the chromogens. The antibodies used were the PharmDx EGFR kit (Dako), the polyclonal rabbit anti-human-c-Kit (CD117, an epitope of KIT) (A4502, Dako) (dilution 1:50) and the polyclonal rabbit anti-human HER-2/neu - c-erbB-2 (A0485, Dako) (dilution 1:1500) in one slide for each antibody.

Microwave oven antigen retrieval from tissue sections was only performed for HER-2/neu. For EGFR staining, tissue sections were digested with protease K at 21°C for 5 min. Nonspecific binding sites were bound using Protein Block (Dako). The slides were incubated with primary antibody for 60 min (HER-2/neu and c-Kit) and 30 min for EGFR. Staining was revealed using the following chromogens: biotinylated secondary antibody, peroxidase-labelled streptavidin and 3,3'-diaminobenzidine tetrahydrochloride-hydrogen peroxide.

Dilution and incubation periods were optimised for each antibody by using positive controls. Specimens whose incubation with the primary antibody had been omitted were used as a negative control. In each experiment, tumours with defined alterations in EGFR, HER-2/neu, or c-Kit were used as positive controls. Each case was scored by a pathologist blinded to patient identity, and scoring system for EGFR, Her2/neu and Kit is based on College of American Pathologists Cell Markers Committee guidelines (Arch Pathol Lab Med 2002–2006).

EGFR positivity revealed by chromogenic staining was localised mainly in the cell membrane. Three staining categories were defined before assessing EGFR: specimens with no or less than 1% of positive cells were defined as 'negative', specimens with 1–20%

of positive cells were defined as 'positive +' and specimens with more than 20% of positive cells were defined as 'positive +++'. These definitions of positive and negative results are in accordance with the published literature (Goldstein and Armin, 2001) but may require modification in specific context.

HER-2/neu expression was confirmed when staining was localised in the membrane. Specimens were scored as 0 (no staining or less than 10% of positive cells), 1+ (weak and focal staining of more than 10% of positive cells), 2+ (weak to moderate and complete membrane staining of more than 10% of positive cells), and 3+ (strong and complete membrane staining of more than 10% of positive cells). Specimens scored as 3+ are Her-2/neu overexpression.

C-Kit expression was scored as negative when there was no staining for c-Kit or when staining was observed in less than 1% of the cells and as positive when the cytoplasm was strongly stained with or without membrane staining in 10% or more cancer cells. C-Kit expression was controlled with mast cells known to express c-Kit.

Statistical analysis

Fischer's exact test was used to evaluate differences in clinical characteristics between positive and negative EGFR, Her-2/neu, and c-Kit expression.

The following variables were studied: age, sex, pathology review, ECOG, serum LDH, disease site, chemotherapy, and response to chemotherapy. Differences in survival duration between subgroups were analysed using a two-sided log rank.

All *P*-values were two-sided. *P* < 0.05 was considered statistically significant. All analyses were carried out with the SPSS package (SPSS program stat).

RESULTS

Fourteen centres enrolled 80 patients onto the GEFCAPI 01 trial. A pathological review of 75 specimens was performed. Tissue from the original biopsy specimens was available in 54 patients for the present immunohistochemical analysis. Patient characteristics are summarised in Table 1.

Eighteen (33%) cases showed no EGFR expression. EGFR was expressed in 36 (66%) of 54 available samples, displaying moderate staining in 17 (31%) cases and strong EGFR positivity in 19 (35%) cases.

In the univariate analysis, EGFR expression correlated with response to chemotherapy (Table 2), but not with age, gender, tumour site, and pathological differentiation. Response to cisplatin-based chemotherapy was 50 and 22% in patients with EGFR-positive and -negative tumours respectively (*P* = 0.046).

Her-2 overexpression was found in only 2 (4%) of 54 tumour specimens. There was no statistically significant correlation between Her-2/neu staining and clinical characteristics, as summarised in Table 1.

Only six cases (10%) showed at least some c-Kit expression. There was no statistically significant correlation between c-Kit staining and clinical characteristics, as summarised in Table 1.

DISCUSSION

In the present study, we showed that EGFR is expressed in a majority (66%) of CUP. Moreover, EGFR expression correlated with response to cisplatin-based chemotherapy. This study is relatively small but it is one of the largest randomised trials in CUP. Treating patients with CUP remains a challenge for physicians and a better understanding of the biology of this neoplasm with the identification of reliable tumour markers that reflect tumour aggressiveness or of predictive factors for new therapeutics might lead to targeted therapies. Growth factor

Table 1 Clinical and pathological characteristics of patients with and without Her-2/neu overexpression, c-Kit expression and EGFR expression

Characteristics	All patients (n = 54)	EGFR- (n = 18)	EGFR+ (n = 36)	HER2/neu- (n = 52)	HER2/neu+ (n = 2)	c-KIT- (n = 48)	c-KIT+ (n = 6)
Age (years)							
Median	58	58	55.5	58	52	55.5	66.5
Range	33-73	33-68	41-73	33-73	45-59	33-73	48-71
Sex							
Male	31 (57%)	10 (55%)	21 (58%)	29 (56%)	2 (100%)	29 (60%)	2 (33%)
Female	23 (43%)	8 (45%)	15 (42%)	23 (44%)	0	19 (40%)	4 (67%)
Pathology							
Adenocarcinoma	29 (53%)	10 (55%)	19 (53%)	27 (52%)	2 (100%)	27 (56%)	2 (33%)
Poorly differentiated	18 (33%)	7 (38%)	11 (30%)	18 (35%)	0	16 (33%)	2 (33%)
Undifferentiated	7 (13%)	1 (5%)	6 (17%)	7 (13%)	0	5 (11%)	2 (33%)
Treatment							
1	28 (51%)	7 (38%)	21 (58%)	27 (52%)	1 (50%)	27 (56%)	1 (17%)
2	26 (49%)	11 (62%)	15 (42%)	25 (48%)	1 (50%)	21 (44%)	5 (83%)
Serum LDH							
Normal	26 (48%)	9 (50%)	17 (47%)	26 (50%)	0	23 (48%)	3 (50%)
Elevated	22 (40%)	8 (45%)	14 (39%)	21 (40%)	1	19 (40%)	3 (50%)
Not available	6 (12%)	1 (5%)	5 (14%)	5	1	6	0
ECOG PS							
0	43 (79.5%)	14 (78%)	29 (80%)	41 (79%)	2	38 (79%)	5 (83%)
1	10 (18.5%)	4 (22%)	6 (17%)	10 (19%)	0	9 (19%)	1 (17%)
Not available	1	1	1	1	0	1	0
Tumour site							
Lymph nodes	32 (60%)	10 (55%)	25 (70%)	35 (67%)	2 (100%)	36 (75%)	4 (66%)
Liver	27 (50%)	11 (60%)	16 (44%)	27 (52%)	0	25 (52%)	2 (33%)
Lung	28 (52%)	11 (60%)	17 (47%)	26 (50%)	2 (100%)	26 (54%)	2 (33%)
Bone	19 (35%)	9 (50%)	10 (28%)	18 (35%)	1 (50%)	18 (37%)	0
Peritoneum	3 (5%)	3 (16%)	9 (25%)	3 (6%)	0	3 (6%)	0
Adrenal gland	6 (11%)	1 (5%)	5 (14%)	12 (23%)	0	6 (12%)	0
Pleura	4 (7%)	1 (5%)	3 (8%)	4 (8%)	0	4 (8%)	0
Brain	2 (4%)	0	2 (5%)	2 (4%)	0	1 (2%)	1 (17%)
Cutaneous	1 (2%)	0	1 (2.5%)	1 (2%)	0	1 (2%)	0
Others	10 (18%)	4 (22%)	6 (17%)	10 (19%)	0	10 (21%)	0

EGFR-negative tumour patient: EGFR score 1+. EGFR-positive tumour patient: EGFR score 2+ or 3+. Treatment 1: cisplatin and gemcitabine. Treatment 2: cisplatin and irinotecan.

Table 2 EGFR expression and response to chemotherapy

Response	EGFR negative (n = 18)	EGFR positive (n = 36)	P-value
Complete response	0	1 (3%)	
Partial response	4 (22%)	17 (47%)	
Stable disease	5 (28%)	9 (25%)	
Progressive disease	7 (39%)	5 (14%)	
Not evaluable	2 (1%)	4 (11%)	
Objective response rate	4 (22%)	18 (50%)	<0.05

EGFR-negative tumour patient: EGFR score 1+. EGFR-positive tumour patient: EGFR score 2+ or 3+.

receptors like EGFR, HER-2, and c-Kit have recently emerged as promising targets for novel therapeutic agents in malignancies such as lung cancer (Shepherd *et al*, 2005), breast cancer (Slamon *et al*, 2001), and gastrointestinal stromal tumour (GIST) (Heinrich *et al*, 2002). This study provides evidence for tyrosine kinase receptor expression/lack of expression in CUP and may allow the study of targeted therapies in these patients. Moreover, immunohistochemistry is an indispensable tool in diagnosis and manage-

ment of CUPs. The role of pathologist is to determine the histopathological type rather than the origin of the primary tumour in view of a suitable therapy based on clinical, morphological, and phenotypic data (Fizazi, 2006).

In the last decade, EGFR has emerged as one of the most important targets for drug development in oncology. Because EGFR is expressed and associated with poor prognosis and a more malignant phenotype in many neoplasms, it has been investigated as a potential target for cancer therapy (Baselga and Arteaga, 2005; Mendelsohn and Baselga, 2006). There are currently two treatment options using anti-EGFR agents under clinical development: a monoclonal antibody directed at the extracellular domain of the receptor and small molecule inhibitors of the EGFR tyrosine kinase. Clinical trials with a humanised murine chimeric monoclonal antibody to EGFR (C225, Cetuximab) in combination with chemotherapy or radiotherapy have shown significant clinical activity in advanced colorectal carcinomas (Cunningham *et al*, 2004) squamous cell carcinomas of the head and neck (Vermorke *et al*, 2007), and non-small cell lung cancer (NSCLC) (Janne *et al*, 2004; Giaccone, 2005). On the other hand, a large number of inhibitors of the EGFR tyrosine kinase are active against NSCLC (Thatcher *et al*, 2005) and pancreatic cancer. It is well known that lung cancer is one of the most frequent origin of unknown primary cancer (Fizazi, 2006). The BR.21 study evaluated the effectiveness

of erlotinib in second-line treatment in NSCLC. The results of BR.21 study demonstrated that erlotinib has the potential to improve overall survival to unselected chemotherapy refractory NSCLC patients (Shepherd *et al*, 2005). A subset of NSCLC patients achieve impressive responses with TKI. Several authors have shown that the mutation or amplification of EGFR are associated with dramatic and sustained response to TKI in lung cancer (Taron *et al*, 2005; Tsao *et al*, 2005; Shigematsu and Gazdar, 2006). Compounds such as cetuximab, gefitinib, and erlotinib could prove valid for targeting tumours expressing EGFR in patients selected according to this marker and may represent a novel therapeutic strategy in patients with CUP (Sequist *et al*, 2007). There is a need for strategies with anti-EGFR agents alone or in combination with conventional chemotherapies and to explore combinations with other molecularly targeted therapies such as angiogenesis inhibitors. These new therapeutic approaches may represent an exciting and promising way of improving the unfavourable prognosis in CUP patients.

Response to cisplatin-based chemotherapy was better in patients with EGFR-positive as compared to those with EGFR-negative tumours whereas there is no difference in overall survival. Several studies suggest that while gefitinib or erlotinib are not effective in the general NSCLC population (Shepherd *et al*, 2005), these targeted therapies have activity in selected patients, and never-smokers and patients of Asian origin and women with adenocarcinoma might expect improvement in survival (Brown and Shepherd, 2005). Moreover, EGFR mutation or EGFR expression may be a positive prognostic factor for survival in advanced NSCLC patients treated with chemotherapy with or without erlotinib, and may predict greater likelihood of response to chemotherapy with or without erlotinib. However, our results must be viewed cautiously since the number of patients was relatively small.

A study by Hainsworth *et al* (2000) showed that 11% of tumour specimens overexpressed Her-2/neu using the Dako immunohistochemical method in a large group of patients who had received uniform treatment with cisplatin-based chemotherapy. In that study, Her-2/neu overexpression occurred mainly in patients with poorly differentiated carcinoma or patients whose tumour was predominantly located above the diaphragm, in the mediastinum or lungs. Another study demonstrated that patients with CUP have a high overexpression of Her-2/neu (Pavlidis *et al*, 1995). In our study, Her-2/neu overexpression was observed in 4% of tumour specimens but 60% of patients had adenocarcinoma and only 30% patients had poorly undifferentiated adenocarcinoma and undifferentiated carcinoma. In these studies, the difference in Her-2/neu overexpression could be explained by the patient population (poorly differentiated vs well-differentiated carcinoma). Moreover, it might be useful to standardise staining procedures used or to compare Her-2/neu overexpression with an immunohistochemical test and FISH amplification.

C-Kit is a tyrosine kinase receptor, which is a target for imatinib mesylate (Gleevec; Novartis Pharma, Basel, Switzerland). Imatinib therapy of c-Kit-positive tumours is an example of rationally targeted cancer therapy, like trastuzumab for the treatment of Her-2/neu-positive breast cancer. Certain malignancies such as chronic myeloid leukaemia (O'Hare *et al*, 2006) and GISTs (Joensuu *et al*, 2002) express c-Kit and respond favourably to imatinib therapy. In a recent study (Went *et al*, 2004), c-Kit expression was shown to occur infrequently in most tumour types. In our study, only 11% of CUP expressed c-Kit, and its expression was not correlated with any clinical patient characteristics.

Anti-EGFR agents may be evaluated in patients with CUP, which overexpress EGFR immunohistochemically. It would be interesting to also study EGFR mutations and EGFR amplification in tumour samples from these patients. Tumour tissues from GEFCAPI 01 are not available to study other molecular markers such as Her-2 amplification, c-Kit mutations, or EGFR mutations or amplification. This analysis could be performed on another




Figure 1 Immunohistochemical patterns of EGFR expression in a carcinomas of an unknown primary site (CUP) specimen showing strong EGFR immunostaining (magnification $\times 400$).




Figure 2 Immunohistochemical patterns of Her-2/neu expression in a carcinomas of an unknown primary site (CUP) specimen showing strong Her-2/neu immunostaining (magnification $\times 400$).




Figure 3 Immunohistochemical patterns of c-Kit expression in a carcinomas of an unknown primary site (CUP) specimen showing strong c-Kit immunostaining (magnification $\times 400$).

clinical trial and offers more extensive information for selecting targeted therapies. In particular, in lung cancer EGFR mutations or EGFR amplification could predict response to EGFR tyrosine

kinases inhibitors (Shigematsu and Gazdar, 2006). Anti-EGFR agents may be evaluated in second- or first-line therapy, in combination with chemotherapy or alone (Figures 1–3).

REFERENCES

- Baselga J, Arteaga CL (2005) Critical update and emerging trends in epidermal growth factor receptor targeting in cancer. *J Clin Oncol* **23**(11): 2445–2459. E-pub 2005 Mar 7. Review
- Brown ER, Shepherd FA (2005) Erlotinib in the treatment of non-small cell lung cancer. *Expert Rev Anticancer Ther* **5**(5): 767–775. Review
- Culine S, Kramar A, Saghachian M, Bugat R, Lesimple T, Lortholary A, Merrouche Y, Laplanche A, Fizazi K, French Study Group on Carcinomas of Unknown Primary (2002) Development and validation of a prognostic model to predict the length of survival in patients with carcinomas of an unknown primary site. *J Clin Oncol* **20**(24): 4679–4683
- Culine S, Lortholary A, Voigt JJ, Bugat R, Theodore C, Priou F, Kaminsky MC, Lesimple T, Pivot X, Coudert B, Douillard JY, Merrouche Y, Allouache J, Goupil A, Negrier S, Viala J, Petrow P, Bouzy J, Laplanche A, Fizazi K, The Trial for the French Study Group on Carcinomas of Unknown Primary (GEFCAPI 01) (2003) Cisplatin in combination with either gemcitabine or irinotecan in carcinomas of unknown primary site: results of a randomized phase II study – trial for the French Study Group on Carcinomas of Unknown Primary (GEFCAPI 01). *J Clin Oncol* **21**(18): 3479–3482
- Cunningham D, Humblet Y, Siena S, Khayat D, Bleiberg H, Santoro A, Bets D, Mueser M, Harstrick A, Verslype C, Chau I, Van Cutsem E (2004) Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med* **351**(4): 337–345
- Fizazi K (2006) *Carcinoma of an Unknown Primary Site*. Taylor and Francis: New York, NY 10016, USA
- Giaccone G (2005) Epidermal growth factor receptor inhibitors in the treatment of non-small-cell lung cancer. *J Clin Oncol* **23**(14): 3235–3242. Review
- Goldstein NS, Armin M (2001) Epidermal growth factor receptor immunohistochemical reactivity in patients with American Joint Committee on Cancer Stage IV colon adenocarcinoma: implications for a standardized scoring system. *Cancer* **92**(5): 1331–1346
- Greco FA, Burris III HA, Litchy S, Barton JH, Bradof JE, Richards P, Scullin Jr DC, Erland JB, Morrissey LH, Hainsworth JD (2002) Gemcitabine, carboplatin, and paclitaxel for patients with carcinoma of unknown primary site: a Minnie Pearl Cancer Research Network study. *J Clin Oncol* **20**(6): 1651–1656
- Hainsworth JD, Lenington WJ, Greco FA (2000) Overexpression of Her-2 in patients with poorly differentiated carcinoma or poorly differentiated adenocarcinoma of unknown primary site. *J Clin Oncol* **18**(3): 632–635
- Heinrich MC, Blanke CD, Druker BJ, Corless CL (2002) Inhibition of KIT tyrosine kinase activity: a novel molecular approach to the treatment of KIT-positive malignancies. *J Clin Oncol* **20**(6): 1692–1703. Review
- Janne PA, Gurubhagavatula S, Yeap BY, Lucca J, Ostler P, Skarin AT, Fidias P, Lynch TJ, Johnson BE (2004) Outcomes of patients with advanced non-small cell lung cancer treated with gefitinib (ZD1839, 'Iressa') on an expanded access study. *Lung Cancer* **44**(2): 221–230
- Joensuu H, Fletcher C, Dimitrijevic S, Silberman S, Roberts P, Demetri G (2002) Management of malignant gastrointestinal stromal tumours. *Lancet Oncol* **3**(11): 655–664. Review
- Krause DS, Van Etten RA (2005) Tyrosine kinases as targets for cancer therapy. *N Engl J Med* **353**(2): 172–187. Review. No abstract available
- Mendelsohn J, Baselga J (2006) Epidermal growth factor receptor targeting in cancer. *Semin Oncol* **33**(4): 369–385. Review
- O'Hare T, Corbin AS, Druker BJ (2006) Targeted CML therapy: controlling drug resistance, seeking cure. *Curr Opin Genet Dev* **16**(1): 92–99. E-pub 2005 Dec 15. Review
- Pavlidis N, Briassoulis E, Bai M, Fountzilias G, Agnantis N (1995) Overexpression of C-myc, Ras and C-erbB-2 oncoproteins in carcinoma of unknown primary origin. *Anticancer Res* **15**(6B): 2563–2567
- Pavlidis N, Fizazi K (2005) Cancer of unknown primary (CUP). *Crit Rev Oncol Hematol* **54**(3): 243–250. Review
- Sequist LV, Bell DW, Lynch TJ, Haber DA (2007) Molecular predictors of response to epidermal growth factor receptor antagonists in non-small-cell lung cancer. *J Clin Oncol* **25**(5): 587–595. Review
- Shepherd FA, Rodrigues Pereira J, Ciuleanu T, Tan EH, Hirsh V, Thongprasert S, Campos D, Maoleekoonpiroj S, Smylie M, Martins R, van Kooten M, Dediu M, Findlay B, Tu D, Johnston D, Bezjak A, Clark G, Santabarbara P, Seymour L, National Cancer Institute of Canada Clinical Trials Group (2005) Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med* **353**(2): 123–132
- Shigematsu H, Gazdar AF (2006) Somatic mutations of epidermal growth factor receptor signaling pathway in lung cancers. *Int J Cancer* **118**(2): 257–262. Review
- Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, Fleming T, Eiermann W, Wolter J, Pegram M, Baselga J, Norton L (2001) Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* **344**(11): 783–792
- Taron M, Ichinose Y, Rosell R, Mok T, Massuti B, Zamora L, Mate JL, Manegold C, Ono M, Queralt C, Jahan T, Sanchez JJ, Sanchez-Ronco M, Hsue V, Jablons D, Sanchez JM, Moran T (2005) Activating mutations in the tyrosine kinase domain of the epidermal growth factor receptor are associated with improved survival in gefitinib-treated chemorefractory lung adenocarcinomas. *Clin Cancer Res* **11**(16): 5878–5885
- Thatcher N, Chang A, Parikh P, Rodrigues Pereira J, Ciuleanu T, von Pawel J, Thongprasert S, Tan EH, Pemberton K, Archer V, Carroll K (2005) Gefitinib plus best supportive care in previously treated patients with refractory advanced non-small-cell lung cancer: results from a randomised, placebo-controlled, multicentre study (Iressa Survival Evaluation in Lung Cancer). *Lancet* **366**(9496): 1527–1537
- Tsao MS, Sakurada A, Cutz JC, Zhu CQ, Kamel-Reid S, Squire J, Lorimer I, Zhang T, Liu N, Daneshmand M, Marrano P, da Cunha Santos G, Lagarde A, Richardson F, Seymour L, Whitehead M, Ding K, Pater J, Shepherd FA (2005) Erlotinib in lung cancer – molecular and clinical predictors of outcome. *N Engl J Med* **353**(2): 133–144. Erratum in: *N Engl J Med*. 2006 Oct 19;355(16):1746
- Vermorken JB, Trigo J, Hitt R, Koralewski P, Diaz-Rubio E, Rolland F, Knecht R, Amellal N, Schueler A, Baselga J (2007) Open-label, uncontrolled, multicenter phase II study to evaluate the efficacy and toxicity of cetuximab as a single agent in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck who failed to respond to platinum-based therapy. *J Clin Oncol* **25**(16): 2171–2177
- Went PT, Dirnhofer S, Bundi M, Mirlacher M, Schraml P, Mangialaio S, Dimitrijevic S, Kononen J, Lugli A, Simon R, Sauter G (2004) Prevalence of KIT expression in human tumors. *J Clin Oncol* **22**(22): 4514–4522