

RESEARCH ARTICLE

Amyloid- β , cortical thickness, and subsequent cognitive decline in cognitively normal oldest-old

Wiesje Pelkmans¹ , Nienke Legdeur¹ , Mara ten Kate^{1,2} , Frederik Barkhof^{2,3} ,
Maqsood M. Yaqub² , Henne Holstege^{1,4} , Bart N. M. van Berckel² , Philip Scheltens¹ ,
Wiesje M. van der Flier^{1,5} , Pieter Jelle Visser^{1,6,7}  & Betty M. Tijms¹ 

¹Alzheimer Center Amsterdam, Department of Neurology I Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

²Department of Radiology & Nuclear Medicine, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

³Queen Square Institute of Neurology and Centre for Medical Image Computing, UCL, London, UK

⁴Department of Clinical Genetics, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

⁵Department of Epidemiology & Biostatistics, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

⁶Department of Psychiatry & Neuropsychology, School for Mental Health and Neuroscience, Maastricht University, Maastricht, The Netherlands

⁷Department of Neurobiology, Care Sciences and Society, Division of Neurogeriatrics, Karolinska Institutet, Stockholm, Sweden

Correspondence

Wiesje Pelkmans, Alzheimer Center
Amsterdam, Amsterdam UMC, 1007 MB,
Amsterdam, The Netherlands. Tel: +31 (0)20
4440823; Fax: +31 (0)20 4448529; E-mail:
w.pelkmans@amsterdamumc.nl

Funding Information

This work has received support from the EU/
EPPIA Innovative Medicines Initiative Joint
Undertaking (EMIF grant No. 115372) and
received kind sponsoring of the PET-tracer
 [¹⁸F]flutemetamol from GE Healthcare.

Received: 26 June 2020; Revised: 16
November 2020; Accepted: 19 November
2020

*Annals of Clinical and Translational
Neurology* 2021; 8(2): 348–358

doi: 10.1002/acn3.51273

Abstract

Objective: To investigate the relationship between amyloid- β ($A\beta$) deposition and markers of brain structure on cognitive decline in oldest-old individuals with initial normal cognition. **Methods:** We studied cognitive functioning in four domains at baseline and change over time in fifty-seven cognitively intact individuals from the EMIF-AD 90+ study. Predictors were $A\beta$ status determined by [¹⁸F]-flutemetamol PET (normal = $A\beta$ – vs. abnormal = $A\beta$ +), cortical thickness in 34 regions and hippocampal volume. Mediation analyses were performed to test whether effects of $A\beta$ on cognitive decline were mediated by atrophy of specific anatomical brain areas. **Results:** Subjects had a mean age of 92.7 ± 2.9 years, of whom 19 (33%) were $A\beta$ +. Compared to $A\beta$ –, $A\beta$ + individuals showed steeper decline on memory ($\beta \pm SE = -0.26 \pm 0.09$), and processing speed ($\beta \pm SE = -0.18 \pm 0.08$) performance over 1.5 years ($P < 0.05$). Furthermore, medial and lateral temporal lobe atrophy was associated with steeper decline in memory and language across individuals. Mediation analyses revealed that part of the memory decline observed in $A\beta$ + individuals was mediated through parahippocampal atrophy. **Interpretation:** These results show that $A\beta$ abnormality even in the oldest old with initially normal cognition is not part of normal aging, but is associated with a decline in cognitive functioning. Other pathologies may also contribute to decline in the oldest old as cortical thickness predicted cognitive decline similarly in individuals with and without $A\beta$ pathology.

Introduction

Even though amyloid- β ($A\beta$) plaques are considered the pathological hallmark of Alzheimer's disease (AD),^{1,2} $A\beta$ pathology is observed frequently in cognitively normal (CN) adults. Furthermore, postmortem and in vivo studies have shown that the prevalence of abnormal $A\beta$ in CN individuals increases with age from 16% at the age of

60, up to 44% of CNs in their 90's,^{3,4} illustrative of the complex relationship of $A\beta$ deposition with cognitive functioning. While previous studies have reported that $A\beta$ pathology in CN individuals is related to subtle cognitive deficits,^{5–7} and an increased risk for cognitive decline and dementia.^{8–10} The relationship of $A\beta$ pathology and cognitive decline in the oldest-old, that is, individuals of 90 years and older, is however less clear. Recent

longitudinal studies in the oldest-old suggest a steeper cognitive decline in non-demented $A\beta^+$ compared to $A\beta^-$ individuals aged older than 90,^{11–13} although one study did not find such an association.¹⁴ Another driver of cognitive decline is cortical atrophy, in particular medial temporal lobe atrophy, which has been observed frequently in the oldest-old as well.^{11,13} Medial temporal lobe atrophy is considered a key feature of $A\beta$ pathology, but in the oldest-old other causes of medial temporal lobe atrophy are common, such as hippocampal sclerosis, cerebrovascular disease, TDP-43 pathology, and aging-related tau astroglipathy (ARTAG).^{15,16} It still remains unclear how abnormal $A\beta$ is related to cognitive decline in cognitively normal individuals over age 90 and to what extent such effects are mediated by cortical atrophy. In this study, we investigated if $A\beta$ pathology is associated with cognitive decline in CN oldest-old. Additionally, where this association was present, we further investigated whether the effect of $A\beta$ on cognitive decline was independent or mediated by cortical thickness.

Methods

Participants

Individuals with normal cognition who underwent an amyloid positron emission tomography (PET) were selected from the Innovative Medicine Initiative European Medical Information Framework for AD (EMIF-AD) 90+ Study conducted at the Amsterdam University Medical Center (UMC). Individuals were recruited through general practitioners or via advertisements, see Legdeur *et al.* (2018)¹⁷ for detailed description of this cohort and overall study design. Normal cognition was defined as a Clinical Dementia Rating (CDR) score of zero, and a Mini-Mental State Examination (MMSE) score of ≥ 26 .

MRI acquisition and processing

3D-T1 weighted and 3D sagittal fluid-attenuated inversion recovery (FLAIR) images were acquired on a Philips 3T Achieva scanner using an 8-channel head coil and a sagittal turbo gradient-echo sequence (T1: 1.00 mm³ isotropic voxels, repetition time (TR) = 7.9 msec, echo time (TE) = 4.5 msec, flip angle (FA) = 8 degrees; FLAIR: 1.12 mm³ isotropic voxels, TR = 4800 msec, TE = 279 msec, and inversion time = 1650 msec). Cortical thickness was estimated from 3D T1 MRI using FreeSurfer (v5.3; <https://surfer.nmr.mgh.harvard.edu>). Non-brain tissue was removed, followed by transformation to MNI space, segmentation and creation of cortical surface meshes.¹⁸ The cortical thickness values were summarized in anatomical regions according to the Desikan–Killiany atlas implemented in

FreeSurfer. To reduce dimensionality of the data we averaged cortical thickness values for each brain region across hemispheres, resulting in 34 cortical regions of interest (ROIs). Hippocampal volume was obtained with FMRIB's Software Library (FSL) FIRST (v5.0.1), as reported previously in Patenaude *et al.* (2011).¹⁹ White matter hyperintensities (WMH) segmentation was performed using a previously established algorithm based on a three-level Gaussian mixture model to model healthy tissues and lesions.²⁰ Because of a skewed distribution, WMH volume was log transformed. Resulting images were visually checked for quality, and data from four subjects had to be excluded due to gross registration or segmentation errors.

Amyloid PET

Dynamic [¹⁸F] flutemetamol amyloid-PET scans were performed on a Philips Ingenuity TF PET-MRI scanner (Philips Medical Systems, Cleveland, OH, USA). The tracer was produced by General Electric (GE) Healthcare at the Cyclotron Research Center of the University of Liège (Liège, Belgium). First, a 30 min dynamic emission scan was started simultaneously with a bolus intravenous injection of 185 MBq [¹⁸F] flutemetamol. The second part of the scan was performed from 90 to 110 min post injection. Immediately prior to each part of the PET scan a dedicated MR sequence was performed for attenuation correction. During scanning, the head was immobilized to reduce movement artifacts. Data from the two scan parts were coregistered and combined into a single 4D image using VINCI Software 2.56 (<https://vinci.sf.mpg.de>) and in-house built software for decay correction of the second part. Parametric nondisplaceable binding potential (BP_{ND}) images were generated from the entire image set using the receptor parametric mapping and cerebellar gray matter as reference tissue.^{21,22} Global cortical BP_{ND} was calculated as the volume weighted average BP_{ND} of 22 regions located within frontal, parietal, temporal, posterior cingulate, and medial temporal lobes.²³ Dynamic BP_{ND} images were used for visual assessment of [¹⁸F] flutemetamol as negative ($A\beta^-$) or positive ($A\beta^+$) by the consensus of three readers, who had been trained according to the manufacturers image interpretation methodology and were blinded to the clinical and demographic data.²⁴

Neuropsychological assessment

A trained neuropsychologist administered cognitive tests within the following cognitive domains: memory, language, processing speed, and executive functioning. For each cognitive domain, tests were combined into a

composite score. For memory we included the CERAD 10 words test (delayed recall),²⁵ the Wechsler Logical Memory Test (delayed recall),²⁶ the Rey Complex Figure Test (delayed copy),²⁷ and the Visual Association Test A.²⁸ For language, we used the 2 min Animal Fluency score,²⁹ and the Graded Naming Test.³⁰ For processing speed we included the Digit Symbol Substitution Test from the Wechsler Adult Intelligence Scale-Revised (WAIS-R),³¹ the Trail Making Test A,³² and the WAIS-III Digit span forward.³³ For executive functioning we included the Trail Making Test B³², the WAIS-III Digit span backward,³³ Letter Fluency (one minute per letter, three letters),²⁹ and the Clock drawing test.³⁴ For a subset ($n = 43$; 75.4%), neuropsychological tests were repeated once circa 1.5 years (1.0–2.8 y) after baseline assessment. For each test, we calculated Z-scores using the baseline mean and standard deviation of the total group. We created composite scores by averaging test Z-scores for each cognitive domain. Trail Making Test A & B scores were inverted so that for all cognitive tests lower scores indicated worse performance.

APOE genotyping

For all participants, blood samples were collected for DNA analysis. DNA was extracted using the QIAamp[®] DNA Blood Mini Kit (QIAGEN GmbH, Hilden, Germany). Apolipoprotein e (APOE) genotype was determined using TaqMan assays (ThermoFisher Scientific, Foster City, CA, USA) on a QuantStudio-12K-Flex system. We classified individuals as APOE $\epsilon 4$ carriers or non-carriers according to their genotype status at rs429358. For three participants, APOE data was missing.

Statistical analysis

We compared demographical characteristics of the $A\beta^-$ and $A\beta^+$ groups using X^2 tests for categorical variables and ANOVA for continuous variables. We ran four linear mixed models (LMM) with subject specific intercepts, and fixed slopes: Model 1 tested the effect of baseline $A\beta$ status on cognitive decline with cognitive domain score as the outcome (*Model 1: Cognition \sim Time* $A\beta$*). Model 2 tested the effect of $A\beta$ status on cortical thickness of all FreeSurfer ROIs and on hippocampal volume (*Model 2: ROI \sim $A\beta$*). Model 3 tested the effect of cortical thickness and hippocampal volume on cognitive decline (*Model 3: Cognition \sim Time*ROI*). Model 4 tested the combined effects of $A\beta$ pathology, cortical thickness, and their interactions with each other on cognitive decline (*Model 4: Cognition \sim Time*ROI* $A\beta$*). Interaction terms were removed when not significant ($P > 0.05$). All models included sex, education, WMH, and age as covariates,

and for hippocampal volume, total intracranial volume was added as a covariate. LMM were corrected for multiple testing using a false discovery rate (FDR) procedure.³⁵ Effects that did not survive FDR correction are shown as $P_{uncorrected}$.

When $A\beta$ status was significantly associated with both cognitive decline and thickness and/or volume in certain ROIs, we performed causal mediation analyses to assess whether the association between $A\beta$ status and cognitive decline was mediated by grey matter brain atrophy. Mediation analyses provides us with a better understanding of the complex pathways of $A\beta$ deposition toward cognitive decline. To estimate the average causal mediation effect three linear models were fitted: the first model has cognitive decline as the outcome of interest as the dependent variable and thickness/volume as predictor, while controlling for $A\beta$; the second model that has the mediator variable thickness/volume as the dependent variable and $A\beta$ status as predictor; the third model averages direct and indirect effects of $A\beta$ status on cognitive decline and are estimated based on the quasi-Bayesian Monte Carlo approximation (1000 simulations). Mediation analyses was performed only in individuals who had repeated assessment of cognitive function ($n = 43$). Finally, we further investigated the effect of other factors known to be associated with cognitive decline, including APOE $\epsilon 4$ genotype, education, and vascular damage. Statistical analyses were performed in R (v4.0.2) using the “lme4” package (v1.1), “mediation” package (v4.5),³⁶ and group estimates were obtained using the “emmeans” package (v1.5).

Results

Demographics

Participants ($n = 57$) had an average age of 92.7 years, ranging from 88 to 102 years, were more often female (63%), and 33% had a visually rated abnormal amyloid PET scan (Table 1). The $A\beta^+$ ($n = 19$) and $A\beta^-$ ($n = 38$) groups did not differ in age, sex, APOE $\epsilon 4$ carrier status, vascular burden, or years of education. Moreover the two groups did not differ in availability of follow-up data, nor the time between test assessments. More years of education was associated with better performance on tests related to processing speed, executive functioning, and less steep decline in memory and executive functioning over time ($p_{FDR} < 0.05$; Table S3). Higher WMH volumes were associated with worse performance on language, processing speed, and faster decline on memory ($p_{FDR} < 0.05$). No association between age, sex, or APOE $\epsilon 4$ carrier status and cognitive performance or cognitive decline was observed.

Table 1. Subject characteristics according to A β status.

	A β - (n = 38)	A β + (n = 19)	Total (n = 57)	P-value
Sex, f (%)	23 (60.5)	13 (68.4)	36 (63.2)	0.771
Age, (y)	92.51 (3.13)	93.00 (2.56)	92.67 (2.94)	0.554
Education, (y)	12.78 (4.53)	11.50 (4.30)	12.35 (4.46)	0.312
A β load (BPND)	0.15 (0.12)	0.56 (0.28)	0.29 (0.27)	<0.001*
APOE ϵ 4 allele carrier (%)	2 (5.7)	3 (16.7)	5 (9.4)	0.426
WMH volume	9.59 (0.89)	9.71 (0.94)	9.63 (0.90)	0.635
Time between BL and FU, (y)	1.57 (0.57)	1.82 (0.64)	1.66 (0.60)	0.191
FU availability (%)	27 (71.1)	16 (84.2)	43 (75.4)	0.446
Passed away at FU (%)	10 (26.3)	2 (10.5)	12 (21.1)	0.301

Data are presented as mean (SD), or n (%). A β , amyloid- β ; BPND, nondisplaceable binding potential; APOE, apolipoprotein e; WMH, White matter hyperintensities (log); BL, baseline; FU, follow-up.

* $P < 0.05$.

Associations of amyloid status and cortical thickness with cognitive decline

First, we tested the effects of A β status on cognitive decline over time (Model 1). At baseline, A β + individuals tended to show worse performance on memory and language, although this did not reach significance (Fig. 1). Over time, A β + individuals showed steeper decline in memory ($\beta \pm SE = -0.26 \pm 0.09$), and processing speed ($\beta \pm SE = -0.18 \pm 0.08$) than A β - individuals ($p_{uncorrected} < 0.05$; Fig. 1). Also, a steeper decline in language performance ($\beta \pm SE = -0.15 \pm 0.08$) was observed at trend level (Table S1a).

Next, we tested effects of A β status on cortical thickness (Model 2). Individuals with A β + showed thinner parahippocampal cortex and a thinner medial orbitofrontal cortex compared to A β - individuals ($p_{uncorrected} < 0.05$; Fig. 2). Hippocampal volume did not differ between A β - and A β + individuals (Table S1c). Then we studied effects of cortical thickness on cognitive decline over time (Model 3). A thinner cortex in predominantly anterior cingulate and multiple lateral temporal regions, including the entorhinal, parahippocampal, fusiform, and superior temporal cortex, was associated with a steeper decline in memory ($p_{FDR} < 0.05$; Fig. 3), with no interaction effects of A β status. In addition, smaller hippocampal volume was associated with worse memory and processing speed performance at baseline and a faster decline in memory over time ($p_{uncorrected} < 0.05$), independent of A β status. Moreover thinner superior frontal and lateral temporal regions, including the fusiform, inferior and superior temporal, posterior cingulate, and

supramarginal cortex, were associated with a steeper decline in language performance ($p_{uncorrected} < 0.05$), of which only the posterior cingulate cortex survived the correction for multiple comparisons. Additionally, a few cortical regions showed that thicker cortex was associated with a steeper decline in executive functioning ($p_{uncorrected} < 0.05$; Fig. 3).

Finally, we examined the interactions of A β status and brain structure measures on the rate of cognitive decline (Model 4). A β + individuals with a thinner parahippocampal and a thicker pars triangularis ($\beta \pm SE = 1.90 \pm 0.91$; $\beta \pm SE = 4.04 \pm 1.87$; $p_{uncorrected} < 0.05$; Fig. 3) showed worse language performance. While also thinner postcentral and superior parietal cortex ($\beta \pm SE = 1.01 \pm 0.37$; $\beta \pm SE = 0.94 \pm 0.36$; $p_{uncorrected} < 0.05$) was associated with faster language decline in A β - individuals. Furthermore, we observed that thicker cortex in occipital and parietal regions was associated with slower processing speed and steeper decline over time in A β + individuals ($p_{uncorrected} < 0.05$; Fig. 3; see Table S1 for full LMM results). We repeated all analyses excluding $n = 12$ individuals that passed away before the second neuropsychological assessment took place, and observed overall largely similar results (see Table S2).

Mediation analyses

Next, we investigated whether the effect of A β pathology on cognitive performance was mediated by cortical thickness (indirect effect) or not (independent effect) for cortical regions that were associated with abnormal A β (i.e., parahippocampal gyrus and medial orbitofrontal cortex). Memory decline associated with A β + was fully mediated by parahippocampal thinning (36.4%; $P < 0.05$; Fig. 4). Decline in language performance associated with A β + was partially mediated by parahippocampal thinning (21.6%; $P < 0.05$), and partially independent (78.4% $P < 0.05$) after controlling for the presence of parahippocampal atrophy. The effect of A β on decline in processing speed, and executive functioning were independent of parahippocampal thickness. Moreover A β associated cognitive decline was independent of medial orbitofrontal thickness for all cognitive domains ($P < 0.05$; Fig. 4).

Discussion

In this study of oldest-old with initially intact cognition, we found that abnormal A β was associated with steeper decline in memory and processing speed performance over 1.5 years. Our findings support the notion that both A β pathology and brain atrophy have detrimental effects on cognitive functioning among cognitively normal individuals that are separate from normal ageing. These




Figure 1. Cognitive performance over time in relation to $A\beta$ status. Spaghetti plots showing individual longitudinal change on memory, language, processing speed, and executive performance according to $A\beta$ status (Amyloid- is normal $A\beta$; Amyloid+ is abnormal $A\beta$). All scores were z-scored and for processing speed and executive functioning inverted such that lower scores indicate worse impairment.

results suggest that $A\beta$ abnormality is indicative of a neurodegenerative process, that also in the oldest-old with apparent high reserve and maintenance mechanisms lead

to cognitive decline. In addition, non- $A\beta$ pathologies may also contribute to decline in the oldest-old as a thinner medial and lateral temporal cortex was related to




Figure 2. Amyloid associations with regional cortical thickness. Beta's are provided for regions with thinner cortex in abnormal $A\beta$ individuals compared to normal $A\beta$ individuals at $p_{\text{uncorrected}} < 0.05$.

subsequent decline in memory and language irrespective of $A\beta$ pathology, indicating that other, possibly $A\beta$ independent pathological processes might also be involved in cognitive decline in the oldest-old.

Numerous studies have reported on the role of $A\beta$ pathology in very early cognitive decline,^{37–39} and we further extend on those findings by showing that the detrimental effect of $A\beta$ is also present in nonagenarians with initially intact cognition. Our baseline results are in line with other cross-sectional data showing only a subtle effect of $A\beta$ pathology in preclinical AD on cognition,^{6,7,40} consistent with the hypothesis that changes in cognition occur relatively late in the AD pathophysiological cascade. Over time, the differences between $A\beta+$ and $A\beta-$ individuals became larger, as the $A\beta+$ subjects showed steeper decline on memory as reported previously,^{41,42} and also in the processing speed domain.

Furthermore, at baseline $A\beta+$ individuals showed thinner cortex in parahippocampal and orbitofrontal regions compared to $A\beta-$, consistent with other work in CN individuals demonstrating a relationship between $A\beta$ and reduced grey matter.^{43–46} However, while previously hippocampal atrophy has been closely related to memory functioning and shown to be a strong and early predictor of conversion to dementia,^{47–49} we observed no differences between $A\beta+$ and $A\beta-$ individuals in hippocampal volume. Possibly, other pathological factors, such as hippocampal sclerosis, TDP-43 pathology, ARTAG, argyrophilic grain disease (AGD), primary age-related tauopathy (PART), and cerebrovascular disease, may contribute to atrophy in these regions, which at old ages become increasingly more common.^{15,16,50} Such pathologies may also explain our observation that thinner medial and lateral temporal cortex was associated with subsequent decline in memory and language regardless of $A\beta$ status. Associations of decreased gray matter volume in temporal, frontal, and parietal regions with progression to dementia have been demonstrated in individuals without $A\beta$.^{51,52} Selective sparing of these cortical regions, most notably the anterior cingulate cortex and medial temporal

lobe, is frequently reported in superagers compared to age-matched controls and has been associated with resilience to cognitive decline.^{53–55} Whether the oldest-old with good cognitive health in the present study are protected by relatively preserved cortical regions is unknown from the present analyses without a control group, however the anterior cingulate and medial temporal thickness were still negatively associated with cognition.

Mediation analyses indicated that cortical regions that were associated with abnormal $A\beta$ (i.e., parahippocampal gyrus and medial orbitofrontal cortex) partly mediated the effects of $A\beta$ on downstream memory and language decline. $A\beta$ no longer had a significant association with memory performance over time when parahippocampal thickness was included in the mediation model, supporting that the structural integrity of the parahippocampal gyrus is important for memory functioning.^{56,57} A finding that might be related to the commonly observed neurofibrillary tangle pathology in the medial temporal lobe in CN adults.⁵⁸ No significant mediation of cortical thickness was observed for the other cognitive domains, indicative of a direct effect of $A\beta$ pathology on cognitive decline.

Another finding we observed, which did not survive correction for multiple comparisons, was that thicker parietal cortices in $A\beta+$ individuals were associated steeper decline in processing speed. This finding was unexpected, as usually cortical thinning is a sign of neuronal loss. It is unclear what such thicker cortex may reflect. Possibly, as this effect was only observed in abnormal $A\beta$, it might reflect a tissue reactive response to $A\beta$ pathology, as previous studies have shown a positive correlation of microglial activation and $A\beta$ deposition.⁵⁹ Microglial activation in preclinical disease stages has been associated with initially preserved or increased brain volume.^{60,61} Still, it has to be noted that the number of $A\beta+$ subjects in this study was small, and requires replication in larger samples. Further repeated MRI studies are needed to investigate whether these brain areas may show thinning at a later point in time.




Figure 3. Associations of cortical thickness with baseline and decline in cognitive functioning, by $A\beta$ status. Effect of cortical thickness on baseline (left) and annual change (right) on memory, language, processing speed, and executive functioning across all subjects and by $A\beta$ status. Top left: thinner caudal* and rostral* anterior cingulate, entorhinal*, fusiform*, inferior, superior* and middle* temporal, insula*, isthmus cingulate, parahippocampal*, and temporal pole* was associated with a faster decline in memory. Top right: thinner caudal middle frontal, entorhinal, fusiform, inferior and superior temporal, medial orbitofrontal, posterior cingulate*, superior frontal, and supramarginal cortex was associated with faster decline in language. $A\beta+$ individuals with thinner pars triangularis and parahippocampal cortex showed worse language performance compared to $A\beta-$ individuals. $A\beta-$ individuals with thinner postcentral and a superior parietal cortex showed faster decline in language performance compared to $A\beta+$ individuals. Bottom left: $A\beta+$ individuals with thinner parahippocampal and a thicker inferior parietal, pars triangularis, pars opercularis showed worse processing speed performance compared to $A\beta-$ individuals. $A\beta+$ individuals with thicker cuneus, frontal pole, postcentral, rostral middle frontal, superior frontal, and superior parietal cortex showed a faster decline on processing speed compared to $A\beta-$ individuals. Bottom right: thicker lateral occipital, pars opercularis, and postcentral regions were associated with faster decline in executive functioning. $A\beta+$ individuals with thicker precuneus showed a faster decline in executive functioning compared to $A\beta-$ individuals. Beta estimates in red indicate thinner cortex is associated with steeper decline on cognitive test score, blue indicates a thicker cortex is associated with steeper decline on cognitive test score. Data are presented for regions significant with an uncorrected P -value < 0.05 . *Indicates region significant at $p_{FDR} < 0.05$.

Furthermore, we investigated the influence of APOE genotype on cognitive decline, which is considered the major genetic risk factor for sporadic AD.⁶² Carriership

of the $\epsilon 4$ allele has been associated with the presence and lower age of onset of $A\beta$ deposits, and $A\beta$ -associated cognitive decline.^{3,63} The underrepresentation of $\epsilon 4$ -




Figure 4. Mediation analysis showing how cortical thinning mediates the effect of $A\beta$ on longitudinal decline in cognitive functioning. The total effect of $A\beta$ on memory, language, processing speed, and executive functioning over time (left). Mediation effect of $A\beta$ through PHC thickness on memory, language, processing speed, and executive functioning over time (middle). Mediation effect of $A\beta$ through mOFC thickness on memory, language, processing speed, and executive functioning over time (right). The figure shows regression coefficients with a 95% confidence interval. (A) the effect of $A\beta$ on cortical thickness; (B) the effect of cortical thickness on cognitive decline when controlling for $A\beta$; c, the total effect of $A\beta$ on cognitive decline (without controlling for mediation effects); c', the direct effect of $A\beta$ on cognitive decline when adjusting for mediation; c-c', the mediation effect. * $P < 0.05$; $A\beta$, amyloid-beta; PHC, parahippocampal cortex; mOFC, medial orbitofrontal cortex.

carriers in our sample (9.4%), might have contributed to the intact cognition at a very high age in these individuals, and to the absence of an association between $APOE\epsilon 4$ carriership and cognitive decline as well. An interesting next step may lie in investigating factors associated with cognitive resilience among the oldest-old who are $APOE\epsilon 4$ carriers.

There are some limitations to our study. Subjects only underwent an MRI and PET scan at baseline, and cognitive decline was based on cognitive assessments at two time points on average 1.5 years apart. Future studies with longer follow-up will aid in determining the temporal ordering of the changes in brain structure and cognitive impairments. Also, the relatively small number of individuals with abnormal amyloid might have resulted in limited statistical power to detect differences that are expected to be subtle in individuals who have initially intact cognition, which is also reflected by the notion that only a few relationships survived correction for multiple

testing, and uncorrected results should be interpreted with caution. Furthermore, information on tau levels was unavailable in our sample, and so, it remains unknown to what extent effects of $A\beta$ on cortical thinning and cognitive decline was influenced by tau pathology. Previous studies suggest that the presence of abnormal tau biomarkers, the other pathological hallmark of AD, is related to worse cognitive functioning in the presence of $A\beta$,⁸ and increases with age.^{4,64} Therefore, future studies measuring tau pathology using CSF or PET in this age group could contribute to a better understanding of the complex $A\beta$ and tau interaction, and their associations with cognition. Strengths of our study include the extensive phenotyping of a longitudinal cohort of oldest-old individuals with maintained cognitive health.

In conclusion, our findings contribute to our understanding of the role of $A\beta$ deposition on cognitive decline in the oldest-old, and suggest that also at very high ages $A\beta$ abnormality is not benign.

Acknowledgments

This work has received support from the EU/EFPIA Innovative Medicines Initiative Joint Undertaking (EMIF grant No. 115372) and received kind sponsoring of the PET-tracer [^{18}F]flutemetamol from GE Healthcare.

Conflict of Interest

Nothing to report.

References

- Montine TJ, Phelps CH, Beach TG, et al. National Institute on Aging–Alzheimer’s Association guidelines for the neuropathologic assessment of Alzheimer’s disease: a practical approach. *Acta Neuropathol* 2012;123:1–11.
- Jack CR, Bennett DA, Blennow K, et al. NIA-AA Research Framework: toward a biological definition of Alzheimer’s disease. *Alzheimer’s Dement* 2018;14:535–562.
- Jansen WJ, Ossenkoppele R, Knol DL, et al. Prevalence of cerebral amyloid pathology in persons without dementia: a meta-analysis. *JAMA - J Am Med Assoc* 2015;313:1924–1938.
- Elobeid A, Libard S, Leino M, et al. Altered proteins in the aging brain. *J Neuropathol Exp Neurol* 2016;75:316–325.
- Visser PJ, Verhey F, Knol DL, et al. Prevalence and prognostic value of CSF markers of Alzheimer’s disease pathology in patients with subjective cognitive impairment or mild cognitive impairment in the DESCRIPA study: a prospective cohort study. *Lancet Neurol* 2009;8:619–627.
- Hedden T, Oh H, Younger AP, Patel TA. Meta-analysis of amyloid-cognition relations in cognitively normal older adults. *Neurology* 2013;80:1341–1348.
- Jansen WJ, Ossenkoppele R, Tijms BM, et al. Association of cerebral amyloid- β aggregation with cognitive functioning in persons without dementia. *JAMA Psychiatry* 2018;75:84.
- Vos SJ, Xiong C, Visser PJ, et al. Preclinical Alzheimer’s disease and its outcome: a longitudinal cohort study. *Lancet Neurol* 2013;12:957–965.
- Donohue MC, Sperling RA, Petersen R, et al. Association between elevated brain amyloid and subsequent cognitive decline among cognitively normal persons. *JAMA - J Am Med Assoc* 2017;317:2305–2316.
- Ebenau JL, Timmers T, Wesselman LMP, et al. ATN classification and clinical progression in subjective cognitive decline. *Neurology* 2020;95:e46–e58.
- Lopez OL, Klunk WE, Mathis C, et al. Amyloid, neurodegeneration, and small vessel disease as predictors of dementia in the oldest-old. *Neurology* 2014;83:1804–1811.
- Kawas CH, Greenia DE, Bullain SS, et al. Amyloid imaging and cognitive decline in nondemented oldest-old: the 90+ Study. *Alzheimer’s Dement* 2013;9:199–203.
- Zhao Y, Tudorascu DL, Lopez OL, et al. Amyloid β deposition and suspected non-Alzheimer pathophysiology and cognitive decline patterns for 12 years in oldest old participants without dementia. *JAMA Neurol* 2018;75:88.
- Balasubramanian AB, Kawas CH, Peltz CB, et al. Alzheimer disease pathology and longitudinal cognitive performance in the oldest-old with no dementia. *Neurology* 2012;79:915–921.
- Kawas CH, Kim RC, Sonnen JA, et al. Multiple pathologies are common and related to dementia in the oldest-old: the 90 + Study. *Neurology* 2015;85:535–542.
- Robinson JL, Corrada MM, Kovacs GG, et al. Non-Alzheimer’s contributions to dementia and cognitive resilience in The 90+ Study. *Acta Neuropathol* 2018;136:377–388.
- Legdeur N, Badissi M, Carter SF, et al. Resilience to cognitive impairment in the oldest-old: design of the EMIF-AD 90+ study. *BMC Geriatr* 2018;18:289.
- Fischl FB. *FreeSurfer*. *NeuroImage* 2012;62:774–781.
- Patenaude B, Smith SM, Kennedy DN, Jenkinson M. A Bayesian model of shape and appearance for subcortical brain segmentation. *NeuroImage* 2011;56:907–922.
- Sudre CH, Cardoso MJ, Bouvy WH, et al. Bayesian model selection for pathological neuroimaging data applied to white matter lesion segmentation. *IEEE Trans Med Imaging* 2015;34:2079–2102.
- Gunn RN, Lammertsma AA, Hume SP, Cunningham VJ. Parametric imaging of ligand-receptor binding in PET using a simplified reference region model. *NeuroImage* 1997;6:279–287.
- Wu Y, Carson RE. Noise reduction in the simplified reference tissue model for neuroreceptor functional imaging. *J Cereb Blood Flow Metab* 2002;22:1440–1452.
- Tolboom N, Yaqub M, van der Flier WM, et al. Detection of Alzheimer pathology in vivo using both 11C-PIB and 18F-FDDNP PET. *J Nucl Med* 2009;50:191–197.
- Collij LE, Konijnenberg E, Reimand J, et al. Assessing amyloid pathology in cognitively normal subjects using 18 F-flutemetamol PET: comparing visual reads and quantitative methods. *J Nucl Med* 2019;60:541–547.
- Morris JC, Heyman A, Mohs RC, et al. The consortium to establish a registry for Alzheimer’s Disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimer’s disease. *Neurology* 1989;39:1159–1165.
- Wechsler D, Holdnack J, Wechsler DL. *Memory scale: fourth edition technical and interpretive manual*. San Antonio Pearson, 2009.
- Meyers JE, Bayless JD, Meyers KR. Rey complex figure: memory error patterns and functional abilities. *Appl Neuropsychol* 1996;3:89–92.
- Lindeboom J, Schmand B, Tulner L, et al. Visual association test to detect early dementia of the Alzheimer type. *J Neurol Neurosurg Psychiatry* 2002;73:126–133.

29. Zhao Q, Guo Q, Hong Z. Clustering and switching during a semantic verbal fluency test contribute to differential diagnosis of cognitive impairment. *Neurosci Bull* 2013;29:75–82.
30. McKenna P, Warrington EK. Testing for nominal dysphasia. *J Neurol Neurosurg Psychiatry* 1980;43:781–788.
31. Wechsler D. *Manual for the Wechsler Adult Intelligence Scale - Revised*. 1981.
32. Reitan RM. Validity of the trail making test as an indicator of organic brain damage. *Percept Mot Skills* 1958;8:271–276.
33. Wilde N, Strauss E. Functional equivalence of WAIS-III/WMS-III digit and spatial span under forward and backward recall conditions. *Clin Neuropsychol* 2002;16:322–330.
34. Teunisse S, Derix MMA, van Crevel H. Assessing the severity of dementia. *Arch Neurol* 1991;48:274–277.
35. Benjamini Y, Yekutieli D. The control of the false discovery rate in multiple testing under dependency. *Ann Stat* 2001;29:1165–1188.
36. Tingley D, Yamamoto T, Hirose K, et al. mediation: R package for causal mediation analysis. *J Stat Softw* 2014;59:1–38.
37. Petersen RC, Wiste HJ, Weigand SD, et al. Association of elevated amyloid levels with cognition and biomarkers in cognitively normal people from the community. *JAMA Neurol* 2016;73:85.
38. Clark LR, Berman SE, Norton D, et al. Age-accelerated cognitive decline in asymptomatic adults with CSF β -amyloid. *Neurology* 2018;90:e1306–e1315.
39. Bilgel M, An Y, Helpfrey J, et al. Effects of amyloid pathology and neurodegeneration on cognitive change in cognitively normal adults. *Brain* 2018;141:2475–2485.
40. Legdeur N, Tijms BM, Konijnenberg E, et al. Associations of Brain Pathology Cognitive and Physical Markers With Age in Cognitively Normal Individuals Aged 60–102 Years. *Gerontol A Biol Sci Med Sci* 2020;75:1609–1617.
41. Snitz BE, Weissfeld LA, Lopez OL, et al. Cognitive trajectories associated with β -amyloid deposition in the oldest-old without dementia. *Neurology* 2013;80:1378–1384.
42. Storandt M, Mintun MA, Head D, Morris JC. Cognitive decline and brain volume loss as signatures of cerebral amyloid- β peptide deposition identified with pittsburgh compound B. *Arch Neurol* 2009;66:1476–1481.
43. van Bergen JMG, Li X, Quevenco FC, et al. Low cortical iron and high entorhinal cortex volume promote cognitive functioning in the oldest-old. *Neurobiol Aging* 2018;64:68–75.
44. Doherty BM, Schultz SA, Oh JM, et al. Amyloid burden, cortical thickness, and cognitive function in the Wisconsin Registry for Alzheimer's Prevention. *Alzheimer's Dement Diagnosis, Assess Dis Monit* 2015;1:160–169.
45. Doré V, Villemagne VL, Bourgeat P, et al. Cross-sectional and longitudinal analysis of the relationship between A β deposition, cortical thickness, and memory in cognitively unimpaired individuals and in Alzheimer disease. *JAMA Neurol* 2013;70:903.
46. Becker JA, Hedden T, Carmasin J, et al. Amyloid- β associated cortical thinning in clinically normal elderly. *Ann Neurol* 2011;69:1032–1042.
47. Mormino EC, Kluth JT, Madison CM, et al. Episodic memory loss is related to hippocampal-mediated β -amyloid deposition in elderly subjects. *Brain* 2009;132:1310–1323.
48. Huijbers W, Mormino EC, Schultz AP, et al. Amyloid- β deposition in mild cognitive impairment is associated with increased hippocampal activity, atrophy and clinical progression. *Brain* 2015;138:1023–1035.
49. Legdeur N, Visser PJ, Woodworth DC, et al. White matter hyperintensities and hippocampal atrophy in relation to cognition: the 90+ Study. *J Am Geriatr Soc* 2019;67:1827–1834.
50. Beach TG, Monsell SE, Phillips LE, Kukull W. Accuracy of the clinical diagnosis of Alzheimer disease at National Institute on Aging Alzheimer disease centers, 2005–2010. *J Neuropathol Exp Neurol* 2012;71:266–273.
51. Verfaillie SCJ, Tijms B, Versteeg A, et al. Thinner temporal and parietal cortex is related to incident clinical progression to dementia in patients with subjective cognitive decline. *Alzheimer's Dement Diagnosis Assess Dis Monit* 2016;5:43–52.
52. ten Kate M, Barkhof F, Visser PJ, et al. Amyloid-independent atrophy patterns predict time to progression to dementia in mild cognitive impairment. *Alzheimers Res Ther* 2017;9:73.
53. Arenaza-Urquijo EM, Przybelski SA, Lesnick TL, et al. The metabolic brain signature of cognitive resilience in the 80+: beyond Alzheimer pathologies. *Brain* 2019;142:1134–1147.
54. de Godoy LL, Alves CAPF, Saavedra JSM, et al. Understanding brain resilience in superagers: a systematic review. *Neuroradiology* 2020; <https://doi.org/10.1007/s00234-020-02562-1>
55. Harrison TM, Maass A, Baker SL, Jagust WJ. Brain morphology, cognition, and β -amyloid in older adults with superior memory performance. *Neurobiol Aging* 2018;67:162–170.
56. Ward AM, Schultz AP, Huijbers W, et al. The parahippocampal gyrus links the default-mode cortical network with the medial temporal lobe memory system. *Hum Brain Mapp* 2014;35:1061–1073.
57. Köhler S, Black SE, Sinden M, et al. Memory impairments associated with hippocampal versus parahippocampal-gyrus atrophy: an MR volumetry study in Alzheimer's disease. *Neuropsychologia* 1998;36:901–904.
58. Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol* 1991;82:239–259.

59. Meyer-Luehmann M, Spire-Jones TL, Prada C, et al. Rapid appearance and local toxicity of amyloid- β plaques in a mouse model of Alzheimer's disease. *Nature* 2008;451:720–724.
60. Hamelin L, Lagarde J, Dorothée G, et al. Early and protective microglial activation in Alzheimer's disease: a prospective study using 18 F-DPA-714 PET imaging. *Brain* 2016;139:1252–1264.
61. Femminella GD, Dani M, Wood M, et al. Microglial activation in early Alzheimer trajectory is associated with higher gray matter volume. *Neurology* 2019;92:e1331–e1343.
62. Corder EH, Saunders AM, Strittmatter WJ, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 1993;261:921–923.
63. Yamazaki Y, Zhao N, Caulfield TR, et al. Apolipoprotein E and Alzheimer disease: pathobiology and targeting strategies. *Nat Rev Neurol* 2019;15:501–518.
64. Lowe VJ, Wiste HJ, Senjem ML, et al. Widespread brain tau and its association with ageing, Braak stage and Alzheimer's dementia. *Brain* 2018;141:271–287.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1A. Baseline and annual change effect of $A\beta$ status on cognitive functioning.

Table S1B. Cortical thickness (mm) according to amyloid status.

Table S1C. Hippocampal volume according to amyloid status.

Table S1D. Associations of cortical thickness with baseline and decline on memory performance by $A\beta$ status.

Table S1E. Associations of hippocampal volume with baseline and decline on memory performance by $A\beta$ status.

Table S1F. Associations of cortical thickness with baseline and decline on language performance by $A\beta$ status.

Table S1G. Associations of hippocampal volume with baseline and decline on language performance by $A\beta$ status.

Table S1H. Associations of cortical thickness with baseline and decline on processing speed performance by $A\beta$ status.

Table S1I. Associations of hippocampal volume with baseline and decline on processing speed performance by $A\beta$ status.

Table S1J. Associations of cortical thickness with baseline and decline on executive functioning performance by $A\beta$ status.

Table S1K. Associations of hippocampal volume with baseline and decline on executive functioning performance by $A\beta$ status.

Table S2A. Baseline and annual change effect of $A\beta$ status on cognitive functioning in a subset without terminal decline.

Table S2B. Cortical thickness (mm) according to amyloid status in a subset without terminal decline.

Table S2C. Hippocampal volume according to amyloid status in a subset without terminal decline.

Table S2D. Associations of cortical thickness with baseline and decline on memory performance by $A\beta$ status in a subset without terminal decline.

Table S2E. Associations of hippocampal volume with baseline and decline on memory performance by $A\beta$ status in a subset without terminal decline.

Table S2F. Associations of cortical thickness with baseline and decline on language performance by $A\beta$ status in a subset without terminal decline.

Table S2G. Associations of hippocampal volume with baseline and decline on language performance by $A\beta$ status in a subset without terminal decline.

Table S2H. Associations of cortical thickness with baseline and decline on processing speed performance by $A\beta$ status in a subset without terminal decline.

Table S2I. Associations of hippocampal volume with baseline and decline on processing speed performance by $A\beta$ status in a subset without terminal decline.

Table S2J. Associations of cortical thickness with baseline and decline on executive functioning performance by $A\beta$ status in a subset without terminal decline.

Table S2K. Associations of hippocampal volume with baseline and decline on executive functioning performance by $A\beta$ status in a subset without terminal decline.

Table S3. Baseline and annual change of other effects on cognitive functioning.