



Simultaneous confidence intervals for all pairwise differences between the coefficients of variation of rainfall series in Thailand

Noppadon Yosboonruang, Sa-Aat Niwitpong and Suparat Niwitpong

Department of Applied Statistics, King Mongkut's University of Technology North Bangkok, Bangkok, Thailand

ABSTRACT

The delta-lognormal distribution is a combination of binomial and lognormal distributions, and so rainfall series that include zero and positive values conform to this distribution. The coefficient of variation is a good tool for measuring the dispersion of rainfall. Statistical estimation can be used not only to illustrate the dispersion of rainfall but also to describe the differences between rainfall dispersions from several areas simultaneously. Therefore, the purpose of this study is to construct simultaneous confidence intervals for all pairwise differences between the coefficients of variation of delta-lognormal distributions using three methods: fiducial generalized confidence interval, Bayesian, and the method of variance estimates recovery. Their performances were gauged by measuring their coverage probabilities together with their expected lengths via Monte Carlo simulation. The results indicate that the Bayesian credible interval using the Jeffreys' rule prior outperformed the others in virtually all cases. Rainfall series from five regions in Thailand were used to demonstrate the efficacies of the proposed methods.

Subjects Statistics, Computational Science, Natural Resource Management, Ecohydrology

Keywords Bayesian, Coefficient of variation, Fiducial generalized confidence interval, Rainfall, Delta-lognormal distribution

Submitted 31 December 2020

Accepted 31 May 2021

Published 23 June 2021

Corresponding author

Suparat Niwitpong,
suparat.n@sci.kmutnb.ac.th

Academic editor

María Ángeles Esteban

Additional Information and
Declarations can be found on
page 23

DOI 10.7717/peerj.11651

© Copyright
2021 Yosboonruang et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

INTRODUCTION

Thailand is located in Southeast Asia and is classed as a tropical area. It is influenced by both the southwest and northeast monsoons. The southwest monsoon crosses Thailand between mid-May to mid-October (the rainy season) and brings moist air from the Indian Ocean that causes clouds and heavy rain. The northeast monsoon crosses Thailand from mid-October to mid-February (the winter season) causing cold and dry weather. Moreover, the South receives additional heavy rainfall due to moisture coming in from the Gulf of Thailand. The season changes from mid-February to mid-May (the summer season) due to uncertainty in the weather and is influenced by tropical cyclones in the South China Sea, and thus, the weather is generally hot and dry but often with heavy rain and thunderstorms (*Thai Meteorological Department, 2015*). Thailand often endures flooding due to thunderstorms, which can take lives and damage property, especially on farms due to Thailand being an agricultural country. Thailand is divided into five regions according to its

Table 1 The provinces of each regions in Thailand.

Regions	Provinces
Northern	Chiang Rai, Mae Hong Son, Chiang Mai, Phayao, Lamphun, Lampang, Phrae, Nan, Uttaradit, Phitsanulok, Sukhothai, Tak, Phichit, Kamphaeng Phet, Phetchabun
Northeastern	Nong Khai, Bueng Kan, Loei, Udon Thani, Nong Bua Lam Phu, Nakhon Phanom, Sakon Nakhon, Mukdahan, Khon Kaen, Kalasin, Maha Sarakham, Roi Et, Chaiyaphum, Yasothon, Amnat Charoen, Ubon Ratchathani, Sri Sa Ket, Nakhon Ratchasima, Buri Ram, Surin
Central	Nakhon Sawan, Uthai Thani, Chai Nat, Sing Buri, Lop Buri, Ang Thong, Sara buri, Suphan Buri, Ayutthaya, Pathum Thani, Kanchanaburi, Ratchaburi, Nakhon Pathom, Nonthaburi, Bangkok Metropolitan, Samut Prakan, Samut Sakhon, Samut Songkhram
Eastern	Nakhon Nayok, Prachin Buri, Sra Kaeo, Chachoeng Sao, Chon Buri, Rayong, Chanthaburi, Trat
Southern	Phetchaburi, Prachuap Khiri Khan, Chumphon, Surat Thani, Nakhon Si Thammarat, Phatthalung, Songkhla, Pattani, Yala, Narathiwat, Ranong, Phang Nga, Krabi, Phuket Trang, Satun

climate pattern and meteorological conditions (Table 1) (*Thai Meteorological Department, 2015*). Therefore, it is important to investigate rainfall dispersion in each area to gain preliminary information for formulating policies to mitigate such incidents.

There have been numerous studies on rainfall series that have used the delta-lognormal distribution. *Fukuchi (1988)* derived the distribution of correlation coefficients of rainfall rates from two areas as bivariate lognormal (delta-lognormal). *Kedem (1990)* showed that the average rain rate over an area follows a delta-lognormal distribution. *Shimizu (1993)* and *Kong et al. (2012)* presented the maximum likelihood estimation of the parameters of rainfall series containing zeros that followed a bivariate lognormal distribution. Moreover, examples of rainfall series that conform to a delta-lognormal distribution can be founded in various studies by *Maneerat, Niwitpong & Niwitpong (2019a)*; *Maneerat, Niwitpong & Niwitpong (2019b)*; *Maneerat, Niwitpong & Niwitpong (2020a)*; *Maneerat, Niwitpong & Niwitpong (2020b)*; *Yosboonruang, Niwitpong & Niwitpong (2019b)*; *Yosboonruang, Niwitpong & Niwitpong (2020)*, and *Yosboonruang & Niwitpong (2020)*. In addition, a delta-lognormal distribution has been applied in other fields, such as *Ingram Jr. et al. (2010)*; *Owen & DeRouen (1980)*; *Fletcher (2008)*; *Wu & Hsieh (2014)*, and *Zhou & Tu (2000)*. Constructing the confidence intervals for several parameters of a delta-lognormal distribution used in statistical inference has been of interest to many researchers. Confidence intervals for the delta-lognormal mean were proposed by *Owen & DeRouen (1980)*; *Kvanli, Shen & Deng (1998)*; *Zhou & Tu (2000)*; *Tian (2005)*; *Chen & Zhou (2006)*; *Tian & Wu (2006)*; *Fletcher (2008)*; *Li, Zhou & Tian (2013)*; *Wu & Hsieh (2014)*; *Hasan & Krishnamoorthy (2018)*, and *Maneerat, Niwitpong & Niwitpong (2018)*; *Maneerat, Niwitpong & Niwitpong (2019a)*; *Maneerat, Niwitpong & Niwitpong (2019b)*. Furthermore, the confidence intervals for variance and the coefficient of variation (CV) of a delta-lognormal distribution were presented by *Buntao & Niwitpong (2012)*; *Buntao & Niwitpong (2013)*; *Yosboonruang, Niwitpong & Niwitpong (2018)*; *Yosboonruang, Niwitpong*

et al. (2019a); *Yosboonruang, Niwitpong et al.* (2019b); *Yosboonruang, Niwitpong et al.* (2020); *Yosboonruang et al.* (2020), and *Maneerat, Niwitpong et al.* (2020a); *Maneerat, Niwitpong et al.* (2020b).

For statistical inference, the CV, the ratio of the standard deviation to the mean, is a good tool for investigating rainfall dispersion. The advantage of using the CV is that it is unitless and thus, is useful for measuring dispersion in data series with different units or drastically different means. Focusing on inferential statistics, the confidence intervals and functions of the CV for several distributions have been presented. *Wong et al.* (2002) suggested a small-sample asymptotic method for constructing the confidence intervals for the CV of normal and non-normal distributions when the sample size is very small. *Mahmoudvand et al.* (2009) proposed two new methods for constructing the confidence intervals for the CV of a normal distribution and compared them with Miller's, Makay's, Vangel's, and Sharma-Krishna's methods; they found that their proposed methods are more appropriate than the others. *Buntao et al.* (2012) proposed the generalized pivotal approach (GPA) and a closed-form method for variance estimation for the difference between the CVs of lognormal and delta-lognormal distributions; their results show that the GPA is the most suitable. After that, they constructed the confidence intervals for the ratio of the CVs of delta-lognormal distributions using GPA and the method of variance estimates recovery (MOVER) (*Buntao et al.*, 2013); their results were similar to the confidence intervals for the difference between the CVs. *Wongkhao, Niwitpong et al.* (2015) presented the generalized confidence interval (GCI) and MOVER to construct the confidence intervals for the ratio of CVs of normal distributions and then compared their methods with the Verrill and Johnson and bootstrapping methods; they found that GCI and MOVER performed better than the others. *Sangnawakij et al.* (2017a) proposed MOVER, GCI, and the asymptotic confidence interval (ACI) for constructing the confidence interval for the CV and difference between the CVs of two-parameter exponential distributions; their results show that GCI was appropriate for a single CV and ACI worked well for the difference between the CVs. In addition, confidence intervals were extended by *Sangnawakij et al.* (2017b) based on the score and Wald intervals for the difference between and ratio of CVs of two gamma distributions; their proposed methods performed well in a comparative study. Recently, *Yosboonruang, Niwitpong et al.* (2018) proposed GCI and a modified Fletcher method to construct the confidence intervals for the CV of a delta-lognormal distribution and found that GCI was the best. Afterward, they introduced the fiducial GCI (FGCI) and MOVER to construct the confidence intervals for the CV of a delta-lognormal distribution (*Yosboonruang, Niwitpong et al.*, 2019a). Moreover, they compared the confidence intervals based on FGCI and a Bayesian method for the CV of a delta-lognormal distribution (*Yosboonruang, Niwitpong et al.*, 2019b); their results indicate that the Bayesian method outperformed FGCI. *Yosboonruang et al.* (2020) constructed confidence intervals using GCI and MOVER based on variance stabilizing transformation, the Wilson score, and Jeffreys' method for the ratio of the CVs of delta-lognormal distributions; their results show that GCI was the most suitable. *Yosboonruang, Niwitpong et al.* (2020) presented FGCI and a Bayesian

method to construct the confidence interval for the difference of CVs of delta-lognormal distributions; they concluded that the Bayesian method was the most appropriate.

Since dispersion in the precipitation series for different areas can be the same or different, simultaneous estimation of this for multiple areas has been investigated using various distributions and parameters. *Mandel & Betensky (2008)* introduced an algorithm for simultaneous confidence interval (SCI) construction and then compared bootstrapped and normal-based SCIs in which the limits of the bootstrap intervals were smaller than the normal-based intervals. *Donner & Zou (2011)* used a two-step MOVER approach for constructing SCIs for multiple contrasts of binomial proportions; their proposed method was reasonable for small-to-moderate sample sizes. *Abdel-Karim (2015)* considered three methods: FGCI-MOVER, MOVER-MOVER, and simultaneous FGCI to construct SCIs for the ratio of means of lognormal distributions; they reported that the MOVER-MOVER method outperformed the others. *Li, Song & Shi (2015)* suggested parametric bootstrapping to construct SCIs for all pairwise differences between the means of two-parameter exponential distributions. *Thangjai, Niwitpong & Niwitpong (2019)* presented three methods: MOVER, a computational approach, and FGCI to construct SCIs for all of the differences between the CVs of lognormal distributions; their results show that MOVER was the best and the computational approach performed similarly to MOVER when the sample size was large. In addition, *Thangjai & Niwitpong (2020)* used parametric bootstrapping, GCI, and MOVER for SCI construction for all of the differences between CVs in two-parameter exponential distributions; their results indicate that GCI was the most appropriate in most cases, while MOVER was the best for large sample sizes.

As mentioned above, rainfall series data follow a delta-lognormal distribution. Since our focus is on comparing the dispersion of rainfall from five regions in Thailand, the pairwise differences between the CVs of their rainfall data distributions are an interesting topic to study. Although there have been numerous methods published for constructing SCIs for the differences between the parameters of several types of distributions, constructing SCIs for all of the pairwise differences between the CVs of delta-lognormal distributions has not yet been reported. GCI is a general method that is often used to construct confidence intervals, but FGCI is stronger than GCI since it provides asymptotically correct frequentist coverage (*Hannig, Abdel-Karim & Iyer, 2006*). Moreover, previous researchers have reported that MOVER is an appropriate method for constructing the SCIs for various parameters of several types of distributions. Therefore, one of our aims was to construct SCIs for this scenario based on FGCI and compare them with ones based on MOVER and Bayesian methodology. The coverage probability, the probability that the confidence interval of the estimate covers the value of the parameter, together with the expected length were used to estimate the performance of the confidence intervals.

METHODS

Let $X_i = (X_{i1}, X_{i2}, \dots, X_{im_i})$, $i = 1, 2, \dots, k$ be a random sample from k independent delta-lognormal distributions, denoted by $X_{ij} \sim \Delta(\mu_i, \sigma_i^2, \delta_{i(0)})$, where $\delta_{i(0)} = P(X_{ij} = 0)$. Since this distribution contains zero and positive values, then the zero values follow

a binomial distribution and the positive values a lognormal distribution denoted by $X_{ij} = 0 \sim \text{Bin}(n_i, \delta_{i(0)})$ and $Y_{ij} = \ln(X_{ij}) \sim N(\mu_i, \sigma_i^2)$, respectively. Moreover, let $n_{i(0)}$ and $n_{i(1)}$ be the numbers of zero and positive values, respectively, where $n_i = n_{i(0)} + n_{i(1)}$. The distribution function of a delta-lognormal distribution is given by

$$f(x_{ij}; \mu_i, \sigma_i^2, \delta_{i(0)}) = \begin{cases} \delta_{i(0)} & ; x_{ij} = 0 \\ \delta_{i(1)} \frac{1}{\sqrt{2\pi}\sigma_i} \left(\frac{1}{x_{ij}}\right) \exp\left\{-\frac{[\ln(x_{ij}) - \mu_i]^2}{2\sigma_i^2}\right\} & ; x_{ij} > 0, \end{cases} \quad (1)$$

where $\delta_{i(1)} = 1 - \delta_{i(0)}$. Following [Aitchison \(1955\)](#), the respective population mean and variance of X_i are

$$E(X_i) = \mu_{X_i} = \delta_{i(1)} \exp\left(\mu_i + \frac{\sigma_i^2}{2}\right) \quad (2)$$

and

$$\text{Var}(X_i) = \sigma_{X_i}^2 = \delta_{i(1)} \exp(2\mu_i + \sigma_i^2) [\exp(\sigma_i^2) - \delta_{i(1)}]. \quad (3)$$

Following this, the CV of X_i can be expressed as

$$\text{CV}(X_i) = v_i = \sqrt{\frac{\exp(\sigma_i^2) - \delta_{i(1)}}{\delta_{i(1)}}}. \quad (4)$$

Since we are interested in constructing the SCIs for all pairwise differences between the CVs, then

$$v_{il} = v_i - v_l = \sqrt{\frac{\exp(\sigma_i^2) - \delta_{i(1)}}{\delta_{i(1)}}} - \sqrt{\frac{\exp(\sigma_l^2) - \delta_{l(1)}}{\delta_{l(1)}}}, \quad (5)$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$. The maximum likelihood estimators of $\delta_{i(1)}$ and μ_i are $\hat{\delta}_{i(1)} = n_{i(1)}/n_i$ and $\hat{\mu}_i = \sum_{j=1}^{n_{i(1)}} \ln(x_{ij})/n_{i(1)}$, respectively. Furthermore, the unbiased estimator for σ_i^2 is $\hat{\sigma}_i^2 = \sum_{j=1}^{n_{i(1)}} [\ln(x_{ij}) - \hat{\mu}_i]^2 / (n_{i(1)} - 1)$.

Assume that $\hat{\delta}_{i(1)}$ and $\hat{\sigma}_i^2$ are independent, then the maximum likelihood estimator of v_i can be defined as

$$\hat{v}_i = \sqrt{\frac{\exp(\hat{\sigma}_i^2) - \hat{\delta}_{i(1)}}{\hat{\delta}_{i(1)}}}. \quad (6)$$

Similarly,

$$\hat{v}_{il} = \hat{v}_i - \hat{v}_l = \sqrt{\frac{\exp(\hat{\sigma}_i^2) - \hat{\delta}_{i(1)}}{\hat{\delta}_{i(1)}}} - \sqrt{\frac{\exp(\hat{\sigma}_l^2) - \hat{\delta}_{l(1)}}{\hat{\delta}_{l(1)}}}, \quad (7)$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$.

According to *Yosboonruang, Niwitpong & Niwitpong (2020)*, the estimated variance of $\hat{v}_i - \hat{v}_l$ can be expressed as

$$\begin{aligned} \hat{Var}(\hat{v}_i - \hat{v}_l) = & \frac{\left\{ \left[\ln(\hat{\delta}_{i(1)}) + \ln\left(\frac{\exp(\hat{\sigma}_i^2) - \hat{\delta}_{i(1)}}{\hat{\delta}_{i(1)}} + 1\right) \right] \left[\frac{\exp(\hat{\sigma}_i^2) - \hat{\delta}_{i(1)}}{\hat{\delta}_{i(1)}} + 1 \right] \right\}^2}{2n_i \left[\frac{\exp(\hat{\sigma}_i^2) - \hat{\delta}_{i(1)}}{\hat{\delta}_{i(1)}} \right]} \\ & + \frac{\left\{ \left[\ln(\hat{\delta}_{l(1)}) + \ln\left(\frac{\exp(\hat{\sigma}_l^2) - \hat{\delta}_{l(1)}}{\hat{\delta}_{l(1)}} + 1\right) \right] \left[\frac{\exp(\hat{\sigma}_l^2) - \hat{\delta}_{l(1)}}{\hat{\delta}_{l(1)}} + 1 \right] \right\}^2}{2n_l \left[\frac{\exp(\hat{\sigma}_l^2) - \hat{\delta}_{l(1)}}{\hat{\delta}_{l(1)}} \right]}, \end{aligned} \quad (8)$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$.

The simultaneous FGCI

To construct the simultaneous FGCI, a fiducial generalized pivotal quantity (FGPQ), which is a subclass of the generalized pivotal quantity (GPQ) (*Hannig, Iyer & Patterson, 2006*), is presented as follows.

Definition 1 Let $X_i = (X_{i1}, X_{i2}, \dots, X_{ini}), i = 1, 2, \dots, k$ be a random sample from k independent delta-lognormal distributions with parameters of interest $(\sigma_i^2, \delta_{i(1)})$ and nuisance parameter μ_i . Let $x_i = (x_{i1}, x_{i2}, \dots, x_{ini}), i = 1, 2, \dots, k$ be an observed value of X_i . GPQ $R(X_i; x_i, \mu_i, \sigma_i^2, \delta_{i(1)})$ is called an FGPQ if it corresponds with the following two conditions (*Weerahandi, 1993; Hannig, Iyer & Patterson, 2006*): 1. For a given x_i , the conditional distribution of $R(X_i; x_i, \mu_i, \sigma_i^2, \delta_{i(1)})$ is free of μ_i . 2. The observed value of $R(X_i; x_i, \mu_i, \sigma_i^2, \delta_{i(1)})$ at $X_i = x_i, r(x_i; x_i, \mu_i, \sigma_i^2, \delta_{i(1)})$ is the parameter of interest.

The FGPQs for σ_i^2 and $\delta_{i(1)}$ can be constructed by applying [Definition 1](#). According to *Hannig, Iyer & Patterson (2006)* and *Li, Zhou & Tian (2013)*, the respective FGPQs for $\delta_{i(1)}$ and σ_i^2 are

$$R_{\delta_{i(1)}} \sim \frac{1}{2} \text{Beta}(n_{i(1)}, n_{i(0)} + 1) + \frac{1}{2} \text{Beta}(n_{i(1)} + 1, n_{i(0)}) \quad (9)$$

and

$$R_{\sigma_i^2} = \frac{(n_{i(1)} - 1)\hat{\sigma}_i^2}{U_i}, \quad (10)$$

where $U_i \sim \chi_{n_{i(1)} - 1}^2$. Following this, the FGPQ for v_i is simply

$$R_{v_i} = \sqrt{\frac{\exp(R_{\sigma_i^2}) - R_{\delta_{i(1)}}}{R_{\delta_{i(1)}}}}. \quad (11)$$

Hence, the FGPQ for the differences between two independent CVs can be expressed as

$$R_{v_{il}} = R_{v_i} - R_{v_l} = \sqrt{\frac{\exp(R_{\sigma_i^2}) - R_{\delta_{i(1)}}}{R_{\delta_{i(1)}}}} - \sqrt{\frac{\exp(R_{\sigma_l^2}) - R_{\delta_{l(1)}}}{R_{\delta_{l(1)}}}}, \quad (12)$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$.

Therefore, the $100(1-\alpha)\%$ two-sided SCI for $v_i - v_l$ based on the FGCI method can be written as $L_{il} \leq v_{il} \leq U_{il}$, where L_{il} and U_{il} are the $\alpha/2$ -th and $(1-\alpha/2)$ -th quantiles of $R_{v_{il}}$, respectively.

Theorem 1 Let $X_i = (X_{i1}, X_{i2}, \dots, X_{in_i}), i = 1, 2, \dots, k$ be a random sample from k independent delta-lognormal distributions with mean μ_i , variance σ_i^2 , and probability of zero values $\delta_{i(0)}$. Let $v_i = \sqrt{[\exp(\sigma_i^2) - \delta_{i(1)}] / \delta_{i(1)}}$ and $v_l = \sqrt{[\exp(\sigma_l^2) - \delta_{l(1)}] / \delta_{l(1)}}$ for $i, l = 1, 2, \dots, k$ and $i \neq l$ be the CV of X_i and X_l , respectively. Furthermore, let \hat{v}_i and \hat{v}_l be the estimators of v_i and v_l , respectively. The estimator for the variance of the difference between v_i and v_l is $\hat{V}ar(\hat{v}_i - \hat{v}_l)$. Let n_i be the sample size of the i -th random sample and $n = n_1 + n_2 + \dots + n_k$. Assume that $n_i/n \rightarrow r_i$ as $n \rightarrow \infty$ where $0 < r_i < 1$. Therefore,

$$P[R_{v_{il}}(\alpha/2) \leq R_{v_{il}} \leq R_{v_{il}}(1-\alpha/2), \forall i \neq l] \rightarrow 1-\alpha. \quad (13)$$

Proof Since

$$P[R_{v_{il}}(\alpha/2) \leq R_{v_{il}} \leq R_{v_{il}}(1-\alpha/2), \forall i \neq l] = P[L_{il} \leq v_i - v_l \leq U_{il}, \forall i \neq l],$$

where $[L_{il}, U_{il}] = \hat{v}_i - \hat{v}_l \pm d_{1-\alpha} \sqrt{\hat{V}ar(\hat{v}_i - \hat{v}_l)}$ and $d_{1-\alpha}$ denotes the $(1-\alpha)$ -th quantile of $R_{v_{il}}$. Thus,

$$\begin{aligned} P[L_{il} \leq v_i - v_l \leq U_{il}, \forall i \neq l] &= P\left[\max_{i \neq l} \left| \frac{\hat{v}_i - \hat{v}_l - (v_i - v_l)}{\sqrt{\hat{V}ar(\hat{v}_i - \hat{v}_l)}} \right| \leq d_{1-\alpha}\right] \\ &= P[D_n \leq d_{1-\alpha}]. \end{aligned}$$

Accordingly,

$$P[L_{il} \leq v_i - v_l \leq U_{il}, \forall i \neq l] \rightarrow 1-\alpha.$$

This implies that

$$P[R_{v_{il}}(\alpha/2) \leq R_{v_{il}} \leq R_{v_{il}}(1-\alpha/2), \forall i \neq l] \rightarrow 1-\alpha. \quad \square$$

The Bayesian method

According to the distributions of X_i for $i = 1, 2, \dots, k$ with the unknown parameters μ_i, σ_i^2 , and $\delta_{i(0)}$, where $\delta_{i(0)} = 1 - \delta_{i(1)}$, the joint likelihood function of k independent delta-lognormal distributions is

$$L(\mu_i, \sigma_i^2, \delta_{i(1)} | x_{ij}) \propto \prod_{i=1}^k (1 - \delta_{i(1)})^{n_{i(0)}} \delta_{i(1)}^{n_{i(1)}} (\sigma_i^2)^{-\frac{n_{i(1)}}{2}} \exp\left\{-\frac{1}{2\sigma_i^2} \sum_{j=1}^{n_{i(1)}} [\ln(x_{ij}) - \mu_i]^2\right\}. \quad (14)$$

By applying the second-order partial derivative of the log-likelihood function with respect to the unknown parameters, the Fisher information matrix of the unknown parameters can be written as

$$I(\mu_i, \sigma_i^2, \delta_{i(1)}) =$$

$$\text{diag} \left[\frac{n_1}{(1-\delta_{1(1)})\delta_{1(1)}} \quad \frac{n_1\delta_{1(1)}}{\sigma_1^2} \quad \frac{n_1\delta_{1(1)}}{2(\sigma_1^2)^2} \quad \cdots \quad \cdots \quad \cdots \quad \frac{n_k}{(1-\delta_{k(1)})\delta_{k(1)}} \quad \frac{n_k\delta_{k(1)}}{\sigma_k^2} \quad \frac{n_k\delta_{k(1)}}{2(\sigma_k^2)^2} \right]. \quad (15)$$

In this paper, we constructed both of equal-tailed SCIs based on simulation data and simultaneous credible intervals based on information from a simulation study of their prior distributions using two forms of Bayesian prior; the suitability of the Jeffreys' rule and uniform priors was determined by considering the values of a random variable of their posterior distributions that correspond to those for a delta-lognormal distribution. See also, [Yosboonruang, Niwitpong & Niwitpong \(2019b\)](#) and [Yosboonruang, Niwitpong & Niwitpong \(2020\)](#).

The Jeffreys' rule prior

The Jeffreys' rule prior is obtained from the square root of the determinant of the Fisher information matrix ([Jeffreys, 1946](#)). It is well known that a delta-lognormal distribution comprises lognormal and binomial distributions. From the CVs in [Eq. \(4\)](#), the parameters of interest are σ_i^2 and $\delta_{i(1)}$, and the Jeffreys' rule priors for these parameters are $p(\sigma_i^2) \propto \sigma_i^{-3}$ and $p(\delta_{i(1)}) \propto (1-\delta_{i(1)})^{-\frac{1}{2}}\delta_{i(1)}^{\frac{1}{2}}$, respectively. Assuming that σ_i^2 and $\delta_{i(1)}$ are independent, the prior distribution for a delta-lognormal distribution can be defined as $p(\sigma_i^2, \delta_{i(1)}) \propto \sigma_i^{-3}(1-\delta_{i(1)})^{-\frac{1}{2}}\delta_{i(1)}^{\frac{1}{2}}$. By combining the likelihood function and the prior distribution of a delta-lognormal distribution, the joint posterior density function can be written as

$$\begin{aligned} p(\sigma_i^2, \delta_{i(1)} | x_{ij}) &= \prod_{i=1}^k \frac{1}{\text{Beta}(n_{i(0)} + \frac{1}{2}, n_{i(1)} + \frac{3}{2})} (1-\delta_{i(1)})^{(n_{i(0)} + \frac{1}{2})-1} \delta_{i(1)}^{(n_{i(1)} + \frac{3}{2})-1} \\ &\quad \times \frac{1}{\sqrt{2\pi} \frac{\sigma_i}{\sqrt{n_{i(1)}}}} \exp \left[-\frac{1}{2 \frac{\sigma_i^2}{n_{i(1)}}} (\mu_i - \hat{\mu}_i)^2 \right] \frac{\left(\frac{n_{i(1)} \hat{\sigma}_i^2}{2} \right)^{\frac{n_{i(1)}}{2}}}{\Gamma\left(\frac{n_{i(1)}}{2}\right)} (\sigma_i^2)^{-\frac{n_{i(1)}}{2}-1} \\ &\quad \times \exp \left(-\frac{n_{i(1)} \hat{\sigma}_i^2}{2 \sigma_i^2} \right), \end{aligned} \quad (16)$$

where $\hat{\mu}_i = \sum_{j=1}^{n_{i(1)}} \ln(x_{ij}) / n_{i(1)}$, and $\hat{\sigma}_i^2 = \sum_{j=1}^{n_{i(1)}} [\ln(x_{ij}) - \hat{\mu}_i]^2 / (n_{i(1)} - 1)$.

By integrating [Eq. \(16\)](#), the respective posterior distributions of σ_i^2 and $\delta_{i(1)}$ are derived as

$$p(\sigma_i^2 | x_{ij}) \propto \prod_{i=1}^k \frac{\left(\frac{n_{i(1)} \hat{\sigma}_i^2}{2} \right)^{\frac{n_{i(1)}}{2}}}{\Gamma\left(\frac{n_{i(1)}}{2}\right)} (\sigma_i^2)^{-\frac{n_{i(1)}}{2}-1} \exp \left(-\frac{n_{i(1)} \hat{\sigma}_i^2}{2 \sigma_i^2} \right), \quad (17)$$

and

$$p(\delta_{i(1)} | x_{ij}) \propto \prod_{i=1}^k \frac{1}{\text{Beta}(n_{i(0)} + \frac{1}{2}, n_{i(1)} + \frac{3}{2})} (1-\delta_{i(1)})^{(n_{i(0)} + \frac{1}{2})-1} \delta_{i(1)}^{(n_{i(1)} + \frac{3}{2})-1}. \quad (18)$$

It should be noted that $p(\sigma_i^2|x_{ij})$ follows an inverse gamma distribution and $p(\delta_{i(1)}|x_{ij})$ follows a beta distribution, denoted by $\sigma_i^2|x_{ij} \sim \text{Inv} - \text{Gamma}(n_{i(1)}/2, n_{i(1)}\hat{\sigma}_i^2/2)$ and $\delta_{i(1)}|x_{ij} \sim \text{Beta}(n_{i(0)} + 1/2, n_{i(1)} + 3/2)$, respectively. Consequently, $\sigma_i^2|x_{ij}$ and $\delta_{i(1)}|x_{ij}$ can be substituted into (5) to construct the equal-tailed SCI and the simultaneous credible interval, respectively.

The uniform prior

Since the uniform prior has a constant function for the prior probability (Stone, 2013), then the uniform priors of σ_i^2 and $\delta_{i(1)}$ are 1, denoted by $p(\sigma_i^2) \propto 1$ and $p(\delta_{i(1)}) \propto 1$, respectively. Afterward, the uniform prior for a delta-lognormal distribution becomes $p(\sigma_i^2, \delta_{i(1)}) \propto 1$. Similar to Eq. (16), the joint posterior density function is obtained by combining $p(\sigma_i^2, \delta_{i(1)})$ with the likelihood function from Eq. (14). Subsequently, we obtain the posterior of σ_i^2 and $\delta_{i(1)}$ by integrating the joint posterior density function with respect to the others. Thus, the posterior distribution is $\sigma_i^2|x_{ij} \sim \text{Inv} - \text{Gamma}[(n_{i(1)} - 2)/2, (n_{i(1)} - 2)\hat{\sigma}_i^2/2]$ for σ_i^2 and $\delta_{i(1)}|x_{ij} \sim \text{Beta}(n_{i(0)} + 1, n_{i(1)} + 1)$ for $\delta_{i(1)}$.

Therefore, the $100(1 - \alpha)\%$ equal-tailed SCI and simultaneous credible interval for v_{il} based on the Bayesian method are $L_{il} \leq v_{il} \leq U_{il}$, where L_{il} and U_{il} are the lower and upper bounds of the intervals, respectively.

Theorem 2 Let $X_i = (X_{i1}, X_{i2}, \dots, X_{in_i}) \sim \Delta(\mu_i, \sigma_i^2, \delta_{i(1)})$, where $i = 1, 2, \dots, k$ and $\delta_{i(0)} = 1 - \delta_{i(1)}$, with sample sizes n_1, n_2, \dots, n_k and $n = n_1 + n_2 + \dots + n_k$. Let $r_i = n_i/n$ as $n \rightarrow \infty$, where $0 < r_i < 1$. For $i, l = 1, 2, \dots, k$ and $i \neq l$, let $v_i = \sqrt{[\exp(\sigma_i^2) - \delta_{i(1)}]/\delta_{i(1)}}$ and $v_l = \sqrt{[\exp(\sigma_l^2) - \delta_{l(1)}]/\delta_{l(1)}}$ be the CVs of X_i and X_l , respectively. Let \hat{v}_i and \hat{v}_l be the estimators of v_i and v_l , respectively. An estimator for the variance of the difference between v_i and v_l is $\hat{\text{Var}}(\hat{v}_i - \hat{v}_l)$. Let $p(\sigma_i^2, \delta_{i(1)})$ and $p(\sigma_i^2, \delta_{i(1)}|x_{ij})$ be the prior distribution and the joint posterior density function for delta-lognormal distribution, respectively. Therefore,

$$P[L_{il} \leq v_{il} \leq U_{il}, \forall i \neq l] \rightarrow 1 - \alpha. \quad \square \quad (19)$$

Proof The proof is similar to Theorem 1.

Algorithm 1: For the FGCI and Bayesian methods

- Step 1. Generate random samples $X_i, i = 1, 2, \dots, k$, with sample sizes n_1, n_2, \dots, n_k and calculate $\hat{\delta}_{i(1)}$ and $\hat{\sigma}_i^2$.
- Step 2. Generate $U_i \sim \chi_{n_{i(1)}-1}^2, \text{Beta}(n_{i(1)}, n_{i(1)} + 1), \text{Beta}(n_{i(1)} + 1, n_{i(1)}), \text{Beta}(n_{i(0)} + 1/2, n_{i(1)} + 3/2), \text{Beta}(n_{i(0)} + 1, n_{i(1)} + 1), \text{Inv} - \text{Gamma}(n_{i(1)}/2, n_{i(1)}\hat{\sigma}_i^2/2)$, and $\text{Inv} - \text{Gamma}[(n_{i(1)} - 2)/2, (n_{i(1)} - 2)\hat{\sigma}_i^2/2]$.
- Step 3. Calculate $R_{\delta_{i(1)}}, R_{\sigma_i^2}, R_{v_i}, R_{v_l}, v_i$, and v_l .
- Step 4. Repeat Steps 2–3 5,000 times.
- Step 5. Compute the 95% SCIs for v_{il} .
- Step 6. Repeat Steps 1–5 15,000 times.

MOVER

The concept of MOVER proposed by Donner & Zou (2012) can be applied to construct the $100(1 - \alpha)\%$ two-sided confidence interval of $v_i - v_l$ for $i, l = 1, 2, \dots, k$ and $i \neq l$, for

which $L_{il} \leq v_{il} \leq U_{il}$ where L_{il} and U_{il} denote the lower and upper limits of the confidence interval, respectively, expressed as

$$L_{il} = \hat{v}_i - \hat{v}_l - \sqrt{(\hat{v}_i - l_i)^2 + (u_l - \hat{v}_l)^2} \quad (20)$$

and

$$U_{il} = \hat{v}_i - \hat{v}_l + \sqrt{(u_i - \hat{v}_i)^2 + (\hat{v}_l - l_l)^2}, \quad (21)$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$. From (4), the parameters of interest are $\delta_{i(1)}$ and σ_i^2 , and so the confidence intervals for these parameters can be constructed.

Since the unbiased estimator of σ_i^2 is given by $\hat{\sigma}_i^2 = \sum_{j=1}^{n_{i(1)}} [\ln(x_{ij}) - \hat{\mu}_i]^2 / (n_{i(1)} - 1)$, for $i = 1, 2, \dots, k$ and where $(n_{i(1)} - 1) \hat{\sigma}_i^2 / \sigma_i^2 \sim \chi_{n_{i(1)} - 1}^2$. Consequently, the respective lower and upper bounds for σ_i^2 are defined as

$$l_{\sigma_i^2} = \frac{(n_{i(1)} - 1) \hat{\sigma}_i^2}{\chi_{1 - \frac{\alpha}{2}, n_{i(1)} - 1}^2} \quad (22)$$

and

$$u_{\sigma_i^2} = \frac{(n_{i(1)} - 1) \hat{\sigma}_i^2}{\chi_{\frac{\alpha}{2}, n_{i(1)} - 1}^2}. \quad (23)$$

The score method proposed by [Wilson \(1927\)](#) is used to construct the confidence limits for $\delta_{i(1)}$. According to [Brown, Cai & DasGupta \(2001\)](#) and [Donner & Zou \(2011\)](#), the respective lower and upper limits of $\delta_{i(1)}$ are given by

$$l_{\delta_{i(1)}} = \frac{n_{i(1)} + \frac{Z_{i(\alpha/2)}^2}{2}}{n_i + Z_{i(\alpha/2)}^2} - Z_{i(\alpha/2)} \frac{\sqrt{\frac{n_{i(0)} n_{i(1)}}{n_i} + \frac{Z_{i(\alpha/2)}^2}{4}}}{n_i + Z_{i(\alpha/2)}^2} \quad (24)$$

and

$$u_{\delta_{i(1)}} = \frac{n_{i(1)} + \frac{Z_{i(\alpha/2)}^2}{2}}{n_i + Z_{i(\alpha/2)}^2} + Z_{i(\alpha/2)} \frac{\sqrt{\frac{n_{i(0)} n_{i(1)}}{n_i} + \frac{Z_{i(\alpha/2)}^2}{4}}}{n_i + Z_{i(\alpha/2)}^2}, \quad (25)$$

where $Z_i, i = 1, 2, \dots, k$ follow a standard normal distribution. This approach is similar to constructing the confidence limits for σ_i^2 and $\delta_{l(1)}$.

Therefore, the $100(1 - \alpha)\%$ two-sided SCIs for $v_i - v_l$ based on the MOVER method are

$$SCI_{il} = [L_{il}, U_{il}], \quad (26)$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$.

Theorem 3 Let $X_i = (X_{i1}, X_{i2}, \dots, X_{in_i}), i = 1, 2, \dots, k$, be random samples from k independent delta-lognormal distributions with mean μ_i , variance σ_i^2 , and probability of zero values $\delta_{i(0)}$. Furthermore, let the sample size of the i -th random sample be n_i , where $n = n_1 + n_2 + \dots + n_k$ and $r_i = n_i/n$ as $n \rightarrow \infty$, for which $0 < r_i < 1$. Let

$v_i = \sqrt{[\exp(\sigma_i^2) - \delta_{i(1)}] / \delta_{i(1)}}$ and $v_l = \sqrt{[\exp(\sigma_l^2) - \delta_{l(1)}] / \delta_{l(1)}}$, for $i, l = 1, 2, \dots, k$ and $i \neq l$, be the CVs of X_i and X_l , respectively. In addition, let \hat{v}_i and \hat{v}_l be the estimators of v_i and v_l , respectively. Let $L_{il} = \hat{v}_i - \hat{v}_l - \sqrt{(\hat{v}_i - l_i)^2 + (u_l - \hat{v}_l)^2}$ and $U_{il} = \hat{v}_i - \hat{v}_l + \sqrt{(u_i - \hat{v}_i)^2 + (\hat{v}_l - l_l)^2}$, where $i, l = 1, 2, \dots, k$ and $i \neq l$, be the respective lower and upper limits of the confidence interval for $v_{il} = v_i - v_l$. Therefore,

$$P(L_{il} \leq v_{il} \leq U_{il}, \forall i \neq l) \rightarrow 1 - \alpha. \quad (27)$$

Proof Suppose that the respective lower and upper limits of the confidence interval for $v_{il} = v_i - v_l$ are

$$L_{il} = \hat{v}_i - \hat{v}_l - \sqrt{(\hat{v}_i - l_i)^2 + (u_l - \hat{v}_l)^2} = \hat{v}_{il} - \sqrt{(\hat{v}_i - l_i)^2 + (u_l - \hat{v}_l)^2}$$

and

$$U_{il} = \hat{v}_i - \hat{v}_l + \sqrt{(u_i - \hat{v}_i)^2 + (\hat{v}_l - l_l)^2} = \hat{v}_{il} + \sqrt{(u_i - \hat{v}_i)^2 + (\hat{v}_l - l_l)^2},$$

where $i, l = 1, 2, \dots, k$ and $i \neq l$. Thus, the respective estimators of variance for \hat{v}_i and \hat{v}_l at $v_i = l_i$ and $v_l = l_l$ are

$$\hat{Var}(\hat{v}_i) = \frac{(\hat{v}_i - l_i)^2}{z_{\alpha/2}^2}$$

and

$$\hat{Var}(\hat{v}_l) = \frac{(\hat{v}_l - l_l)^2}{z_{\alpha/2}^2},$$

where $z_{\alpha/2}$ is the $\alpha/2$ -th quantile of the standard normal distribution. Similarly, the respective estimators of variance for \hat{v}_i and \hat{v}_l at $v_i = u_i$ and $v_l = u_l$ are

$$\hat{Var}(\hat{v}_i) = \frac{(u_i - \hat{v}_i)^2}{z_{\alpha/2}^2}$$

and

$$\hat{Var}(\hat{v}_l) = \frac{(u_l - \hat{v}_l)^2}{z_{\alpha/2}^2}.$$

Hence, the respective lower and upper limits can be expressed as

$$\begin{aligned} L_{il} &= \hat{v}_{il} - z_{\alpha/2} \sqrt{\frac{(\hat{v}_i - l_i)^2}{z_{\alpha/2}^2} + \frac{(u_l - \hat{v}_l)^2}{z_{\alpha/2}^2}} \\ &= \hat{v}_{il} - z_{\alpha/2} \sqrt{\hat{Var}(\hat{v}_i) + \hat{Var}(\hat{v}_l)} \end{aligned}$$

and

$$\begin{aligned} U_{il} &= \hat{v}_{il} + z_{\alpha/2} \sqrt{\frac{(u_i - \hat{v}_i)^2}{z_{\alpha/2}^2} + \frac{(\hat{v}_l - l_l)^2}{z_{\alpha/2}^2}} \\ &= \hat{v}_{il} + z_{\alpha/2} \sqrt{\hat{Var}(\hat{v}_i) + \hat{Var}(\hat{v}_l)}. \end{aligned}$$

Therefore,

$$\begin{aligned} P(L_{il} \leq v_{il} \leq U_{il}) &= P\left[v_{il} \in \left(\hat{v}_{il} \pm z_{\alpha/2} \sqrt{\hat{V}\text{ar}(\hat{v}_i) + \hat{V}\text{ar}(\hat{v}_l)}\right), \forall i \neq l\right] \\ &= P\left[\max_{i \neq l} \left| \frac{\hat{v}_{il} - v_{il}}{\sqrt{\hat{V}\text{ar}(\hat{v}_i) + \hat{V}\text{ar}(\hat{v}_l)}} \right| \leq z_{\alpha/2}\right] \\ &= P[D'_n \leq z_{\alpha/2}]. \end{aligned}$$

Suppose that $n_i/n \rightarrow r_i \in (0, 1)$ as $n \rightarrow \infty, i = 1, 2, \dots, k$ where $n = n_1 + n_2 + \dots + n_k$. From the central limit theorem, $n(\hat{v}_i - v_i) \xrightarrow{d} Z_i, i = 1, 2, \dots, k$, where $Z_i \stackrel{iid}{\sim} N(0, \sigma_i^2/r_i)$, while from Slutsky's theorem, $D'_n \rightarrow D'$, where $D' = \max_{i \neq l} \left| (Z_i - Z_l) / \sqrt{\sigma_i^2/r_i + \sigma_l^2/r_l} \right|$.

Following Skorokhod's theorem, let Y_n and Y be random variables from the common probability space with distributions D'_n and D' , respectively. Thus, Y_n converges to Y almost surely, denoted by $Y_n \xrightarrow{a.s.} Y$, and D'_n converges to D' almost surely, denoted by $D'_n \xrightarrow{a.s.} D'$. Assume that Z_i and Z_i^* are independent and identically distributed random variables. Thus, $T(X, X^*, \mu, \sigma^2) \rightarrow D'^*$, where $D'^* = \max_{i \neq l} \left| (Z_i^* - Z_l^*) / \sqrt{\sigma_i^2/r_i + \sigma_l^2/r_l} \right|$, for $i, l = 1, 2, \dots, k$, and $i \neq l$. Since the limiting distribution of $T(X, X^*, \mu, \sigma^2)$ is continuous and $z_{\alpha/2}(X) \rightarrow q_{\alpha/2}$, where $q_{\alpha/2}$ is the $\alpha/2$ -th quantile of the distribution of D'^* , we can obtain

$$P(D'_n \leq z_{\alpha/2}) \rightarrow P(D' \leq q_{\alpha/2}) = P(D'^* \leq q_{\alpha/2}) = 1 - \alpha, \text{ as } n \rightarrow \infty. \text{ Therefore,}$$

$$P\left[v_{il} \in \left(\hat{v}_{il} \pm z_{\alpha/2} \sqrt{\hat{V}\text{ar}(\hat{v}_i) + \hat{V}\text{ar}(\hat{v}_l)}, \forall i \neq l\right)\right] \rightarrow 1 - \alpha,$$

which implies that

$$P(L_{il} \leq v_{il} \leq U_{il}, \forall i \neq l) \rightarrow 1 - \alpha. \quad \square$$

Algorithm 2: For MOVER

- Step 1. Generate random samples $X_i, i = 1, 2, \dots, k$ with sample size n_1, n_2, \dots, n_k and calculate $\hat{\delta}_{i(1)}$ and $\hat{\sigma}_i^2$.
- Step 2. Generate $\chi_{1-\alpha/2, n_{i(1)}-1}^2, \chi_{\alpha/2, n_{i(1)}-1}^2$, and $Z_i \sim N(0, 1)$.
- Step 3. Calculate $l_{\sigma_i^2}, l_{\sigma_i^2}, u_{\sigma_i^2}, u_{\sigma_i^2}, l_{\delta_{i(1)}}, l_{\delta_{i(1)}}, u_{\delta_{i(1)}},$ and $u_{\delta_{i(1)}}$.
- Step 4. Repeat Steps 2–3 5,000 times.
- Step 5. Compute the 95% SCIs for v_{il} .
- Step 6. Repeat Steps 1–5 15,000 times.

SIMULATION RESULTS

Here, the performances of the proposed methods via Monte Carlo simulation with the R statistical program are presented. The best method attains a coverage probability equal to or greater than the nominal simultaneous confidence level of 0.95 together with the

shortest expected length. The simulations were conducted with 15,000 iterations for each combination of parameters. Furthermore, 5,000 replications for the FGCI and Bayesian methods for each case of parameter combination were carried out. Sample sizes were set as 25, 50, and 100; $\delta_{i(1)} = 0.2, 0.5, 0.8$; and $\sigma_i^2 = 0.5, 1.0, 2.0$.

The results in Tables 2–4 and Figs. 1–3 show that the coverage probabilities of FGCI and the equal-tailed Bayesian using Jeffreys' rule prior were close to or greater than the nominal confidence level for almost all k values. Similarly, the coverage probabilities of the equal-tailed Bayesian using the uniform prior, the Bayesian credible intervals using Jeffreys' rule and uniform priors, and MOVER were close to or greater than the nominal confidence level for all cases. For most cases, the Bayesian credible interval using Jeffreys' rule prior attained the shortest expected length, except for $n_i = 50$; $\delta_{i(1)} = 0.5, 0.8$; and $\sigma_i^2 = 0.5, 1.0$, for which the expected lengths of FGCI were the shortest.

EMPIRICAL STUDY

Thailand is generally divided into five areas by topography, i.e., Northern (A1), Northeastern (A2), Central (A3), Eastern (A4), and Southern (A5). The daily rainfall data from these areas in August 2020 were used to assess the performances of the proposed methods for SCI construction. The distributions of these data are presented in Fig. 4, which shows right-skewness for all of the datasets. Thus, the minimum Akaike information criterion (AIC) and the lowest Bayesian information criterion (BIC) were used to test the fitting of the distributions to such data. From AIC and BIC results in Table 5, it is evident that the positive values in the rainfall datasets from the five areas conform to lognormal distributions. Moreover, normal Q-Q plots were constructed to show the distributions of the log-transformed positive rainfall data from the five areas (Fig. 5), which verified the AIC and BIC results that these datasets follow lognormal distributions. A summary of these data are

$$\begin{aligned} n_1 &= 31, \hat{\delta}_1 = 0.7097, \hat{\mu}_1 = 0.7715, \hat{\sigma}_1^2 = 3.4565, \hat{\eta}_1 = 6.6088, \\ n_2 &= 31, \hat{\delta}_2 = 0.6774, \hat{\mu}_2 = 1.4332, \hat{\sigma}_2^2 = 2.9550, \hat{\eta}_2 = 5.2294, \\ n_3 &= 31, \hat{\delta}_3 = 0.6452, \hat{\mu}_3 = 1.5512, \hat{\sigma}_3^2 = 2.8638, \hat{\eta}_3 = 5.1154, \\ n_4 &= 31, \hat{\delta}_4 = 0.4839, \hat{\mu}_4 = 1.4178, \hat{\sigma}_4^2 = 2.1487, \hat{\eta}_4 = 4.0888, \\ n_5 &= 31, \hat{\delta}_5 = 0.4839, \hat{\mu}_5 = 1.8040, \hat{\sigma}_5^2 = 2.1962, \hat{\eta}_5 = 4.1930. \end{aligned}$$

Table 6 reports the 95% SCIs and credible intervals for all pairwise differences between the CVs of the daily rainfall series from five areas in Thailand. The results show that the expected length of the Bayesian credible interval using the Jeffreys' rule prior was the shortest, which corresponds with the simulation results. Therefore, it is a good choice for constructing the SCI for all of the pairwise differences between the CVs of the precipitation series from the five areas in Thailand.

DISCUSSION

The simulation results indicate that the Bayesian credible interval using Jeffreys' rule prior outperformed the other methods in virtually all cases. Although the coverage probabilities

Table 2 The coverage probabilities and expected lengths for the 95% SCIs and credible intervals for all pairwise differences between the CVs of delta-lognormal distributions for $k = 3$.

$n_1 : n_2 : n_3$	$\delta_{1(1)} : \delta_{2(1)} : \delta_{3(1)}$	$\sigma_1^2 : \sigma_2^2 : \sigma_3^2$	Coverage probabilities (Expected lengths)					
			FGCI	B.Jrule-E	B.Uni-E	B.Jrule-C	B.Uni-C	MOVER
25:25:25	0.5:0.5:0.5	0.5:0.5:0.5	0.9642	0.9788	0.9842	0.9956	0.9980	0.9986
			(2.1698)	(2.1537)	(2.5030)	(2.0848)	(2.4026)	(3.4408)
		1.0:1.0:1.0	0.9573	0.9605	0.9718	0.9957	0.9984	0.9941
			(6.8818)	(6.2442)	(7.9188)	(5.7177)	(7.1021)	(9.6034)
		2.0:2.0:2.0	0.9516	0.9465	0.9631	0.9978	0.9991	0.9838
			(68.0142)	(53.9752)	(88.1169)	(37.9389)	(55.1280)	(85.1444)
	0.5:1.0:2.0	0.9557	0.9540	0.9678	0.9708	0.9820	0.9878	
		(24.7584)	(20.5209)	(31.5389)	(13.1644)	(17.7557)	(31.4612)	
	0.8:0.8:0.8	0.5:0.5:0.5	0.9542	0.9632	0.9725	0.9834	0.9894	0.9954
			(1.2558)	(1.2613)	(1.3684)	(1.2412)	(1.3444)	(1.8437)
		1.0:1.0:1.0	0.9533	0.9526	0.9636	0.9857	0.9915	0.9868
			(3.2363)	(3.1117)	(3.4638)	(2.9940)	(3.3202)	(4.2113)
2.0:2.0:2.0		0.9499	0.9458	0.9570	0.9930	0.9963	0.9756	
		(16.6440)	(15.4579)	(18.2078)	(13.7933)	(16.0213)	(20.0584)	
0.5:1.0:2.0	0.9513	0.9496	0.9598	0.9659	0.9750	0.9797		
	(7.3206)	(6.8777)	(7.9629)	(5.5921)	(6.3072)	(8.7800)		
50:50:50	0.2:0.2:0.2	0.5:0.5:0.5	0.9692	0.9869	0.9906	0.9991	0.9997	0.9994
			(4.3383)	(4.1487)	(5.2706)	(3.9321)	(4.8731)	(7.2083)
		1.0:1.0:1.0	0.9593	0.9672	0.9778	0.9988	0.9996	0.9965
			(17.7495)	(14.6338)	(23.0519)	(12.3745)	(17.8689)	(25.8720)
		2.0:2.0:2.0	0.9525	0.9489	0.9668	0.9984	0.9996	0.9882
			(813.5319)	(360.7209)	(3.75E+03)	(130.6376)	(338.9741)	(1.00E+03)
	0.5:1.0:2.0	0.9560	0.9567	0.9719	0.9742	0.9846	0.9910	
		(131.2015)	(80.4363)	(238.4643)	(36.1680)	(67.2572)	(169.2385)	
	0.5:0.5:0.5	0.5:0.5:0.5	0.9609	0.9797	0.9827	0.9904	0.9926	0.9990
			(1.2086)	(1.2989)	(1.3657)	(1.2849)	(1.3499)	(1.9562)
		1.0:1.0:1.0	0.9536	0.9613	0.9678	0.9870	0.9907	0.9934
			(3.0015)	(2.9770)	(3.1957)	(2.8997)	(3.1064)	(4.2360)
		2.0:2.0:2.0	0.9496	0.9488	0.9579	0.9910	0.9942	0.9831
			(13.0784)	(12.5308)	(13.8315)	(11.6629)	(12.8048)	(16.6567)
	0.5:1.0:2.0	0.9510	0.9541	0.9619	0.9644	0.9712	0.9872	
		(6.2117)	(6.0214)	(6.5869)	(5.1934)	(5.6079)	(7.9402)	
	0.8:0.8:0.8	0.5:0.5:0.5	0.9545	0.9652	0.9702	0.9764	0.9804	0.9960
			(0.7563)	(0.7866)	(0.8128)	(0.7810)	(0.8069)	(1.1146)
1.0:1.0:1.0		0.9512	0.9530	0.9582	0.9744	0.9788	0.9870	
		(1.7222)	(1.7159)	(1.7838)	(1.6925)	(1.7587)	(2.2244)	
2.0:2.0:2.0		0.9489	0.9476	0.9528	0.9832	0.9863	0.9746	
		(6.2458)	(6.1250)	(6.4310)	(5.9199)	(6.2078)	(7.4204)	
0.5:1.0:2.0	0.9534	0.9532	0.9592	0.9644	0.9690	0.9816		

(continued on next page)

Table 2 (continued)

$n_1 : n_2 : n_3$	$\delta_{1(1)} : \delta_{2(1)} : \delta_{3(1)}$	$\sigma_1^2 : \sigma_2^2 : \sigma_3^2$	Coverage probabilities (Expected lengths)					
			FGCI	B.Jrule-E	B.Uni-E	B.Jrule-C	B.Uni-C	MOVER
100:100:100	0.2:0.2:0.2	0.5:0.5:0.5	(3.2059)	(3.1646)	(3.3108)	(2.8957)	(3.0156)	(3.8254)
			0.9655	0.9862	0.9876	0.9955	0.9969	0.9995
			(2.1032)	(2.2970)	(2.4455)	(2.2649)	(2.4065)	(3.5694)
			0.9572	0.9696	0.9746	0.9935	0.9959	0.9966
			(5.5368)	(5.4606)	(6.0091)	(5.2547)	(5.7570)	(8.2473)
			0.9528	0.9546	0.9629	0.9951	0.9973	0.9891
		1.0:1.0:1.0	(28.1449)	(26.2788)	(30.4653)	(23.6063)	(27.0268)	(37.4768)
			0.9541	0.9585	0.9665	0.9672	0.9744	0.9903
			(12.6988)	(12.0766)	(13.7614)	(9.9027)	(11.0231)	(16.9562)
			0.9541	0.9585	0.9665	0.9672	0.9744	0.9903
			(12.6988)	(12.0766)	(13.7614)	(9.9027)	(11.0231)	(16.9562)
			0.9541	0.9585	0.9665	0.9672	0.9744	0.9903
	0.5:0.5:0.5	0.5:0.5:0.5	0.9595	0.9793	0.9803	0.9852	0.9866	0.9988
			(0.7728)	(0.8596)	(0.8774)	(0.8547)	(0.8723)	(1.2568)
			0.9558	0.9646	0.9673	0.9806	0.9824	0.9949
			(1.7356)	(1.7761)	(1.8252)	(1.7576)	(1.8058)	(2.4523)
			0.9501	0.9513	0.9555	0.9803	0.9831	0.9832
			(6.1101)	(6.0586)	(6.2714)	(5.9140)	(6.1174)	(7.6906)
		1.0:1.0:1.0	0.9518	0.9565	0.9603	0.9613	0.9653	0.9889
			(3.1854)	(3.1976)	(3.2977)	(2.9846)	(3.0716)	(4.0742)
			0.9518	0.9565	0.9603	0.9613	0.9653	0.9889
			(3.1854)	(3.1976)	(3.2977)	(2.9846)	(3.0716)	(4.0742)
			0.9518	0.9565	0.9603	0.9613	0.9653	0.9889
			(3.1854)	(3.1976)	(3.2977)	(2.9846)	(3.0716)	(4.0742)
0.8:0.8:0.8	0.5:0.5:0.5	0.9542	0.9668	0.9691	0.9721	0.9739	0.9966	
		(0.5013)	(0.5298)	(0.5376)	(0.5273)	(0.5350)	(0.7395)	
		0.9495	0.9526	0.9554	0.9654	0.9678	0.9864	
		(1.0816)	(1.0919)	(1.1103)	(1.0843)	(1.1025)	(1.3892)	
		0.9517	0.9516	0.9546	0.9739	0.9756	0.9766	
		(3.4703)	(3.4545)	(3.5212)	(3.4088)	(3.4741)	(4.0656)	
	1.0:1.0:1.0	0.9496	0.9518	0.9552	0.9579	0.9608	0.9814	
		(1.8867)	(1.8883)	(1.9231)	(1.8062)	(1.8377)	(2.2430)	
		0.9496	0.9518	0.9552	0.9579	0.9608	0.9814	
		(1.8867)	(1.8883)	(1.9231)	(1.8062)	(1.8377)	(2.2430)	
		0.9496	0.9518	0.9552	0.9579	0.9608	0.9814	
		(1.8867)	(1.8883)	(1.9231)	(1.8062)	(1.8377)	(2.2430)	
25:50:100	0.5:0.5:0.5	0.9614	0.9782	0.9840	0.9829	0.9885	0.9984	
		(1.4118)	(1.4647)	(1.6172)	(1.4069)	(1.5307)	(2.2374)	
		0.9525	0.9572	0.9670	0.9764	0.9838	0.9941	
		(3.9253)	(3.7120)	(4.3712)	(3.3455)	(3.7992)	(5.4323)	
		0.9525	0.9502	0.9609	0.9827	0.9875	0.9826	
		(27.5125)	(23.2226)	(33.9099)	(16.5621)	(20.9132)	(34.6995)	
	1.0:1.0:1.0	0.9551	0.9600	0.9648	0.9793	0.9827	0.9901	
		(3.8830)	(3.8512)	(4.0951)	(3.6913)	(3.9292)	(5.2078)	
		0.9572	0.9679	0.9740	0.9754	0.9825	0.9958	
		(0.8600)	(0.8810)	(0.9323)	(0.8589)	(0.9050)	(1.2513)	
		0.9516	0.9525	0.9596	0.9709	0.9775	0.9869	
		(2.0776)	(2.0320)	(2.1907)	(1.9232)	(2.0528)	(2.6582)	
25:50:100	0.8:0.8:0.8	0.9513	0.9486	0.9558	0.9776	0.9826	0.9751	
		(8.8667)	(8.4294)	(9.4811)	(7.3245)	(8.0262)	(10.4756)	
		0.9532	0.9545	0.9592	0.9702	0.9735	0.9833	
		(2.2424)	(2.2348)	(2.3100)	(2.1757)	(2.2505)	(2.7807)	
		0.9532	0.9545	0.9592	0.9702	0.9735	0.9833	
		(2.2424)	(2.2348)	(2.3100)	(2.1757)	(2.2505)	(2.7807)	

Notes.

B.Jrule-E, B.Uni-E represented the equal-tailed Bayesian confidence intervals using Jeffreys' rule and uniform priors, respectively, and B.Jrule-C and B.Uni-C represented the Bayesian credible intervals using Jeffrey's rule and uniform priors.

Table 3 The coverage probabilities and expected lengths for the 95% SCIs and credible intervals for all pairwise differences between the CVs of delta-lognormal distributions for $k = 5$.

$n_1 : \dots : n_5$	$\delta_{1(1)} : \dots : \delta_{5(1)}$	$\sigma_1^2 : \dots : \sigma_5^2$	Coverage probabilities (Expected lengths)						
			FGCI	B.Jrule-E	B.Uni-E	B.Jrule-C	B.Uni-C	MOVER	
25 ⁵	0.5 ⁵	0.5 ⁵	0.9643 (2.1555)	0.9794 (2.1428)	0.9854 (2.4893)	0.9958 (2.0756)	0.9981 (2.3915)	0.9986 (3.4245)	
		1.0 ⁵	0.9558 (6.9019)	0.9597 (6.2584)	0.9717 (7.9374)	0.9954 (5.7313)	0.9982 (7.1150)	0.9935 (9.6234)	
		2.0 ⁵	0.9519 (62.1565)	0.9470 (50.5751)	0.9625 (80.8805)	0.9973 (37.1890)	0.9993 (53.9109)	0.9834 (79.5531)	
		0.5 ² : 1.0 : 2.0 ²	0.9545 (25.5380)	0.9538 (23.5178)	0.9676 (37.1670)	0.9715 (15.2344)	0.9818 (20.9148)	0.9870 (36.4956)	
	0.8 ⁵	0.5 ⁵	0.9548 (1.2560)	0.9642 (1.2612)	0.9728 (1.3683)	0.9833 (1.2413)	0.9895 (1.3445)	0.9958 (1.8432)	
		1.0 ⁵	0.9529 (3.2430)	0.9523 (3.1186)	0.9627 (3.4726)	0.9857 (3.0017)	0.9915 (3.3297)	0.9875 (4.2211)	
		2.0 ⁵	0.9495 (16.3407)	0.9447 (15.2264)	0.9567 (17.9101)	0.9929 (13.5908)	0.9963 (15.7761)	0.9757 (19.7367)	
		0.5 ² : 1.0 : 2.0 ²	0.9525 (7.2308)	0.9508 (7.5393)	0.9614 (8.7262)	0.9696 (6.1994)	0.9775 (7.0193)	0.9807 (9.6191)	
	50 ⁵	0.2 ⁵	0.5 ⁵	0.9676 (4.3044)	0.9864 (4.1252)	0.9896 (5.2361)	0.9990 (3.9094)	0.9996 (4.8405)	0.9994 (7.1615)
			1.0 ⁵	0.9610 (17.6359)	0.9688 (14.5147)	0.9792 (22.9054)	0.9988 (12.3017)	0.9997 (17.7946)	0.9969 (25.7023)
			2.0 ⁵	0.9532 (428.000)	0.9499 (253.198)	0.9673 (996.537)	0.9988 (122.265)	0.9997 (272.000)	0.9882 (578.397)
			0.5 ² : 1.0 : 2.0 ²	0.9548 (191.069)	0.9550 (122.152)	0.9707 (490.477)	0.9735 (47.626)	0.9849 (101.477)	0.9903 (276.993)
0.5 ⁵		0.5 ⁵	0.9602 (1.2089)	0.9789 (1.2989)	0.9818 (1.3660)	0.9899 (1.2850)	0.9921 (1.3502)	0.9987 (1.9561)	
		1.0 ⁵	0.9548 (3.0083)	0.9618 (2.9852)	0.9677 (3.2034)	0.9876 (2.9072)	0.9911 (3.1133)	0.9941 (4.2457)	
		2.0 ⁵	0.9506 (13.0787)	0.9496 (12.5396)	0.9578 (13.8536)	0.9913 (11.6771)	0.9944 (12.8287)	0.9839 (16.6737)	
		0.5 ² : 1.0 : 2.0 ²	0.9537 (6.1486)	0.9574 (6.5705)	0.9644 (7.1952)	0.9695 (5.7079)	0.9752 (6.1752)	0.9881 (8.6478)	
0.8 ⁵		0.5 ⁵	0.9541 (0.7579)	0.9654 (0.7881)	0.9700 (0.8142)	0.9763 (0.7825)	0.9799 (0.8083)	0.9958 (1.1163)	
		1.0 ⁵	0.9516 (1.7233)	0.9536 (1.7170)	0.9589 (1.7855)	0.9755 (1.6937)	0.9790 (1.7603)	0.9874 (2.2261)	
		2.0 ⁵	0.9508 (6.2622)	0.9495 (6.1471)	0.9548 (6.4545)	0.9838 (5.9443)	0.9871 (6.2338)	0.9760 (7.4475)	
		0.5 ² : 1.0 : 2.0 ²	0.9512	0.9522	0.9574	0.9646	0.9691	0.9808	

(continued on next page)

Table 3 (continued)

$n_1 : \dots : n_5$	$\delta_{1(1)} : \dots : \delta_{5(1)}$	$\sigma_1^2 : \dots : \sigma_5^2$	Coverage probabilities (Expected lengths)					
			FGCI	B.Jrule-E	B.Uni-E	B.Jrule-C	B.Uni-C	MOVER
100^5	0.2^5	0.5^5	(3.1366)	(3.3900)	(3.5463)	(3.1199)	(3.2508)	(4.0931)
			0.9660	0.9867	0.9883	0.9954	0.9966	0.9994
		1.0^5	(2.1027)	(2.2959)	(2.4445)	(2.2638)	(2.4055)	(3.5676)
			0.9566	0.9680	0.9732	0.9935	0.9958	0.9968
		2.0^5	(5.5583)	(5.4771)	(6.0303)	(5.2701)	(5.7768)	(8.2755)
			0.9521	0.9526	0.9617	0.9951	0.9971	0.9883
	$0.5^2 : 1.0 : 2.0^2$	(27.9967)	(26.1432)	(30.2853)	(23.4497)	(26.8163)	(37.2629)	
		0.9550	0.9598	0.9676	0.9697	0.9763	0.9915	
	0.5^5	0.5^5	(12.6571)	(13.2932)	(15.2030)	(11.0107)	(12.3119)	(18.6781)
			0.9594	0.9791	0.9807	0.9848	0.9862	0.9987
		1.0^5	(0.7720)	(0.8589)	(0.8767)	(0.8540)	(0.8717)	(1.2557)
			0.9546	0.9635	0.9664	0.9793	0.9816	0.9947
2.0^5		(1.7331)	(1.7739)	(1.8229)	(1.7555)	(1.8035)	(2.4495)	
		0.9527	0.9540	0.9582	0.9825	0.9852	0.9848	
$0.5^2 : 1.0 : 2.0^2$	(6.0963)	(6.0429)	(6.2535)	(5.9008)	(6.1028)	(7.6733)		
	0.9527	0.9579	0.9613	0.9645	0.9681	0.9884		
100^5	0.8^5	0.5^5	(3.1078)	(3.4067)	(3.5167)	(3.1961)	(3.2916)	(4.3313)
			0.9542	0.9670	0.9691	0.9721	0.9742	0.9966
		1.0^5	(0.5010)	(0.5295)	(0.5374)	(0.5269)	(0.5348)	(0.7392)
			0.9509	0.9544	0.9570	0.9665	0.9688	0.9875
		2.0^5	(1.0812)	(1.0918)	(1.1102)	(1.0842)	(1.1024)	(1.3890)
			0.9506	0.9507	0.9533	0.9730	0.9752	0.9755
	$0.5^2 : 1.0 : 2.0^2$	(3.4721)	(3.4569)	(3.5238)	(3.4111)	(3.4762)	(4.0692)	
		0.9500	0.9524	0.9552	0.9593	0.9619	0.9806	
	$25^2 : 50 : 100^2$	0.5^5	(1.8380)	(2.0041)	(2.0416)	(1.9233)	(1.9575)	(2.3760)
			0.9608	0.9770	0.9830	0.9830	0.9886	0.9985
		1.0^5	(1.5163)	(1.4923)	(1.6614)	(1.4314)	(1.5704)	(2.2909)
			0.9549	0.9595	0.9688	0.9785	0.9852	0.9940
2.0^5		(4.4104)	(3.8914)	(4.6543)	(3.4894)	(4.0239)	(5.7364)	
		0.9511	0.9484	0.9600	0.9813	0.9872	0.9826	
$0.5^2 : 1.0 : 2.0^2$	(32.7160)	(25.3223)	(37.4111)	(17.8405)	(23.1773)	(37.9900)		
	0.9552	0.9603	0.9655	0.9801	0.9835	0.9901		
0.8^5	0.5^5	(3.8324)	(4.0083)	(4.2595)	(3.8460)	(4.0906)	(5.3939)	
		0.9555	0.9656	0.9719	0.9753	0.9817	0.9959	
	1.0^5	(0.9159)	(0.8965)	(0.9522)	(0.8736)	(0.9239)	(1.2760)	
		0.9506	0.9512	0.9583	0.9707	0.9774	0.9856	
	2.0^5	(2.2438)	(2.0817)	(2.2545)	(1.9682)	(2.1113)	(2.7304)	
		0.9496	0.9469	0.9551	0.9774	0.9824	0.9744	
$0.5^2 : 1.0 : 2.0^2$	(10.1646)	(8.9783)	(10.2014)	(7.7693)	(8.6055)	(11.2231)		
	0.9515	0.9536	0.9582	0.9704	0.9737	0.9832		
			(2.2063)	(2.3189)	(2.3964)	(2.2595)	(2.3367)	(2.8768)

Notes.

 25^5 represents 25:25:25:25:25.

Table 4 The coverage probabilities and expected lengths for the 95% SCIs and credible intervals for all pairwise differences between the CVs of delta-lognormal distributions for $k = 10$.

$n_1 : \dots : n_{10}$	$\delta_{1(1)} : \dots : \delta_{10(1)}$	$\sigma_1^2 : \dots : \sigma_{10}^2$	Coverage probabilities (Expected lengths)					
			FGCI	B.Jrule-E	B.Uni-E	B.Jrule-C	B.Uni-C	MOVER
25 ¹⁰	0.5 ¹⁰	0.5 ¹⁰	0.9644	0.9796	0.9852	0.9958	0.9981	0.9987
			(2.1718)	(2.1549)	(2.5017)	(2.0864)	(2.4056)	(3.4437)
		1.0 ¹⁰	0.9559	0.9596	0.9717	0.9952	0.9982	0.9935
			(6.9095)	(6.2680)	(7.9343)	(5.7342)	(7.1201)	(9.6381)
		2.0 ¹⁰	0.9513	0.9466	0.9627	0.9970	0.9990	0.9832
			(64.2109)	(51.6243)	(83.5403)	(37.5819)	(54.7163)	(81.8119)
	0.8 ¹⁰	0.5 ¹⁰	0.9558	0.9645	0.9733	0.9836	0.9899	0.9958
			(1.2509)	(1.2566)	(1.3621)	(1.2369)	(1.3398)	(1.8373)
		1.0 ¹⁰	0.9518	0.9513	0.9621	0.9855	0.9914	0.9871
			(3.2511)	(3.1246)	(3.4760)	(3.0068)	(3.3358)	(4.2290)
		2.0 ¹⁰	0.9511	0.9468	0.9581	0.9931	0.9962	0.9759
			(16.2516)	(15.1258)	(17.7445)	(13.5223)	(15.6914)	(19.6085)
50 ¹⁰	0.2 ¹⁰	0.5 ¹⁰	0.9678	0.9859	0.9892	0.9988	0.9995	0.9993
			(4.3466)	(4.1581)	(5.2771)	(3.9368)	(4.8806)	(7.2207)
		1.0 ¹⁰	0.9600	0.9680	0.9784	0.9988	0.9997	0.9966
			(17.6759)	(14.5839)	(22.8034)	(12.3672)	(17.8805)	(25.7964)
		2.0 ¹⁰	0.9525	0.9491	0.9671	0.9988	0.9997	0.9881
			(584.809)	(314.782)	(1958.586)	(130.579)	(314.105)	(825.014)
	0.5 ¹⁰	0.5 ¹⁰	0.9614	0.9800	0.9828	0.9905	0.9927	0.9989
			(1.2078)	(1.2981)	(1.3643)	(1.2842)	(1.3493)	(1.9553)
		1.0 ¹⁰	0.9543	0.9614	0.9676	0.9874	0.9909	0.9942
			(2.9985)	(2.9750)	(3.1898)	(2.8975)	(3.1028)	(4.2323)
		2.0 ¹⁰	0.9505	0.9497	0.9577	0.9915	0.9944	0.9834
			(13.0586)	(12.5061)	(13.7967)	(11.6493)	(12.7980)	(16.6309)
0.8 ¹⁰	0.5 ¹⁰	0.9543	0.9653	0.9699	0.9761	0.9799	0.9960	
		(0.7572)	(0.7873)	(0.8131)	(0.7817)	(0.8075)	(1.1153)	
	1.0 ¹⁰	0.9510	0.9530	0.9584	0.9747	0.9791	0.9872	
		(1.7261)	(1.7192)	(1.7869)	(1.6958)	(1.7624)	(2.2287)	
	2.0 ¹⁰	0.9500	0.9485	0.9541	0.9835	0.9866	0.9749	
		(6.2757)	(6.1553)	(6.4591)	(5.9506)	(6.2399)	(7.4567)	
100 ¹⁰	0.2 ¹⁰	0.5 ¹⁰	0.9658	0.9868	0.9885	0.9957	0.9968	0.9994
			(2.1038)	(2.2979)	(2.4450)	(2.2660)	(2.4078)	(3.5707)
		1.0 ¹⁰	0.9573	0.9684	0.9738	0.9937	0.9959	0.9968
	0.5 ¹⁰	0.9515	0.9523	0.9612	0.9950	0.9972	0.9881	
		(27.8994)	(26.0553)	(30.1289)	(23.3884)	(26.7565)	(37.1355)	
		1.0 ¹⁰	0.9595	0.9799	0.9813	0.9856	0.9869	0.9989
100 ¹⁰	0.5 ¹⁰	0.9595	0.9799	0.9813	0.9856	0.9869	0.9989	
		(0.7720)	(0.8589)	(0.8767)	(0.8540)	(0.8718)	(1.2559)	
		1.0 ¹⁰	0.9536	0.9622	0.9653	0.9784	0.9810	0.9945

(continued on next page)

Table 4 (continued)

$n_1 : \dots : n_{10}$	$\delta_{1(1)} : \dots : \delta_{10(1)}$	$\sigma_1^2 : \dots : \sigma_{10}^2$	Coverage probabilities (Expected lengths)					
			FGCI	B.Jrule-E	B.Uni-E	B.Jrule-C	B.Uni-C	MOVER
0.8^{10}		2.0^{10}	(1.7364)	(1.7768)	(1.8256)	(1.7582)	(1.8066)	(2.4531)
			0.9505	0.9518	0.9557	0.9815	0.9841	0.9838
		0.5^{10}	(6.1121)	(6.0620)	(6.2702)	(5.9173)	(6.1188)	(7.6931)
			0.9541	0.9668	0.9689	0.9720	0.9741	0.9964
		1.0^{10}	(0.5013)	(0.5298)	(0.5376)	(0.5272)	(0.5351)	(0.7395)
			0.9508	0.9542	0.9569	0.9665	0.9689	0.9874
2.0^{10}	(1.0804)	(1.0907)	(1.1089)	(1.0832)	(1.1014)	(1.3878)		
		0.9503	0.9502	0.9530	0.9726	0.9747	0.9753	
		(3.4656)	(3.4492)	(3.5157)	(3.4036)	(3.4692)	(4.0610)	

Notes.
 25^{10} represents 25:25:25:25:25:25:25:25:25:25.




Figure 1 Comparison of the performances of the proposed methods in terms of their coverage probabilities and expected lengths with various sample sizes: (A) $k = 3$ (B) $k = 5$ (C) $k = 10$.

Full-size DOI: 10.7717/peerj.11651/fig-1

in some cases were close to 1.00, suggesting that overestimation may have occurred, the expected lengths were the shortest. Therefore, the Bayesian credible interval using Jeffreys' rule prior can be used to construct the SCIs for all of the pairwise differences between the CVs of delta-lognormal distributions. Since constructing SCIs concerns the differences between the parameters of interest for all pairwise comparisons, our findings correspond




Figure 2 Comparison of the performances of the proposed methods in terms of their coverage probabilities and expected lengths with various probabilities of non-zero values: (A) $k = 3$ (B) $k = 5$ (C) $k = 10$.

Full-size  DOI: [10.7717/peerj.11651/fig-2](https://doi.org/10.7717/peerj.11651/fig-2)

with *Yosboonruang, Niwitpong & Niwitpong (2020)* who found that the highest posterior density Bayesian using Jeffreys' rule prior is appropriate for constructing the confidence interval for the difference between two independent CVs of delta-lognormal distributions. However, *Abdel-Karim (2015)* and *Thangjai, Niwitpong & Niwitpong (2019)* reported that MOVER is the most suitable for constructing SCIs for the mean or CV of a lognormal distribution, but this is not in agreement with our findings for the data and scenario used in this study since the range of intervals for its SCI was wider than when using the Bayesian methods. In addition, the SCI range between the CVs of the daily rainfall data series from the five different areas of Thailand was too wide, and so this demonstrates that it is different in rainfall dispersion from five areas in Thailand.

CONCLUSIONS

Herein, we proposed methods to construct the SCIs for all pairwise differences between the CVs of delta-lognormal distributions, including FGCI, two Bayesian methods constructed under the equal-tailed confidence intervals and credible intervals using the Jeffreys' rule and uniform priors, and MOVER. The performances of the proposed methods were determined via their coverage probabilities together with their expected lengths under




Figure 3 Comparison of the performances of the proposed methods in terms of their coverage probabilities and expected lengths with various variances: (A) $k = 3$ (B) $k = 5$ (C) $k = 10$.

Full-size DOI: [10.7717/peerj.11651/fig-3](https://doi.org/10.7717/peerj.11651/fig-3)

Table 5 AIC and BIC results for testing the distributions of the positive daily rainfall data from the five areas of Thailand in August 2020.

Regions	AIC				BIC			
	Normal	Lognormal	Cauchy	Exponential	Normal	Lognormal	Cauchy	Exponential
Northern	200.1677	126.6431	154.5509	143.7143	202.3498	128.8252	156.7329	144.7143
Northeastern	186.9685	145.5208	170.8114	151.8661	189.0576	147.6098	172.9005	152.9106
Central	187.4002	142.8220	159.7491	148.7169	189.3916	144.8135	161.7405	149.7126
Eastern	129.2900	99.5405	110.7899	100.7356	130.7061	100.9566	112.2060	101.4437
Southern	140.0174	111.4550	124.1323	114.2896	141.4335	112.8711	125.5484	114.9977

various circumstances. The results indicate that the Bayesian credible interval using the Jeffreys' rule prior was suitable for constructing the SCIs for all pairwise differences between the CVs of delta-lognormal distributions in terms of the coverage probability together with the expected length. Furthermore, FGCI is appropriate for constructing these SCIs in cases of the variances equal to 0.5 and 1.0 with the proportion of non-zero values equal to 0.5 and 0.8 for the sample sizes of 50 and 100. In addition, the results of using daily rainfall data from five regions in Thailand coincided with those from the simulation study.




Figure 4 The density of daily rainfall data in the five areas of Thailand in August 2020.

Full-size [DOI: 10.7717/peerj.11651/fig-4](https://doi.org/10.7717/peerj.11651/fig-4)




Figure 5 Normal Q-Q plots of the log-transformed positive daily rainfall data from the five areas of Thailand in August 2020.

Full-size [DOI: 10.7717/peerj.11651/fig-5](https://doi.org/10.7717/peerj.11651/fig-5)

Table 6 The 95% two-sided confidence intervals and credible intervals for all pairwise differences between the CVs of daily rainfall data from the five areas of Thailand in August 2020.

Regions	CI_{FGCI}	$CI_{B, Jrule-E}$	$CI_{B, Uni-E}$	$CI_{B, Jrule-C}$	$CI_{B, Uni-C}$	CI_{MOVER}
A1-A2	[-18.6001,34.6174]	[-18.9230,29.7356]	[-17.4271,40.6785]	[-20.0641,27.7802]	[-22.5574,31.0130]	[-24.4099,41.5271]
A1-A3	[-17.9489,34.8373]	[-14.7811,30.1761]	[-19.4179,40.0759]	[-18.8783,24.1259]	[-20.8181,36.4352]	[-24.6036,41.6372]
A1-A4	[-12.0747,36.3137]	[-11.4058,31.0863]	[-13.2535,43.2976]	[-15.6054,24.9219]	[-18.9824,33.3564]	[-19.2874,42.6266]
A1-A5	[-14.6835,36.5005]	[-12.4631,30.6305]	[-15.3529,42.0874]	[-15.9005,25.3713]	[-18.3991,35.6191]	[-20.8273,42.5263]
A2-A3	[-19.1684,19.8007]	[-15.4068,20.8296]	[-21.9396,19.5370]	[-16.9424,18.6778]	[-18.6853,21.7661]	[-25.8643,25.7772]
A2-A4	[-14.2284,20.9574]	[-12.1077,21.4176]	[-15.2508,20.4711]	[-12.3215,21.1200]	[-16.0818,19.4744]	[-20.5246,26.7455]
A2-A5	[-15.1643,20.7719]	[-14.3124,21.0636]	[-16.9309,20.4638]	[-15.1223,19.4770]	[-15.7284,20.9116]	[-22.0733,26.6474]
A3-A4	[-14.0533,20.3410]	[-12.8066,16.4067]	[-16.1177,22.6669]	[-13.4979,15.6852]	[-17.3305,21.2102]	[-20.6315,26.9413]
A3-A5	[-15.2305,20.4287]	[-14.3490,16.8954]	[-17.6976,22.7643]	[-14.9963,16.0516]	[-16.0804,23.8267]	[-22.1806,26.8432]
A4-A5	[-17.1536,14.8658]	[-15.0784,13.8954]	[-18.1792,16.4151]	[-14.9530,14.0968]	[-18.1763,16.4173]	[-23.1424,21.4920]

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research was funding by King Mongkut's University of Technology North Bangkok. Grant number: KMUTNB-BasicR-64-32. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

King Mongkut's University of Technology North Bangkok: KMUTNB-BasicR-64-32.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Noppadon Yosboonruang and Suparat Niwitpong conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Sa-Aat Niwitpong performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

All the data and R code are available as [Supplemental File](#).

The data from northern, northeastern, and southern regions of Thailand were collected by the Northern, Upper Northeastern, and Southern East Coast Meteorological Centers and published at:

- Northern Meteorological Center: Available at <http://www.cmnet.tmd.go.th/>,

- Upper Northeastern Meteorological Center: <http://www.khonkaen.tmd.go.th/Home.php>,

- Southern-East Coast Meteorological Center: <http://www.songkhla.tmd.go.th/>.

The data from central Thailand was collected by the Thai Meteorological Department at: <https://www.tmd.go.th/en/>.

The data from eastern Thailand was obtained from the Eastern Region Irrigation Hydrology Center at: <http://hydro-6.com/>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.11651#supplemental-information>.

REFERENCES

- Abdel-Karim AH. 2015.** Construction of simultaneous confidence intervals for ratios of means of lognormal distributions. *Communications in Statistics - Simulation and Computation* **44**(2):271–283 DOI [10.1080/03610918.2013.767911](https://doi.org/10.1080/03610918.2013.767911).
- Aitchison J. 1955.** On the distribution of a positive random variable having a discrete probability mass at the origin. *Journal of the American Statistical Association* **50**:901–908.
- Brown LD, Cai TT, DasGupta A. 2001.** Interval estimation for a binomial proportion. *Statistical Science* **16**(2):101–117 DOI [10.1214/ss/1009213285](https://doi.org/10.1214/ss/1009213285).
- Buntao N, Niwitpong S-A. 2012.** Confidence intervals for the difference of coefficients of variation for lognormal distributions and delta-lognormal distributions. *Applied Mathematical Sciences* **6**(134):6691–6704.
- Buntao N, Niwitpong S-A. 2013.** Confidence intervals for the ratio of coefficients of variation of delta-lognormal distribution. *Applied Mathematical Sciences* **7**(77):3811–3818 DOI [10.12988/ams.2013.35248](https://doi.org/10.12988/ams.2013.35248).
- Chen Y-H, Zhou X-H. 2006.** Generalized confidence intervals for the ratio or difference of two means for lognormal populations with zeros. *UW Biostatistics Working Paper Series*.
- Donner A, Zou GY. 2011.** Estimating simultaneous confidence intervals for multiple contrasts of proportions by the method of variance estimates recovery. *Statistics in Biopharmaceutical Research* **3**(2):320–335 DOI [10.1198/sbr.2010.09050](https://doi.org/10.1198/sbr.2010.09050).
- Donner A, Zou GY. 2012.** Closed-form confidence intervals for functions of the normal mean and standard deviation. *Statistical Methods in Medical Research* **21**(4):347–359 DOI [10.1177/0962280210383082](https://doi.org/10.1177/0962280210383082).
- Fletcher D. 2008.** Confidence intervals for the mean of the delta-lognormal distribution. *Environmental and Ecological Statistics* **15**(2):175–189 DOI [10.1007/s10651-007-0046-8](https://doi.org/10.1007/s10651-007-0046-8).
- Fukuchi H. 1988.** Correlation properties of rainfall rates in the United Kingdom. *Antennas and Propagation IEE Proceedings H - Microwaves* **135**(2):83–88 DOI [10.1049/ip-h-2.1988.0018](https://doi.org/10.1049/ip-h-2.1988.0018).
- Hannig JEL, Abdel-Karim A, Iyer H. 2006.** Simultaneous fiducial generalized confidence intervals for ratios of means of lognormal distributions. *Austrian Journal of Statistics* **35**(2&3):261–269.

- Hannig J, Iyer H, Patterson P. 2006.** Fiducial generalized confidence intervals. *Journal of the American Statistical Association* **101**(473):254–269
[DOI 10.1198/016214505000000736](https://doi.org/10.1198/016214505000000736).
- Hasan MS, Krishnamoorthy K. 2018.** Confidence intervals for the mean and a percentile based on zero-inflated lognormal data. *Journal of Statistical Computation and Simulation* **88**(8):1499–1514 [DOI 10.1080/00949655.2018.1439033](https://doi.org/10.1080/00949655.2018.1439033).
- Ingram Jr. GW, Richards WJ, Lamkin JT, Muhling B. 2010.** Annual indices of Atlantic bluefin tuna (*Thunnus thynnus*) larvae in the Gulf of Mexico developed using delta-lognormal and multivariate models. *Aquatic Living Resources* **23**:35–47
[DOI 10.1051/alr/2009053](https://doi.org/10.1051/alr/2009053).
- Jeffreys H. 1946.** An invariant form for the prior probability in estimation problems. *Proceedings of the Royal Society of London. Series a. Mathematical and Physical Sciences* **186**(1007):453–461 [DOI 10.1098/rspa.1946.0056](https://doi.org/10.1098/rspa.1946.0056).
- Kedem B. 1990.** Estimation of mean rain rate: application to satellite observations. *Journal of Geophysical Research* **95**(D2):1965–1972 [DOI 10.1029/JD095iD02p01965](https://doi.org/10.1029/JD095iD02p01965).
- Kong CY, Jamaludin S, Yusof F, Foo HM. 2012.** Parameter estimation for bivariate mixed lognormal distribution. *Journal of Science and Technology* **4**(1):41–48.
- Kvanli AH, Shen YK, Deng LY. 1998.** Construction of confidence intervals for the mean of a population containing many zero values. *Journal of Business & Economic Statistics* **16**(3):362–368 [DOI 10.2307/1392513](https://doi.org/10.2307/1392513).
- Li J, Song W, Shi J. 2015.** Parametric bootstrap simultaneous confidence intervals for differences of means from several two-parameter exponential distributions. *Statistics and Probability Letters* **106**:39–45 [DOI 10.1016/j.spl.2015.07.002](https://doi.org/10.1016/j.spl.2015.07.002).
- Li X, Zhou X, Tian L. 2013.** Interval estimation for the mean of lognormal data with excess zeros. *Statistics and Probability Letters* **83**(11):2447–2453
[DOI 10.1016/j.spl.2013.07.004](https://doi.org/10.1016/j.spl.2013.07.004).
- Mahmoudvand R, Hassani H. 2009.** Two new confidence intervals for the coefficient of variation in a normal distribution. *Journal of Applied Statistics* **36**(4):429–442
[DOI 10.1080/02664760802474249](https://doi.org/10.1080/02664760802474249).
- Mandel M, Betensky RA. 2008.** Simultaneous confidence intervals based on the percentile bootstrap approach. *Computational Statistics & Data Analysis* **52**(4):2158–2165 [DOI 10.1016/j.csda.2007.07.005](https://doi.org/10.1016/j.csda.2007.07.005).
- Maneerat P, Niwitpong S-A, Niwitpong S. 2018.** Confidence intervals for the ratio of means of delta-lognormal distribution. Cham: Springer International Publishing, 161–174.
- Maneerat P, Niwitpong S-A, Niwitpong S. 2019a.** Bayesian confidence intervals for a single mean and the difference between two means of delta-lognormal distributions. *Communications in Statistics - Simulation and Computation* 1–29
[DOI 10.1080/03610918.2019.1616095](https://doi.org/10.1080/03610918.2019.1616095).
- Maneerat P, Niwitpong S-A, Niwitpong S. 2019b.** Confidence intervals for the mean of delta-lognormal distribution. Cham: Springer International Publishing, 264–274
[DOI 10.1007/978-3-030-04263-9_20](https://doi.org/10.1007/978-3-030-04263-9_20).

- Maneerat P, Niwitpong S-A, Niwitpong S. 2020a.** A Bayesian approach to construct confidence intervals for comparing the rainfall dispersion in Thailand. *PeerJ* **8**:e8502 DOI [10.7717/peerj.8502](https://doi.org/10.7717/peerj.8502).
- Maneerat P, Niwitpong S-A, Niwitpong S. 2020b.** Bayesian confidence intervals for the difference between variances of delta-lognormal distributions. *Biometrical Journal* **62**:1769–1790 DOI [10.1002/bimj.201900079](https://doi.org/10.1002/bimj.201900079).
- Owen WJ, DeRouen TA. 1980.** Estimation of the mean for lognormal data containing zeroes and left-censored values, with applications to the measurement of worker exposure to air contaminants. *Biometrics* **36**(4):707–719 DOI [10.2307/2556125](https://doi.org/10.2307/2556125).
- Sangnawakij P, Niwitpong S-A. 2017a.** Confidence intervals for coefficients of variation in two-parameter exponential distributions. *Communications in Statistics - Simulation and Computation* **46**(8):6618–6630 DOI [10.1080/03610918.2016.1208236](https://doi.org/10.1080/03610918.2016.1208236).
- Sangnawakij P, Niwitpong S-A. 2017b.** Confidence intervals for functions of coefficients of variation with bounded parameter spaces in two gamma distributions. *Songklanakarin Journal of Science and Technology* **39**(1):27–39.
- Shimizu K. 1993.** A bivariate mixed lognormal distribution with an analysis of rainfall data. *Journal of Applied Meteorology* **32**(2):161–171 DOI [10.1175/1520-0450\(1993\)032<0161:ABMLDW>2.0.CO;2](https://doi.org/10.1175/1520-0450(1993)032<0161:ABMLDW>2.0.CO;2).
- Stone JV. 2013.** Bayes' Rule: a tutorial introduction to Bayesian analysis. Sheffield: Sebtel Press.
- Thai Meteorological Department. 2015.** The climate of Thailand. Available at https://www.tmd.go.th/en/archive/thailand_climate.pdf (accessed on 30 August 2020).
- Thangjai W, Niwitpong S. 2020.** Simultaneous confidence intervals for all differences of coefficients of variation of two-parameter exponential distributions. *Thailand Statistician* **18**(2):135–149.
- Thangjai W, Niwitpong S-A, Niwitpong S. 2019.** Simultaneous confidence intervals for all differences of coefficients of variation of log-normal distributions. *Hacetatepe Journal of Mathematics & Statistics* **48**(5):1505–1521 DOI [10.15672/hujms.454804](https://doi.org/10.15672/hujms.454804).
- Tian L. 2005.** Inferences on the mean of zero-inflated lognormal data: the generalized variable approach. *Statistics in Medicine* **24**(20):3223–3232 DOI [10.1002/sim.2169](https://doi.org/10.1002/sim.2169).
- Tian L, Wu J. 2006.** Confidence intervals for the mean of lognormal data with excess zeros. *Biometrical Journal. Biometrische Zeitschrift* **48**(1):149–156 DOI [10.1002/bimj.200510155](https://doi.org/10.1002/bimj.200510155).
- Weerahandi S. 1993.** Generalized confidence interval. *Journal of the American Statistical Association* **88**(423):899–905 DOI [10.1080/01621459.1993.10476355](https://doi.org/10.1080/01621459.1993.10476355).
- Wilson EB. 1927.** Probable inference, the law of succession, and statistical inference. *Journal of the American Statistical Association* **22**(158):209–212 DOI [10.1080/01621459.1927.10502953](https://doi.org/10.1080/01621459.1927.10502953).
- Wong ACM, Wu J. 2002.** Small sample asymptotic inference for the coefficient of variation: normal and nonnormal models. *Journal of Statistical Planning and Inference* **104**(1):73–82 DOI [10.1016/S0378-3758\(01\)00241-5](https://doi.org/10.1016/S0378-3758(01)00241-5).

- Wongkhao A, Niwitpong S-A, Niwitpong S. 2015.** Confidence intervals for the ratio of two independent coefficients of variation of normal distribution. *Far East Journal of Mathematical Sciences* **98(6)**:741–757 DOI [10.17654/FJMSNov2015_741_757](https://doi.org/10.17654/FJMSNov2015_741_757).
- Wu W-H, Hsieh H-N. 2014.** Generalized confidence interval estimation for the mean of delta-lognormal distribution: an application to New Zealand trawl survey data. *Journal of Applied Statistics* **41(7)**:1471–1485 DOI [10.1080/02664763.2014.881780](https://doi.org/10.1080/02664763.2014.881780).
- Yosboonruang N, Niwitpong S. 2020.** Statistical inference on the ratio of delta-lognormal coefficients of variation. *Applied Science and Engineering Progress* **8** DOI [10.14416/j.asep.2020.06.003](https://doi.org/10.14416/j.asep.2020.06.003).
- Yosboonruang N, Niwitpong S, Niwitpong S-A. 2019a.** Confidence intervals for coefficient of variation of three parameters delta-lognormal distribution. Cham: Springer International Publishing, 352–363.
- Yosboonruang N, Niwitpong S-A, Niwitpong S. 2018.** Confidence intervals for the coefficient of variation of the delta-lognormal distribution. Cham: Springer International Publishing, 327–337.
- Yosboonruang N, Niwitpong S-A, Niwitpong S. 2019b.** Measuring the dispersion of rainfall using Bayesian confidence intervals for coefficient of variation of delta-lognormal distribution: a study from Thailand. *PeerJ* **7**:e7344 DOI [10.7717/peerj.7344](https://doi.org/10.7717/peerj.7344).
- Yosboonruang N, Niwitpong S-A, Niwitpong S. 2020.** The Bayesian confidence intervals for measuring the difference between dispersions of rainfall in Thailand. *PeerJ* **8**:e9662 DOI [10.7717/peerj.9662](https://doi.org/10.7717/peerj.9662).
- Zhou XH, Tu W. 2000.** Confidence intervals for the mean of diagnostic test charge data containing zeros. *Biometrics* **56(4)**:1118–1125 DOI [10.1111/j.0006-341X.2000.01118.x](https://doi.org/10.1111/j.0006-341X.2000.01118.x).