




The Role of Neuroglobin in the Sleep-Wake Cycle

Fabio García-García¹ Mario Eduardo Acosta-Hernández¹ Luis Beltrán-Parrazal² 
 Juan Carlos Rodríguez-Alba¹

¹Instituto de Ciencias de la Salud, Departamento de Biomedicina, Universidad Veracruzana, Xalapa, Veracruz, Mexico

²Instituto de Investigaciones Cerebrales, Universidad Veracruzana, Xalapa, Veracruz, Mexico

Address for correspondence Fabio García-García, PhD, Universidad Veracruzana, Instituto de Ciencias de la Salud, Departamento de Biomedicina, Xalapa, Veracruz, Mexico (email: fgarcia@uv.mx).

Sleep Sci 2023;16(3):e362–e367.

Abstract

Neuroglobin (Ngb) is a protein expressed in the central and peripheral nervous systems of the vertebrate. The Ngb has different functions in neurons, including regulating O₂ homeostasis, oxidative stress, and as a neuroprotector after ischemia/hypoxia events. The Ngb is a hemoprotein of the globin family, structurally like myoglobin and hemoglobin. Ngb has higher expression in the cortex, hypothalamus, thalamus, brainstem, and cerebellum in mammals. Interestingly, Ngb immunoreactivity oscillates according to the sleep-wake cycle and decreases after 24 hours of sleep deprivation, suggesting that sleep homeostasis regulates Ngb expression. In addition, Ngb expresses in brain areas related to REM sleep regulation. Therefore, in the present review, we discuss the potential role of the Ngb in the sleep-wake regulation of mammals.

Keywords

- ▶ sleep deprivation
- ▶ REM sleep
- ▶ oxidative stress
- ▶ neuroprotection

Introduction

Sleep is a state of the conscience regulated by the brain. Evidence shows that different brain areas and neurotransmitters participate in sleep induction and regulation. For example, non-rapid eye movements (NREM) sleep depends on hypothalamic nuclei activity. In contrast, rapid eye movements (REM) sleep is generated by several brainstem nuclei. Both sleep stages are characterized by a specific pattern in electroencephalogram and muscle activity.

NREM sleep is induced by activating the GABAergic neurons from the ventrolateral preoptic (VLPO) and the medial preoptic (MnPO) nucleus. These GABAergic neurons increase firing activity exclusively during a NREM sleep period. Furthermore, the activity of GABAergic neurons inhibits neurons from nuclei that participate in wakefulness induction. In contrast, the orexin (ORX) released by the neurons from the lateral hypothalamus stimulates neurons from the ventral tegmental area (VTA) to release dopamine and induce wakefulness. In addition, the tuberomammillary

hypothalamic neurons also are encouraged by ORX to produce histamine release. Finally, ORX stimulates neurons from the locus coeruleus and dorsal raphe nucleus in the brainstem to release noradrenaline and serotonin. In these complex interactions between brain areas and neurotransmitters to induce sleep and wakefulness, diverse intracellular molecules are also involved.

The neural firing rates, synaptic markers, molecular pathways, and gene expression show differential up and down-regulation across brain areas and sleep stages.¹ In this sense, the neuroglobin (Ngb) is a protein expressed in specific brain regions related to the sleep-wake cycle, and its expression change after sleep deprivation. Therefore, in the present review, we discuss the potential role of the Ngb in the sleep-wake regulation of mammals.

Neuroglobin

The Ngb was described in mammals' brains in 2000.² It has been reported widely in vertebrates from humans, mice, and

received
 May 30, 2022
 accepted
 November 23, 2022

DOI <https://doi.org/10.1055/s-0043-1772806>.
 ISSN 1984-0659.

© 2023. Brazilian Sleep Association. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

rats to blowfish and zebrafish, suggesting that Ngb protein is highly conserved through evolution.³⁻⁵ The Ngb is a monomeric globular hemoprotein of about 150 amino acids with a relative molecular mass of 17 kDa. A single copy gene codifies Ngb in all species, and the coding region in the mammalian Ngb gene carries three introns located at B12.2, E11.0, and G7.0. Despite the introns B12.2 and G7.0 being present in other globin genes, the intron E11.0 is unique to Ngb.^{6,7} Furthermore, all vertebrate contains this intron in a conserved position within the coding regions except zebrafish, including a fourth intron.⁶ The human Ngb gene is located on the long arm of chromosome 14 at position q24.3.⁶ Although, Ngb shares about 20% to 25% sequence identity compared to the homologous globins hemoglobin and myoglobin,^{3,7} despite the difference, Ngb adopts a 3D folding identical to the rest of the globins, which is the main characteristic of the family (►Figure 1).

For example, hemoglobin and myoglobin have six coordination bonds for six ligands; four are nitrogen atoms part of the porphyrin molecule. The fifth bonds bind to the porphyrin with the histidine's nitrogen atom at position 93, while the sixth bond with O₂. In contrast, Ngb forms six bonds, but all of them with O₂.⁸ The occupancy of the six-coordinating bonds implies that exogenous ligands must displace the distal histidine to bond to the prosthetic group. Therefore, the mechanism of exogenous ligand binding to Ngb is more

complex than the other globins.⁸ Thus, Ngb may enhance O₂ to high metabolic neurons and may also be involved in detoxifying reactive oxygen species.⁹

Neuroglobin Functions

Ngb gene and protein are highly conserved throughout evolution.¹⁰ A single copy gene codifies Ngb protein in all species except trout. In the last decade, research on the role that Ngb plays in the brain has intensified. Most studies have been focused on understanding the potential chemical reactions that Ngb regulates and its role as an O₂ transporter in neurons and glial cells.

Experimental evidence indicates that Ngb plays multiple roles in health and disease.^{6,11,12} For example, it has been documented that Ngb is expressed at concentrations between 1-100 μM exclusively in the brain, both in neurons and astrocytes and in Müller cells^{13,14}; however, Ngb concentration is significantly increased in Müller cells after an ocular injury.¹² Furthermore, biochemical studies support that Ngb can facilitate oxygen transport only if it is at a high concentration. In contrast, low concentrations of Ngb (~1 μM) probably participate only in enzymatic activities and cell signaling.^{6,12,15} Furthermore, it has been suggested that high concentrations of Ngb in tissues such as the retina prevent hypoxia and increase O₂ uptake by up to 40%.¹⁶

A





Fig. 1 Crystal structure of murine Neuroglobin (Ngb)⁵² and photomicrographic shows basal immunoreactivity to Ngb in the rat's anteroventral periventricular nucleus (avPe). Panel A. Ngb is a hexacoordinated protein that shares 20 to 25% of the amino acid structure with the rest of all globins. Ngb adopts a 3D-folding like other globins, eight alpha-subunits named A-H. Have no external ligands; therefore, any gaseous ligand must compete for binding to the heme iron. Panel B. Coronal section of the avPe illustrates the bilateral immunoreactivity of Ngb cells (4x). Panel C. The density of Ngb positive cells is higher in this brain region (10x). Panel D. Amplification shows the Ngb immunoreactivity in all somas, including the nucleus (100x). 3V third ventricle. Bregma – 0.12 mm. ac anterior commissure, och optic chiasm.

A significant increase in the Ngb immunoreactivity is reported in the cerebellum of animals exposed to moderate concentrations of CO₂ in vivo.¹⁷ In addition, it has been shown that an increase in the expression levels of Ngb in the eye can prevent hypoxia caused by induced ocular hypertension in rats.¹⁸ This result suggests that Ngb could be essential in metabolic adaptation to hypoxic events. Moreover, a decrease in Ngb expression decreases the number and viability of cortical neurons. At the same time, the overexpression of Ngb in the same neuronal type increases the viability of neurons in an in vitro hypoxia model.¹⁹

Recently, it has been suggested that Ngb regulates the levels of inflammatory cytokines such as Interleukin-6 (IL-6), Tumor Necrosis Factor-alpha (TNF α), Interleukin 1-beta (IL-1B), and Vascular Endothelial Growth Factor (VEGF) and the microglia activation during a hypoxic environment.²⁰ This effect seems to be related to the regulatory impact of Ngb on the Wnt/ β -catenin and Nuclear Factor kappa-light-chain-enhancer of activated B cells (NF- κ B) signaling pathways, which induce the proteasomal degradation of Disheveled-1.²¹

Although most of the details of the mechanisms through which Ngb exerts its neuroprotective activity are unknown, colocalization studies in cells of the nervous system suggest a direct interaction between Ngb and mitochondria. It has been suggested that Ngb could participate in regulating energy metabolism and apoptosis mechanisms through its interaction with ferric cytochrome c. This antiapoptotic function of ferric cytochrome c requires the participation of Ngb-reducing activity on ferric cytochrome C, which prevents the activation of caspase-9.^{22,23}

In the last decade, it has been described that some natural and synthetic compounds with small chemical structures can upregulate the expression of the Ngb due to their ability to cross the blood-brain barrier. For example, in hypoxic cerebral neurons, boosted Ngb expression was observed in cultures treated for 24 h with the iron chelator deferoxamine (Dfx). This effect could be due to Dfx enhanced expression of hypoxia-inducible genes, which are closely related to Ngb expression.^{19,24} Besides, valproic and cinnamic acids showed induction of Ngb in vitro. In this sense, valproic acid prompts Ngb expression, like Dfx, whereas cinnamic acid has a more significant effect. However, other short-fatty acids tested did not significantly impact Ngb expression, suggesting that chemical structures and functional groups have a crucial role in this biological effect.²⁵

Furthermore, hormones can also influence the upregulation of Ngb. For instance, in the SK-N-BE human neuroblastoma cell line and mouse hippocampal primary neurons, the treatment with 17 β -estradiol (E₂) induces an increase in the Ngb protein levels in a time- and dose-dependent manner. Nevertheless, male steroid hormones did not affect Ngb protein expression. Interestingly, polyphenol compounds with estrogen-like properties such as daidzein, genistein, polydatin, biochanin A, and fucosterol provided promising results to induce Ngb upregulation.^{26,27} Thus, this evidence suggests exploring new therapeutic approaches for neurodegenerative disorders by establishing in vivo effective doses and adequate treatment periods to prevent adverse effects.

In summary, there is a considerable volume of experimental evidence indicating that exposure to natural or synthetic molecules, hormones or CO₂, the induction of oxidative stress, hypoxia, epilepsy, and ischemia, and even sleep deprivation cause an increase in Ngb levels in the brain and cerebellum that potentially allows it to carry out neuroprotective functions.²⁸

The Role of the Neuroglobin in Sleep-Wake Cycle Regulation

The Ngb is expressed in brain nuclei that participate in the sleep-wake cycle. For example, the locus coeruleus nucleus (LC), pedunculopontine tegmental nucleus (PPTg), laterodorsal tegmental nucleus (LDTg), and preoptic area (POA) are brain regions with specific Ngb expression.²⁹ Also, the neurons from these nuclei modify their firing frequency patterns across the sleep-wake cycle.³⁰ Recently, we reported reducing the number of Ngb-positive cells in the PPTg, LDTg, and periventricular area (Pe) in rats after sleep deprivation for 24 hours.³¹ The corticosterone plasma levels in sleep-deprived rats did not increase compared to non-sleep-deprived rats, suggesting that stress was not responsible for the decrease in Ngb immunoreactivity.³¹ Therefore, sleep loss could significantly reduce the Ngb cell-positive cells; perhaps the protein expression is sensitive to disruptions in sleep homeostasis.

In addition, as Ngb exerts antioxidant effects,³² we evaluated whether sleep deprivation produces oxidative stress in brain tissue since sleep loss induces oxidative stress. However, the results showed that sleep deprivation did not cause oxidative stress in the brain tissue. The above was measured by generating malondialdehyde, a metabolite produced by membrane lipid peroxidation.³¹ Altogether, the results support the hypothesis that the expression of Ngb depends on sleep homeostasis because sleep deprivation reduces its immunoreactivity independent of physiological and oxidative stress.

A recent report shows that 6 hours of sleep rebound after 24 hours of sleep deprivation were sufficient to restore the number of Ngb-positive cells in the PPTg and LDTg nuclei. These results suggest that sleep recovery after sleep deprivation promotes Ngb expression.³³ Furthermore, it is well documented that the total time spent in NREM and REM during the initial 6 hours of sleep recovery is increased. Particularly, REM sleep time is increased almost twice after 24-h of sleep deprivation. Moreover, it is reported that EEG slow-wave activity (SWA; mean power density 0.75-4.0 Hz) during NREM sleep is elevated relative to a non-sleep-deprived condition, and the number of brief awakenings is reduced.^{34,35} Therefore, it is crucial to characterize whether sleep time after sleep deprivation is associated with Ngb expression.

In addition, in the preliminary study, our group found that the intracerebral administration of the ORX (10 μ g/5 μ L, Human orexin, Cat. 06012, Sigma Aldrich, St Louis, MO, USA) in rats results in a significant increase in the total number of the Ngb positive cells in MnPO, Perifornical area, PPTg, and LC. These brain nuclei receive projections

from the lateral hypothalamus and express orexin receptors.³⁶ ORX is a peptide synthesized by neurons located through the medial-lateral extent of the hypothalamus and responsible for inducing wakefulness.³⁶ Furthermore, prolonged wakefulness causes cellular damage.^{37,38} This cellular damage is related to neurodegenerative disease development.³⁹ In this sense, *in vitro* studies suggest that cytosolic Ngb could react with mitochondrial cytochrome c, interfering with apoptotic pathways to prevent cellular damage.^{23,28,40} Therefore, Ngb could participate as an endogenous neuroprotector in situations of sleep loss and prolonged shift work. For example, patients with obstructive sleep apnea (OSA) and that who suffer sleep fragmentation during the night report higher serum levels of Ngb, suggesting that Ngb has a protective role in conditions of low oxygen levels.⁴¹

In addition, lower Ngb expression after orexin administration was observed in the LDTg nucleus. Several studies in cats and rats show that the administration of ORX into the pons, particularly in the LDTg nucleus, promotes wakefulness and suppresses REM sleep.^{42,43} Furthermore, an optogenetic study reported that activating cholinergic neurons in the PPTg during the NREM sleep period induces REM sleep. However, this effect was less evident when stimulated LTDg neurons.⁴⁴ Together these results suggest that PPTg could participate more actively than the LDTg nucleus in controlling the onset of REM sleep. These results also support the hypothesis that Ngb is increased to mediate the metabolic rate induced by sustained neuronal activity (► **Figure 2**).

As aforementioned, sleep loss associated with OSA disorder increase Ngb serum levels. The OSA is a condition that generates intermittent hypoxia (IH), sleep disruption,

and increased apoptotic processes due to oxidative stress. Interestingly, plasma levels of Ngb increase in proportion to the severity of the OSA.⁴¹ Furthermore, Ngb over-expression in IH animal models protects against lipid peroxidation.^{45,46} In a murine model of ischemia-reperfusion, Ngb positive cells notably increased 24 h post-reperfusion in the cerebral cortex and remained higher than controls 48 h after ischemia-reperfusion.⁴⁶ These findings provide evidence for the neuroprotective role of Ngb during cellular stress.

Sleep is also associated with neuroprotective functions since sleep loss is a risk factor for developing Alzheimer's and other neurological and psychiatric disorders. Furthermore, Ngb is also related to neurodegenerative diseases such as Alzheimer's.⁴⁷⁻⁴⁹ In this sense, it is plausible that Ngb acts as the first line of defense in developing neurodegenerative pathologies. Therefore, even when it sounds logical, it is necessary to investigate whether chronic sleep restriction may affect Ngb expression and if the reduction of the Ngb induces the neurodegenerative disease's progress.

Ngb is highly expressed in neurons of the suprachiasmatic nucleus (SCN) in the hypothalamus that serves as the master clock in the mammal's brain. SCN neurons co-express Ngb and PER-1, a clock gene involved in the circadian regulation and maintenance of SCN activity.⁵⁰ SCN receives a monosynaptic input from the retina, where Ngb is highly expressed. Interestingly, Ngb mRNA expression in SCN neurons of mice increases during the light period; meanwhile, during the dark period decreases, showing an evident circadian pattern.⁵¹ Then, Ngb-deficient mice disrupt the SCN light responses and increase the expression of PER1 after a light stimulus during the dark period.⁵¹ It also correlated with




Fig. 2 Neuronal pathways that regulate sleep. The ventrolateral preoptic nucleus (VLPO) and median preoptic nucleus (MnPO) send inhibitory projections to the brain nucleus in the hypothalamus and brainstem involved in wake to induce sleep. The activation of the neurons hypothetically induces neuroglobin (Ngb) expression in the nuclei related to sleep. Ngb can participate in the control of metabolic rate induced by oxidative stress. Abbreviations: TMN, Tuberomammillary nucleus; VTA, ventral tegmental area; DRn, dorsal raphe nucleus; PBn, parabrachial nucleus; PZ, parafacial zone.

preliminary results of our laboratory that show an increase in the number of Ngb-positive cells during sleep (light phase) and a decrease during the wake phase (dark phase). Since sleep is regulated in a circadian fashion directly from SCN, it could be one of the main signals for Ngb synthesis and regulation during sleep.

Conclusions

The evidence discussed in the present review suggests that Ngb is susceptible to response to sleep loss and sleep rebound restored. Potentially, Ngb could participate as an endogenous neuroprotector in situations of sleep loss and prolonged shift work. These results also support the hypothesis that Ngb is increased to mediate the metabolic rate induced by sustained neuronal activity. However, more studies are necessary to understand the role of the Ngb in sleep and other behaviors and the consequences of its deficiencies.

Financial Disclosure

None declared.

Non-financial Disclosure

None declared.

Conflict of Interest Statement

None declared.

References

- Navarro-Lobato I, Genzel L. The up and down of sleep: From molecules to electrophysiology. *Neurobiol Learn Mem* 2019; 160:3–10
- Burmester T, Weich B, Reinhardt S, Hankeln T. A vertebrate globin expressed in the brain. *Nature* 2000;407(6803):520–523
- Xie LK, Yang SH. Brain globins in physiology and pathology. *Med Gas Res* 2016;6(03):154–163
- Roesner A, Fuchs C, Hankeln T, Burmester T. A globin gene of ancient evolutionary origin in lower vertebrates: evidence for two distinct globin families in animals. *Mol Biol Evol* 2005;22(01):12–20
- Qiu XY, Chen XQ. Neuroglobin - recent developments. *Biomol Concepts* 2014;5(03):195–208
- Ascenzi P, di Masi A, Leboffe L, et al. Neuroglobin: From structure to function in health and disease. *Mol Aspects Med* 2016;52:1–48
- Burmester T, Hankeln T. Neuroglobin: a respiratory protein of the nervous system. *News Physiol Sci* 2004;19(03):110–113
- Pesce A, Dewilde S, Nardini M, et al. The human brain hexacoordinated neuroglobin three-dimensional structure. *Micron* 2004; 35(1-2):63–65
- Alekseeva OS, Grigor'ev IP, Korzhhevskii DE. Neuroglobin, an oxygen-binding protein in the mammalian nervous system (localization and putative functions). *J Evol Biochem Physiol* 2017; 53:249–258
- Wystub S, Laufs T, Schmidt M, et al. Localization of neuroglobin protein in the mouse brain. *Neurosci Lett* 2003;346(1-2):114–116
- Chen XQ, Qin LY, Zhang CG, et al. Presence of neuroglobin in cultured astrocytes. *Glia* 2005;50(02):182–186
- Lechauve C, Augustin S, Roussel D, Sahel J-A, Corral-Debrinski M. Neuroglobin involvement in visual pathways through the optic nerve. *Biochim Biophys Acta* 2013;1834(09):1772–1778
- Brunori M, Vallone B. A globin for the brain. *FASEB J* 2006;20(13): 2192–2197
- Schmidt M, Giessl A, Laufs T, Hankeln T, Wolfrum U, Burmester T. How does the eye breathe? Evidence for neuroglobin-mediated oxygen supply in the mammalian retina. *J Biol Chem* 2003;278 (03):1932–1935
- Cwerman-Thibault H, Lechauve C, Augustin S, et al. Neuroglobin Can Prevent or Reverse Glaucomatous Progression in DBA/2J Mice. *Mol Ther Methods Clin Dev* 2017;5:200–220
- Roberts PA, Gaffney EA, Luthert PJ, Foss AJE, Byrne HM. Retinal oxygen distribution and the role of neuroglobin. *J Math Biol* 2016; 73(01):1–38
- Beltran-Parral L, Acuna D, Ngan AM, et al. Neuroglobin, cytochrome c, and transcriptional profiling of hypoxia-related genes in the rat cerebellum after prenatal chronic very mild carbon monoxide exposure (25 ppm). *Brain Res* 2010;1330:61–71
- Shi S-Y, Feng X-M, Li Y, Li X, Chen X-L. Expression of neuroglobin in ocular hypertension induced acute hypoxic-ischemic retinal injury in rats. *Int J Ophthalmol* 2011;4(04):393–395
- Sun Y, Jin K, Mao XO, Zhu Y, Greenberg DA. Neuroglobin is up-regulated by and protects neurons from hypoxic-ischemic injury. *Proc Natl Acad Sci U S A* 2001;98(26):15306–15311
- Tun SBB, Barathi VA, Luu CD, Lynn MN, Chan ASY. Effects of Exogenous Neuroglobin (Ngb) on retinal inflammatory chemokines and microglia in a rat model of transient hypoxia. *Sci Rep* 2019;9(01):18799
- Xun Y, Li Z, Tang Y, et al. Neuroglobin Regulates Wnt/ β -Catenin and NF κ B Signaling Pathway through Dvl1. *Int J Mol Sci* 2018;19 (07):2133
- Bønding SH, Henty K, Dingley AJ, Brittain T. The binding of cytochrome c to neuroglobin: a docking and surface plasmon resonance study. *Int J Biol Macromol* 2008;43(03):295–299
- Fago A, Mathews AJ, Moens L, Dewilde S, Brittain T. The reaction of neuroglobin with potential redox protein partners cytochrome b5 and cytochrome c. *FEBS Lett* 2006;580(20):4884–4888
- Ciccone L, Nencetti S, Socci S, Orlandini E. Neuroglobin and neuroprotection: the role of natural and synthetic compounds in neuroglobin pharmacological induction. *Neural Regen Res* 2021;16(12):2353–2358
- Jin K, Mao XO, Xie L, John V, Greenberg DA. Pharmacological induction of neuroglobin expression. *Pharmacology* 2011;87(1-2):81–84
- De Marinis E, Ascenzi P, Pellegrini M, et al. 17 β -estradiol—a new modulator of neuroglobin levels in neurons: role in neuroprotection against H₂O₂-induced toxicity. *Neurosignals* 2010;18(04): 223–235
- Barreto GE, McGovern AJ, Garcia-Segura LM. Role of Neuroglobin in the Neuroprotective Actions of Estradiol and Estrogenic Compounds. *Cells* 2021;10(08):1907
- De Simone G, Sbardella D, Oddone F, Pesce A, Coletta M, Ascenzi P. Structural and (Pseudo-)Enzymatic Properties of Neuroglobin: Its Possible Role in Neuroprotection. *Cells* 2021;10(12): 3366
- Hundahl CA, Allen GC, Nyengaard JR, et al. Neuroglobin in the rat brain: localization. *Neuroendocrinology* 2008;88(03):173–182
- Szymusiak R, Alam N, McGinty D. Discharge patterns of neurons in cholinergic regions of the basal forebrain during waking and sleep. *Behav Brain Res* 2000;115(02):171–182
- Melgarejo-Gutiérrez M, Acosta-Peña E, Venebra-Muñoz A, Escobar C, Santiago-García J, García-García F. Sleep deprivation reduces neuroglobin immunoreactivity in the rat brain. *Neuroreport* 2013;24(03):120–125
- Amri F, Ghouili I, Amri M, Carrier A, Masmoudi-Kouki O. Neuroglobin protects astroglial cells from hydrogen peroxide-induced oxidative stress and apoptotic cell death. *J Neurochem* 2017;140 (01):151–169
- Melgarejo-Gutiérrez M, García-García F, Hernández-Márquez G, Morgado-Valle C, Acosta-Hernández ME, Rodríguez-Alba JC. Sleep Recovery Restored Neuroglobin Immunoreactivity in Rat LDTg-PPTg Nuclei. *Sleep Disord* 2020;2020:8353854

- 34 Lancel M, Kerkhof GA. Effects of repeated sleep deprivation in the dark- or light-period on sleep in rats. *Physiol Behav* 1989;45(02): 289–297
- 35 Franken P, Dijk DJ, Tobler I, Borbély AA. Sleep deprivation in rats: effects on EEG power spectra, vigilance states, and cortical temperature. *Am J Physiol* 1991;261(1 Pt 2):R198–R208
- 36 Peyron C, Tighe DK, van den Pol AN, et al. Neurons containing hypocretin (orexin) project to multiple neuronal systems. *J Neurosci* 1998;18(23):9996–10015
- 37 Ramanathan L, Gulyani S, Nienhuis R, Siegel JM. Sleep deprivation decreases superoxide dismutase activity in rat hippocampus and brainstem. *Neuroreport* 2002;13(11):1387–1390
- 38 de Souza L, Smaili SS, Ureshino RP, et al. Effect of chronic sleep restriction and aging on calcium signaling and apoptosis in the hippocampus of young and aged animals. *Prog Neuropsychopharmacol Biol Psychiatry* 2012;39(01):23–30
- 39 Qiu H, Zhong R, Liu H, Zhang F, Li S, Le W. Chronic Sleep Deprivation Exacerbates Learning-Memory Disability and Alzheimer's Disease-Like Pathologies in A β PP(swe)/PS1(Δ E9) Mice. *J Alzheimers Dis* 2016;50(03):669–685
- 40 Flocchetti M, De Marinis E, Ascenzi P, Marino M. Neuroglobin and neuronal cell survival. *Biochim Biophys Acta* 2013;1834(09): 1744–1749
- 41 Xu M, Yang Y, Zhang J. Levels of neuroglobin in serum and neurocognitive impairments in Chinese patients with obstructive sleep apnea. *Sleep Breath* 2013;17(02):573–582
- 42 Xi M-C, Morales FR, Chase MH. Effects on sleep and wakefulness of the injection of hypocretin-1 (orexin-A) into the laterodorsal tegmental nucleus of the cat. *Brain Res* 2001; 901(1-2):259–264
- 43 de Lecea L, Sutcliffe JG. The hypocretins and sleep. *FEBS J* 2005; 272(22):5675–5688
- 44 Van Dort CJ, Zachs DP, Kenny JD, et al. Optogenetic activation of cholinergic neurons in the PPT or LDT induces REM sleep. *Proc Natl Acad Sci U S A* 2015;112(02):584–589
- 45 Haines B, Demaria M, Mao X, et al. Hypoxia-inducible factor-1 and neuroglobin expression. *Neurosci Lett* 2012;514(02):137–140
- 46 Shang A, Zhou D, Wang L, et al. Increased neuroglobin levels in the cerebral cortex and serum after ischemia-reperfusion insults. *Brain Res* 2006;1078(01):219–226
- 47 Chen L-M, Xiong Y-S, Kong F-L, et al. Neuroglobin attenuates Alzheimer-like tau hyperphosphorylation by activating Akt signaling. *J Neurochem* 2012;120(01):157–164
- 48 Sun F, Mao X, Xie L, Greenberg DA, Jin K. Neuroglobin protein is upregulated in Alzheimer's disease. *J Alzheimers Dis* 2013;36 (04):659–663
- 49 Szymanski M, Wang R, Fallin MD, Bassett SS, Avramopoulos D. Neuroglobin and Alzheimer's dementia: genetic association and gene expression changes. *Neurobiol Aging* 2010;31(11):1835–1842
- 50 Hundahl CA, Hannibal J, Fahrenkrug J, Dewilde S, Hay-Schmidt A. Neuroglobin expression in the rat suprachiasmatic nucleus: colocalization, innervation, and response to light. *J Comp Neurol* 2010;518(09):1556–1569
- 51 Hundahl CA, Fahrenkrug J, Hay-Schmidt A, Georg B, Faltoft B, Hannibal J. Circadian behaviour in neuroglobin deficient mice. *PLoS One* 2012;7(04):e34462
- 52 Avella G, Ardiccioni C, Scaglione A, et al. Engineering the internal cavity of neuroglobin demonstrates the role of the haem-sliding mechanism. *Acta Crystallogr D Biol Crystallogr* 2014;70(Pt 6): 1640–1648