

Ramadan Fasting and Risk of Covid-19

Abstract

Almost all religions recommend periods of fasting. Many adult Muslims fast during the holy month of Ramadan each year. Ramadan fasting as a type of intermittent fasting is a non-pharmacological intervention refining the overall health. This year, Ramadan is coincided with coronavirus disease 2019 (COVID-19) outbreak making it one of the most challenging fasting periods for Muslims in the world. There is no solid direct evidence to suggest any adverse effect of Ramadan fasting during the COVID-19 pandemic in healthy individuals. However, there are exemptions in Ramadan Fasting and those at risk of health issues should not fast. COVID-19 is a new disease and there is limited studies concerning its risk factors. The purpose of this review was shedding more light on the potential mechanisms involved in influence of practice of fasting in all forms, including Ramadan fasting on the vulnerability to infection.

Keywords: COVID-19, pandemic, Ramadan fasting

Introduction

Muslims fast from sunrise to sunset during the month of Ramadan is a compulsory act for all healthy Muslim adults. While fasting is an important part of Ramadan, it is also a time of self-reflection and a wonderful opportunity to quit harmful habits for Muslims.

Ramadan is the 9th lunar month migrates throughout the seasons. The 2020 Ramadan coincides with the coronavirus disease 2019 (COVID-19) outbreak, making it one of the most challenging fasting periods for Muslims in the world.

In recent years, the effect of intermittent fasting (IF) as a way of calorie restriction (CR) on health outcome has been considered in many studies. Indeed a lessening in caloric intake without malnutrition, has frequently been found to improve health in a variety of species,^[1] including non-human primates.^[2]

In medical terms, Ramadan fasting (RF) can be considered a unique model of IF with periods of food, drink and smoking abstinence daily for 1 month. Knowledge about the effects of Ramadan fasting on the viral respiratory disease is poor, mainly because of a scarcity of scientific research in this area. So we focus on the

association of Ramadan fasting/IF and potential preventive/susceptive mechanisms on viral disease. IF influences several body systems, including the immune system, and metabolic state which plays a critical role in regulating and maintaining body response to pathogens. The pandemics caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) raised growing interest concerning the effect of RF on COVID-19.^[3] The purpose of this review was shedding more light on the potential mechanisms involved in influence of Ramadan fasting on the vulnerability to infection.

Fasting and Metabolic Changes

Fasting is associated with shifting the primary energy source of the body from glucose to free fatty acid (FFA). In a fed human, glucose is used for energy and fat is stored in adipose tissue as triglycerides. During fasting, the liver converts fatty acids derived from triglycerides to ketone bodies to supply new source of energy for many tissues.^[4] This states are associated with altered immune cell function in the body.^[5] In a fast human, ketone bodies levels increase within 8 to 12 hours after the beginning of fasting and could reach to 6-8 mm after prolonged fasting.^[6,7]

The brain relies on the ketone bodies β -hydroxybutyrate and acetoacetate as fuel, besides glucose, in fasting.^[8]

How to cite this article: Javanmard SH, Otraj Z. Ramadan fasting and risk of Covid-19. *Int J Prev Med* 2020;11:60.

Shaghayegh Haghjooy Javanmard, Zahra Otraj¹

Applied Physiology Research Center, Cardiovascular Research Institute, Isfahan University of Medical Science, Isfahan, Iran, ¹Vice-Chancellery for Research and Technology, Isfahan University of Medical Sciences, Isfahan, Iran

Address for correspondence:

Dr. Shaghayegh Haghjooy Javanmard, Applied Physiology Research Center, Cardiovascular Research Institute, Isfahan University of Medical Science, Isfahan, Iran. E-mail: sh_haghjoo@med.mui.ac.ir

Access this article online

Website:
www.ijpvmjournal.net/www.ijpm.ir

DOI:
10.4103/ijpvm.IJPVM_236_20

Quick Response Code:



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

β -hydroxybutyrate can inhibit innate immune responses, sparing ATP for the functioning of ketone-dependent organs such as the brain and heart.

Ketone bodies are effective signaling molecules acting on several cell and organ functions.^[9] They have major effects on body metabolic state through peroxisome proliferator-activated receptor γ coactivator 1 α (PGC-1 α), poly (adenosine diphosphate [ADP]-ribose) polymerase 1 (PARP1), and ADP ribosyl cyclase pathways.^[10]

Fasting also has significant effect on metabolic hormones concentration and function including insulin, IGF-I, adiponectin, and ghrelin. It also has a robust role in tissue homeostasis by suppressing cell growth, and enhancing apoptosis of damaged cells.^[11]

IF also arise adaptive stress response that leads to increased expression of several defenses mechanisms including DNA repair, protein quality control, autophagy, oxidative stress response pathways

Fasting and Respiratory System Function

Several studies reported no significant change in the lung function parameters during Ramadan as compared to the pre-Ramadan period. However, some groups reported significant increase of vital capacity and mean/peak expiratory flow rate it seems that RF did not affect expiratory flow rates in healthy subjects. FVC was decreased significantly in the post-Ramadan period compared to Ramadan.

Poor hydration and inadequate water intake is one of the concerns raised around susceptibility to COVID-19 infection by Ramadan fasting. There is no evidence for association of RF with severe water deprivation and negative water balance in healthy subjects.^[12] It has been shown that total body water remained within the normal range of 30-46 kg.^[13] In addition, although fasting causes some changes in urinary volume and metabolites concentration, there is no significant change in maximum urinary concentration.^[14]

The adverse effects of dehydration on mucociliary function as a key defense mechanisms against infection is previously known.^[15] Indeed, RF research denies the harmful effect of fasting on mucociliary clearance. In a study conducted to evaluate the difference in a mucociliary clearance among volunteers who underwent RF, there was no significant difference in the mucociliary clearance time between the RF and control.^[16]

Fasting and Immune System

Several studies highlighted the mild inflammatory state associated with obesity and metabolic syndrome. RF have been shown to significantly damp down the inflammation of the body by suppressing pro-inflammatory cytokine expression. Indeed, intermittent fasting modulates the levels

of visceral fat and several additional adipokines, including leptin, IL-6, TNF- α and IGF-1.^[17,18]

Since, leptin raises neutrophil chemotaxis and macrophage phagocytosis, and triggers the maturation and function of T-cells, some changes of immune functions during fasting could be in part the result of a drop in the plasma leptin.^[19,20]

The number of total leukocytes, granulocytes, lymphocytes and monocytes were significantly decreased during Ramadan in comparison to pre-Ramadan in some reports,^[17] while in infected patients RF was associated with increasing the numbers of macrophages. It has been shown that fasting could activate a regenerative switch in stem cells, leading to the generation of new immune cells.^[21]

Interestingly, in Muslim fasting HIV patients, the change of twice-daily dose of antivirals to once-daily dosed had no significant changes on CD4 cell counts, viral load, or disease state in comparison with the twice-daily dose therapy for non-fasting patients.^[22]

The results of a comprehensive systematic review showed that fasting during Ramadan was associated with mild transient changes in the immune system, which return to the basal pre-Ramadan status shortly afterward.^[23]

Fasting also has some effects on pathogen defense mechanisms and adaptive/innate immune responses. Since immune cells also rely on glucose to sustain their function and proliferation reduced glucose availability as well as hypoinsulinemia during fasting may have negative effect on immune defense mechanisms. However, their function can be restored after re-feeding.^[24]

On the other hand, it has been shown that innate immunity response for intracellular infection factors during RF was not decreased. In fact, it increased, indicating the beneficial effect of fasting to protect against bacterial infection.^[23]

Conclusion

In conclusion, scientific evidence indicates that all types of IF including RF has several health benefits in healthy people without malnutrition. In addition, RF might not be harmful for many patients with controlled and stable disease.^[25] However, undernourishment could be dangerous in acute viral diseases.^[26] We couldn't find any evidence of adverse effect of metabolic changes associated with RF on the immune and respiratory system related to COVID-19.

Although Ramadan fasting is safe for all healthy individuals, the Qur'an exempts ill patients and those at risk of various diseases from fasting. COVID-19 is a novel disease and there is not enough information about its risk factors. However, older people (more than 65 years) and those who have serious comorbidity such as cardiovascular disease, diabetes, cancer and chronic lung and kidney

disease might be at higher risk for severe illness from COVID-19.^[27]

It has been shown that Muslims aware of their religious exemption from fasting decide not to take that exemption and fast.^[28] In respect of patient-centered care, it seems that risk assessment, organized education and choice of appropriate medication can reduce the possible hazards associated with fasting among at risk people.

Stress of RF could lead to a transient increased level of cortisol and potentially prolonged the viral shedding duration in COVID-19 patients,^[29,30] so national health authorities should be considered the regulations and advice regarding physical distancing and other public health measures related to COVID-19 during Ramadan.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 02 May 2020 **Accepted:** 02 May 2020

Published: 25 May 20

References

- Anton SD, Moehl K, Donahoo WT, Marosi K, Lee SA, Mainous AG, *et al.* Flipping the metabolic switch: Understanding and applying the health benefits of fasting. *Obesity* (Silver Spring) 2018;26:254-68.
- Colman RJ, Anderson RM, Johnson SC, Kastman EK, Kosmatka KJ, Beasley TM, *et al.* Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science* 2009;325:201-4.
- Paraskevis D, Kostaki EG, Magiorkinis G, Panayiotakopoulos G, Sourvinos G, Tsiodras S. Full-genome evolutionary analysis of the novel corona virus (2019-nCoV) rejects the hypothesis of emergence as a result of a recent recombination event. *Infect Genet Evol* 2020;79:104212.
- de Cabo R, Mattson MP. Effects of intermittent fasting on health, aging, and disease. *N Engl J Med* 2019;381:2541-51.
- Youm YH, Nguyen KY, Grant RW, Goldberg EL, Bodogai M, Kim D, *et al.* The ketone metabolite beta-hydroxybutyrate blocks NLRP3 inflammasome-mediated inflammatory disease. *Nat Med* 2015;21:263-9.
- Cahill GF Jr. Starvation in man. *N Engl J Med* 1970;282:668-75.
- Newman JC, Verdin E. Ketone bodies as signaling metabolites. *Trends Endocrinol Metab* 2014;25:42-52.
- Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G, *et al.* The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: A randomized trial in young overweight women. *Int J Obes (Lond)* 2011;35:714-27.
- Newman JC, Verdin E. β -Hydroxybutyrate: A signaling metabolite. *Annu Rev Nutr* 2017;37:51-76.
- Puchalska P, Crawford PA. Multi-dimensional roles of ketone bodies in fuel metabolism, signaling, and therapeutics. *Cell Metab* 2017;25:262-84.
- Longo VD, Mattson MP. Fasting: Molecular mechanisms and clinical applications. *Cell Metab* 2014;19:181-92.
- Leiper JB, Molla AM, Molla AM. Effects on health of fluid restriction during fasting in Ramadan. *Eur J Clin Nutr* 2003;57(Suppl 2):S30-8.
- Chumlea WC, Guo SS, Zeller CM, Reo NV, Siervogel RM. Total body water data for white adults 18 to 64 years of age: The Fels Longitudinal Study. *Kidney Int* 1999;56:244-52.
- Shirreffs SM. Markers of hydration status. *Eur J Clin Nutr* 2003;57(Suppl 2):S6-9.
- Lewis BW, Patial S, Saini Y. Immunopathology of airway surface liquid dehydration disease. *J Immunol Res* 2019;2019:2180409.
- Develioglu ON, Sirazi S, Topak M, Purisa S, Kulekci M. Differences in mucociliary activity of volunteers undergoing Ramadan versus Nineveh fasting. *Eur Arch Otorhinolaryngol* 2013;270:1655-9.
- Faris MA, Kacimi S, Al-Kurd RA, Fararjeh MA, Bustanji YK, Mohammad MK, *et al.* Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects. *Nutr Res* 2012;32:947-55.
- Davies AR, Efthimiou E. Curing type 2 diabetes mellitus with bariatric surgery - Reality or delusion? *Br J Diabetes Vasc Dis* 2012;12:173-6.
- La Cava A, Matarese G. The weight of leptin in immunity. *Nat Rev Immunol* 2004;4:371-9.
- Xu DL, Wang DH. Fasting suppresses T cell-mediated immunity in female Mongolian gerbils (*Meriones unguiculatus*). *Comp Biochem Physiol A Mol Integr Physiol* 2010;155:25-33.
- Mihaylova MM, Cheng CW, Cao AQ, Tripathi S, Mana MD, Bauer-Rowe KE, *et al.* Fasting activates fatty acid oxidation to enhance intestinal stem cell function during homeostasis and aging. *Cell Stem Cell* 2018;22:769-78.e4.
- Yakasai AM, Muhammad H, Babashani M, Jumare J, Abdulmumini M, Habib AG. Once-daily antiretroviral therapy among treatment-experienced Muslim patients fasting for the month of Ramadan. *Trop Doct* 2011;41:233-5.
- Adawi M, Watad A, Brown S, Aazza K, Aazza H, Zouhir M, *et al.* Ramadan fasting exerts immunomodulatory effects: Insights from a systematic review. *Front Immunol* 2017;8:1144.
- Walrand S, Moreau K, Caldefie F, Tridon A, Chassagne J, Portefaix G, *et al.* Specific and nonspecific immune responses to fasting and refeeding differ in healthy young adult and elderly persons. *Am J Clin Nutr* 2001;74:670-8.
- Bragazzi NL, Briki W, Khabbache H, Rammouz I, Mnadla S, Demaj T, *et al.* Ramadan fasting and infectious diseases: A systematic review. *J Infect Dev Ctries* 2015;9:1186-94.
- Wang A, Huen SC, Luan HH, Yu S, Zhang C, Gallezot JD, *et al.* Opposing effects of fasting metabolism on tissue tolerance in bacterial and viral inflammation. *Cell* 2016;166:1512-25.e12.
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: A retrospective cohort study. *Lancet* 2020;395:1054-62.
- Amin MEK, Abdelmageed A. Clinicians' perspectives on caring for Muslim patients considering fasting during Ramadan. *J Relig Health* 2019. doi: 10.1007/s10943-019-00820-y.
- Roky R, Houti I, Moussamih S, Qotbi S, Aadil N. Physiological and chronobiological changes during Ramadan intermittent fasting. *Ann Nutr Metab* 2004;48:296-303.
- Janicki-Deverts D, Cohen S, Turner RB, Doyle WJ. Basal salivary cortisol secretion and susceptibility to upper respiratory infection. *Brain Behav Immun* 2016;53:255-61.