

Triamcinolone acetonide and anecortave acetate do not stimulate uveal melanoma cell growth

Mariam el Filali, Irene Homminga, Willem Maat, Pieter A. van der Velden, Martine J. Jager

Department of Ophthalmology, Leiden University Medical Centre, Leiden, the Netherlands

Purpose: Radiotherapy-induced radiation retinopathy can develop in over 40% of eyes treated for uveal melanoma. Triamcinolone acetonide (TA) and anecortave acetate (AA) can be used to treat radiation retinopathy. It is not known whether TA or AA has any effect on potentially still viable uveal melanoma cells in the choroid after radiotherapy. We therefore studied the effect of these drugs on the proliferation of uveal melanoma cell lines in vitro. Furthermore, as these drugs are supposed to counteract vascular leakage, we determined their effect on the expression and production of the proangiogenic vascular endothelial growth factor-A (VEGF-A), the antiangiogenic pigment epithelium-derived factor (PEDF), and thrombospondin-1 (TSP-1) in uveal melanoma cells.

Methods: Three uveal melanoma cell lines were treated in vitro with TA or AA. Cell proliferation was measured by counting cells and using the Water-Soluble Tetrazolium Salt-1 (WST-1) assay. VEGF-A and PEDF production was measured by ELISA, and intracellular expression of angiogenic-associated genes including VEGF-A, PEDF, and TSP-1 was determined by real-time quantitative RT-PCR.

Results: We found no effect of TA or AA on tumor cell growth or production of VEGF-A and PEDF in any of the three uveal melanoma cell lines tested. Regarding expression as measured by RT-PCR, TA had an inhibiting effect on TSP-1 in only one cell line, and no effect on VEGF-A or PEDF. AA showed a similar lack of effect.

Conclusions: Since TA and AA do not stimulate uveal melanoma cell growth, it seems to be safe to use these drugs to treat radiation retinopathy after irradiation for uveal melanoma. Additional experiments using more cell lines or primary tumor cell cultures are needed to validate this conclusion. Furthermore, the results of our study suggest that TA does not exert its antileakage effect through downregulation of VEGF-A or upregulation of TSP-1 or PEDF in uveal melanoma cell lines. It is possible that TA and AA influence these pro- and antiangiogenic factors only under hypoxic circumstances. Further investigation is needed.

Uveal melanoma (UM) is the most common primary intraocular tumor in adults, with an incidence of approximately 5–7 per 1,000,000 in Caucasian populations [1-4]. The five-year tumor-related survival rate is about 70%–80% [5,6], and eventually about 50% of patients will die from the disease [7,8]. Because the eye lacks lymphatic vessels, the route of dissemination is almost exclusively hematogenous. The most common site of metastasis formation is the liver [9,10]. Predictors of survival for UM patients are identified by histologic cell type, tumor diameter, tumor location, age, gender [11], and cytogenetic parameters. Loss of chromosome 3 is one of the most significant predictors for UM-related deaths [12,13]. There are different modes of therapy available for treating a primary UM. Enucleation remains a common treatment for large tumors. Small and medium-sized tumors with a prominence <8 mm can be treated with proton beam therapy, local resection, or radiation [14,15]. Radiotherapy is often combined with transpupillary thermotherapy, which may also allow treatment of larger tumors [16,17]. Radiotherapy, however, not only damages UM cells, but also

normal healthy cells located in the immediate area surrounding the tumor. Capillary endothelial cells are especially sensitive to radiation and presumably die directly or later by mitotic death [18]. Capillary endothelial cell loss causes changes in the structure and permeability of the vessels in the affected area. This leads to an occlusive microangiopathy, clinically known as radiation retinopathy. A variety of sight-threatening manifestations are associated with radiation retinopathy, such as macular edema and neovascularization [19]. The onset of radiation retinopathy occurs on average 26 months after radioactive plaque therapy [20]. The cause of this delay is not quite clear, although duration of the endothelial cell cycle and delayed mitotic cell death might be an explanation [18]. Gunduz et al. [21] found that approximately 5% of patients treated by radioactive plaque developed radiation retinopathy after one year. After five years this percentage rose to about 40%.

At the moment, there is no established effective treatment for radiation retinopathy and its complications, although laser photocoagulation has shown some beneficial effect [20]. Finger and Chin [22] demonstrated that antiangiogenic agents, such as Avastin, have a beneficial effect.

Triamcinolone acetonide (TA) is a glucocorticoid that has already been shown to improve vision for a few months in

Correspondence to: M.J. Jager, Department of Ophthalmology, Leiden University Medical Centre, P.O. Box 9600, 2300 RC Leiden, the Netherlands; Phone: +31715263938; FAX: +31715248222; email: m.j.jager@lumc.nl

patients with macular edema associated with diabetic retinopathy [23-25] and in patients with exudative age-related macular degeneration (ARMD) [26]. Although the causal mechanisms underlying these diseases differ, diabetic retinopathy, exudative ARMD, and radiation retinopathy share the same sight-threatening complications, such as macular edema and neovascularization. A recent study reported a temporary positive effect of intravitreal TA injections in a group of 31 patients with radiation maculopathy [27].

While TA may be a suitable drug to treat complications of radiation retinopathy, its effect on UM cells is unknown. The possibility that there are living UM cells in the choroid of eyes treated with radiotherapy cannot be excluded [28]; local recurrences do develop [29]. We therefore set out to study the effect of TA on the proliferation of UM cells before using this drug in patients with radiation retinopathy. To our knowledge, this has not been investigated previously.

TA has been shown to have an antiangiogenic effect [30-32], though the mechanism by which this effect comes about is not clear. The extent of tumor angiogenesis is determined by the balance between proangiogenic and antiangiogenic molecules, released by both tumor cells and surrounding cells [33]. The formation of new vessels is a process involving endothelial cells; tumor cells that lack a sufficient blood supply are most likely the incentive of angiogenesis. Vascular endothelial growth factor-A (VEGF-A) plays an important role in angiogenesis, regulating vasopermeability and the proliferation and migration of endothelial cells [34]. VEGF-A seems also to be the key mediator in ocular vessel diseases as well as in tumor angiogenesis [35].

We also studied two inhibitors of angiogenesis, pigment epithelium-derived factor (PEDF) and thrombospondin-1 (TSP-1). Various studies have documented that PEDF inhibits angiogenesis in the eye [36,37], whereas TSP-1 has been shown to inhibit cutaneous melanoma progression by suppressing tumor vessel formation [38].

One of the disadvantages of TA, however, is that it causes ocular hypertension in about 30% of the cases [39] and may lead to glaucomatous damage. A new steroid-derived substance, anecortave acetate (AA), an angiostatic cortisone, has been developed to be devoid of corticosteroid side effects such as ocular hypertension [40]. Hence, AA may be a good alternative to TA. A first study of AA administered in a juxtasceral depot to treat subfoveal choroidal neovascularization in ARMD showed good results in the prevention of further vessel development [41]. We therefore analyzed the effect of both TA as well as AA on proliferation, expression and production of pro- and antiangiogenic factors.

METHODS

Cell lines: Three well described cell lines were selected on the basis of known differences in VEGF-A production (low,

medium, high) to perform our experiments: Mel 285, 92-1 and OCM-1. Cell line OCM-1 was kindly provided by Dr. J. Kan-Mitchell (Karmanos Cancer Institute, Detroit, MI) [43]. Cell line 92-1 was established in our laboratory [42]. Mel 285 was a generous gift of Dr. B.R. Ksander (Schepens Eye Research Institute, Boston, MA) [44]. Cell lines Mel 285 and 92-1 were cultured in Roswell Park Memorial Institute (RPMI) 1640 (Gibco Life Technologies, Breda, the Netherlands) to which 10% fetal calf serum, 2% glutamine (Gibco), and 2% penicillin and streptomycin (Gibco) were added. OCM-1 was cultured in Dulbecco's Modified Eagle's Medium (DMEM; Gibco) with 10% fetal calf serum and 2% penicillin and streptomycin (Gibco). Cultures were passaged every three to four days.

Triamcinolone acetonide and AL-4940 preparations: We used triamcinolone acetonide T6501 (Sigma-Aldrich, St. Louis, MO). The recommended intravitreal dose of TA is 4 mg (0.1 ml) [27]. The dose to which the UM cells are expected to be exposed *in vivo* after injection of TA is 1 mg/ml. Unfortunately, due to massive TA crystal deposits the maximal tolerated dose *in vitro* was 0.04 mg/ml. Ten mg TA was suspended in 230 μ l methanol, producing a 100 mM stock suspension. This stock was serially diluted with medium to concentrations of 10 μ M and 100 μ M. Extra methanol was added to the 10 μ M suspension to obtain a concentration of 0.1% methanol, similar to the 100 μ M suspension. As a control, we used solutions of 0.1% methanol in medium. We used 9(11)-dehydrocortisol (AL-4940-06; Alcon Laboratories, Fort Worth, TX) instead of AA. *In vivo* AA is rapidly deacetylated into AL-4940, and AL-4940 is therefore the predominantly present active form [45]. A 10 mM stock solution was prepared by dissolving 5.4 mg AL-4940 in 1.6 ml dimethyl sulfoxide (DMSO). This stock was serially diluted with medium to concentrations of 0.1, 1.0, and 10 μ M. A solution of 0.1% DMSO was used as control.

Triamcinolone acetonide administration: Growth assays were performed in triplicate. For each assay, 64 wells per cell line were filled with 1×10^4 UM cells per well in 1 ml medium, using 24 well plates. Cells were allowed to settle for 24 h. After 24 h, the regular medium was replaced by 1 ml of one of the following media: RPMI/DMEM (control), RPMI/DMEM with methanol (second control), 10 μ M TA suspension, or 100 μ M TA suspension; for the AL-4940 assay: RPMI (control), RPMI with DMSO (second control), 0.1 μ M AL-4940 solution, 1.0 μ M AL-4940 solution, or 10 μ M AL-4940 solution. Cells were cultured in an incubator at 37 °C and 5.0% CO₂.

Cell proliferation and cytotoxicity: Four, six, eight, and ten days after medium replacement, cells were counted or replaced with fresh specific media. Cells from four wells of each cell line at one time point were pooled to collect a high enough number of cells to obtain a reliable count. Cell death was determined by trypan blue dye-exclusion using a Bürker

TABLE 1. PRIMER SEQUENCES OF THE GENES STUDIED IN THE RT-PCR ASSAY

Primer	Gene	Forward primer (5'-3')	Reverse primer (5'-3')
Endogenous controls	<i>RPS-11</i>	AAGCAGCCGACCATCTTTCA	CGGGAGCTTCTCCTTGCC
	<i>HRPT</i>	CGAGATGTGATGAAGGAGATGG	GCAGGTCAGCAAAGAATTTATAGC
	<i>β-ACTIN</i>	CGGGACCTGACTGACTACCTC	CTCCTTAATGTACGCACGATTT
Genes under study	<i>VEGF-A</i>	GCCCTTGCCTTGCTGCTCTACC	GTGATGATTCTGCCTCCTCTTC
	<i>PEDF</i>	AGCAATCTCCTTCTCGGTGTGG	CCTCACGGTCTCTCTTCATCC
	<i>TSP-1</i>	AGGTCTTCAGCGTGGTGTCC	ACAAACAGGGTGATGCTCTTCC

Primer sequences of the control and studied genes that were used in real-time quantitative RT-PCR assay to determine level of expression in UM cell lines. Abbreviations: β -ACTIN is beta-actin; HPRT is hypoxanthine phosphoribosyltransferase; RPS-11 is ribosomal protein S11.

counting chamber (Omnilabo, Breda, the Netherlands). In addition, cell proliferation was measured by mitochondrial function using the WST-1 assay (Roche Diagnostics, Indianapolis, IN), as previously described [46]. In short, 96 well plates were allotted 1250 UM cells per well and either filled with regular medium (control) or 100 μ M TA suspension. Absorbance was measured at 450 nm (n=8) on a multiwell spectrophotometer (Perkin Elmer, Wellesley, MA). *VEGF-A and PEDF production*: In the supernatant obtained on day eight, VEGF-A and PEDF concentrations were determined using commercial solid phase sandwich enzyme-linked immunosorbent assay kits. For VEGF-A, we used a human VEGF-A ELISA immunoassay kit (Biosource, Camarillo, CA), and for PEDF, we used a PEDF Sandwich ELISA antigen detection kit (BioProducts MD, Middletown, MD).

Real-time Quantitative RT-PCR analysis: Several different gene mRNA expressions were analyzed by real-time quantitative reverse transcriptase-polymerase chain reaction (RT-PCR). RNA was isolated on day 8 in one TA assay, using an Rneasy® Mini Kit (Qiagen, Valencia, CA). RNA samples were stored at -80 °C until further processing, when approximately 1 μ g of RNA per sample was reverse-transcribed using the iScript cDNA synthesis kit (Bio-Rad, Hercules, CA). The solutions (20 μ l) were diluted by adding sterile water until the volume reached a total of 100 μ l. In 96 wells, 2 μ l of this solution was added to a 15 μ l solution of iQ SYBR Green Supermix, forward and reverse primers (10 μ M solutions) for VEGF-A, PEDF, TSP-1, beta-actin (β -ACTIN), hypoxanthine phosphoribosyltransferase (HRPT), or ribosomal protein S11 (RPS-11), and sterile water (volume ratio respectively 10:1:1:8). The primers for all genes under study were designed with the Primer Express software (PE Applied Biosystems; Foster city, CA; Table 1) [47]. A quantitative analysis of the samples was then performed for all genes by real-time quantitative RT-PCR in a MyiQ iCycler real-time PCR system (Bio-Rad). An accepted method to correct the sample-to-sample variation when determining gene expression is to select a cellular housekeeping gene that serves as an endogenous control, against which the target gene

expression levels can be normalized [48]. RPS-11 is a housekeeping gene that has been recently used to normalize gene expression in UM cells [47]. β -actin is a relatively stable cytoskeletal protein generally thought to be present at a constant level in cells, regardless of experimental treatment or technical procedure [49]. HPRT, an enzyme in purine metabolism, is reported as a constitutively expressed housekeeping gene [49,50].

The PCR reaction settings were 95 °C for 3 min, then 40 cycles at 95 °C for 30 s and 60 °C for 30 s, then 95 °C for 1 min and 60 °C for 1 min. Using a control, we calculated assigned relative quantity for any sample for all genes as follows:

$$\text{Relative Quantity}_{\text{sample (Gene X)}} = E_{\text{Gene X}}^{[C_T(\text{Control}) - C_T(\text{sample})]}$$

where E represents efficiency of primer or (probe) set, $C_T(\text{Control})$ equals the average C_T for the sample which has been assigned as a control, and $C_T(\text{sample})$ is the average C_T for the sample. This is referred to as normalized expression [51].

Statistical analysis: Data are expressed as the mean±standard deviation. Student's *t*-test was used to determine whether there were statistically significant differences between treatment groups in the cell viability assay. A $p < 0.05$ was considered statistically significant.

RESULTS

Influence of TA and AA on UM cell proliferation: TA and AA may be potential drugs to treat radiation retinopathy. However harmful side effects such as proliferation of UM cells have to be ruled out.

Treatment of UM cells with TA did not induce cell death as measured by the Bürker counting chamber (data not shown). The WST-1 assay demonstrated no significant effect of TA treatment on cell proliferation of the UM cell lines 92.1 ($p=0.755$), OCM-1 ($p=0.844$), and Mel285 ($p=0.487$; Figure 1). AA showed similar results. TA and AA had no toxic effect on UM cells, as there were no differences in the percentages of dead cells between the different treatments (data not shown).

VEGF-A and PEDF production: To investigate any direct pro- or antiangiogenic effects of TA and AA on UM cells, we studied the influence of these drugs on VEGF-A and PEDF production of three UM cell lines using an ELISA on culture supernatant. Since cell numbers per well differed considerably between some treatments and experiments, we corrected for the number of cells by dividing the concentration of VEGF-A or PEDF measured in the supernatant by the amount of cells present in the same well. All cell lines produced VEGF-A and

PEDF, but clear differences were observed in the amounts: OCM-1 produced the highest levels of VEGF-A (50 pg/ml/ 10^4 cells); 92-1 produced approximately one-fifth of this amount (10 pg/ml/ 10^4 cells); and Mel 285 produced the lowest amount of VEGF-A (5 pg/ml/ 10^4 cells).

Cell line 92-1 produced large amounts of PEDF (2.000 pg/ml/ 10^4 cells), around 100 times the amount produced by cell line OCM-1 (20 pg/ml/ 10^4 cells). Mel 285 produced very little PEDF (3 pg/ml/ 10^4 cells). Addition of TA or AA to the cell cultures had no effect on either VEGF-A or PEDF production (Figure 2A,B).

VEGF-A, PEDF, and TSP-1 expression: We determined by real-time quantitative RT-PCR whether TA or AA had any effect on several genes in UM cell lines that were associated with angiogenesis. VEGF-A was expressed in all cell lines at variable levels. TA treatment had no major (>1.0 time fold difference) effect on VEGF-A expression. On the contrary, TA gave a dose-dependent reduction of TSP-1 expression in Mel285 (time fold difference=1.5–2.9). TSP-1 was hardly expressed in 92.1 (Figure 3). AA had no significant effect either on VEGF-A or TSP-1 expression. PEDF is highly expressed in cell line 92-1 and only to a minimal extend in the other cell lines. PEDF expression in cell line 92-1 was not affected by TA or AA (data not shown).

DISCUSSION

Our results showed no apparent stimulating or inhibiting effect of TA on UM cell proliferation in the concentrations used in this study [27]. The effect of TA on UM cell lines has not yet been investigated, but the effect of TA on other cell types has been the object of some studies. For instance, TA was found to inhibit the growth of retinal pigment epithelium (RPE) when ARPE 19 cells were treated with concentrations ranging from 1 μ M to 2.3 mM, but stimulated growth at a concentration of 1 nM [52,53]. Another study, using bovine retinal endothelial cells, found that TA at a concentration of 115 μ M had no effect on growth; concentrations ranging from 230 μ M to 4.6 mM had an inhibiting effect, and concentrations of 6.8 mM and higher were cytotoxic [54]. Likewise, in vitro incubation of RPE with corticosteroids induced a specific and dose-dependent reduction of cell viability. These toxic events were not associated with the anti-inflammatory activity of these compounds but depended on the hydrosolubility of their formulation [55].

While previous studies on RPE cells demonstrated an inhibition of VEGF production and expression by TA, we found no effect of TA on VEGF-A production or expression in UM cells. For instance, TA reduced VEGF expression of ARPE19 cells under oxidative stress [53] and inhibited cobalt-stimulated VEGF production in Müller cells, probably by a destabilization of VEGF mRNA [56]. The lack of inhibition of TA on VEGF-A production and expression in our study could perhaps be explained by the absence of specific stimuli




Figure 1. Effect of TA on uveal melanoma cell proliferation. Cells were treated with four different suspensions: Park Memorial Institute/Dulbecco's Modified Eagle Medium (RPMI/DMEM; control), RPMI/DMEM with methanol (second control), 10 mM TA suspension, or 100 mM TA suspension. Cell proliferation was determined by a WST-1 assay (absorbance (OD) and the mean \pm SD is provided for each time point (n=8)). No substantial difference was seen in proliferation of uveal melanoma cell lines between TA treatment and control (cell line 92.1 (p=0.755), OCM-1 (p=0.844), and Mel285 (p=0.487)). Abbreviations: OD is optical density; TA is triamcinolone acetonide; CO is control.

that would upregulate VEGF-A production and expression, such as oxidative stress or cobalt administration. In the study by Sears et al. [56], only cobalt-stimulated VEGF production was inhibited by TA, while basal VEGF production was not affected. Similarly, another study showed no alteration in VEGF expression by TA in a rat retina when no special stress was applied [57]. We are currently performing experiments to determine the effect of TA and other drugs on UM cells under stressful conditions, such as hypoxia.

PEDF inhibits angiogenesis by inducing apoptosis in endothelial cells that try to form new vessels [58,59]. Besides the separate levels of VEGF-A and PEDF, the balance between the two seems to be decisive in whether or not angiogenesis takes place [60]. Yang et al. previously demonstrated expression of PEDF in UM cell lines. Furthermore, they observed an inhibitory effect of angiostatin on the ratio of VEGF:PEDF mRNA levels in vitro [59]. Likewise, Mel 285, 92-1, and OCM-1 all produced PEDF in our experiments. We observed no effect of TA on the production of PEDF by any of the UM cell lines. Therefore, stimulation of production of PEDF by UM cells seems not to be the antiangiogenic working mechanism of this drug under normal circumstances. The high expression of PEDF by cell line 92-1 was unexpected, since PEDF has been shown to work as a tumor growth inhibitor through antiangiogenic mechanisms [61-63] and cell line 92-1 was derived from a highly malignant tumor [42]. However, there are reports that melanocytic tumors have a relatively high expression of PEDF [64]. Also, PEDF might have a biphasic activity, inhibiting angiogenesis at normal amounts, but stimulating angiogenesis at high concentrations [65].

The balance between VEGF-A and PEDF, as we have described, seems also to be influenced by TSP-1 in that it inhibits endothelial cell migration and proliferation and also induces apoptosis [66]. In cell line Mel 285, TSP-1 expression was about twofold inhibited in cells grown in TA compared to the control cells. Previous reports on the effect of TA on the expression of TSP-1 could not be found, though some effects of other corticosteroids have been reported.




Figure 2. Production of VEGF-A and PEDF in three different UM cell lines exposed to various TA and AA doses. Cells were treated with four or five different suspensions. **A:** For triamcinolone acetonide (TA) experiments, we used either Park Memorial Institute/Dulbecco's Modified Eagle Medium (RPMI/DMEM; control), RPMI/DMEM with methanol (second control), a 10 mM TA suspension, or a 100 mM TA suspension; **B:** For anecortave acetate (AA) experiments, we used either RPMI (control), RPMI with dimethyl sulfoxide (DMSO; second control), 0.1 mM AL-4940 solution, 1.0 mM AL-4940 solution, or 10 mM AL-4940 solution. The level of vascular endothelial growth factor-A (VEGF-A) protein (n=2) is shown in pg/ml (mean±SD). All cell lines produced VEGF-A and pigment epithelium-derived factor (PEDF), but clear differences were observed: OCM-1 produced the highest levels of VEGF-A, while cell line Mel 285 produced the lowest amount of VEGF-A. Cell line 92-1 produced large amounts of PEDF (>2000 pg/ml). Mel 285 produced scant PEDF. Addition of TA or AA to the cell cultures had no effect on either VEGF-A or PEDF production.




Figure 3. The effect of TA (two doses) on gene expression of VEGF-A and TSP-1 in three different uveal melanoma cell lines. Cells were treated with four different suspensions: Park Memorial Institute/Dulbecco's Modified Eagle Medium (RPMI/DMEM; control), RPMI/DMEM with methanol (second control), 10 mM triamcinolone acetonide (TA) suspension, or 100 mM TA suspension. Expression levels of vascular endothelial growth factor-A (VEGF-A) and thrombospondin-1 (TSP-1) are normalized against two endogenous controls (normalized fold; mean±SD). Experiments were performed twice. VEGF-A was expressed in all cell lines at variable levels. TA treatment had no major (>1.0 fold) effect on VEGF-A expression. TSP-1 was not expressed in 92.1. Mel 285 showed a decreased TSP-1 expression (time-fold difference=1.5-2.9) after treatment with TA.

Hydrocortisone upregulated TSP-1 expression in a glioma cell line [67], and dexamethasone increased TSP-1 expression in a murine trophoblast-like cell culture [68]. The results of Mel 285 do not agree with this, and may indicate an unexpected effect of TA. TSP-1 expression of cell line 92-1 was very low, possibly due to inactivation of this gene in the same manner as described for hypermethylated p16 (INK4a) [69].

Parallel experiments with AA, designed to eliminate any possible side effects of the glucocorticoid fraction, showed similar results as with TA. There seems to be no differential effect of this angiostatic cortisone compared to TA on potentially still viable UM cells in the choroid after radiotherapy; both drugs have hardly any effect at all.

The risk of an increase in local recurrences of UM by TA or AA in the treatment of radiation retinopathy by direct stimulation of UM cells seems to be low. We based these conclusions on a restricted number of cell lines, and extrapolation to the in vivo situation is by definition limited. In addition, the clinical dose of TA could not be investigated due to massive crystal deposits. Perhaps a certain threshold has to be reached before TA can demonstrate any effect.

Nonetheless, our results suggest that TA does not exert any antiangiogenic effect through influencing basal VEGF-A, PEDF, TSP-1 production or expression in UM cells. However, influence of hypoxia and stimulation of these cell lines by VEGF or FGF2 or by antiangiogenic factors is possible and is under investigation.

Furthermore, there are many other pro- and antiangiogenic factors that could be altered by TA. Plasminogen activator inhibitor type-1 for instance, has been implicated in metastasis formation and tumor-associated angiogenesis in UM [70], and would be an interesting candidate for further studies. These factors will need further investigation.

ACKNOWLEDGMENTS

This study was supported by an unrestricted grant from Alcon. Anecortave Acetate was kindly supplied by Alcon.

REFERENCES

- Egan KM, Seddon JM, Glynn RJ, Gragoudas ES, Albert DM. Epidemiologic aspects of uveal melanoma. *Surv Ophthalmol* 1988; 32:239-51. [PMID: 3279559]
- Singh AD, Topham A. Incidence of uveal melanoma in the United States: 1973-1997. *Ophthalmology* 2003; 110:956-61. [PMID: 12750097]
- Hu DN, Yu GP, McCormick SA, Schneider S, Finger PT. Population-based incidence of uveal melanoma in various races and ethnic groups. *Am J Ophthalmol* 2005; 140:612-7. [PMID: 16226513]
- Virgili G, Gatta G, Ciccolallo L, Capocaccia R, Biggeri A, Crocetti E, Lutz JM, Paci E. EURO CARE Working Group. Incidence of uveal melanoma in Europe. *Ophthalmology* 2007; 114:2309-15. [PMID: 17498805]
- Bergman L, Seregard S, Nilsson B, Lundell G, Ringborg U, Ragnarsson-Olding B. Uveal melanoma survival in Sweden from 1960 to 1998. *Invest Ophthalmol Vis Sci* 2003; 44:3282-7. [PMID: 12882771]
- Singh AD, Topham A. Survival rates with uveal melanoma in the United States: 1973-1997. *Ophthalmology* 2003; 110:962-5. [PMID: 12750098]
- Augsburger JJ, Correa ZM, Freire J, Brady LW. Long-term survival in choroidal and ciliary body melanoma after enucleation versus plaque radiation therapy. *Ophthalmology* 1998; 105:1670-8. [PMID: 9754176]
- Kujala E, Makitie T, Kivela T. Very long-term prognosis of patients with malignant uveal melanoma. *Invest Ophthalmol Vis Sci* 2003; 44:4651-9. [PMID: 14578381]
- Kath R, Hayungs J, Bornfeld N, Sauerwein W, Hoffken K, Seeber S. Prognosis and treatment of disseminated uveal melanoma. *Cancer* 1993; 72:2219-23. [PMID: 7848381]
- Rietschel P, Panageas KS, Hanlon C, Patel A, Abramson DH, Chapman PB. Variates of survival in metastatic uveal melanoma. *J Clin Oncol* 2005; 23:8076-80. [PMID: 16258106]
- Mooy CM, De Jong PT. Prognostic parameters in uveal melanoma: a review. *Surv Ophthalmol* 1996; 41:215-28. [PMID: 8970236]
- Prescher G, Bornfeld N, Hirche H, Horsthemke B, Jockel KH, Becher R. Prognostic implications of monosomy 3 in uveal melanoma. *Lancet* 1996; 347:1222-5. [PMID: 8622452]
- Sisley K, Rennie IG, Parsons MA, Jacques R, Hammond DW, Bell SM, Potter AM, Rees RC. Abnormalities of chromosomes 3 and 8 in posterior uveal melanoma correlate with prognosis. *Genes Chromosomes Cancer* 1997; 19:22-8. [PMID: 9135991]
- Damato B. Developments in the management of uveal melanoma. *Clin Experiment Ophthalmol* 2004; 32:639-47. [PMID: 15575836]
- Pötter R, Janssen K, Prott FJ, Widder J, Haverkamp U, Busse H, Müller RP. Ruthenium-106 eye plaque brachytherapy in the conservative treatment of uveal melanoma: evaluation of 175 patients treated with 150 Gy from 1981-1989. *Front Radiat Ther Oncol* 1997; 30:143-9. [PMID: 9205894]
- Bartlema YM, Oosterhuis JA, Jounnee-De Korver JG, Tjho-Heslinga RE, Keunen JE. Combined plaque radiotherapy and transpupillary thermotherapy in choroidal melanoma: 5 years' experience. *Br J Ophthalmol* 2003; 87:1370-3. [PMID: 14609837]
- Desjardins L, Lumbroso-Le Rouic L, Levy-Gabriel C, Dendale R, Delacroix S, Nauraye C, Estève M, Plancher C, Asselain B. Combined proton beam radiotherapy and transpupillary thermotherapy for large uveal melanomas: a randomized study of 151 patients. *Ophthalmic Res* 2006; 38:255-60. [PMID: 16888407]
- Archer DB, Amoaku WM, Gardiner TA. Radiation retinopathy—clinical, histopathological, ultrastructural and experimental correlations. *Eye* 1991; 5:239-51. [PMID: 2070883]
- Zamber RW, Kinyoun JL. Radiation retinopathy. *West J Med* 1992; 157:530-3. [PMID: 1441494]
- Finger PT, Kurli M. Laser photocoagulation for radiation retinopathy after ophthalmic plaque radiation therapy. *Br J Ophthalmol* 2005; 89:730-8. [PMID: 15923510]
- Gunduz K, Shields CL, Shields JA, Cater J, Freire JE, Brady LW. Radiation retinopathy following plaque radiotherapy for

- posterior uveal melanoma. *Arch Ophthalmol* 1999; 117:609-14. [PMID: 10326957]
22. Finger PT, Chin K. Anti-vascular endothelial growth factor bevacizumab (avastin) for radiation retinopathy. *Arch Ophthalmol* 2007; 125:751-6. [PMID: 17562985]
 23. Er H, Yilmaz H. Intravitreal cortisone injection for refractory diffuse diabetic macular edema. *Ophthalmologica* 2005; 219:394-400. [PMID: 16286802]
 24. Patelli F, Fasolino G, Radice P, Russo S, Zumbo G, Di Tizio FM, Frisone G, Marchi S. Time course of changes in retinal thickness and visual acuity after intravitreal triamcinolone acetonide for diffuse diabetic macular edema with and without previous macular laser treatment. *Retina* 2005; 25:840-5. [PMID: 16205561]
 25. Sorensen TL, Haamann P, Villumsen J, Larsen M. Intravitreal triamcinolone for macular oedema: efficacy in relation to aetiology. *Acta Ophthalmol Scand* 2005; 83:67-70. [PMID: 15715560]
 26. Jonas JB, Spandau UH, Kampeter BA, Harder B. Follow-up after intravitreal triamcinolone acetonide for exudative age-related macular degeneration. *Eye* 2007; 21:387-94. [PMID: 16410809]
 27. Shields CL, Demirci H, Dai V, Marr BP, Mashayekhi A, Materin MA, Manquez ME, Shields JA. Intravitreal triamcinolone acetonide for radiation maculopathy after plaque radiotherapy for choroidal melanoma. *Retina* 2005; 25:868-74. [PMID: 16205566]
 28. Pe'er J, Stefani FH, Seregard S, Kivela T, Lommatzsch P, Prause JU, Sobottka B, Damato B, Chowder I. Cell proliferation activity in posterior uveal melanoma after Ru-106 brachytherapy: an EORTC ocular oncology group study. *Br J Ophthalmol* 2001; 85:1208-12. [PMID: 11567966]
 29. Harbour JW, Char DH, Kroll S, Quivey JM, Castro J. Metastatic risk for distinct patterns of postirradiation local recurrence of posterior uveal melanoma. *Ophthalmology* 1997; 104:1785-92. [PMID: 9373108]
 30. Murata M, Shimizu S, Horiuchi S, Taira M. Inhibitory effect of triamcinolone acetonide on corneal neovascularization. *Graefes Arch Clin Exp Ophthalmol* 2006; 244:205-9. [PMID: 16044325]
 31. Hashimoto I, Nakanishi H, Shono Y, Toda M, Tsuda H, Arase S. Angiostatic effects of corticosteroid on wound healing of the rabbit ear. *J Med Invest* 2002; 49:61-6. [PMID: 11901762]
 32. Antoszyk AN, Gottlieb JL, Machemer R, Hatchell DL. The effects of intravitreal triamcinolone acetonide on experimental pre-retinal neovascularization. *Graefes Arch Clin Exp Ophthalmol* 1993; 231:34-40. [PMID: 8428678]
 33. Folkman J. The role of angiogenesis in tumor growth. *Semin Cancer Biol* 1992; 3:65-71. [PMID: 1378311]
 34. Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med* 2003; 9:669-76. [PMID: 12778165]
 35. Carmeliet P. VEGF as a key mediator of angiogenesis in cancer. *Oncology* 2005; 69:4-10. [PMID: 16301830]
 36. Dawson DW, Volpert OV, Gillis P, Crawford SE, Xu H, Benedict W, Bouck NP. Pigment epithelium-derived factor: a potent inhibitor of angiogenesis. *Science* 1999; 285:245-8. [PMID: 10398599]
 37. Matsuoka M, Ogata N, Otsuji T, Nishimura T, Takahashi K, Matsumura M. Expression of pigment epithelium derived factor and vascular endothelial growth factor in choroidal neovascular membranes and polypoidal choroidal vasculopathy. *Br J Ophthalmol* 2004; 88:809-15. [PMID: 15148217]
 38. Trotter MJ, Colwell R, Tron VA. Thrombospondin-1 and cutaneous melanoma. *J Cutan Med Surg* 2003; 7:136-41. [PMID: 12447621]
 39. Jaissle GB, Szurman P, Bartz-Schmidt KU. Ocular side effects and complications of intravitreal triamcinolone acetonide injection.
 40. Clark AF, Mellon J, Li XY, Ma D, Leher H, Apte R, Alizadeh H, Hegde S, McLenaghan A, Mayhew E, D'Orazio TJ, Niederkorn JY. Inhibition of intraocular tumor growth by topical application of the angiostatic steroid anecortave acetate. *Invest Ophthalmol Vis Sci* 1999; 40:2158-62. [PMID: 10440274]
 41. Schmidt-Erfurth U, Michels S, Michels R, Aue A. Anecortave acetate for the treatment of subfoveal choroidal neovascularization secondary to age-related macular degeneration. *Eur J Ophthalmol* 2005; 15:482-5. [PMID: 16001382]
 42. De Waard-Siebinga I, Blom DJ, Griffioen M, Schrier PI, Hoogendoorn E, Beverstock G, Danen EH, Jager MJ. Establishment and characterization of an uveal-melanoma cell line. *Int J Cancer* 1995; 62:155-61. [PMID: 7622289]
 43. Kan-Mitchell J, Mitchell MS, Rao N, Liggett PE. Characterization of uveal melanoma cell lines that grow as xenografts in rabbit eyes. *Invest Ophthalmol Vis Sci* 1989; 30:829-34. [PMID: 2722439]
 44. Nareyeck G, Zeschnigk M, Prescher G, Lohmann DR, Anastassiou G. Establishment and characterization of two uveal melanoma cell lines derived from tumors with loss of one chromosome 3. *Exp Eye Res* 2006; 83:858-64. [PMID: 16750193]
 45. Clark AF. AL-3789: a novel ophthalmic angiostatic steroid. *Expert Opin Investig Drugs* 1997; 6:1867-77. [PMID: 15989587]
 46. Narayanan R, Kenney MC, Kamjoo S, Trinh TH, Seigel GM, Resende GP, Kuppermann BD. Trypan blue: effect on retinal pigment epithelial and neurosensory retinal cells. *Invest Ophthalmol Vis Sci* 2005; 46:304-9. [PMID: 15623789]
 47. Zuidervaart W, van der Velden PA, Hurks MH, van Nieuwpoort FA, Out-Luiting CJ, Singh AD, Frants RR, Jager MJ, Gruijs NA. Gene expression profiling identifies tumour markers potentially playing a role in uveal melanoma development. *Br J Cancer* 2003; 89:1914-9. [PMID: 14612903]
 48. Bustin SA. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol* 2000; 25:169-93. [PMID: 11013345]
 49. Gilsbach R, Kouta M, Bonisch H, Bruss M. Comparison of in vitro and in vivo reference genes for internal standardization of real-time PCR data. *Biotechniques* 2006; 40:173-7. [PMID: 16526406]
 50. de Kok JB, Roelofs RW, Giesendorf BA, Pennings JL, Waas ET, Feuth T, Swinkels DW, Span PN. Normalization of gene expression measurements in tumor tissues: comparison of 13 endogenous control genes. *Lab Invest* 2005; 85:154-9. [PMID: 15543203]
 51. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. qBase relative quantification framework and software for management and automated analysis of real-

- time quantitative PCR data. *Genome Biol* 2007; 8:R19. [PMID: 17291332]
52. Yeung CK, Chan KP, Chan CK, Pang CP, Lam DS. Cytotoxicity of triamcinolone on cultured human retinal pigment epithelial cells: comparison with dexamethasone and hydrocortisone. *Jpn J Ophthalmol* 2004; 48:236-42. [PMID: 15175915]
53. Matsuda S, Gomi F, Oshima Y, Tohyama M, Tano Y. Vascular endothelial growth factor reduced and connective tissue growth factor induced by triamcinolone in ARPE19 cells under oxidative stress. *Invest Ophthalmol Vis Sci* 2005; 46:1062-8. [PMID: 15728566]
54. Spandau UH, Sauder G, Schubert U, Hammes HP, Jonas JB. Effect of triamcinolone acetonide on proliferation of retinal endothelial cells in vitro and in vivo. *Br J Ophthalmol* 2005; 89:745-7. [PMID: 15923512]
55. Valamanesh F, Torriglia A, Savoldelli M, Gandolphe C, Jeany JC, BenEzra D, Behar-Cohen F. Glucocorticoids induce retinal toxicity through mechanisms mainly associated with paraptosis. *Mol Vis* 2007; 13:1746-57. [PMID: 17960113]
56. Sears JE, Hoppe G. Triamcinolone acetonide destabilizes VEGF mRNA in Muller cells under continuous cobalt stimulation. *Invest Ophthalmol Vis Sci* 2005; 46:4336-41. [PMID: 16249516]
57. Gao H, Qiao X, Gao R, Mieler WF, McPherson AR, Holz ER. Intravitreal triamcinolone does not alter basal vascular endothelial growth factor mRNA expression in rat retina. *Vision Res* 2004; 44:349-56. [PMID: 14659961]
58. Bouck N. PEDF: anti-angiogenic guardian of ocular function. *Trends Mol Med* 2002; 8:330-4. [PMID: 12114112]
59. Yang H, Xu Z, Iuvone PM, Grossniklaus HE. Angiostatin decreases cell migration and vascular endothelium growth factor (VEGF) to pigment epithelium derived factor (PEDF) RNA ratio in vitro and in a murine ocular melanoma model. *Mol Vis* 2006; 12:511-7. [PMID: 16735992]
60. Tong JP, Yao YF. Contribution of VEGF and PEDF to choroidal angiogenesis: a need for balanced expressions. *Clin Biochem* 2006; 39:267-76. [PMID: 16409998]
61. Hosomichi J, Yasui N, Koide T, Soma K, Morita I. Involvement of the collagen I-binding motif in the anti-angiogenic activity of pigment epithelium-derived factor. *Biochem Biophys Res Commun* 2005; 335:756-61. [PMID: 16102727]
62. Streck CJ, Zhang Y, Zhou J, Ng C, Nathwani AC, Davidoff AM. Adeno-associated virus vector-mediated delivery of pigment epithelium-derived factor restricts neuroblastoma angiogenesis and growth. *J Pediatr Surg* 2005; 40:236-43. [PMID: 15868591]
63. Garcia M, Fernandez-Garcia NI, Rivas V, Carretero M, Escamez MJ, Gonzalez-Martin A, Medrano EE, Volpert O, Jorcano JL, Jimenez B, Larcher F, Del Rio M. Inhibition of xenografted human melanoma growth and prevention of metastasis development by dual antiangiogenic/antitumor activities of pigment epithelium-derived factor. *Cancer Res* 2004; 64:5632-42. [PMID: 15313901]
64. Tsuru M, Arima N, Toyozumi Y, Kato S. Pigment epithelium-derived factor as a new diagnostic marker for melanocytic tumors. *Kurume Med J* 2005; 52:81-7. [PMID: 16422173]
65. Apte RS, Barreiro RA, Duh E, Volpert O, Ferguson TA. Stimulation of neovascularization by the anti-angiogenic factor PEDF. *Invest Ophthalmol Vis Sci* 2004; 45:4491-7. [PMID: 15557459]
66. Ren B, Yee KO, Lawler J, Khosravi-Far R. Regulation of tumor angiogenesis by thrombospondin-1. *Biochim Biophys Acta* 2006; 1765:178-88. [PMID: 16406676]
67. Vedoy CG, Sogayar MC. Isolation and characterization of genes associated with the anti-tumor activity of glucocorticoids. *Brain Res Mol Brain Res* 2002; 106:57-69. [PMID: 12393265]
68. Leclerc N, Luppen CA, Ho VV, Nagpal S, Hacia JG, Smith E, Frenkel B. Gene expression profiling of glucocorticoid-inhibited osteoblasts. *J Mol Endocrinol* 2004; 33:175-93. [PMID: 15291752]
69. van der Velden PA, Metzelaar-Blok JA, Bergman W, Monique H, Hurks H, Frants RR, Gruis NA, Jager MJ. Promoter hypermethylation: a common cause of reduced p16(INK4a) expression in uveal melanoma. *Cancer Res* 2001; 61:5303-6. [PMID: 11431374]
70. Ma D, Gerard RD, Li XY, Alizadeh H, Niederkorn JY. Inhibition of metastasis of intraocular melanomas by adenovirus-mediated gene transfer of plasminogen activator inhibitor type 1 (PAI-1) in an athymic mouse model. *Blood* 1997; 90:2738-46. [PMID: 9326241]