

# Organocatalytic Synthesis of $\alpha$ -Trifluoromethyl Allylboronic Acids by Enantioselective 1,2-Borotropic Migration

Sybrand J. T. Jonker,<sup>§</sup> Ramasamy Jayarajan,<sup>§</sup> Tautvydas Kireilis, Marie Deliaival, Lars Eriksson, and Kálmán J. Szabó\*



Cite This: *J. Am. Chem. Soc.* 2020, 142, 21254–21259



Read Online

ACCESS |



Metrics & More



Article Recommendations




Supporting Information

**ABSTRACT:** Chiral  $\alpha$ -substituted allylboronic acids were synthesized by asymmetric homologation of alkenylboronic acids using  $\text{CF}_3/\text{TMS}$ -diazomethanes in the presence of BINOL catalyst and ethanol. The chiral  $\alpha$ -substituted allylboronic acids were reacted with aldehydes or oxidized to alcohols in situ with a high degree of chirality transfer. The oxygen-sensitive allylboronic acids can be purified via their isolated diaminonaphthalene (DanH)-protected derivatives. The highly reactive purified allylboronic acids reacted in a self-catalyzed reaction at room temperature with ketones, imines, and indoles to give congested trifluoromethylated homoallylic alcohols/amines with up to three contiguous stereocenters.

Chiral allylboronic acids<sup>1</sup> are ideal reagents for asymmetric synthesis because of their high reactivity in self-catalyzed allylboration reactions that occur with high stereochemical fidelity. However, the synthesis of chiral allylboronic acids has been an unmet challenge in organic synthesis. Our experience with Pd-catalyzed synthesis of (achiral) allylboronic acids<sup>2</sup> (Figure 1a) and conclusions based on related mechanistic studies<sup>3</sup>

suggested that a metal-free approach would be rewarding for effective control of the stereoselectivity. We hypothesized that the synthesis of chiral allylboronic acids may be devised by using an organocatalytic homologation strategy. The first methods for asymmetric homologation of organoboron compounds were reported by the Matteson group.<sup>4,5</sup> Aggarwal and co-workers<sup>6</sup> applied a useful lithiation–borylation method (Figure 1b) for the synthesis of chiral allyl-Bpin species,<sup>7</sup> including an example of an  $\alpha$ -trifluoromethyl allylboronate derivative.<sup>8</sup> This method is based on stoichiometric formation of chiral lithium carbenoid intermediates, and therefore, it is not suitable for the direct synthesis of allylboronic acids. The Ley<sup>9–11</sup> and Wang<sup>12</sup> groups (Figure 1c) reported a homologation method based on diazo carbenoid reagents. This method was suitable for the synthesis of (achiral) allylboronic acids, which were used in one-pot allylboration<sup>9,11</sup> or converted to their Bpin derivatives.<sup>12</sup> A similar approach was employed by Molander and co-workers (Figure 1c) for the synthesis of benzylboronic acids from trifluoromethyl diazomethane.<sup>13,14</sup> Arnold and co-workers presented a method for the synthesis of chiral  $\alpha$ - $\text{CF}_3$  alkyl- and benzylboron compounds by directed evolution of enzymes.<sup>15,16</sup> Fluorinated organoboronates are useful reagents for selective synthesis of organofluorines.<sup>13–21</sup> The  $\text{CF}_3$  motif very often occurs<sup>22–24</sup> in pharmaceuticals and agrochemical products (Figure 1d).<sup>25–29</sup>

Here we present a new methodology for the synthesis of chiral  $\alpha$ - $\text{CF}_3$  allylboronic acids (Figure 1e). Our concept (Figure 2) is based on reacting alkenylboroxine 2, trifluor-




**Figure 1.** Synthesis of organoboronates and boronic acids as well as examples of bioactive molecules with a  $\text{CF}_3$  group.

Received: September 16, 2020

Published: December 3, 2020





**Figure 2.** Concept of catalytic asymmetric homologation with 1,2-borotropic migration.

omethyl diazomethane **3**, catalytic amounts of BINOL (**4**),<sup>30–36</sup> and stoichiometric amounts of EtOH. Alkenylboroxine **2** readily reacts with diazo compound **3**.<sup>9–11</sup> However, this reaction results in racemic product, such as *rac*-**1-OR**. The racemic background reaction can be avoided by addition of EtOH to the reaction mixture, which forms unreactive alkenylboronic esters **2-OEt**, which are weaker Lewis acids<sup>37</sup> than the corresponding boroxines **2**.<sup>30,31</sup> Because of the dynamic covalent bonding<sup>38</sup> ability of boron, BINOL **4** undergoes transesterification with **2-OEt** to form chiral alkenyl boronate **A**. Exchange of the alkyl group to an aromatic moiety leads to a substantial increase in the Lewis acidity of boron,<sup>37</sup> and therefore, **A** and **3** form ate complex **B** in the stereoselection step of the process (see Figure S3). Then the alkenyl group undergoes stereoselective 1,2-migration<sup>10,39</sup> to afford **C**. Subsequently, ethanolysis of **C** gives product **1-OEt**.

The optimal conditions for the homologation involved using **2a** with an excess of **3**, 20 mol % **4** and 2 equiv of EtOH (Table 1, entry 1). The oxygen-sensitive allylboronic ester **1a-OEt** was protected with diaminonaphthalene (DanH)<sup>40</sup> to give **5a** with 98% ee in 69% yield. When the reaction was repeated with 10 mol % catalyst **4**, the yield was substantially lowered (12%), but the enantioselectivity was practically unchanged (96% ee) (entry 2). Replacement of iodo-BINOL **4** with bromo-BINOL (entry 3) led to decreases in the yield (9%) and the enantioselectivity (88% ee). Interestingly, increasing the loading of bromo-BINOL (entry 4) to 30 mol % led to a high yield (73%) and selectivity (94% ee). When bulky  $\gamma$ -substituents were employed in the BINOL catalyst (entry 5), both the yield and selectivity strongly declined. Application of the parent BINOL as the catalyst gave a low yield (4%) and relatively low selectivity (72% ee). When a commercially available alkenylboronic acid was used as the substrate (entry 7), the reaction proceeded in poor yield (18%) but with excellent selectivity (97% ee). When EtOH was replaced by *i*-PrOH (entry 8), the yield dropped (44%) but the selectivity was still high (96% ee). In the absence of EtOH (entry 9), a complex reaction mixture was obtained, from which **5a** was isolated in 4% yield with 47% ee. The poor enantioselectivity

**Table 1. Optimal Conditions for Synthesis of  $\alpha$ -CF<sub>3</sub> Allylboronic Acids<sup>a</sup>**


Entry	Change	NMR yield <sup>b</sup> (%)	Yield <sup>c</sup> (%)	ee (%)
1	No change	>95	69	98
2	10 mol% of catalyst <b>4</b>	22	12	96
3	20 mol% (R)-3,3'-Dibromo-1,1'-bi-2-naphthol instead of <b>4</b>	34	9	88
4	30 mol% (R)-3,3'-Dibromo-1,1'-bi-2-naphthol instead of <b>4</b>	>95	73	94
5	20 mol% (R)-3,3'-Bis(3,5-bis(trifluoromethyl)phenyl)-1,1'-bi-2-naphthol instead of <b>4</b>	8	trace	43
6	20 mol% (R)-1,1'-Bi(2-naphthol) instead of <b>4</b>	8	4	72
7	Boronic acid <b>2</b> was used as received	54	18	97
8	2 equiv. <i>i</i> -PrOH instead of EtOH	81	44	96
9	No alcohol added	14 <sup>d</sup>	4	47
10	No alcohol added and no catalyst	18 <sup>d</sup>	trace	0
11	No molecular sieves added	7	trace	84
12	Reaction at room temperature instead of 40 °C	9	3	96
13	Solvent is toluene instead of DCM	33	14	92

<sup>a</sup>Boroxine **2a** (0.033 mmol, equivalent to 0.1 mmol of the boronic acid), **3** (0.3 mmol), **4** (0.02 mmol, 20 mol %), and ethanol (0.2 mmol) were reacted in DCM (0.8 mL) for 48 h at 40 °C, and then DanH (0.15 mmol) was added. <sup>b</sup>Yields of **5a** determined by <sup>19</sup>F NMR spectroscopy. <sup>c</sup>Isolated yields. <sup>d</sup>A complex reaction mixture was obtained.

can be rationalized by the racemic background reaction (**2** + **3** → *rac*-**1-OR** in Figure 2). The complex reaction mixture is a consequence of the poor stability of **1** and its boroxine in the absence of EtOH. Simple aliphatic alcohols esterify the boronic acids/boroxines and thus protect them from decomposition under the reaction conditions of the borylation (Figure 1a).<sup>2,32,41</sup> When both EtOH and the BINOL catalyst were omitted (entry 10), a complex reaction mixture was obtained again. Without molecular sieves (entry 11), the yield was poor, probably because the slow formation of chiral alkenyl-BINOL-type intermediate **A** (Figure 2). At room temperature, changing dichloromethane (DCM) to toluene leads to lowering the yield and a slight decrease of the ee (entries 12–13).

Under the optimal conditions, alkyl-substituted alkenylboronic acids **2a–c** reacted readily to give the corresponding  $\alpha$ -CF<sub>3</sub> allylboronic acid esters **1(a–c)-OEt** and Bdan derivatives **5a–c** (Figure 3a). Aryl-substituted alkenylboronic acids (**2d–g**) reacted somewhat slower than the aliphatic ones. Cinnamyl derivative **5d** was formed in 54% yield (93% ee) when 20 mol % catalyst was used. However, with 20 mol % catalyst, **5e** formed only in 26% yield (89% ee). Therefore, the catalyst loading was increased to 30 mol % to obtain acceptable yields of **5e–g** (50–70%). The absolute configuration of crystalline **5e** was determined to be *S* by X-ray diffraction. On the basis of the structural similarities of the substrates and the reaction conditions, we assumed that the absolute configuration of the other species (**5a–d**, **5f**, and **5g**) was the same. The reactions can be easily scaled up. For example, the synthesis of **5a** on 1 and 2 mmol scales occurred with 98 and 96% ee in 78 and 68% yield, respectively.

The transient allylboron compounds **1-OEt** reacted with aldehyde **6a** in situ (Figure 1b). The enantioselectivity for the formation of **7a–d** varied between 90 and 98% ee. In addition, only one of the four possible diastereomers was formed in each case. We did not detect any *Z* isomer of **7a–e** in the crude product of the reaction. Usually,  $\alpha$ -substituted allylboron



**Figure 3.** Synthesis and applications of chiral  $\alpha$ -substituted allylboronic acids. <sup>a</sup>1 mmol scale. <sup>b</sup>2 mmol scale. <sup>c</sup>30 mol % 4 was used. <sup>d</sup>At 30 °C. <sup>e</sup>6b (0.15 mmol) was used.

compounds with bulky protecting groups (e.g., pinacol or 9-BBN) react with poor *E/Z* selectivity in allylboration reactions.<sup>42,43</sup>

These selectivity issues can often be solved by application of additives, but in the presented processes, poor *E/Z* selectivity was avoided by the small size of the B(OEt)<sub>2</sub> group. Notably, small molecules with alkenyl-CF<sub>3</sub> motifs<sup>44–48</sup> are very important drugs, such as in anticancer agents,<sup>25</sup> pesticides,<sup>26</sup> and herbicides<sup>27</sup> (Figure 1d). Formation of 7e from 1f proceeded with 75% ee. The relatively low enantioselectivity is a consequence of the fact that 1f-OEt is formed with lower selectivity (86% ee, 5f) than other allylboronic acids. The

yields are in the range of 41–60% based on alkenylboronic acid monomers after a two-step process. Another useful reaction is the stereoselective in situ oxidation of the chiral allylboron compounds to the corresponding  $\alpha$ -CF<sub>3</sub> allylic alcohols 8a–c (Figure 3c), which were obtained in 50–78% yield with 90–99% ee. The corresponding trifluoroethanol motif<sup>49,50</sup> occurs for example in antitumor agent Z<sup>28</sup> and the monoamine oxidase inhibitor bexloxtone (Figure 1d).<sup>29</sup>

The asymmetric homologation concept can also be extended to the synthesis of chiral  $\alpha$ -silyl allylboronic acids, such as 1h-OEt (Figure 3d). In this reaction, 3 was replaced with TMS-diazomethane. Dan protection of 1h-OEt failed, and therefore, we isolated pinane derivative 9. The homologation affording 9 proceeded with high selectivity (99:1 d.r., corresponding to 98% ee for 1h-OEt) in 51% yield. In situ allylboration of 6a gave homoallylic alcohol 10 with high selectivity.

We were able to obtain purified oxygen-sensitive allylboronic acids such as 1a and 1d by hydrolysis of the corresponding isolated Dan-protected products (5a and 5d) (Table 2). The increased reactivity of the purified products unleashed the outstanding synthetic potential of chiral allylboronic acids. As we reported previously, in the presence of molecular sieves (or other drying agents), pure allylboronic acids form very reactive allylboroxines.<sup>2,30,31,37</sup> Purified 1a in the presence of molecular sieves reacted with 6a in just 10 min to afford 7a (Table 2, entry 1). Notably, the enantioselectivities with purified and in situ-formed 1a were identical. This was also confirmed by the reaction of cinnamyl analogue 1d with 6a (entry 2). Allylboration of 6b with in situ-generated 1a-OEt failed to give 7f (Figure 3b). However, purified 1a in the presence of molecular sieves gave 1a-boroxine (see the Supporting Information), which reacted with 6b to afford 7f (entry 3) with excellent selectivity (98% ee) in 67% yield. The purification (1a-OEt → 5a → 1a sequence) is essential to obtain 7f, as demonstrated by a control experiment (entry 4). When 2 equiv of EtOH was added to 1a prior to addition of 6b, formation of 7f was not observed. Likewise, 1a-Bpin did not react with 6b under the reaction conditions applied for 1a (entry 5). Aliphatic ketones (6c–e) also reacted smoothly with allylboronic acids. Cyclohexanones 6c and 6d gave the corresponding products 7g and 7h with 91–97% ee in 50–72% yield (entries 6 and 7). The reaction of racemic methyl cyclohexanone 6d with 1d is spectacular, as in this reaction the major enantiomer (97% ee) 7h was formed with three contiguous stereocenters in a single reaction step. Acyclic aliphatic ketone 6e reacted in high yield (72%) but with only 82% ee, affording densely functionalized tertiary homoallyl alcohol 7i. The synthetic utility of purified chiral allylboronic acids was further demonstrated by allylboration of indoles<sup>11,31,51,52</sup> 6f and 6g with 1d to afford 7j and 7k with high selectivities (entries 9 and 10). From skatole 6g, the *E*-alkenyl-CF<sub>3</sub> product 7k with three adjacent stereocenters was formed with 89% ee. Isoquinoline derivative<sup>31,53</sup> 6h reacted with purified 1d to afford 7m with 93% ee in 54% yield. Allylboration of 6i<sup>54,55</sup> with 1a gave  $\alpha$ -amino acid derivative 7n with 98% ee in 72% yield.

In summary, we have presented a new methodology for the catalytic synthesis of chiral  $\alpha$ -CF<sub>3</sub> or  $\alpha$ -SiMe<sub>3</sub> allylboronic acids using stabilized diazomethane derivatives. The basic concept of stereoselective 1,2-borotropic migration can certainly be extended to nonstabilized diazoalkanes as well by solving the issues of electrophilic side reactions (e.g., protonation of the diazoalkanes) competing with the formation of the ate complex

**Table 2. Showcase for Application of Purified  $\alpha$ -CF<sub>3</sub> Allylboronic Acids in Stereoselective Synthesis<sup>a</sup>**

Entry	Boronic acid	Electrophile	Product	Time (h)	Yield <sup>b</sup> (%)	ee (%)
1				0.13	43	98
2				4	53	93
3				18	67	98
4				18	no reaction	
5				18	no reaction	
6				18	72	91
7				18	50	97
8				18	72	82
9				18	62	93
10				18	48	89
11				18	54	93
12				18	72	98

<sup>a</sup>Unless otherwise stated, **5** (0.1 mmol) was hydrolyzed with 3 N H<sub>2</sub>SO<sub>4</sub> in DME, and then **1** was extracted with toluene under Ar. <sup>b</sup>Isolated yields. <sup>c</sup><sup>1</sup>H NMR spectra of **1a** and **1d** are given in the Supporting Information. <sup>d</sup>Using 2 equiv of EtOH without molecular sieves. <sup>e</sup>**1a-Bpin** and **6b** were stirred at 40 °C without molecular sieves.

(B). The enantioenriched  $\alpha$ -CF<sub>3</sub> and  $\alpha$ -SiMe<sub>3</sub> allylboronic acids readily undergo in situ allylboration with aldehydes or can be converted to the corresponding allylic alcohols with high levels of chirality transfer. The purified chiral allylboronic acids are very reactive and highly stereoselective reagents in the allylation of aldehydes, ketones, imines, and indoles. Very promising application areas for these types of allylboronic acids are in drug design (Figure 1d) and natural product synthesis.<sup>56–60</sup>

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.0c09923>.

Materials and methods, characterization data, and NMR spectra (PDF)

Crystallographic data for **5e** (CIF)

## AUTHOR INFORMATION

### Corresponding Author

Kálmán J. Szabó – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91 Stockholm, Sweden; [orcid.org/0000-0002-9349-7137](https://orcid.org/0000-0002-9349-7137); Email: [kalman.j.szabo@su.se](mailto:kalman.j.szabo@su.se)

### Authors

Sybrand J. T. Jonker – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91 Stockholm, Sweden; [orcid.org/0000-0002-5402-8418](https://orcid.org/0000-0002-5402-8418)

Ramasamy Jayarajan – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91 Stockholm, Sweden; [orcid.org/0000-0002-1848-1434](https://orcid.org/0000-0002-1848-1434)

Tautvydas Kireilis – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91 Stockholm, Sweden

Marie Deliaval – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91 Stockholm, Sweden; [orcid.org/0000-0003-1396-2818](https://orcid.org/0000-0003-1396-2818)

Lars Eriksson – Department of Materials and Environmental Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91 Stockholm, Sweden

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.0c09923>

### Author Contributions

<sup>§</sup>S.J.T.J. and R.J. contributed equally.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

The authors thank the Knut and Alice Wallenberg Foundation (2018.0066) and the Swedish Research Council (2017-04235) for financial support. A postdoctoral fellowship for R.J. from the Wenner-Gren Foundation (UPD2019-0149) is gratefully acknowledged. We thank Dr. D. Wang for performing some preliminary experiments.

## REFERENCES

- (1) Diner, C.; Szabó, K. J. Recent Advances in the Preparation and Application of Allylboron Species in Organic Synthesis. *J. Am. Chem. Soc.* **2017**, *139*, 2–14.
- (2) Raducan, M.; Alam, R.; Szabó, K. J. Palladium-Catalyzed Synthesis and Isolation of Functionalized Allylboronic Acids: Selective, Direct Allylboration of Ketones. *Angew. Chem., Int. Ed.* **2012**, *51*, 13050–13053.
- (3) Larsson, J. M.; Szabó, K. J. Mechanistic Investigation of the Palladium-Catalyzed Synthesis of Allylic Silanes and Boronates from Allylic Alcohols. *J. Am. Chem. Soc.* **2013**, *135*, 443–455.
- (4) Matteson, D. S.; Sadhu, K. M. Boronic ester homologation with 99% chiral selectivity and its use in syntheses of the insect pheromones (3*S*,4*S*)-4-methyl-3-heptanol and exo-brevicomin. *J. Am. Chem. Soc.* **1983**, *105*, 2077–2078.

- (5) Matteson, D. S. Boronic Esters in Asymmetric Synthesis. *J. Org. Chem.* **2013**, *78*, 10009–10023.
- (6) Stymiest, J. L.; Bagutski, V.; French, R. M.; Aggarwal, V. K. Enantiodivergent conversion of chiral secondary alcohols into tertiary alcohols. *Nature* **2008**, *456*, 778–782.
- (7) Hesse, M.; Essafi, S.; Watson, C.; Harvey, J.; Hirst, D.; Willis, C.; Aggarwal, V. K. Highly Selective Allylboration of Aldehydes Using  $\alpha,\alpha$ -Disubstituted Allylic Pinacol Boronic Esters. *Angew. Chem., Int. Ed.* **2014**, *53*, 6145–6149.
- (8) Nandakumar, M.; Rubial, B.; Noble, A.; Myers, E. L.; Aggarwal, V. K. Ring-Opening Lithiation–Borylation of 2-Trifluoromethyl Oxirane: A Route to Versatile Tertiary Trifluoromethyl Boronic Esters. *Angew. Chem., Int. Ed.* **2020**, *59*, 1187–1191.
- (9) Poh, J.-S.; Lau, S.-H.; Dykes, I. G.; Tran, D. N.; Battilocchio, C.; Ley, S. V. A multicomponent approach for the preparation of homoallylic alcohols. *Chem. Sci.* **2016**, *7*, 6803–6807.
- (10) Bomio, C.; Kabeshov, M. A.; Lit, A. R.; Lau, S.-H.; Ehlert, J.; Battilocchio, C.; Ley, S. V. Unveiling the role of boroxines in metal-free carbon–carbon homologies using diazo compounds and boronic acids. *Chem. Sci.* **2017**, *8*, 6071–6075.
- (11) Forni, J. A.; Lau, S.-H.; Poh, J.-S.; Battilocchio, C.; Ley, S. V.; Pastre, J. C. Diastereoselective synthesis of functionalized indolines using in situ generated allyl boronic species. *Synlett* **2018**, *29*, 825–829.
- (12) Wu, C.; Bao, Z.; Xu, X.; Wang, J. Metal-free synthesis of gem-silylboronate esters and their Pd(0)-catalyzed cross-coupling with aryl iodides. *Org. Biomol. Chem.* **2019**, *17*, 5714–5724.
- (13) Argintaru, O. A.; Ryu, D.; Aron, I.; Molander, G. A. Synthesis and Applications of  $\alpha$ -Trifluoromethylated Alkylboron Compounds. *Angew. Chem., Int. Ed.* **2013**, *52*, 13656–13660.
- (14) Molander, G. A.; Ryu, D. Diastereoselective Synthesis of Vicinally Bis(trifluoromethylated) Alkylboron Compounds through Successive Insertions of 2,2,2-Trifluoroethane. *Angew. Chem., Int. Ed.* **2014**, *53*, 14181–14185.
- (15) Kan, S. B. J.; Huang, X.; Gumulya, Y.; Chen, K.; Arnold, F. H. Genetically programmed chiral organoborane synthesis. *Nature* **2017**, *552*, 132.
- (16) Huang, X.; Garcia-Borràs, M.; Miao, K.; Kan, S. B. J.; Zutshi, A.; Houk, K. N.; Arnold, F. H. A Biocatalytic Platform for Synthesis of Chiral  $\alpha$ -Trifluoromethylated Organoborons. *ACS Cent. Sci.* **2019**, *5*, 270–276.
- (17) Akiyama, S.; Kubota, K.; Mikus, M. S.; Paioti, P. H. S.; Romiti, F.; Liu, Q.; Zhou, Y.; Hoveyda, A. H.; Ito, H. Catalytic Enantioselective Synthesis of Allylic Boronates Bearing a Trisubstituted Alkenyl Fluoride and Related Derivatives. *Angew. Chem., Int. Ed.* **2019**, *58*, 11998–12003.
- (18) Morrison, R. J.; van der Mei, F. W.; Romiti, F.; Hoveyda, A. H. A Catalytic Approach for Enantioselective Synthesis of Homoallylic Alcohols Bearing a Z-Alkenyl Chloride or Trifluoromethyl Group. A Concise and Protecting Group-Free Synthesis of Mycothiazole. *J. Am. Chem. Soc.* **2020**, *142*, 436–447.
- (19) Kojima, R.; Akiyama, S.; Ito, H. A Copper(I)-Catalyzed Enantioselective  $\gamma$ -Boryl Substitution of Trifluoromethyl-Substituted Alkenes: Synthesis of Enantioenriched  $\gamma,\gamma$ -gem-Difluoroallylboronates. *Angew. Chem., Int. Ed.* **2018**, *57*, 7196–7199.
- (20) Fager, D. C.; Morrison, R. J.; Hoveyda, A. H. Regio- and Enantioselective Synthesis of Trifluoromethyl-Substituted Homoallylic  $\alpha$ -Tertiary  $\text{NH}_2$ -Amines by Reactions Facilitated by a Threonine-Based Boron-Containing Catalyst. *Angew. Chem., Int. Ed.* **2020**, *59*, 11448–11455.
- (21) Jiang, Q.; Guo, T.; Yu, Z. Copper-Catalyzed Asymmetric Borylation: Construction of a Stereogenic Carbon Center Bearing Both  $\text{CF}_3$  and Organoboron Functional Groups. *J. Org. Chem.* **2017**, *82*, 1951–1960.
- (22) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. Next Generation of Fluorine-Containing Pharmaceuticals, Compounds Currently in Phase II–III Clinical Trials of Major Pharmaceutical Companies: New Structural Trends and Therapeutic Areas. *Chem. Rev.* **2016**, *116*, 422–518.
- (23) Preshlock, S.; Tredwell, M.; Gouverneur, V. 18F-Labeling of Arenes and Heteroarenes for Applications in Positron Emission Tomography. *Chem. Rev.* **2016**, *116*, 719–766.
- (24) Jeschke, P. The Unique Role of Fluorine in the Design of Active Ingredients for Modern Crop Protection. *ChemBioChem* **2004**, *5*, 570–589.
- (25) Kung, P. P.; Meng, J. J. 2-Amino Pyridine Compounds. WO 2008/059368 A2, 2008.
- (26) Stubbs, V. K.; Wilshire, C.; Webber, L. G. Cyhalothrin—a novel acaricidal and insecticidal synthetic pyrethroid for the control of the cattle tick (*Boophilus microplus*) and the buffalo fly (*Haematobia irritans exigua*). *Aust. Vet. J.* **1982**, *59*, 152–155.
- (27) Hemelaere, R.; Desroches, J.; Paquin, J.-F. Introduction of the 4,4,4-Trifluorobut-2-ene Chain Exploiting a Regioselective Tsuji–Trost Reaction Catalyzed by Palladium Nanoparticles. *Org. Lett.* **2015**, *17*, 1770–1773.
- (28) Bernardon, J. M.; Biadatti, T. Vitamin D Analogues. WO 03/050067 A2, 2003.
- (29) Warot, D.; Berlin, I.; Patat, A.; Durrieu, G.; Zieleniuk, I.; Puech, A. J. Effects of Befloxatone, a Reversible Selective Monoamine Oxidase-A Inhibitor, on Psychomotor Function and Memory in Healthy Subjects. *J. Clin. Pharmacol.* **1996**, *36*, 942–950.
- (30) Alam, R.; Vollgraff, T.; Eriksson, L.; Szabó, K. J. Synthesis of Adjacent Quaternary Stereocenters by Catalytic Asymmetric Allylboration. *J. Am. Chem. Soc.* **2015**, *137*, 11262–11265.
- (31) Alam, R.; Diner, C.; Jonker, S.; Eriksson, L.; Szabó, K. J. Catalytic Asymmetric Allylboration of Indoles and Dihydroisoquinolines with Allylboronic Acids: Stereodivergent Synthesis of up to Three Contiguous Stereocenters. *Angew. Chem., Int. Ed.* **2016**, *55*, 14417–14421.
- (32) Zhao, J.; Jonker, S. J. T.; Meyer, D. N.; Schulz, G.; Tran, C. D.; Eriksson, L.; Szabó, K. J. Copper-Catalyzed Synthesis of Allenylboronic Acids. Access to Sterically Encumbered Homopropargylic Alcohols and Amines by Propargylboration. *Chem. Sci.* **2018**, *9*, 3305–3312.
- (33) Wu, T. R.; Shen, L.; Chong, J. M. Asymmetric Allylboration of Aldehydes and Ketones Using 3,3'-Disubstitutedbinaphthol-Modified Boronates. *Org. Lett.* **2004**, *6*, 2701–2704.
- (34) Wu, T. R.; Chong, J. M. Ligand-Catalyzed Asymmetric Alkynylboration of Enones: A New Paradigm for Asymmetric Synthesis Using Organoboranes. *J. Am. Chem. Soc.* **2005**, *127*, 3244–3245.
- (35) Lou, S.; Moquist, P. N.; Schaus, S. E. Asymmetric Allylboration of Acyl Imines Catalyzed by Chiral Diols. *J. Am. Chem. Soc.* **2007**, *129*, 15398–15404.
- (36) Jiang, Y.; Schaus, S. E. Asymmetric Petasis Borono-Mannich Allylation Reactions Catalyzed by Chiral Biphenols. *Angew. Chem., Int. Ed.* **2017**, *56*, 1544–1548.
- (37) Huang, G.; Diner, C.; Szabó, K. J.; Himo, F. Mechanism and Stereoselectivity of the BINOL-Catalyzed Allylboration of Skatoles. *Org. Lett.* **2017**, *19*, 5904–5907.
- (38) Nishiyabu, R.; Kubo, Y.; James, T. D.; Fossey, J. S. Boronic acid building blocks: tools for self assembly. *Chem. Commun.* **2011**, *47*, 1124–1150.
- (39) Wang, H.; Jing, C.; Noble, A.; Aggarwal, V. K. Stereospecific 1,2-Migrations of Boronate Complexes Induced by Electrophiles. *Angew. Chem., Int. Ed.* **2020**, *59*, 16859–16872.
- (40) Noguchi, H.; Hojo, K.; Suginome, M. Boron-Masking Strategy for the Selective Synthesis of Oligoarenes via Iterative Suzuki–Miyaura Coupling. *J. Am. Chem. Soc.* **2007**, *129*, 758–759.
- (41) Selander, N.; Kipke, A.; Sebelius, S.; Szabó, K. J. Petasis Borono-Mannich Reaction and Allylation of Carbonyl Compounds via Transient Allyl Boronates Generated by Palladium Catalyzed Substitution of Allyl Alcohols. An Efficient One-Pot Route to Stereodefined  $\alpha$ -Amino Acids and Homoallyl Alcohols. *J. Am. Chem. Soc.* **2007**, *129*, 13723–13731.
- (42) Fasano, V.; McFord, A. W.; Butts, C. P.; Collins, B. S. L.; Fey, N.; Alder, R. W.; Aggarwal, V. K. How big is the pinacol boronic ester as a substituent? *Angew. Chem., Int. Ed.* **2020**, *59*, 22403–22407.

(43) Althaus, M.; Mahmood, A.; Suárez, J. R.; Thomas, S. P.; Aggarwal, V. K. Application of the Lithiation–Borylation Reaction to the Preparation of Enantioenriched Allylic Boron Reagents and Subsequent In Situ Conversion into 1,2,4-Trisubstituted Homoallylic Alcohols with Complete Control over All Elements of Stereochemistry. *J. Am. Chem. Soc.* **2010**, *132*, 4025–4028.

(44) Koh, M. J.; Nguyen, T. T.; Lam, J. K.; Torker, S.; Hyvl, J.; Schrock, R. R.; Hoveyda, A. H. Molybdenum chloride catalysts for Z-selective olefin metathesis reactions. *Nature* **2017**, *542*, 80–85.

(45) Straathof, N. J. W.; Cramer, S. E.; Hessel, V.; Noël, T. Practical Photocatalytic Trifluoromethylation and Hydrotrifluoromethylation of Styrenes in Batch and Flow. *Angew. Chem., Int. Ed.* **2016**, *55*, 15549–15553.

(46) Hafner, A.; Bräse, S. Efficient Trifluoromethylation of Activated and Non-Activated Alkenyl Halides by Using (Trifluoromethyl)-trimethylsilane. *Adv. Synth. Catal.* **2011**, *353*, 3044–3048.

(47) Imhof, S.; Randl, S.; Blechert, S. Ruthenium catalysed cross metathesis with fluorinated olefins. *Chem. Commun.* **2001**, 1692–1693.

(48) Hong, H.; Li, Y.; Chen, L.; Li, B.; Zhu, Z.; Chen, X.; Chen, L.; Huang, Y. Electrochemical Synthesis Strategy for C<sub>vinyl</sub>-CF<sub>3</sub> Compounds through Decarboxylative Trifluoromethylation. *J. Org. Chem.* **2019**, *84*, 5980–5986.

(49) Chen, X.; Gong, X.; Li, Z.; Zhou, G.; Zhu, Z.; Zhang, W.; Liu, S.; Shen, X. Direct transfer of tri- and di-fluoroethanol units enabled by radical activation of organosilicon reagents. *Nat. Commun.* **2020**, *11*, 2756.

(50) Xu, Q.; Zhou, H.; Geng, X.; Chen, P. Nonenzymatic kinetic resolution of racemic 2,2,2-trifluoro-1-aryl ethanol via enantioselective acylation. *Tetrahedron* **2009**, *65*, 2232–2238.

(51) Ullrich, P.; Schmauck, J.; Brauns, M.; Mantel, M.; Breugst, M.; Pietruszka, J. Enantioselective Allylation of Indoles: A Surprising Diastereoselectivity. *J. Org. Chem.* **2020**, *85*, 1894–1905.

(52) Nowrouzi, F.; Batey, R. A. Regio- and Stereoselective Allylation and Crotylation of Indoles at C2 Through the Use of Potassium Organotrifluoroborate Salts. *Angew. Chem., Int. Ed.* **2013**, *52*, 892–895.

(53) Wu, T. R.; Chong, J. M. Asymmetric Allylboration of Cyclic Imines and Applications to Alkaloid Synthesis. *J. Am. Chem. Soc.* **2006**, *128*, 9646–9647.

(54) Jonker, S. J. T.; Diner, C.; Schulz, G.; Iwamoto, H.; Eriksson, L.; Szabó, K. J. Catalytic asymmetric propargyl- and allylboration of hydrazonoesters: a metal-free approach to sterically encumbered chiral  $\alpha$ -amino acid derivatives. *Chem. Commun.* **2018**, *54*, 12852–12855.

(55) Fujita, M.; Nagano, T.; Schneider, U.; Hamada, T.; Ogawa, C.; Kobayashi, S. Zn-Catalyzed Asymmetric Allylation for the Synthesis of Optically Active Allylglycine Derivatives. Regio- and Stereoselective Formal  $\alpha$ -Addition of Allylboronates to Hydrazono Esters. *J. Am. Chem. Soc.* **2008**, *130*, 2914.

(56) Touré, B. B.; Hall, D. G. Natural Product Synthesis Using Multicomponent Reaction Strategies. *Chem. Rev.* **2009**, *109*, 4439–4486.

(57) Daniels, B. E.; Ni, J.; Reisman, S. E. Synthesis of Enantioenriched Indolines by a Conjugate Addition/Asymmetric Protonation/Aza-Prins Cascade Reaction. *Angew. Chem., Int. Ed.* **2016**, *55*, 3398–3402.

(58) Lu, Z.; Yang, M.; Chen, P.; Xiong, X.; Li, A. Total Synthesis of Hapalindole-Type Natural Products. *Angew. Chem., Int. Ed.* **2014**, *53*, 13840–13844.

(59) Tang, Q.; Fu, K.; Ruan, P.; Dong, S.; Su, Z.; Liu, X.; Feng, X. Asymmetric Catalytic Formal 1,4-Allylation of  $\beta,\gamma$ -Unsaturated  $\alpha$ -Ketoesters: Allylboration/Oxy-Cope Rearrangement. *Angew. Chem., Int. Ed.* **2019**, *58*, 11846–11851.

(60) Trost, B. M.; Cregg, J. J.; Hohn, C.; Bai, W.-J.; Zhang, G.; Tracy, J. S. Ruthenium-catalysed multicomponent synthesis of the 1,3-dienyl-6-oxy polyketide motif. *Nat. Chem.* **2020**, *12*, 629–637.