




Article

Religious Affiliations Influence Health-Related and General Decision Making: A Brazilian Nationwide Survey

Marcelo Borges ^{1,*} , Giancarlo Lucchetti ², Frederico C. Leão ¹, Homero Vallada ^{1,3,4} and Mario F. P. Peres ^{1,4,5}

¹ PROSER, Programa de Espiritualidade e Religiosidade, Instituto de Psiquiatria, Hospital das Clinicas HCFMUSP, Faculdade de Medicina, Universidade de Sao Paulo, Sao Paulo 05403-903, Brazil; leaofc@gmail.com (F.C.L.); hvallada@usp.br (H.V.); mariop3r3s@gmail.com (M.F.P.P.)

² School of Medicine, Federal University of Juiz de Fora, Juiz de Fora 36038-330, Brazil; g.lucchetti@yahoo.com.br

³ Departamento & Instituto de Psiquiatria (LIM-23) HCFMUSP, Sao Paulo 05403-903, Brazil

⁴ Faculdade de Medicina FMUSP, Universidade de Sao Paulo, Sao Paulo 05403-903, Brazil

⁵ Hospital Israelita Albert Einstein, São Paulo 05403-903, Brazil

* Correspondence: borgespsi@icloud.com

Abstract: Approximately 90% of the world's population is involved in some spiritual/religious practice, and this dimension has a relevant role in life. Many studies demonstrate the associations between spirituality/religiosity (S/R), and physical, mental, and social health. Systematic reviews have indicated positive associations; however, the mechanisms behind religious coping are not fully understood. The present study aimed to examine the role of religious affiliation in general (ordinary) and health-related decisions. A nationwide, population-based, cross-sectional study was conducted in Brazil using a self-administered online survey. How much religious affiliation influences decision making was investigated. A total of 1133 participants were included, who were classified as Catholics (43.9%), Evangelicals (18.7%), spiritualists (12.8%), non-religious (11.9%), and others (12.7%). Most participants (66.5%) believed that their religious affiliations had moderate to high influences on their decisions. Participants rated the influence as high in marriage (62.7%), in donations (60.1%), in volunteering (55%), in friendships (53.9%), and in work (50.5%). Concerning health-related decisions, the influence was rated as high in drug use (45.2%), in accepting medical recommendations (45%), and in smoking (43.2%). The influence of religious affiliation on general decision making was significantly correlated for dietary restrictions ($r = 0.462$), alcohol consumption ($r = 0.458$), drug use ($r = 0.441$), tobacco consumption ($r = 0.456$), and refusal of medical recommendations ($r = 0.314$). Improving the understanding of how a patient's beliefs, practices, and experiences affect their health may help healthcare practitioners to take into account religious considerations, not only regarding influences on habits but also regarding adherence to medical treatment advice.

Keywords: nationwide; online survey; religion; sensitive information; public health



Citation: Borges, M.; Lucchetti, G.; Leão, F.C.; Vallada, H.; Peres, M.F.P. Religious Affiliations Influence Health-Related and General Decision Making: A Brazilian Nationwide Survey. *Int. J. Environ. Res. Public Health* **2021**, *18*, 2873. <https://doi.org/10.3390/ijerph18062873>

Academic Editor: Paul B. Tchounwou

Received: 12 February 2021

Accepted: 5 March 2021

Published: 11 March 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Approximately 90% of the world's population is involved in some spiritual/religious practice, and this dimension plays a relevant role in life [1]. Evidence of spirituality and religiousness's effect on health has been constantly growing. Approximately 30,000 articles were published in the last 15 years in this field of research according to a bibliometric analysis [2].

Many studies have demonstrated associations between spirituality/religiosity (S/R) and physical, mental, and social health [2]. Systematic reviews have shown that most studies indicated some positive association in which the S/R was a protective factor, but some studies found negative or neutral associations [3,4]. Many professional organizations recognize that spiritual care is an important component of healthcare, such the World

Health Organization (WHO). The Joint Commission on the Accreditation of Healthcare Organizations (JCAHO) [5] recommended including S/R in clinical care and education. Concerning mental health, the World Psychiatric Association, American Psychological Association [6], and Royal College of Psychiatrists [7] have sections dedicated to S/R issues. The way patients experience the S/R dimension can be either protective or harmful, depending on how these resources are used in the health/disease process (religious coping). The mechanisms behind religious coping are not fully understood, but religion is a relevant aspect of an individual's cultural background, important not only in daily decisions (e.g., dating, friendship, and clothing choices) but also health-related attitudes (e.g., diet, smoking, and drug use). Beliefs, practices, and experiences can influence decisions involving medical treatment. Religiousness is associated with adherence, giving meaning to an illness [8], and care to one's body [9]. Religiousness can be associated with acceptance, refusal, or rejection of medical recommendations [10]. Religions significantly influence important health habits (use/abuse of alcohol [11], drugs, and smoking [12]). Since patient care is influenced by cultural and religious beliefs, assessing how much a particular religious affiliation influences decision making could contribute to a comprehensive understanding of the amount or the magnitude of this relationship.

On the one hand, studies in this field have not been designed to consider patients' perceptions about religion [13], and the basis may start with data being extracted from nationwide censuses not primarily designed to assess religion's scientific aspects in a broader way [14]. On the other hand, most studies were conducted in specific populations [15]. Sometimes, religious affiliation was compared with non-religiousness [16]. The majority of studies have been carried out in the United States and maybe are not fully applicable in other contexts and cultures [17]. Previous international studies have already used online surveys to investigate spirituality and religion in the general population [18,19], and the online approach is still mostly used for health professionals [20,21]. On the specific subject of religious affiliation, no study was found that investigates the patient's perception of its influence on daily or health-related decision making.

Therefore, this study aimed to evaluate how much religious affiliation influences daily life and health-related decisions, thereby providing to healthcare providers more information to address religious considerations and demonstrate how a patient's beliefs, practices, and experiences may affect his/her health.

2. Materials and Methods

This is a nationwide, cross-sectional, quota sampled, self-administrated online survey study. Invitations were sent by Qualtrics® Panels, one of the main companies offering online instrument services [22], for the project "Spiritual and Religious Beliefs, Practices and Experiences in the General Population." It was sponsored by the Interfaith Coalition on Spirituality and Health (coalizaointerfe.org, accessed on 10 September 2020), a Brazilian institution composed of representative members of several religious affiliations, and non-religious faith and professional healthcare practices in Brazil. Data were collected between 28 June 2016 and 22 August 2016, through a self-administered online survey. Respondents were invited to participate and complete the online survey, and those who agreed signed an online consent term.

The sample size was determined by non-probability quota samples, thereby assuring that distinct subgroups of the population and their characteristics were represented. Quotas were set to limit the respondents according to social class distribution, age, gender, and geographic location, so the population surveyed could meet the same profile of the general adult population in Brazil, according to the 2010 census. A previous article [14] showed that using this sampling procedure could result in accurate representative surveys for Brazil as a whole. Inclusion criteria were 18 years old or more, and online access to a website, email, or social media network. Missing background data, invalid email addresses, incomplete answers, and missing responses were exclusion criteria. The survey took less than 30 min and attention filters and quality check questions were used.

A question about religious affiliation was asked: "What is the best option below to define your current religious affiliation?" Response options included 16 options: "Catholic," "Evangelical," "Kardec Spiritist," "Jewish," "Buddhist," "Umbanda," "Candomble," "atheist," "agnostic," "spiritualist," "Christian," "Jehovah's Witness," "Seicho-no-ie," "Wicca," "no religion," and "others." Due to the limited number of responders in some affiliations, we re-allocated the sample into 6 groups: "Catholics," "Evangelicals," "spiritualists" (Kardec spiritists and spiritualists), "other Christians," "non-religious" (agnostic, atheist, and no religion), and "other denominations" (Jewish, Buddhist, Umbanda, Candomble, Jehovah's Witness, Seicho-no-ie, Wicca, and others). Kardec Spiritists and spiritualists supposedly behave similarly. Although "others" seems heterogeneous, their sample size was still small, justifying their being analyzed together.

The self-reported influence of religious affiliation on general life was assessed by the question: "How much does religious affiliation influence your decisions?" The self-reported influence of religious affiliation on various attitudes was examined by: "How much does your religious affiliation influence your": "dating," "marriage," "friendships," "work," "clothing choices," "volunteering," "donations," and "politics." The self-reported influence of religious affiliation on health-related attitudes was examined by: "How much does your religious affiliation influence" your decisions regarding "diet," "alcohol consumption," "drug use," "tobacco consumption," "refusal of medical recommendations," and "acceptance of medical recommendations"?

All questions presented 5 alternatives: 1 (very little), 2 (a little bit), 3 (moderately), 4 (quite a lot), and 5 (a lot). Analysis was performed with 95% confidence intervals. For correlations, we used Pearson tests; for comparisons, paired *t*-tests for normal samples. The Wilcoxon test was used for non-parametric comparisons, and Fisher and chi-square tests were used to compare proportions (IBM SPSS Statistics version 25). The study followed international ethical regulations (Declaration of Helsinki principles) and was approved by the local Ethics Research Committee of IPq HCFMUSP (CAAE 64956717.4.0000.0065). All participants gave informed consent online.

3. Results

From 1252 participants, 1131 (90.3%) completed the selected questions. The sample was composed of 536 (47.4%) men and 595 (52.6%) women, with a mean age of 40.5; 57.9% of the participants were married and 37.8% had a monthly family income between US\$800.00 and US\$2000.00 (Table 1).

The sample was categorized into six religious groups: Catholics (43.9%), Evangelicals (18.7%), spiritualists (12.8%), other Christians (6.6%), other denominations (6.1%), and non-religious (11.9%). Regarding how much their religious affiliations influenced their decision making, 21.4% of the participants reported "very little," 12.1% "a little bit," 23.2% "moderately," 26.1% "quite a lot," and 17.2% reported "a lot."

Religious affiliation correlated with daily decisions to various degrees: "dating" ($r = 0.530$), "marriage" ($r = 0.551$), "friendships" ($r = 0.565$), "job" ($r = 0.571$), "clothing choices" ($r = 0.507$), "volunteering" ($r = 0.568$), "donations" ($r = 0.590$), and "politics" ($r = 0.349$). Likewise, health-related decisions correlated as follows: "diet" ($r = 0.462$), "alcohol consumption" ($r = 0.458$), "drug use" ($r = 0.441$), and "tobacco consumption" ($r = 0.456$). "Acceptance of medical recommendations" has a correlation of 0.560. "Refusal of medical recommendations" presented as $r = 0.314$, demonstrating that the influences of religious affiliation were lower on health-related aspects, as detailed by religious affiliation categories (Table 2).

Table 1. Descriptive characteristics of the sample.

Variables	Categories	Catholics	Evangelicals	Spiritualists	Other	Other Christians	Non-Religious
		Total (n = 496) n(%)	Total (n = 411) n(%)	Total (n = 145) n(%)	Total (n = 69) n(%)	Total (n = 75) n(%)	Total (n = 135) n(%)
Gender	Female	235 (47.4)	122 (57.8)	94 (64.8)	36 (56.2)	45 (60)	63 (43.7)
	Male	261 (52.6)	89 (42.2)	51 (35.2)	33 (47.8)	30 (40.0)	72 (53.3)
Age Range (years)	18–25	116 (23.4)	48 (22.7)	28 (19.3)	10 (14.5)	24 (32.0)	30 (22.2)
	26–35	93 (18.8)	48 (22.7)	31 (21.4)	14 (20.3)	18 (24.0)	27 (20.0)
	36–45	103 (20.8)	46 (21.8)	37 (25.5)	16 (23.2)	8 (10.7)	24 (17.8)
	46–55	80 (16.2)	22 (10.4)	23 (15.9)	18 (26.1)	12 (16.0)	22 (16.3)
	56–65	72 (14.5)	34 (16.1)	14 (9.7)	7 (10.1)	9 (12.0)	18 (13.3)
	>65	31 (6.3)	13 (6.2)	12 (8.3)	4 (5.8)	4 (5.3)	14 (10.4)
Educational Level	Incomplete high school	18 (3.6)	16 (7.6)	5 (3.4)	9 (13)	2 (2.7)	7 (5.2)
	Completed high school	192 (38.7)	105 (49.8)	41 (28.3)	33 (47.8)	29 (38.7)	52 (38.5)
	Completed university	193 (38.9)	61 (28.9)	60 (41.4)	19 (27.5)	30 (40.0)	62 (45.9)
	Postgraduated	93 (18.8)	29 (13.7)	39 (26.9)	8 (11.6)	14 (18.7)	14 (10.4)
Marital Status	Single	149 (30.0)	56 (26.5)	43 (29.7)	24 (34.8)	29 (38.7)	74 (54.8)
	Judicially separated	12 (2.4)	6 (2.8)	3 (2.1)	0 (0)	0 (0)	4 (3.0)
	Divorced	28 (5.6)	7 (3.3)	6 (4.1)	6 (8.7)	3 (4.0)	7 (5.2)
	Married	300 (60.5)	138 (65.4)	88 (60.7)	38 (55.1)	43 (57.3)	48 (35.6)
	Widowed	7 (1.4)	4 (1.9)	5 (3.4)	1 (1.4)	0 (0)	2 (1.5)
Monthly income range (USD) *	>4114	46 (10.6)	14 (8.1)	12 (10.2)	5 (9.3)	5 (7.7)	8 (6.8)
	2058–4114	132 (30.5)	36 (20.8)	36 (30.5)	10 (18.5)	14 (21.5)	31 (26.5)
	823–2057	158 (36.5)	61 (35.3)	48 (40.7)	18 (33.3)	28 (43.1)	50 (42.7)
	411–822	71 (16.4)	40 (23.1)	16 (13.6)	17 (31.5)	10 (15.4)	20 (17.1)
	<411	26 (6.0)	22 (12.7)	6 (5.1)	4 (7.4)	8 (12.3)	8 (6.8)
Race/Color	White	325 (65.5)	107 (50.7)	95 (65.5)	38 (55.1)	34 (45.3)	94 (69.6)
	Black	19 (3.8)	15 (7.1)	9 (6.2)	4 (5.8)	3 (4.0)	11 (8.1)
	Asian	12 (2.4)	6 (2.8)	2 (1.4)	3 (4.3)	2 (2.7)	4 (3.0)
	Pardo (mixed)	138 (27.8)	81 (38.4)	38 (26.2)	22 (31.9)	36 (48)	25 (18.5)
	Indigenous	2 (0.4)	2 (0.9)	1 (0.7)	2 (2.9)	0 (0)	1 (0.7)
Region	Northeast	144 (29)	51 (24.2)	44 (30.3)	18 (26.1)	18 (24)	33 (24.4)
	South	66 (13.3)	34 (16.1)	23 (15.9)	11 (15.9)	9 (12)	18 (13.3)
	Midwest	37 (7.5)	13 (6.2)	10 (6.9)	4 (5.8)	9 (12)	16 (11.9)
	Southeast	213 (42.9)	93 (44.1)	62 (42.8)	27 (39.1)	35 (46.7)	60 (44.4)
	North	36 (7.3)	20 (9.5)	6 (4.1)	9 (13)	4 (5.3)	8 (5.9)

* Conversion from Brazilian Real using the mid-price for the dollar in July 2018 (1 USD = BRL 3.83).

Table 2. Correlations between generic questions, daily decisions, and health-related questions.


	Catholics	Evangelicals	Spiritualists	Other	Other Christians	Non Religious
Dating	0.50 *	0.41 *	0.36 *	0.55 *	0.49 *	0.34 *
Marriage	0.51 *	0.44 *	0.39 *	0.46 *	0.53 *	0.37 *
Friendship	0.59 *	0.41 *	0.37 *	0.48 *	0.43 *	0.50 *
Job	0.62 *	0.34 *	0.41 *	0.42 *	0.48 *	0.55 *
Dress	0.50 *	0.40 *	0.26 *	0.44 *	0.46 *	0.60 *
Volunteers	0.57 *	0.34 *	0.43 *	0.61 *	0.64 *	0.52 *
Donation	0.62 *	0.37 *	0.43 *	0.56 *	0.56 *	0.52 *
Politics	0.35 *	0.22 *	0.29 *	0.20 *	0.44 *	0.30 *
Diet	0.47 *	0.32 *	0.32 *	0.49 *	0.32 *	0.53 *
Alcohol	0.39 *	0.34 *	0.43 *	0.52 *	0.34 *	0.44 *
Drug use	0.38 *	0.29 *	0.40 *	0.37 *	0.45 *	0.33 *
Tobacco	0.36 *	0.36 *	0.49 *	0.43 *	0.50 *	0.33 *
Refuse Medical	0.30 *	0.19 *	0.29 *	0.27 **	0.33 *	0.54 *
Accept Medical	0.61 *	0.31 *	0.50 *	0.59 *	0.46 *	0.50 *

* Pearson’s correlation is significant at the 0.01 level (2-tailed); ** Pearson’s correlation is significant at the 0.01 level (1-tailed). Correlation Level: High, [dark grey]; Medium, [medium grey]; Weak, [light grey]; Very weak, [white].

Most participants’ (66.5%) believe that their religious affiliations have moderate to high influences on their general decisions. Participants rated the influence of religious affiliation as high on marriage (62.7%), on donations (60.1%), on volunteering (55%), on friendship (53.9%), on work (50.5%), on dating (49.6%), on choice of clothing (42%) and on politics (27.2%) (Figure 1).

Concerning health-related decisions, the influence of religious affiliation was high on drug use (45.2%), on accepting medical recommendations (45%), on smoking (43.2%), on alcohol consumption (39.7%), on dietary restrictions (37.1%), and on refusing medical advice (22.6%) (Figure 2).

Evangelicals had the highest levels of religious influence within daily life and health-related decisions, and as expected, the non-religious group had the least influence from religion (though they were referring to the influences of beliefs on life decisions).



Low = “very little” + “little bit”
 Moderate = “moderately”
 High = “quite a lot”+ “very much”
 Self-reported % of How much religious affiliation influences specific daily life decision making

Figure 1. Religious affiliation’s influences in daily decisions.




Figure 2. Religious affiliation’s influences in health-related decisions.

4. Discussion

Our data show religious affiliation influences daily life and health-related decisions. Overall, the influence of religious affiliation on general decisions was considered high by the participants, particularly in areas such as marriage, donations, friendships, and work. However, this influence is also considerable in health-related decisions, such as the refusal of medical advice, diet, smoking, and alcohol use. These findings provide further support to the scientific literature, which has been showing that religion is a relevant aspect of individuals’ cultural background [1], important not only in daily life but also in health-related decisions and attitudes [2]. Additionally, the way patients experience the S/R dimension can be either protective or harmful, depending on how these resources are used [23,24].

Historically, religions incorporated forms of fasting and self-starvation. Many important religious figures, such as Moses, Buddha, and Jesus abstained from eating as part of ascetic/mystic practices. A controlled trial study found evidence of treatment via spirituality-oriented groups based on the eating attitude test (EAT), and eating disorder specific symptoms, such as bingeing, purging, restricting, and attitudes about food and dieting [25]. That was corroborated in a previous correlational study [26]. In our study, 63% of participants reported a small or very small influence of religion on diet, as opposed to 37% reporting a moderate influence or above. Although the majority said religion does not affect their dietary patterns, it is very important to consider that one-third of the general population says the opposite. Several health problems are sensitive to dietary patterns—obesity, hypertension, diabetes, and dyslipidemia, not to mention less prevalent disorders. Clinicians should be aware of a patient’s religiosity so a treatment plan could be tailored and would not be compromised by their adherence to dietary recommendations.

Studies have demonstrated a relationship between abstinence and alcohol and use/abuse [27], but also pointed out the difficulty of identifying which factors of the

spiritual dimension are present in this mediation. Increased S/R involvement is related to lower alcohol consumption and better outcomes in alcohol abuse treatments. Although religious affiliation does not define alcohol-related attitudes, it may contribute to abuse [28]. Attitudes toward alcohol consumption in this study showed 39.7% of individuals rating religious influence as moderate and above. This finding supports the idea that religion is a mediator, suppressing alcohol consumption. One may hypothesize that this is due to the social conduct rules of one's religion, but it could be related to spiritual aspects too. Concerning drug use, an epidemiological study conducted in the U.S. presented a higher risk of having started the use of cannabis, cocaine, or other extra-medical drugs for the non-religious. Additionally, those who declared less importance for religion in life were more likely to use drugs [29]. A causal-comparative drug abuse study demonstrated neutral effects of religion on addiction; however, the sample had high religiosity and commitment to beliefs [30]. Studies with minorities showed religiosity to be a protective factor [31]. Regarding the influence on drug use, this study demonstrated 45.2% of participants rating it as moderately important or more so.

For tobacco consumption, a nationwide study performed in the United Kingdom [12] indicated that the prevalence of tobacco use is higher among people who identified themselves to be without religion (66.2%) than those who declared themselves to be Christians (60.0%) and relatively lower among those who declared themselves to be Muslims (35.2%). In this study, only 9.6% of non-religious people rated their beliefs as having an influence of moderate and above that; 57.3% of Christians and 62.1% of Evangelicals declared an influence of moderate or above. However, the mechanisms involved in this process still require further study. Among the possible reasons is the fact that many religious denominations prohibit the use of substances such as tobacco; the time spent in activities related to the S/R dimension has decreased risk behaviors; religious affiliation can increase the likelihood of confrontations regarding one's character; and participation in a religious institution has the characteristic of social involvement, which can discourage such behaviors [32].

Concerning the influence of religion on following or refusing medical advice, one study showed trust in physicians and the health system. A nationwide U.S. study showed a positive association with adherence to treatment and strength of religious affiliation [33]. A case series study investigated patients, noting that the S/R dimension helped adherence to treatment, indicating a positive association [9]. Another nationwide study with 41-year-olds and older showed that high religious attendance (more than a weekly) granted a 40% reduced hazard of mortality compared with those who never attended, and there was a positive association between religious attendance and the importance of religion to the respondent. However, higher levels of religiousness are by no means always associated with better health. Our findings show amounts of influence of "very little," (40.5%), "a little bit" (14.5%), "moderately" (22.2%), "quite a lot" (15.1%), and "a lot" (7.7%) on accepting medical recommendations. However, for denying or refusal of medical recommendations, the results were: "very little" (63.2%), "a little bit" (14.1%), "moderately" (13.0%), "quite a lot" (5.7%), and "a lot" (4.0%).

Thus, awareness about the global role of S/R in health has been increasing [34]. However, nationwide studies frequently do not have a primary design to assess religious issues as a main topic [14]. This study demonstrated how religious affiliation influences attitudes toward daily life and health-related decisions. It may have clinical relevance for healthcare professionals. A good approach needs awareness of religions to avoid unhelpful judgment and preconceptions [35], to increase trust with each patient.

This study had some limitations that should be considered. First, this was a cross-sectional survey, avoiding inferences about cause and effect. Second, data collection was carried out only for persons 18 years old or older who had online access to email, social media, and websites; it is possible that some groups of the population, such as residents from rural areas, those with lower socioeconomic levels, and older adults, were included at a lower frequency. Finally, we used self-reported measures to investigate the influences of religious affiliation on daily decisions, instead of evaluating their behavior in real-life

scenarios. However, previous international studies have already used online surveys to investigate spirituality and religion in the general population [18,19] and the online approach is still mostly used by health professionals [20,21].

5. Conclusions

In conclusion, religious affiliation influences daily life and health-related decisions to different degrees. Healthcare providers need to consider religious aspects of patients' lives to consider how a patient's beliefs, practices, and experiences may affect their health, not only because of religion's influences on habits such as diet, smoking, drinking, and drug use, but also regarding adherence to medical advice.

Author Contributions: M.B.—substantial contributions to the conceptualization, design of the work, analysis, interpretation of data, drafting of the manuscript, and revision of the paper for important intellectual content. G.L.—substantial contributions to the conceptualization, analysis, interpretation of data, drafting of the manuscript, and revision of the paper for important intellectual content. F.C.L.—substantial contributions to the conceptualization, drafting of the manuscript, and revision of the paper for important intellectual content. H.V.—substantial contributions to the conceptualization, drafting of the manuscript, and revision of the paper for important intellectual content. M.F.P.—substantial contributions to the conceptualization, data acquisition, analysis, interpretation of the data for the work, drafting of the manuscript, and revision of the paper for important intellectual content. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study followed international ethical regulations (Declaration of Helsinki principles) and was approved by local Ethics Research Committee of IPq HCFMUSP. All participants signed an online informed consent form.

Informed Consent Statement: Patient consent was waived because it is a public opinion survey with unidentified participants according to Brazilian policies resolution N° 510 (<http://conselho.saude.gov.br/resolucoes/2016/Reso510.pdf>, accessed on 10 September 2020).

Data Availability Statement: Project “Spiritual and Religious Beliefs, Practices and Experiences in the General Population,” was sponsored by Interfaith Coalition on Spirituality and Health (coalizaointerfe.org, accessed on 10 September 2020), a Brazilian institution composed of representative members of several religious affiliations, and non-religious faith and healthcare practices in Brazil.

Acknowledgments: Financial contribution from IHPV (Instituto Homero Pinto Vallada), São Paulo, Brazil.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Moreira-Almeida, A.; Neto, F.L.; Koenig, H.G. Religiousness and mental health: A review. *Rev. Bras. Psiquiatr.* **2006**, *36*, 176–182. [[CrossRef](#)]
2. Lucchetti, G.; Lucchetti, A.L. Spirituality, religion, and health: Over the last 15 years of field research (1999–2013). *Int. J. Psychiatry Med.* **2014**, *48*, 199–215. [[CrossRef](#)]
3. Koenig, H.G. Religion, spirituality, and health: The research and clinical implications. *ISRN Psychiatry* **2012**, *2012*, 278730. [[CrossRef](#)] [[PubMed](#)]
4. Goncalves, J.P.B.; Lucchetti, G.; Menezes, P.R.; Vallada, H. Complementary religious and spiritual interventions in physical health and quality of life: A systematic review of randomized controlled clinical trials. *PLoS ONE* **2017**, *12*, e0186539. [[CrossRef](#)]
5. Lucchetti, G.; Lucchetti, A.L.; Puchalski, C.M. Spirituality in medical education: Global reality? *J. Relig. Health* **2012**, *51*, 3–19. [[CrossRef](#)]
6. Vieten, C.; Scammell, S.; Pilato, R.; Ammondson, I.; Pargament, K.I.; Lukoff, D. Spiritual and religious competencies for psychologists. *Psychol. Relig. Spirit.* **2013**, *5*, 129–144. [[CrossRef](#)]
7. Moreira-Almeida, A.; Koenig, H.G.; Lucchetti, G. Clinical implications of spirituality to mental health: Review of evidence and practical guidelines. *Rev. Bras. Psiquiatr.* **2014**, *36*, 176–182. [[CrossRef](#)] [[PubMed](#)]
8. Borrás, L.; Mohr, S.; Brt, P.Y.; Gillieron, C.; Eytan, A.; Huguelet, P. Religious beliefs in schizophrenia: Their relevance for adherence to treatment. *Schizophr. Bull.* **2007**, *33*, 1238–1246. [[CrossRef](#)] [[PubMed](#)]
9. Grosseohme, D.H.; Ragsdale, J.R.; Cotton, S.; Meyers, M.A.; Clancy, J.P.; Seid, M.; Joseph, P.M. Using spirituality after an adult CF diagnosis: Cognitive reframing and adherence motivation. *J. Health Care Chaplain.* **2012**, *18*, 110–120. [[CrossRef](#)] [[PubMed](#)]

10. Koenig, H.G. *Spirituality and Health Research: Methods, Measurements, Statistics, and Resources*; Templeton Press: Conshohocken, PA, USA, 2012.
11. Kogan, S.M.; Luo, Z.; Murry, V.M.; Brody, G.H. Risk and protective factors for substance use among African American high school dropouts. *Psychol. Addict. Behav.* **2005**, *19*, 382–391. [[CrossRef](#)]
12. Hussain, M.; Walker, C.; Moon, G. Smoking and Religion: Untangling Associations Using English Survey Data. *J. Relig. Health* **2019**, *58*, 2263–2276. [[CrossRef](#)]
13. Garcia, G.; Ellison, C.G.; Sunil, T.S.; Hill, T.D. Religion and selected health behaviors among Latinos in Texas. *J. Relig. Health* **2013**, *52*, 18–31. [[CrossRef](#)]
14. Peres, M.F.P.; de Oliveira, A.B.; Leão, F.C.; Filho, H.P.V.; Moreira-Almeida, A.; Lucchetti, G. Religious landscape in Brazil: Comparing different representative nationwide approaches to obtain sensitive information in healthcare research. *SSM Popul. Health* **2018**, *6*, 85–90. [[CrossRef](#)] [[PubMed](#)]
15. Kim, K.K.; Yu, E.S.; Chen, E.H.; Kim, J.; Brintnall, R.; Vance, S. Smoking behavior, knowledge, and beliefs among Korean Americans. *Cancer Pract.* **2000**, *8*, 223–230. [[CrossRef](#)] [[PubMed](#)]
16. Craig, B.A.; Morton, D.P.; Kent, L.M.; Gane, A.B.; Butler, T.L.; Rankin, P.M.; Price, K.R. Religious Affiliation Influences on the Health Status and Behaviours of Students Attending Seventh-Day Adventist Schools in Australia. *J. Relig. Health* **2017**, *57*, 994–1009. [[CrossRef](#)]
17. VanderWeele, T.J. Religion and health in Europe: Cultures, countries, context. *Eur. J. Epidemiol.* **2017**, *32*, 857–861. [[CrossRef](#)] [[PubMed](#)]
18. Connor, K.M.; Davidson, J.R.; Lee, L.C. Spirituality, resilience, and anger in survivors of violent trauma: A community survey. *J. Trauma Stress.* **2003**, *16*, 487–494. [[CrossRef](#)] [[PubMed](#)]
19. Peres, M.F.P.; Kamei, H.H.; Tobo, P.R.; Lucchetti, G. Mechanisms Behind Religiosity and Spirituality's Effect on Mental Health, Quality of Life and Well-Being. *J. Relig. Health* **2018**, *57*, 1842–1855. [[CrossRef](#)] [[PubMed](#)]
20. McSherry, W.; Jamieson, S. The qualitative findings from an online survey investigating nurses' perceptions of spirituality and spiritual care. *J. Clin. Nurs.* **2013**, *22*, 3170–3182. [[CrossRef](#)] [[PubMed](#)]
21. Selman, L.; Young, T.; Vermandere, M.; Stirling, I.; Leget, C. Research priorities in spiritual care: an international survey of palliative care researchers and clinicians. *J. Pain Symptom Manag.* **2014**, *48*, 518–531. [[CrossRef](#)]
22. Brandon, D.M.; Long, J.H.; Loraas, T.M.; Mueller-Phillips, J.; Vansant, B. Online Instrument Delivery and Participant Recruitment Services: Emerging Opportunities for Behavioral Accounting Research. *Behav. Res. Account.* **2013**, *26*, 1–23. [[CrossRef](#)]
23. Lucchetti, G.; de Almeida, L.G.; Lucchetti, A.L. Religiousness, mental health, and quality of life in Brazilian dialysis patients. *Hemodial Int.* **2012**, *16*, 89–94. [[CrossRef](#)]
24. Vitorino, L.M.; Lucchetti, G.; Leao, F.C.; Vallada, H.; Peres, M.F.P. The association between spirituality and religiousness and mental health. *Sci. Rep.* **2018**, *8*, 17233. [[CrossRef](#)]
25. Richards, P.S.; Berrett, M.E.; Hardman, R.K.; Eggett, D.L. Comparative efficacy of spirituality, cognitive, and emotional support groups for treating eating disorder inpatients. *Eat Disord.* **2006**, *14*, 401–415. [[CrossRef](#)]
26. Smith, F.T.; Hardman, R.K.; Richards, P.S.; Fischer, L. Intrinsic religiousness and spiritual well-being as predictors of treatment outcome among women with eating disorders. *Eat Disord.* **2003**, *11*, 15–26. [[CrossRef](#)]
27. Lucchetti, G.; Koenig, H.G.; Pinsky, I.; Laranjeira, R.; Vallada, H. Religious beliefs and alcohol control policies: A Brazilian nationwide study. *Rev. Bras. Psiquiatr.* **2014**, *36*, 4–10. [[CrossRef](#)]
28. Witkiewitz, K.; McCallion, E.; Kirouac, M. Religious Affiliation and Spiritual Practices: An Examination of the Role of Spirituality in Alcohol Use and Alcohol Use Disorder. *Alcohol. Res. Curr. Rev.* **2016**, *38*, 55–58.
29. Degenhardt, L.; Chiu, W.T.; Sampson, N.; Kessler, R.C.; Anthony, J.C. Epidemiological patterns of extra-medical drug use in the United States: Evidence from the National Comorbidity Survey Replication, 2001–2003. *Drug Alcohol. Depend.* **2007**, *90*, 210–223. [[CrossRef](#)]
30. Sajjadi, H.; Ghaedamini Harouni, G.; Sharifian, S.M. Personal, familial and environmental determinants of drug abuse: A causal-comparative study. *Glob. J. Health Sci.* **2015**, *7*, 367–374. [[CrossRef](#)]
31. Drabble, L.; Trocki, K.F.; Klinger, J.L. Religiosity as a protective factor for hazardous drinking and drug use among sexual minority and heterosexual women: Findings from the National Alcohol Survey. *Drug Alcohol. Depend.* **2016**, *161*, 127–134. [[CrossRef](#)]
32. Garrusi, B.; Nakhaee, N. Religion and Smoking: A Review of Recent Literature. *Int. J. Psychiatry Med.* **2012**, *43*, 279–292. [[CrossRef](#)] [[PubMed](#)]
33. Benjamins, M.R. Religious influences on trust in physicians and the health care system. *Int. J. Psychiatry Med.* **2006**, *36*, 69–83. [[CrossRef](#)]
34. Summerskill, W.; Horton, R. Faith-based delivery of science-based care. *Lancet* **2015**, *386*, 1709–1710. [[CrossRef](#)]
35. Lucchetti, G.; Bassi, R.M.; Lucchetti, A.L. Taking spiritual history in clinical practice: A systematic review of instruments. *Explore* **2013**, *9*, 159–170. [[CrossRef](#)]