

The Autophagy Protein CsATG8 is Involved in Asexual Development and Virulence in the Pepper Anthracnose Fungus *Colletotrichum scovillei*

Kwang Ho Lee , Adiyantara Gumilang , Teng Fu , Sung Wook Kang  and Kyoung Su Kim 

Division of Bio-Resource Sciences and Interdisciplinary Program in Smart Agriculture, Kangwon National University, Chuncheon, Korea

ABSTRACT

Autophagy serves as a survival mechanism and plays important role in nutrient recycling under conditions of starvation, nutrient storage, and differentiation of plant pathogenic fungi. However, autophagy-related genes have not been investigated in *Colletotrichum scovillei*, a causal agent of pepper fruit anthracnose disease. *ATG8* is involved in autophagosome formation and is considered a marker of autophagy. Therefore, we generated an *ATG8* deletion mutant, $\Delta Csatg8$, via homologous recombination to determine the functional roles of *CsATG8* in the development and virulence of *C. scovillei*. Compared with the wild-type, the deletion mutant $\Delta Csatg8$ exhibited a severe reduction in conidiation. Conidia produced by $\Delta Csatg8$ were defective in survival, conidial germination, and appressorium formation. Moreover, conidia of $\Delta Csatg8$ showed reduced lipid amount and PTS1 selectivity. A virulence assay showed that anthracnose development on pepper fruits was reduced in $\Delta Csatg8$. Taken together, our results suggest that *CsATG8* plays various roles in conidium production and associated development, and virulence in *C. scovillei*.

ARTICLE HISTORY

Received 29 August 2022
Revised 11 November 2022
Accepted 12 November 2022

KEYWORDS

Colletotrichum scovillei;
pepper anthracnose;
autophagy; *CsATG8*

1. Introduction

Pepper (*Capsicum annuum* L.) is one of the most economically important crops, with the production of dry and green peppers reaching approximately 42.2 million tons worldwide in 2019 [1,2]. Anthracnose disease, caused by fungal species from the *Colletotrichum* genus, is one of the most devastating fungal diseases in pepper [3,4]. Anthracnose disease on pepper fruit and leaves is characterized by sunken lesions and spots [5]. *Colletotrichum scovillei*, belonging to the *Colletotrichum acutatum* species complex, is the main fungal pathogen responsible for pepper yield losses in countries in subtropical and temperate zones [6].

The term autophagy encompasses a number of cellular phenomena [7], including macroautophagy (i.e., nonspecific engulfment of cytosolic components by double-membraned vesicles, which subsequently fuse with the vacuole/lysosome, where the contents are degraded) and pexophagy (i.e., degradation of targeted peroxisomes in the vacuole/lysosome) [8]. Autophagy is a nonselective degradation process that is conserved from yeast to humans [8,9]. Fungal autophagy is involved in the degradation and recycling of unnecessary organelles and long-lived proteins in eukaryotic cells [10–12].

The functional roles of autophagy genes have been well characterized in *Saccharomyces cerevisiae* [8,13–15]. Among the 41 autophagy genes in the genome of *S. cerevisiae*, 19 are essential for autophagosome formation [16]. Autophagy proteins have been classified into five major groups according to their functions: the Atg1 kinase complex, the Atg8 conjugation system, the Atg12 conjugation system, the phosphatidylinositol 3-phosphate kinase complex, and the Atg2–Atg18 complex and Atg9 [17]. Atg4 plays a role as a cysteine protease, and Atg5 is a putative lipase involved in the lysis of autophagic bodies [8,16]. Atg8, used as a marker of autophagy, is associated with the function of the autophagosome membrane [9]. The functional roles of Atg8 are associated with the growth, development, and virulence of several filamentous fungi [8,13,15,18]. For example, in *Aspergillus oryzae*, the deletion of *atg8* caused a rapid increase in hyphal vacuolation under nutrient-starved conditions [8]. In *Magnaporthe oryzae*, $\Delta Moatg8$ was completely defective in conidiation and virulence [18]. In *Colletotrichum orbiculare*, $\Delta Coatg8$ exhibited severe defects in conidial germination and appressorium development [13]. However, the functional roles of

CONTACT Kyoung Su Kim  kims@kangwon.ac.kr

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/12298093.2022.2148393>.

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group on behalf of the Korean Society of Mycology. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ATG8 have not been evaluated in the pepper fruit anthracnose pathogen, *C. scovillei*.

This study was performed to investigate the functional roles of CsATG8 in the development and virulence of *C. scovillei* using a deletion mutant (Δ Csatg8). The results showed that Δ Csatg8 was defective in conidiation, conidium survival, conidial germination, appressorium formation, virulence, and tolerance to oxidative stress. Our results suggest that CsATG8 plays important roles in the growth, development, and virulence of *C. scovillei* in anthracnose disease.

2. Materials and methods

2.1. Fungal strains and culture conditions

The transformants were generated from *C. scovillei* wild-type strain KC05. The fungal strains were cultured on minimal medium agar (MMA; glucose 10 g/l, NaNO₃ 2 g/l, KH₂PO₄ 1 g/l, MgSO₄·7H₂O 0.5 g/l, agar 20 g/l, and Trace element solution 0.1 ml/l), oatmeal agar media (OMA; 50 g oatmeal and 25 g agar/l), and V8 media (V8 juice 80 ml/l, 10 N NaOH₃ 10 μ l/l, agar 15 g/l) at 25 °C and for 5-day dark and next 2-day light. Mycelia for DNA and RNA extraction were grown in completed media broth (CM broth; 0.6% yeast extract, 0.6% casamino acids, and 1% sucrose per liter), and shaken at 150 rpm and 25 °C for 3 days.

2.2. Bioinformatics tools

The sequences of CsATG8 and its homologs were downloaded from National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov>) and Comparative Fungal Genomics Platform (CFGP, <http://cfgp.riceblast.snu.ac.kr>) [18]. Phylogenetic relationships among CsATG8 and

other ubiquitin-like domains were analyzed using MEGA 7.0 software. The identities between CsATG8 and its homologs were analyzed using NCBI BLASTP (<https://blast.ncbi.nlm.nih.gov/>). Domain structures were predicted using InterPro Scan (<http://www.ebi.ac.uk/interpro/>) and visualized using Illustrator for Biological Sequences, version 1.0.3. Primers were designed using Primer Quest Design Tool (<http://sg.idtdan.com/site>).

2.3. Targeted gene deletion and complementation

Target gene deletion was performed using a double-joint PCR method with slight modification [19]. The upstream and downstream regions (~1.5 kb) were amplified with primers 5 F/5R and 3 F/3R (Table 1), respectively. The selection marker, the HPH cassette, was amplified using primers HPHF/HPHR (Table 1) from pBCATPH [20]. To generate the deletion constructs, those PCR products were fused and amplified using 5 F/3R and NF/NR, respectively (Table 1). The amplified deletion constructs were transformed into wild-type protoplasts using a previously reported method [21,22,23]. To generate complemented strain *Csatg8c*, a full-length sequence of CsATG8 was co-transformed with geneticin resistant cassette into Δ Csatg8 protoplasts.

2.4. Vector construction

The carboxyl-terminal amino acid sequence serine-lysine-leucine (SKL) was used as a PTS1 [24]. The *dsRED* gene with a PTS1 was amplified from the pDsRed-gen-36 using the primers DsRED_F and DsRED_SKL_R (Table 1). The pIGPAPA vector was linearized by PCR with the primers DsRED_VF/DsRED_VR. The amplified *dsRED*

Table 1. Primers are used in this study.

Primers	Sequence (5'→3')
CsATG8 (CAP 002212)	
CsATG8 5 F	CATCGCCATCGCTCACAGG
CsATG8 5 R	TACATAGCAGACCGAAACGCGAAGGAGTAATCGGTCGTCGAAG
CsATG8 3 F	GGCGAGGGTCAAGATATAGATGAGGTGGATGGCGTTCTA
CsATG8 3 R	CCTAAGTCAAGAGCCACGAAA
CsATG8 NF	AGCAGATGCTGGGTCACATA
CsATG8 NR	ATAACCGACCTCAGGTTCTT
CsATG8 SF/cF	TTGAAGAACGGTCAGGTCAAG
CsATG8 SR/cR	TCTCATCCTCATTGCATCAGC
CsATG8 PF	GGCTGTTTGTGTCTTCA
CsATG8 PR	TAAGTCAAGAGCCACGAAA
CsATG8 RTF	GACCGTATTCCCGTTATCTGC
CsATG8 RTR	AGTCGCCAAAGGTGTTCT C
CsATG8 qRTF	GACCGTATTCCCGTTATC
CsATG8 qRTR	CGACGAAGATGAAGATGG
DsRED SKL F	ATGGCCTCCTCCGAGAACG
DsRED SKL R	TTATAACTTGGACAGGAACAGGTGGTGGCCGG
DsRED VF	CTGTCCAAGTTATAAAGCGGCCCGCCGGCTGCAG
DsRED VR	CTCGGAGGAGGCCATGTTGGCAGGTGTGGTATG
HPH F	GGCTTGGCTGGAGCTAGTGGAGG
HPH R	CTCCGGAGCTGACATCGACACCAAC

gene with a PTS1 was cloned into the linearized pIGPAPA using the overlap DNA Cloning Kit (Elpis Biotech, Daejeon, Korea).

2.5. Nucleic acid manipulation

Genomic DNA used in screening PCR was extracted through a quick and safe method [25]. To perform southern blotting, fungal genomic DNA was extracted using a standard method [26]. The genomic DNA isolated from wild-type and candidate mutants were digested with restriction enzyme *Nco*I, and then hybridized with a biotin-labeled DNA probe (about 500 bp) (Roche, Indianapolis, IN). Total RNA was extracted from frozen fungal tissues using the Easy-Spin Total RNA Extraction Kit (Intron Biotechnology, Seoul, South Korea), according to the manufacturer's protocol. First-strand cDNA was synthesized from 5 µg total RNA using the oligo (dT) primer with the SuperScriptTM III First-Strand Synthesis System Kit (InvitrogenTM Life Technologies, CA). The *β-tubulin* (CAP_007327) gene was used as the control. The primer sets used for RT-PCR are listed in (Table 1). Real-Time quantitative PCR was performed on the StepOne Real-Time PCR System (Applied Biosystems, CA) using HIPI Real-Time PCR 2 × Master Mix (SYBR Green) (Elpis Biotech).

2.6. Phenotypic analysis

Conidiation was measured by counting the number of conidia harvested with 5 ml of distilled water using a hemocytometer. The conidial length was measured using an Axio Image A2 microscope (Carl Zeiss Microscope Division, Oberkochen, Germany) with ZEN imaging software. Conidial germination and appressorium formation were performed by dropping conidial suspensions (5×10^4 conidia/ml), prepared from filtering through 3 layers of miracloth (Calbiochem, San Diego, CA) and wash for 3 times, onto the hydrophobic surface of coverslips and incubating in a humid box at 25 °C. To evaluate mycelial growth under chemical stresses and starvation, mycelial agar plugs from MMA were inoculated onto CMA containing 20 and 30 mM H₂O₂, and incubated at 25 °C without light for 5 days. To evaluate conidium survival and lipid reservoir, the conidial suspension was stained by Phloxine B and Nile red (Sigma, St Louis, MO). Infection assays were performed by inoculating conidial suspension (20×10^4 conidia/ml) onto healthy intact, wounded pepper fruits and incubating them in a humid box at 25 °C for 5–10 days. For the appressorium penetration and invasive growth assays, conidial drops (5×10^4 conidia/ml) were placed on the surface of pepper fruits and incubated in a moistened box at 25 °C. All experiments were repeated 3 times with 3 replicates.

3. Results

3.1. Phylogenetic analysis, domain structure, and targeted gene deletion

Phylogenetic analysis showed that CsATG8 is closely related to its orthologues in fungi. NCBI BLASTP search showed that ATG8 is highly conserved in eukaryotic organisms, with high levels of identity to orthologues from *Colletotrichum gloeosporioides* (98.35%), *Neurospora crassa* (99.17%), *Penicillium rubens* (98.31%), *Botrytis cinerea* (96.58%), *Sclerotinia sclerotiorum* (97.44%), *Fusarium oxysporum* (100%), *Colletotrichum higginsianum* (98.35%), *Pyricularia oryzae* (94.07%), *Yarrowia lipolytica* (90.76%), *Candida albicans* (80.34%), *S. cerevisiae* (79.31%), and lower levels of identity to the more distant *Arabidopsis thaliana* (80.17%) and *Homo sapiens* (59.48%) orthologues [27,28] (Figure S1(A,B)). Domain prediction showed that all ATG8 proteins contain a small ubiquitin-like domain (IPR029071) (Figure S1(B)). These results indicate that CsATG8 homologs are highly conserved in fungi.

To investigate the roles of CsATG8, we generated a targeted gene deletion mutant by homologous recombination (Figure S2(A)). Δ Csatg8 was confirmed by Southern blotting and RT-PCR (Figure S2(B,C)). To verify the phenotypes of Δ Csatg8 caused by the deletion of CsATG8, we generated a complemented strain (*Csatg8c*), which was verified by RT-PCR (Figure S2(C)).

3.2. Roles of CsATG8 in conidiation, conidial morphology, reduced lipid amount, and PTS1 selectivity

Δ Csatg8 showed normal mycelial growth rate but significantly reduced conidiation compared with the wild-type and *Csatg8c* (Figure 1(A,B) and Figure S3), suggesting that CsATG8 is involved in the conidiation of *C. scovillei*. A vacuole-like structure was observed in $29 \pm 3.6\%$ of conidia in Δ Csatg8, compared with $5.3 \pm 2.5\%$ and $6.3 \pm 1.5\%$ of those in the wild-type and *Csatg8c*, respectively (Figure 2(A,B)). It has been reported that peroxisomal matrix proteins are imported into the peroxisome via the PTS1 pathway [24]. Therefore, we expressed the dsRED-SKL fusion protein in wild-type, Δ Csatg8, and *Csatg8c*. The cytoplasmic localized RFP signal was detected in $24 \pm 3.5\%$ of conidia in Δ Csatg8 compared with $2.6 \pm 1.1\%$ and $6.3 \pm 1.5\%$ of those in the wild-type and *Csatg8c*, respectively (Figure 2(C)). Because Lipids droplets are a source of energy in conidia and involved in conidium fitness and germination [29,30], we stained lipid droplets with Nile red and found an accumulation of lipid droplets in




Figure 1. Measurement of conidiation. (A) Observation of conidia developed from conidiophores. Photographs were taken after incubation of 3-day-old mycelial agar plugs in a humid box for 7 h with light; (B) Quantitative evaluation of conidiation. The indicated strains were grown on V8 medium for 5 days in the dark and 2 days in the light. * $p < 0.05$ (Tukey's test). Scale bar = 20 μm .

10.5 \pm 2.6% of conidia in $\Delta Csatg8$, compared with 97.7 \pm 1.5% of those in the wild-type (Figure 2(D,E)). This result suggests that *CsATG8* is involved in a proper lipid reservoir. We further examined conidium survival in $\Delta Csatg8$ by staining with phloxine B [31–34], and the results showed that the proportion of dead conidia was significantly greater in $\Delta Csatg8$ (20.3 \pm 4.5%) than in the wild-type (3 \pm 0.5%) and $Csatg8c$ (3.6 \pm 1.5%) (Figure 3(A,B)). This result suggests that *CsATG8* is involved in conidial survival in *C. scovillei*. Taken together, these results suggest that *CsATG8* is involved in the conidiation, conidium viability, and autophagy-related processes in *C. scovillei*.

3.3. Role of *CsATG8* in appressorium formation

Appressorium development is a prerequisite for anthracnose disease [2]. To investigate the role of *CsATG8* in pre-infection, we evaluated the time course of conidial germination and appressorium

formation on the hydrophobic surface of coverslips. After 6 h, the germination rate of the wild-type was approximately 72 \pm 2.5%, whereas the rate of germ tube formation was only approximately 35 \pm 2.0% in $\Delta Csatg8$. The defect in conidial germination was recovered in $Csatg8c$. In comparison with the wild-type and $Csatg8c$ strains, most conidia of $\Delta Csatg8$ failed to differentiate into appressoria at 16 h (Figure 4(A,B)). These results suggest that *CsATG8* is involved in conidial germination and appressorium formation in *C. scovillei*.

3.4. Roles of *CsATG8* in anthracnose disease formation

To determine the role of *CsATG8* in anthracnose development, we inoculated conidial suspensions onto intact pepper fruits. $\Delta Csatg8$ was greatly reduced in causing anthracnose disease, whereas both the wild-type and $Csatg8c$ caused severe anthracnose disease after 10 days (Figure 5(A)). These observations suggest that *CsATG8* plays an important role in the virulence of *C. scovillei*. We inoculated conidial suspensions onto wounded pepper fruits. After 5 days, $\Delta Csatg8$ caused a similar level of anthracnose disease as did those of wild-type and $Csatg8c$ (Figure 5(B)). This observation reveals that the reduced virulence of $\Delta Csatg8$ did not result from a defect of invasive hyphae growth. These results suggest that *CsATG8* is important for virulence through the regulation of appressorium-mediated penetration in *C. scovillei*.

3.5. Roles of *CsATG8* in tolerance to stress conditions

To determine whether *CsATG8* is involved in tolerance to stress conditions, we evaluated mycelial growth on CMA amended with oxidative stresses (Figure 6(A,B)). Among those stresses, mycelial growth was significantly inhibited by H_2O_2 in $\Delta Csatg8$ (by 56.6 \pm 3.0%) compared with the wild-type (by 44.0 \pm 2.0%) and $Csatg8c$ (by 42.0 \pm 2.0%) (Figure 6(A,B)). This result suggests that *CsATG8* is involved in the tolerance to oxidative stress of *C. scovillei*.

4. Discussion

This study was performed to functionally characterize *ATG8*, one of 23 autophagy-related genes in *C. scovillei*, because *ATG8* is a major autophagy protein involved in the development and virulence of plant pathogenic fungi [8,13,15,18].

The function of *ATG8* is related to conidiation in several phytopathogenic fungi [13,14,18,35]. In




Figure 2. Detection of intracellular vacuoles and lipids. (A) Observation of vacuole-like structures and dsRED:PTS1 localization. Conidia were harvested in MMA medium for 5 days in the dark and 2 days in the light; (B, C) Quantitative examination of vacuole-like structures and the dsRED:PTS1 fusion protein. At least 100 conidia were examined per replicate. * $p < 0.05$ (Tukey's test); (D, E) Measurement of lipid droplets; (D) Observation of lipid localization. The conidia were stained with Nile red and incubated for 1 h in the dark; (E) Quantitative measurement of lipid droplets. At least 100 conidia were examined per replicate. * $p < 0.05$ (Tukey's test). Scale bar = 10 μm .

M. oryzae, the deletion of *ATG8* significantly reduced conidiation and conidiophore development [18]. Our results showed that conidiation was significantly reduced in $\Delta Csatg8$ (Figure 1(A,B)), suggesting that *CsATG8* is involved in conidiation. Interestingly, $\Delta Csatg8$ produced morphologically abnormal conidia, which were smaller than those of the wild-type (Figure 2(A)). Moreover, the phloxine B assay showed that $\Delta Csatg8$ produced dead conidia (Figure 3(A)). Defects in the viability of $\Delta Csatg8$ conidia may reveal a novel function of *ATG8* in plant pathogenic fungi.

ATG8 plays an important role in infection-related morphogenesis [18]. *ATG8* deletion mutants were reported to be defective in conidial germination and appressorium formation in *M. oryzae* and *C. orbiculare* [13,18]. In the present study, conidial germination was significantly lower in $\Delta Csatg8$ ($35 \pm 2.0\%$) than in the wild-type ($72 \pm 2.5\%$) after 6 h of incubation. In the conidial viability assay, $\Delta Csatg8$ showed a conidial survival rate of $79.7 \pm 4.5\%$ (Figure 3(B)), indicating that reduced conidial germination was not only due to low conidial viability. Furthermore, $\Delta Csatg8$ showed reduced appressorium formation

(Figure 4(A,B)). These results suggested that *CsATG8* is involved in conidial germination and appressorium formation in *C. scovillei*.

Pexophagy, a type of selective autophagy, is involved in degrading lipids in peroxisomes [12,36]. Nile red staining assay showed that $\Delta Csatg8$ accumulated fewer lipid droplets than did the wild-type (Figure 2(D,E)), suggesting that *CsATG8* may be involved in the lipid reservoir. Peroxisomal membrane proteins are targeted to the peroxisome membrane by two distinct peroxisomal targeting signals, PTS1 and PTS2 [37,38]. Accompanied by the formation of autophagosomes, peroxisomes are degraded inside vacuoles [13]. RFP:PTS1 signals in $\Delta Csatg8$ exhibited cytoplasmic localization, whereas the wild-type control showed a punctate localization pattern [36]. *CsATG8* may be involved in the formation of intracellular organelles and peroxisomes.

The $\Delta Csatg8$ showed severely reduced anthracnose disease formation on pepper fruits (Figure 5(A,B)). Previous studies demonstrated that *ATG8* deletion mutants were defective in appressorium-mediated penetration [13,18]. Consistently, $\Delta Csatg8$ was significantly reduced in penetration ability to




Figure 3. Conidial viability assay. (A) Observation of conidial survival. Conidia harvested from 10-day-old MMA plugs were stained with phloxine B. Dead conidia were stained red. Scale bar = 20 μ m; (B) Quantitative evaluation of conidial survival. At least 100 conidia were examined per replicate. * $p < 0.05$ (Tukey's test).




Figure 4. Appressorium formation assay. (A) Photographs of appressorium formation. Conidial suspensions (5×10^4 /ml) were placed on the hydrophobic surface of coverslips and incubated in a humid box at 25 $^{\circ}$ C for 16 h. Scale bar = 10 μ m; (B) Quantitative measurement of appressorium formation. At least 100 conidia were examined per replicate. * $p < 0.05$ (Tukey's test).



Figure 5. Virulence assays. Conidial suspensions (20×10^4 /ml) were dropped onto intact and wounded pepper fruits and incubated in a humid plastic box at 25 $^{\circ}$ C for 14 days.




Figure 6. Mycelial growth under conditions of oxidative stress. Mycelial agar plugs were inoculated onto CMA or CMA containing 20 or 30 mM H₂O₂ and then incubated at 25 $^{\circ}$ C in the dark for 5 days. * $p < 0.05$ (Tukey's test).

that of the wild-type and *Csatg8c* in the present study (Figure 5(A,B)), suggesting that *CsATG8* is involved in appressorium-mediated penetration. $\Delta Csatg8$ caused anthracnose disease with similar lesion size on wounded pepper fruits compared to the wild-type. The anthracnose lesions caused by $\Delta Csatg8$ were black in color, which may be caused by increased pigmentation in $\Delta Csatg8$, as mycelia of $\Delta Csatg8$ were more pigmented compared to that of wild-type (Figure S3). Therefore, the marked reduction in anthracnose lesions in $\Delta Csatg8$ should be due to a defect in appressorium-mediated penetration.

Taken together, our results showed that $\Delta Csatg8$ is defective in conidiation, appressorium formation, virulence, lipid reservoir, PTS1 selection, and tolerance to oxidative stress in *C. scovillei*. Our results suggest that *CsATG8* plays important roles in the development and virulence of *C. scovillei* in anthracnose disease.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This study was supported by the Basic Science Research Program through the National Research Foundation of Korea grant [NRF-2020R1A2C100550700] funded by the Ministry of Education, Science and Technology, Republic of Korea.

ORCID

Kwang Ho Lee  <http://orcid.org/0000-0002-5001-2929>

Adiyantara Gumilang  <http://orcid.org/0000-0002-8887-9763>

Teng Fu  <http://orcid.org/0000-0003-1310-6501>

Sung Wook Kang  <http://orcid.org/0000-0002-7049-2499>

Kyoung Su Kim  <http://orcid.org/0000-0002-0827-0844>

References

- [1] Food and Agriculture Organization of the United Nations [Internet]. Rome (Italy): FAOSTAT; 2021 [cited 2021 November 3]. Available from: <http://www.fao.org/faostat/en/#data/QC/>
- [2] Lee NH, Fu T, Shin JH, et al. The small GTPase *CsRAC1* is important for fungal development and pepper anthracnose in *Colletotrichum scovillei*. *Plant Pathol J*. 2021;37(6):607–618.
- [3] Förster H, Adaskaveg JE. Identification of subpopulations of *Colletotrichum acutatum* and epidemiology of almond anthracnose in California. *Phytopathology*. 1999;89(11):1056–1065.
- [4] Giacomini RM, de Fátima Ruas C, Moreira AFP, et al. Inheritance of anthracnose resistance (*Colletotrichum scovillei*) in ripe and unripe *Capsicum annuum* fruits. *J Phytopathol*. 2020; 168(3):184–192.
- [5] Saxena A, Raghuvanshi R, Gupta VK, et al. Chilli anthracnose: the epidemiology and management. *Front Microbiol*. 2016;7:1527.
- [6] Fu T, Han JH, Shin JH, et al. Homeobox transcription factors are required for fungal development and the suppression of host defense mechanisms in the *Colletotrichum scovillei*-pepper pathosystem. *mBio*. 2021;12(4):e01620.
- [7] Yorimitsu T, Klionsky DJ. Autophagy: molecular machinery for self-eating. *Cell Death Differ*. 2005; 12(S2):1542–1552.
- [8] Pollack JK, Harris SD, Marten MR. Autophagy in filamentous fungi. *Fungal Genet Biol*. 2009;46(1): 1–8.
- [9] Klionsky DJ, Cuervo AM, Seglen PO. Methods for monitoring autophagy from yeast to human. *Autophagy*. 2007;3(3):181–206.
- [10] Pinan-Lucarré B, Balguerie A, Clavé C. Accelerated cell death in *Podospora* autophagy mutants. *Eukaryot Cell*. 2005;4(11):1765–1774.
- [11] Pollack JK, Li ZJ, Marten MR. Fungal mycelia show lag time before re-growth on endogenous carbon. *Biotechnol Bioeng*. 2008;100(3):458–465.
- [12] Reggiori F, Klionsky DJ. Autophagy in the eukaryotic cell. *Eukaryot Cell*. 2002;1(1):11–21.
- [13] Asakura M, Ninomiya S, Sugimoto M, et al. Atg26-mediated pexophagy is required for host invasion by the plant pathogenic fungus *Colletotrichum orbiculare*. *Plant Cell*. 2009;21(4): 1291–1304.
- [14] Deng Y, Qu Z, Naqvi NI. Role of macroautophagy in nutrient homeostasis during fungal development and pathogenesis. *Cells*. 2012;1(3):449–463.
- [15] Kershaw MJ, Talbot NJ. Genome-wide functional analysis reveals that infection-associated fungal autophagy is necessary for rice blast disease. *Proc Natl Acad Sci USA*. 2009;106(37):15967–15972.
- [16] Hirata E, Ohya Y, Suzuki K. Atg4 plays an important role in efficient expansion of autophagic isolation membranes by cleaving lipidated Atg8 in *Saccharomyces cerevisiae*. *PLoS One*. 2017;12(7): e0181047.
- [17] Nakatogawa H, Suzuki K, Kamada Y, et al. Dynamics and diversity in autophagy mechanisms: lessons from yeast. *Nat Rev Mol Cell Biol*. 2009; 10(7):458–467.
- [18] Deng YZ, Ramos-Pamplona M, Naqvi NI. Autophagy-assisted glycogen catabolism regulates asexual differentiation in *Magnaporthe oryzae*. *Autophagy*. 2009;5(1):33–43.
- [19] Yu JH, Hamari Z, Han KH, et al. Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi. *Fungal Genet Biol*. 2004;41(11):973–981.
- [20] Choi J, Kim Y, Kim S, et al. MoCRZ1, a gene encoding a calcineurin-responsive transcription factor, regulates fungal growth and pathogenicity of *Magnaporthe oryzae*. *Fungal Genet Biol*. 2009; 46(3):243–254.
- [21] Kim S, Park SY, Kim KS, et al. Homeobox transcription factors are required for conidiation and appressorium development in the rice blast fungus *Magnaporthe oryzae*. *PLOS Genet*. 2009;5(12): e1000757.

- [22] Sweigard JA, Chumley FG, Valent B. Cloning and analysis of *CUT1*, a cutinase gene from *Magnaporthe grisea*. *Mol Gen Genet*. 1992;232(2): 174–182.
- [23] Shin JH, Han JH, Park HH, et al. Optimization of polyethylene glycol-mediated transformation of the pepper anthracnose pathogen *Colletotrichum scovillei* to develop an applied genomics approach. *Plant Pathol J*. 2019;35(6):575–584.
- [24] Wang JY, Wu XY, Zhang Z, et al. Fluorescent colocalization of PTS1 and PTS2 and its application in analysis of the gene function and the peroxisomal dynamic in *Magnaporthe oryzae*. *J Zhejiang Univ Sci B*. 2008;9(10):802–810.
- [25] Chi MH, Park SY, Lee YH. A quick and safe method for fungal DNA extraction. *Plant Pathol J*. 2009;25(1):108–111.
- [26] Sambrook J, Russell DW. 2001. *Molecular cloning: a laboratory manual*. 3rd ed. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press.
- [27] Han JH, Chon JK, Ahn JH, et al. Whole genome sequence and genome annotation of *Colletotrichum acutatum*, causal agent of anthracnose in pepper plants in South Korea. *Genom Data*. 2016;8:45–46.
- [28] Kim JO, Choi KY, Han JH, et al. The complete mitochondrial genome sequence of the ascomycete plant pathogen *Colletotrichum acutatum*. *Mitochondrial DNA A DNA Mapp Seq Anal*. 2016;27(6):4547–4548.
- [29] Bianchi DE, Turian G. Lipid content of conidia of *Neurospora crassa*. *Nature*. 1967;214(5095): 1344–1345.
- [30] Daryaei A. *Conidium “fitness” in Trichoderma* [dissertation]. Oxford (PA): Lincoln University; 2014.
- [31] Fu T, Shin JH, Lee NH, et al. Mitogen-activated protein kinase CsPMK1 is essential for pepper fruit anthracnose by *Colletotrichum scovillei*. *Front Microbiol*. 2022;13:770119.
- [32] Jeon J, Rho H, Kim S, et al. Role of MoAND1-mediated nuclear positioning in morphogenesis and pathogenicity in the rice blast fungus, *Magnaporthe oryzae*. *Fungal Genet Biol*. 2014;69: 43–51.
- [33] Park J, Kong S, Kim S, et al. Roles of forkhead-box transcription factors in controlling development, pathogenicity, and stress response in *Magnaporthe oryzae*. *Plant Pathol J*. 2014;30(2): 136–150.
- [34] Sun CB, Suresh A, Deng YZ, et al. A multidrug resistance transporter in magnaporthe is required for host penetration and for survival during oxidative stress. *Plant Cell*. 2006;18(12):3686–3705.
- [35] Liu XH, Lu JP, Zhang L, et al. Involvement of a *Magnaporthe grisea* serine/threonine kinase gene, MgATG1, in appressorium turgor and pathogenesis. *Eukaryot Cell*. 2007;6(6):997–1005.
- [36] Shin JH, Fu T, Kim KS. Pex7 selectively imports PTS2 target proteins to peroxisomes and is required for anthracnose disease development in *Colletotrichum scovillei*. *Fungal Genet Biol*. 2021; 157:103636.
- [37] Fujiki Y, Matsuzono Y, Matsuzaki T, et al. Import of peroxisomal membrane proteins: the interplay of Pex3p-and Pex19p-mediated interactions. *Biochim Biophys Acta*. 2006;1763(12):1639–1646.
- [38] Otera H, Okumoto K, Tateishi K, et al. Peroxisome targeting signal type 1 (PTS1) receptor is involved in import of both PTS1 and PTS2: studies with PEX5-defective CHO cell mutants. *Mol Cell Biol*. 1998;18(1):388–399.