

SHORT REPORT

Open Access

# DNA barcoding does not separate South American *Triatoma* (Hemiptera: Reduviidae), Chagas Disease vectors

Silvia Andrade Justi<sup>1,2</sup>, Carolina Dale<sup>2</sup> and Cleber Galvão<sup>2\*</sup>

## Abstract

**Background:** DNA barcoding assumes that a biological entity is completely separated from its closest relatives by a *barcoding gap*, which means that intraspecific genetic distance (from COI sequences) should never be greater than interspecific distances. We investigated the applicability of this strategy in identifying species of the genus *Triatoma* from South America.

**Findings:** We calculated intra and interspecific Kimura-2-parameter distances between species from the *infestans*, *matogrossensis*, *sordida* and *rubrovaria* subcomplexes. In every subcomplex examined we observed at least one intraspecific distance greater than interspecific distances.

**Conclusions:** Although DNA barcoding is a straightforward approach, it was not applicable for identifying Southern American *Triatoma* species, which may have diverged recently. Thus, caution should be taken in identifying vector species using this approach, especially in groups where accurate identification of taxa is fundamentally linked to public health issues.

**Keywords:** Triatominae, Chagas disease, DNA barcoding, Molecular identification

## Findings

DNA barcoding, as proposed by Hebert et al. [1] assumes that a biological entity is completely separated from its closest relatives by a *barcoding gap* [2], which means that intraspecific genetic distances (from COI sequences) are never greater than interspecific distances.

*Triatoma* Laporte (Hemiptera: Reduviidae) is the most diverse genus of Chagas Disease vectors, and accurate identification of species is imperative for the efficiency of vector control programs. The *Triatoma* genus is divided into species complexes and subcomplexes according to geographic distribution and morphological similarity [3].

Recently, Justi et al. [4] reported that the relationships between species assigned to South American *Triatoma*-subcomplexes could not be untangled with the data in hand. We were then prompted to investigate whether DNA barcoding would be a useful tool for identifying

the species within the *infestans*, *matogrossensis*, *sordida* and *rubrovaria* subcomplexes [3].

Kimura-2-parameter genetic distances [5] were calculated pairwise within each of the above mentioned subcomplexes (Table 1) using the software MEGA v. 5 [6], and intra and interspecific distances were compared.

In all subcomplexes we observed at least one intraspecific distance greater than interspecific distances (Table 1). To be considered appropriate to identify species within a group, intraspecific distances must always be greater than interspecific ones [2], and therefore DNA barcoding is not accurate for the species-level identification of South American *Triatoma*. Moreover, the method fails to account for hybridization events, which are naturally observed in *Triatoma* [7,8], and introgression, which is frequent in nuclear DNA [9]. These considerations argue that Hebert et al.'s [1] proposal of cataloguing biodiversity based only on DNA barcoding may severely underestimate it.

Besides that, as highlighted by Dujardin et al. [10], the morphological changes observed in closely related "species", or "lineages" as we prefer to call them, may have

\* Correspondence: galvao@ioc.fiocruz.br

<sup>2</sup>Laboratório Nacional e Internacional de Referência em Taxonomia de Triatomíneos, Instituto Oswaldo Cruz, FIOCRUZ, Rio de Janeiro, Brazil  
Full list of author information is available at the end of the article

**Table 1 K2p-distances between species of the *Triatoma* subcomplexes studied**

Subcomplex	GenBank	Number	Geographic Origin		1	2	3	4	5										
<i>infestans</i>	KC249330	1	Chaco Tita, Cochabamba, Bolivia	<i>T. delpontei</i> 53															
	KC249346	2	Chaco Tita, Cochabamba, Bolivia	<i>T. infestans</i> 44	<b>0.021</b>														
	KC249349	3	Cotapachi, Cochabamba, Bolivia	<i>T. infestans</i> 58	<b>0.025</b>	0.018													
	KC249352	4	Mataral, Cochabamba, Bolivia	<i>T. infestans</i> 60	<b>0.025</b>	0.018	0.005												
	KC249354	5	Ilicuni, Cochabamba, Bolivia	<i>T. infestans</i> 63	<b>0.021</b>	0.016	0.000	0.006											
	KC249355	6	Montevideo, Uruguay	<i>T. infestans</i> 69	0.072	<b>0.061</b>	<b>0.064</b>	<b>0.069</b>	<b>0.103</b>										
<i>matogrossensis</i>	KC249327,KC249328	7	Posse, GO, Brazil	<i>T. costalimai</i> 35															
	KC249329	8	Chiquitania, Cochabamba, Bolivia	<i>T. costalimai</i> 42	0.154														
	KC249360	9	São Gabriel D'oeste, MS, Brazil	<i>T. matogrossensis</i> 192	0.134	0.138													
	KC249361	10	Bahia, Brazil	<i>T. matogrossensis</i> 31	0.151	0.152	<b>0.047</b>												
	KC249391	11	Pantanal, MT, Brazil	<i>T. vanda</i> 28	0.156	0.151	0.047	0.040											
	KC249392	12	Rio Verde do MatoGrosso, MT, Brazil	<i>T. vanda</i> 73	0.138	0.146	<b>0.005</b>	0.046	0.045										
	KC249393,KC249394	13	Rondonópolis, MT, Brazil	<i>T. vanda</i> 74	0.158	0.150	0.048	0.059	0.007	0.052									
<i>rubrovaria</i>	KC249322	14	São Gerônimo, RS, Brazil	<i>T. carvalhoi</i> 78															
	KC249323	15	Caçapava do Sul, RS, Brazil	<i>T. circummaculata</i> 120	0.039														
	KC249324	16	Sítio Faxina, Piratini, RS, Brazil	<i>T. circummaculata</i> 121	0.029	0.025													
	KC249325	17	Sítio Faxina, Piratini, RS, Brazil	<i>T. circummaculata</i> 122	0.017	<b>0.039</b>	<b>0.033</b>												
	KC249356	18	Nova Petrópolis, RS, Brazil	<i>T. klugi</i> 75	0.018	<b>0.037</b>	<b>0.031</b>	<b>0.017</b>											
	KC249369	19	Sítio Faxina, Piratini, RS, Brazil	<i>T. rubrovaria</i> 123	<b>0.055</b>	0.023	0.029	0.055	0.057										
	KC249370	20	Sítio venda da Lagoa, Canguçu, RS, Brazil	<i>T. rubrovaria</i> 134	0.065	0.052	0.065	0.065	0.061	0.070									
	KC249372	21	SítioFaxina, Pinheiro Machado, RS, Brazil	<i>T. rubrovaria</i> 136	0.042	0.019	0.027	0.036	0.036	0.031	0.035								
	KC249373	22	Sítio venda da Lagoa, Canguçu, RS, Brazil	<i>T. rubrovaria</i> 140	0.038	0.020	0.019	0.043	0.040	0.029	0.032	0.012							
	KC249374	23	Canguçu, RS, Brazil	<i>T. rubrovaria</i> 156	0.039	0.020	0.019	0.045	0.042	0.029	0.032	0.012	0.000						
	KC249375	24	Caçapava do Sul, RS, Brazil	<i>T. rubrovaria</i> 76	0.021	0.029	0.021	0.016	0.016	0.033	<b>0.074</b>	0.034	0.038	0.038					
	KC249376	25	Quevedos, RS, Brazil	<i>T. rubrovaria</i> 77	0.029	0.030	0.043	0.022	0.029	0.046	<b>0.065</b>	0.031	0.046	0.048	0.026				
	<i>sordida</i>	KC249338	26	Rivadaria, Argentina	<i>T. garciabesi</i> 89														
KC249342		27	Santa Cruz, Bolívia	<i>T. guasayana</i> 55	0.077														
KC249343		28	Santa Cruz, Bolívia	<i>T. guasayana</i> 82	0.065	0.056													

**Table 1 K2p-distances between species of the *Triatoma* subcomplexes studied (Continued)**

KC249379,KC249380	29	Romerillo, Cochabamba, Bolivia	<i>T. sordida</i> 46	0.029	0.060	0.060							
KC249381,KC249382	30	Romerillo, Cochabamba, Bolivia	<i>T. sordida</i> 47	<b>0.030</b>	0.061	0.061	0.000						
KC249383	31	La Paz, Bolivia	<i>T. sordida</i> 83	0.081	0.013	0.063	0.066	0.066					
KC249384	32	Pantanal, MS, Brazil	<i>T. sordida</i> 85	0.069	0.012	0.062	0.065	<b>0.065</b>	0.025				
KC249385	33	Santa Cruz, Bolivia	<i>T. sordida</i> 86	0.043	0.082	0.074	0.035	<b>0.035</b>	0.073	0.082			
KC249387	34	San Miguel Corrientes, Argentina	<i>T. sordida</i> 88	0.061	0.058	0.063	0.070	<b>0.071</b>	0.058	0.055	0.052		
KC249388	35	Poconé, MT, Brazil	<i>T. sordida</i> 90	0.069	0.017	0.058	0.075	<b>0.075</b>	0.031	0.011	0.078	0.051	

Highlighted distances deviate from the DNA barcoding premis that intraspecific distances are smaller than interspecific distances.

led taxonomists to rush into describing subspecies or species, even genera. Molecular phylogenetic studies are in their infancy in unravelling the evolution of Triatominae, and a comprehensive molecular phylogeny, including more than one specimen for most lineages, was published only in 2014 [4], although several analyses were conducted focusing on small species groups. Taken together, these statements make it clear that further investigations of Triatominae evolution are long overdue, preferably integrating morphological, molecular and ecological data.

Lineage evolution has not occurred, but it is happening now. Concerning lineages designated in the *infestans* complex (including the subcomplexes studied here), separation is much clearer in terms of morphology than in molecular systematics. In cases where lineages have not reached reciprocal monophyly, defining taxonomic entities is not a straightforward issue [11]. Therefore caution is necessary, especially in a group where accurate identification of taxa is fundamentally linked to public health issues.

## Conclusions

Although DNA barcoding is a straightforward approach, it was not applicable for identifying Southern American *Triatoma* species, which may have diverged recently. Thus, caution should be taken in identifying vector species using this approach, especially in groups where accurate identification of taxa is fundamentally linked to public health issues.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

SAJ designed the study, acquired data, performed all the analyses, interpreted the results, drafted and reviewed the manuscript. CD designed the study, acquired data, performed all the analyses, interpreted the results, drafted and reviewed the manuscript. CG designed the study and reviewed the manuscript. All authors read and approved the final version of the manuscript.

## Authors' information

CD is PhD student funded by Instituto Oswaldo Cruz and SAJ is a Post Doctoral Fellow funded by CNPq.

## Acknowledgements

We thank Carlos G. Schrago for reviewing our manuscript and for the useful comments.

## Author details

<sup>1</sup>Universidade Federal do Rio de Janeiro, Instituto de Biologia, Laboratório de Biologia Evolutiva Teórica e Aplicada, Rio de Janeiro, Brazil. <sup>2</sup>Laboratório Nacional e Internacional de Referência em Taxonomia de Triatomíneos, Instituto Oswaldo Cruz, FIOCRUZ, Rio de Janeiro, Brazil.

Received: 13 September 2014 Accepted: 4 November 2014

Published online: 21 November 2014

## References

1. Hebert PDN, Cywinska A, Ball SL, DeWaard JR: *Biological identifications through DNA barcodes*. Biol Sci: In Proc Roy Soc Lond B; 2003:313–321.

- Meyer CP, Paulay G: DNA barcoding: error rates based on comprehensive sampling. *PLoS Biol* 2005, **3**(Suppl. 12):e422.
- Schofield CJ, Galvão C: Classification, evolution, and species groups within the Triatominae. *Acta Trop* 2009, **110**(Suppl 2–3):88–100.
- Justi SA, Russo AMC, Santos-Mallet JR, Obara MT, Galvão C: Molecular phylogeny of Triatomini (Hemiptera: Reduviidae: Triatominae). *Parasit Vectors* 2014, **7**:149.
- Kimura M: A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 1980, **16**:111–120.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S: MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony method. *Mol Biol Evol* 2011, **28**:2731–2739.
- Carcavallo RU, Girón IG, Jurberg J, Galvao C, Noireau F, Canale D: Mutations, Hybrids and Teratologies. In *Atlas of Chagas Disease Vectors in the Americas*. Edited by Editora FIOCRUZ. Rio de Janeiro: Carcavallo, Girón, Jurberg, Lent; 1998:515–536.
- Costa J, Peterson AT, Dujardin JP: Morphological evidence suggests homoploid hybridization as a possible mode of speciation in the Triatominae (Hemiptera, Heteroptera, Reduviidae). *Inf Gen Evol* 2009, **9**:263–270.
- Zhang AB, Sota T: Nuclear gene sequences resolve species phylogeny and mitochondrial introgression in *Leptocarabus* beetles showing trans-species polymorphisms. *Mol Phylogenet Evol* 2007, **45**:534–546.
- Dujardin JP, Costa J, Bustamante D, Jaramillo N, Catalá S: Deciphering morphology in Triatominae: the evolutionary signals. *Acta Trop* 2009, **110**(Suppl 2–3):101–111.
- Arnold ML: Natural Hybridization and species concepts. In *Natural Hybridization and Evolution*. Oxford: Arnold: Oxford University Press; 1997:11–22.

doi:10.1186/s13071-014-0519-1

**Cite this article as:** Justi et al.: DNA barcoding does not separate South American *Triatoma* (Hemiptera: Reduviidae), Chagas Disease vectors. *Parasites & Vectors* 2014 **7**:519.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

