

DNA-induced unfolding of the thyroid hormone receptor α A/B domain through allostery

Elias J. Fernandez, Vandna Gahlot, Celeste Rodriguez and Jacob Amburn

Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, TN, USA

Keywords

A/B domain; interdomain allostery; intrinsically disordered protein domain; nuclear receptor; protein–DNA interactions; thyroid receptor

Correspondence

E. J. Fernandez, BCMB Department, The University of Tennessee, Knoxville, TN 37996, USA
Fax: +1-(865) 974-6306
Tel: +1-(865) 974-4090
E-mail: elias.fernandez@utk.edu

The A/B domains of nuclear receptors such as thyroid receptor α (TR α) are considered to be conformationally flexible and can potentially adopt multiple structural conformations. We used intrinsic tryptophan fluorescence quenching and circular dichroism spectroscopy to characterize the unfolding of this A/B domain upon DNA binding to the contiguous DNA-binding domain (DBD). We propose that this allosteric change in A/B domain conformation can allow it to make the multiple interactions with distinct molecular factors of the transcriptional preinitiation complex. We further suggest that by influencing the affinity of the DBD for DNA, A/B domain can fine-tune the recognition of promoter DNA by TR α .

(Received 23 January 2017, revised 10 April 2017, accepted 11 April 2017)

doi:10.1002/2211-5463.12229

The effects of the thyroid hormone (triiodothyronine, T3) are widespread in development, homeostasis and metabolism. The T3 receptors (thyroid hormone receptor, TR) are encoded by two closely related genes (α and β) [1]. The T3R α genes in humans express the T3-binding isoform TR α 1 [2]. The TR β gene expresses TR β 1 and TR β 2, which differ only in their N-terminal A/B regions, and are also distinct from the A/B region of TR α 1 [3]. TR α is mostly expressed in the brain [4] and is associated with the development of the nervous system [5]. TR α is constitutively localized within the nucleus where it interacts with nucleosomal DNA [6,7]. In the absence of T3 ligand, TR α is observed to actively repress transcription through interactions with transcriptional corepressors such as SMRT and NCoR [8–10].

Thyroid hormone receptors are members of the nuclear receptor (NR) superfamily of ligand-mediated transcription factors [2]. NRs have common modular structural features that include an N-terminal domain

(A/B domain, Fig. 1A). This A/B domain is of variable length and amino acid sequence and encompasses a ligand-independent transactivation function (AF1) domain that is critical for regulating transactivation [11,12]. Following the A/B domain is a highly conserved DNA-binding domain (DBD; C domain, Fig. 1A) that binds palindromic DNA sequences called hormone response elements (HRE). A short ‘hinge’ sequence (D domain) connects the DBD (C domain) to a C-terminal ligand-binding domain (LBD; E/F domain, Fig. 1A). Upon binding agonist-ligands, the LBD (E/F domain) undergoes conformational changes which results in the recruitment of coactivator molecules [13–17]. Antagonists and inverse agonists disrupt the ‘active-state’ LBD and the resulting LBD conformation functions as a docking site for corepressors [18–20]. Also, except for the A/B domains, the amino acid sequences of TR α and TR β are over 90% identical. Since TRs differ most significantly in the N-

Abbreviations

CD, circular dichroism; DBD, DNA-binding domain; DR4, direct repeat 4; ITC, isothermal titration calorimetry; LBD, ligand-binding domain; NR, nuclear receptor; T3, triiodothyronine; TRE, thyroid receptor response element; TR, thyroid hormone receptor.




Fig. 1. (A) NR domain topology displaying single-letter domain assignments. The region circled in green (above) is the focus of this study and the structural topology shows the relative orientation of the domains with DR4 TRE DNA (below). (B) The amino acid sequence of the TR α A/B + C domain molecular construct is colour coded (A/B domain in red and C domain in green). The single tryptophan is shown in blue. (C and D) Results from the *Escherichia coli* overexpression and purification of the TR α (A/B + DBD) and TR α (DBD) molecular constructs, respectively. The molecular weight standards are on lane 6 (the positions of the 75 kDa and 25 kDa standards are labelled) and the purified proteins are in lane 7.

terminal A/B domain, it is suggested that this region plays a significant role in mediating the distinct roles of these receptors [21]. It has also been proposed that TR α -mediated transcriptional regulation can also occur through specific interactions of the A/B domain with the PIC, specifically with transcription factor IIB (TFIIB) [21–24] and the TATA-binding protein (TBP) [25]. Transcriptional repression and similar interactions have also been observed between TR β and TFIIB [21,23,26].

By and large, the N-terminal domain of NRs is the least understood. This A/B region is diverse in size, sequence and is conformationally malleable [12,27,28], implying that this domain plays disparate roles in conferring cell type and/or promoter specificity [21]. Moreover, there are no data on the atomic resolution structure of any NR A/B domain conformation to date.

Nuclear receptor structure is strongly affected by the presence and even sequence of the DNA response element [29]. The source of these may result from conformational changes within the DBD as observed in structures of glucocorticoid (GR) bound to multiple

GREs [30,31]. This may explain, in part, the DNA-dependent interactions between the TR α DBD and LBD (E/F domain) reported earlier [32]. DNA binding is also central to allosteric communication between the A/B and C (DBD) domains [27,33–36]. Multiple DNA-binding site sequences have been identified for TR α . TR isoforms and oligomers exhibit preferential binding to specific DNA sequences called thyroid response elements (TRE) [37]. These TRE sequences consist of consensus AGGTCA (half-sites) arranged as direct repeats (DR), palindromic sequences (Pal) or inverted palindromic sequences (IP), each with differing spacing between the half-sites.

Allostery is a recognized regulatory feature within NRs such that ligand binding and even minor perturbations (such as nonbinding-site mutations) are detected at distal regions of NRs [15–17,27,38–41]. With distinct structural changes, allostery has been observed to link ligand, coactivator and the DNA-binding sites [17,32,42]. Furthermore, DNA binding is also central to allosteric communications between the A/B and C (DBD) domains [27,33–36,43]. Increasingly, cooperative interactions between multiple NR

domains have also been reported to modulate transactivation suggesting additional layers of regulation [32,44].

Here, we report a notable conformational change in the TR α A/B domain that is initiated through allostery through the TR α DBD by DNA. The shorter, 50-amino acid A/B domain of TR α encompasses several of the structural motifs that have been identified in NRs with significantly larger A/B domains to be important for ligand-independent activity [24]. Of these, distinct variations of the KRKRK amino acid sequence motif are common to several NRs including TR, progesterone (PR) and the liver X receptor (LXR) [28]. We are able to observe that the TR α A/B domain can allosterically enhance the binding affinity of the receptor for direct repeat 4 (DR4) TRE DNA. Furthermore, using a combination of circular dichroism (CD) and intrinsic tryptophan fluorescence spectroscopy, we can report that the binding of DNA to the TR α DBD (C domain) induces unfolding within the flanking TR α A/B domain. Overall, these observations suggest a structural basis for intramolecular cooperativity within TR α that fine-tunes binding to specific DNA sites.

Experimental procedures

Protein expression and purification

The chicken thyroid hormone receptor $\alpha 1$ gene (cTR $\alpha 1$, NCBI accession #: NP_990644.1) is over 90% identical to human TR $\alpha 1$ (NCBI accession #: NP_955366.1) at the amino acid level and is used for all experiments here. TR α (A/B + DBD, amino acid 1–154), TR α (DBD, amino acids 37–154) [32] and TR α (A/B domain, amino acids 1–50) were cloned into the plasmid pET15b (Life Technologies Inc., Carlsbad, CA, USA) to produce pET15b-TR α (A/B + DBD), pET15b-TR α (DBD) and pET15b-TR α (A/B domain), respectively. Proteins were produced in *Escherichia coli* BL21 (DE3) RIPL cells. Protein synthesis was induced with 0.5 mM isopropyl β -D-thiogalactoside (IPTG) at 20°C. Cells were lysed by sonication in 50 mM Tris, pH 8.0, 500 mM NaCl, 20 mM Imidazole, 10% glycerol, 1 protease inhibitor tablet, 5.7 mM β -mercaptoethanol, 0.5 μ M PMSF, 10 μ M ZnCl₂, 10 mM MgCl₂, recombinant DNase I (10 U). 6XHis-tagged TR (A/B + DBD) and TR (DBD) were purified using Ni-NTA agarose (Qiagen®, Germantown, MD, USA) with 0.3 M Imidazole, 50 mM Tris, pH 8.0, 500 mM NaCl and 10% glycerol. Proteins were further purified by size-exclusion chromatography (SEC) using S200 Superdex 16/60 column (GE Healthcare Life Sciences, Pittsburgh, PA, USA) in buffer consisting 50 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), pH = 7.5 (at 25 °C), 125 mM NaCl, 5 mM

MgCl₂, 1 mM tris (2-carboxyethyl)phosphine hydrochloride; TCEP). Protein was analysed using SDS/PAGE. Protein concentration was determined using the Bradford Assay (BioRad®, Hercules, CA, USA).

Preparation of DR4 TRE DNA adduct

19-mer DNA oligos containing the thyroid hormone response element (TRE) consensus site (DR4: 5'-CCAGGTCATTTTCAGGTCAG-3', where the underlined sequence is the NR binding site) were commercially obtained (Life Technologies Inc.) as single-stranded oligomers [45]. Double-stranded DR4 TRE was prepared by mixing the complementary strands in equimolar ratios to a final concentration of 2 mM, followed by heat denaturation at 95 °C for 5 min and annealing by gradual cooling to room temperature.

Isothermal titration calorimetry (ITC)

Thyroid receptor α (A/B + DBD) and TR α (DBD), purified by SEC, were used for isothermal titration calorimetry (ITC) measurements using VP-ITC MicroCal™ (MicroCal Inc., Northampton, MA, USA). Protein and ligand were prepared in 50 mM HEPES, pH 7.5, 125 mM NaCl, 5 mM MgCl₂ and 1 mM TCEP. For titration experiments, protein concentration ranged from 30 to 45 μ M and ligand DR4 TRE: 5'-CCAGGTCATTTTCAGGTCAG-3' concentration ranged from 300 to 400 μ M. Both protein and ligand were degassed for 5–10 min. The experiments were initiated by injecting 28 \times 10 μ L aliquots of DR4 TRE from the syringe into the calorimetric cell containing 1.5 mL of protein solution. All the titrations were performed at 25 °C and the buffer (pH adjusted to 7.5 at 25 °C). The change in thermal power as a function of each injection was automatically recorded using MICROCAL ORIGIN software and the raw data were further processed to yield binding isotherms of heat released per injection as a function of molar ratio of DR4 TRE to TR α (A/B + DBD) or TR α (C domain). The data were acquired and processed using the MICROCAL ORIGIN (MicroCal Inc.) software. Data were collected in triplicate.

Fluorescence spectroscopy

Fluorescence emission spectra of purified TR α (A/B + DBD) in 50 mM HEPES, pH 7.5, 125 mM NaCl, 5 mM MgCl₂, 1 mM TCEP were recorded at various concentrations of DR4 TRE. A total of 2 mL protein (2 μ M) was used to which 2 μ L of DR4 TRE (0–9.4 μ M) was added for each scan. To monitor the effect of sample dilution due to DR4 TRE titrations into protein, equal volumes of buffer were titrated into 2 mL protein (2 μ M). The spectra were monitored using a PerkinElmer-LS 55 Fluorescence Spectrometer at excitation wavelength of 295 nm at 300 nm \cdot min⁻¹.

Emission wavelength range was set at 310 nm to 450 nm, with slit width of 5.0 nm; 1 cm path length rectangular cuvettes were used to take all measurements at room temperature. The final fluorescence intensity change curve was a result of three averaged curves from individual experiments. The contribution of DR4-TRE to the TR α (A/B + DBD) + DR4 TRE spectrum was corrected by subtracting the spectrum of TR α (A/B + DBD) + buffer. Since multiple studies have shown that two molecules of TR bind a single TRE DNA [32,46], titration data curves were fitted to a two-site binding, nonlinear regression fitting model by PRISM7 (GraphPad software, La Jolla, CA, USA, www.graphpad.com), where change in fluorescence intensity was plotted against increasing concentrations of DR4 TRE ranging from 0.0 μM to 9.4 μM (Fig. 1B).

Circular dichroism (CD)

Circular dichroism spectra of TR α (A/B + DBD) and TR α (DBD; in 50 mM sodium phosphate buffer, pH = 7.5–8.0, 80 mM NaCl, and 5 mM MgCl₂, 1 mM TCEP) in the presence and absence of DR4 TRE DNA were recorded using a JASCO J-815 CD spectrometer. Protein to DNA ratio was 1: 1.1 for all experiments. All spectra were collected at 100 nm·min⁻¹ scan rate in 2 mm cuvettes maintained at 4 °C. The band width was 4 nm with data pitch 1 nm. CD spectra of buffer and DR4 TRE (4–5 μM) were also recorded separately as controls. Each spectrum shown is the result of 30 spectra accumulations, averaged and smoothed. All the spectra were corrected for the contributions of the buffer and TRE DR4 [47]. Mean residue ellipticity ([θ], (deg cm² dmol⁻¹)) was calculated using the CAPITO software [48].

Results

Here, we present data from studies on a 154-amino acid, two-domain molecular construct that encompasses the contiguous A/B (N terminus) and the C domains (DBD) of TR α (Fig. 1A). The TR α A/B domain comprises approximately 50 amino acids with an evolutionary conserved KRKRK motif (Fig. 1B) consisting of multiple charged residues [21,24]. Additionally, this construct contains a single tryptophan residue that is conveniently located within the A/B domain (¹⁹Trp) and adjacent to the KRKRK motif which has enabled us to monitor the local changes in conformation with steady-state intrinsic tryptophan fluorescence spectroscopy. In summary, we present data on the structural conformation of the TR α A/B domain, the conformational changes in this domain that are transmitted by allostery when the DBD (C domain) binds DNA, and the effect of the A/B domain on DNA recognition and binding.

The structural topology of TR α is shown in Fig. 1A. The two TR α constructs – TR α (A/B + DBD) and TR α (DBD), were purified to homogeneity as monomers of TR (A/B + DBD; 20.1 kDa) and TR (DBD; 15.9 kDa; Fig. 1C,D).

The TR α A/B \leftrightarrow DBD allostery influences the binding affinity for DNA

The selectivity for DNA is central to the transcriptional activity of NRs. Here, we provide evidence that allostery between the TR α A/B domain and the DBD also occurs in reverse, i.e. TR α A/B domain can influence the behaviour of the TR α (C domain only) vis-à-vis its DNA-binding affinity. Using ITC, we compare the binding affinity (K_d) of TR α (A/B + DBD) domains and TR α (DBD) for DR4 TRE DNA. We observe a three-fold increase in affinity of the intact TR α (A/B + DBD) domain for DR4 TRE DNA ($K_d = 2.31 \pm 0.21 \mu\text{M}$) over the truncated TR α DBD ($K_d = 6.65 \pm 0.50 \mu\text{M}$; Fig. 2). Also, the stoichiometry (N) of binding by both TR α (A/B + DBD) and TR α (DBD) is approximately $N = 0.5$ for TRE DR4, indicating that a single DR4 TRE binds two protein molecules. This is consistent with previous data showing two TR-interacting half-sites within the DR4 TRE [32,45]. Analyses of the thermodynamic parameters suggest that the TR α (A/B + DBD) \leftrightarrow DR4 TRE interaction is entropically less favourable ($T\Delta S = -2.05$ kcal/mol) than the corresponding entropic contributions to the TR α (DBD) \leftrightarrow DR4 TRE interactions ($T\Delta S = 1.08$ kcal·mol⁻¹). Therefore, it is likely that the higher affinity between TR α (A/B + DBD) and DR4 TRE is directed by the approximately 1.6-fold higher enthalpic contribution ($\Delta H = -9.63 \pm 1.20$ kcal·mol⁻¹) over the corresponding TR α (DBD) \leftrightarrow DR4 TRE interactions ($\Delta H = -5.98 \pm 0.43$ kcal·mol⁻¹; Table 1).

TRE binding to the DBD can influence specific local conformation of the A/B domain

Our studies above indicate that there is an allosteric pathway that links the DNA-binding site within the TR α DBD to the N-terminal TR α A/B domain (Fig. 2). Here, we sought to determine if the DNA-dependent allosteric communication between TR α A/B \leftrightarrow DBD is manifested in measurable conformational changes, specifically within the TR α A/B domain. Fortunately, there exists only a single Trp residue within the entire TR α (A/B + DBD) molecular construct. Furthermore, at position 19 this ¹⁹Trp is also both midway within the TR α A/B domain (residues 1–50) and distal from the DNA-binding




Fig. 2. ITC measurements were performed to measure heat changes upon titrating DR4 TRE DNA into (A). TR α (A/B + DBD) and (B). TR α (DBD). For all titrations, the c values ($c = nK_aM_{tot}$, where n is the stoichiometry parameter, K_a is the association constant = $1/K_d$ and M_{tot} is the concentration of the macromolecule, TR α) range from 6.5 to 9, which is within the ideal range for determining binding constants by ITC [73]. Data obtained are summarized in Table 1.

Table 1. Thermodynamic parameters of TRE DR4 interaction with TR α (A/B + DBD) and TR α DBD. Parameters are determined at 25 °C and pH = 7.5, as described in Experimental Procedures. The reported values are the average of three experiments and the errors are the standard deviation.

Protein complexes	K_d (μ M)	ΔH (kcal·Mol $^{-1}$)	N^a	ΔG (kcal·Mol $^{-1}$)	$T\Delta S$ (kcal·Mol $^{-1}$)
TR α (A/B + DBD) + DR4	2.31 ± 0.21	-9.63 ± 1.20	0.54 ± 0.03	-7.68	-2.05
TR α (DBD) + DR4	6.65 ± 0.50	-5.98 ± 0.43	0.53 ± 0.02	-7.06	1.08

^a The apparent stoichiometry from the curve fitting data.

TR α DBD (residues 51–154; Fig. 1B). Thus, this single *Trp* enables us to directly identify conformational changes within the central region of TR α (A/

B + DBD). *Trp* fluorescence quenching has been a common indicator of local and global conformational changes within the NR A/B domains [35,49,50] and

due to allostery [16]. We monitored the dose-dependent changes in intrinsic steady-state tryptophan fluorescence, accompanied by an approximately 5 nm red-shift in fluorescence maxima, within TR α (A/B + DBD) in the presence of DR4 TRE (Fig. 3A). The measurable decrease in fluorescence suggests a specific change in the ^{19}Trp conformation, and furthermore, the conformational changes within the ^{19}Trp sidechain are more likely from a progressive decrease in its local hydrophobic environment, presumably from an increased exposure to the surrounding buffer [16]. These titrations were also analysed to provide a quantitative measure of binding affinity: since the two DR4 half-sites are indistinguishable for binding TR α [32], the average binding affinity of TR α (A/B + DBD) for DR4 TRE is $K_d = 2.69 \pm 0.22 \mu\text{M}$. This binding constant confirms data obtained by calorimetry.

TRE binding to the DBD results in unfolding of the TR α A/B domain

The spectroscopic analyses above suggest an allosteric conformational change within the TR α A/B domain upon binding DNA at the TR α (DBD). To determine the specific DNA-dependent changes in structure within the TR α A/B domain, we utilized CD spectroscopy. Given that minor changes in the secondary structure of proteins can be detected in the raw CD spectra (θ in rad cm^{-1} vs. wavelength in nm) in the far-UV ($\lambda = 190\text{--}260$ nm) range, we compared the CD spectra of the TR α (A/B + DBD) domains with TR α (DBD) in the absence and when complexed with DR4 TRE (Fig. 3B). For the TR α (DBD), there is a prominent change in the minima at 208 nm and 222 nm of the CD spectrum in the presence of DNA, which suggests a significant increase in α -helical structure of the




Fig. 3. Conformational changes determined by Fluorescence and CD spectroscopy. (A) Change in intrinsic tryptophan fluorescence of TR α (A/B + DBD) is monitored in response to increasing levels of DR4 TRE DNA. The data above are obtained after subtracting buffer and DR4 TRE DNA contributions. In addition, no static quenching of molecular *Trp* was observed by DR4 TRE DNA. (B and C) Raw CD spectra of TR α (A/B + DBD) and TR α (DBD), respectively, \pm DR4 TRE DNA. (D) The CD (θ), ($\text{deg cm}^2 \text{dmol}^{-1}$) vs. wavelength, (nm) spectra of the TR α (A/B domain) was calculated by individually subtracting the θ values for TR α (DBD) from TR α (A/B + DBD), for each corresponding wavelength, \pm DR4 TRE DNA, respectively. The assumption made is that the conformations of the TR α (C domain), \pm DR4 TRE DNA, are the same in both TR α (A/B + DBD) and TR α (DBD). Inset, CD spectra of TR α (A/B domain) measured directly \pm DR4 TRE.

TR(DBD) upon binding DNA (Fig. 3C). Such conformational changes in NR DBDs have been previously observed using NMR spectroscopy confirming a dosage-dependent stabilization of the NR DBD upon binding DNA [51–55]. In this study, the CD spectra of TR α (A/B + DBD) indicates that while the TR α segment is predominantly α -helical, the complexation of TR α (A/B + DBD) with DNA results in a markedly smaller change in secondary structure from the DNA-free protein when compared with the corresponding structural changes within the TR α DBD-only (Fig. 3C). To determine the source of this discrepancy between the TR α (A/B + DBD) and TR α DBD, we subtracted the spectroscopically measured molar ellipticity of CD of the TR α (DBD) from the TR α (A/B + DBD) domain. The resulting spectrum estimates the ‘calculated’ molar ellipticity ($[\theta]$, (deg cm² dmol⁻¹), and therefore the conformational change, of the TR α (A/B domain) within the TR α (A/B + DBD):DNA complex (Fig. 3D). Additionally, we do not detect significant secondary structure changes to the isolated TR α (A/B domain) in the presence of DR4 TRE (Fig. 3D inset). Taken together, these results suggest that the TR α (A/B domain) has partial α -helical secondary structure within the ‘DNA-free’ TR α (A/B + DBD). Upon binding DNA, the contiguous A/B domain and the DBD undergo contrasting conformational changes – while the A/B domain appears to convert from a more structured to a conformationally less-rigid state, the DBD becomes conformationally more stable. Overall, this α -helical-to-random coil unfolding of the TR α A/B domain appears to counteract the propensity for greater α -helicity within the TR α (DBD) upon binding DR4 TRE. This may explain, in part, the smaller overall change in TR α (A/B + DBD) in comparison with the TR α (DBD), upon binding DR4 TRE.

Discussion

Multiple lines of evidence suggest that the NR A/B domains are flexible and can adopt distinct conformations through allostery initiated by DNA:DBD interactions [12,34,43,49,50,56–59]. A common observation is that the A/B domains in all NRs studied to date, the DNA-initiated allostery elicits an increase in secondary structure (mostly α -helicity) of this domain.

Multiple attempts to determine the structures of full-length NRs have failed to identify the conformation of their N-terminal domains [60]. Yet, all these structures have indicated that there is no apparent direct interaction between the A/B domain and the DBD. Our observations suggest that DNA-dependent

conformational changes within the TR α A/B domain are distinct from the corresponding changes within the other NR A/B domains listed above. The implications for the unique mode of TR α A/B domain \leftrightarrow DBD allostery are broad. For instance, the TR α A/B domain is reported to interact with several cellular cofactors including TFIIB [21–24] and TBP [25]. Similar interactions have been observed between NRs and the PIC, such as the androgen (AR) [61,62], COUP-TF [63], oestrogen (ER) [63,64], GR [65], mineralocorticoid (MR) [66] and PR [34,63] receptors, among others. In each of these NRs, and distinct from TR α as reported here, the A/B domain is constrained to a more folded conformation by DNA-allostery. This more-structurally constrained A/B domain is observed to enhance the NR \leftrightarrow cofactor interaction.

In TR α , the sequence of basic residues ²³KRKR²⁷K has been identified to make specific interactions with TFIIB (Fig. 1B) [24]. Adjacent to this basic motif is ¹⁹Trp, which we show here by DR4 TRE DNA dose-dependent fluorescence quenching to undergo conformational changes to a more exposed environment and this would be expected with the unfolding of this region of the TR α A/B domain. From truncation and associated binding studies, the corresponding TR α -interacting domain of TFIIB is identified to be contained within residues 178–201 of an amphipathic α -helix [24]. Curiously, this TR α -interacting TFIIB α -helix has also separately been identified as integral to the binding interface between TFIIB and DNA [67]. Together, these studies suggest that the formation of the TR α :TFIIB and the TFIIB:DNA complexes are mutually exclusive and that binding to TR α can disrupt the TFIIB-DNA complex. In the absence of direct structural data, it is tempting to speculate that the DNA-induced unfolding of the TR α A/B domain plays a role in inserting itself into the TFIIB-DNA complex and the newly created TR α :TFIIB is stabilized by both interactions made by the charged ²³KRKR²⁷K and through the exposed apolar backbone of the TR α A/B domain. Indeed, such DNA-induced unfolding events are less commonly reported in the literature and the Ets-1 transcription factor is a singular prior example of an analogous DNA-induced unfolding within a flanking domain through allostery [68,69]. In Ets-1, this induced unfolding is proposed to ameliorate inhibitory intramolecular interactions and encourage intermolecular interactions that promote gene transcription.

Additionally, this study reinforces the observation that DNA recognition is finely tuned by the domains flanking the NR DBD. In both DNA-bound TR α :

RXR heterodimeric [45] and the TR β monomeric [46] structures, the conformation of the TR DBD is virtually identical, suggesting a generic mechanism for DNA recognition and binding. Yet, using DBD and DBD-LBD constructs of TR α , we have earlier established that the affinity of the DBD for DNA can be modulated through intramolecular allostery [32]. Moreover, even subtle changes within these flanking domains (A/B or E/F domains) such as mutations [70] and interactions with cellular factors [32] or small-molecule ligands [71] can affect DNA binding. Given the distinct unfolding process of the TR α A/B domain, the mechanism by which this domain can allosterically influence DBD \leftrightarrow DNA interactions is likely to be different from those of AR [35] and PR [72].

In summary, our data here suggest a distinct consequence of allostery within TR α . The data from CD spectroscopy show that conformational changes induced within the TR(DBD) are transmitted ‘upstream’ to the flanking A/B domain. The resultant conformation of the TR α A/B domain is less ordered within the intact, DNA-bound TR α (A/B + DBD) than in the absence of DNA. This unfolding results in the repositioning of ¹⁹Trp observed from the quenching of tryptophan fluorescence. The unusual feature of DNA-induced, allosterically driven conformational changes within the TR α A/B domain is the overall loss in secondary structure, quantified as a decrease in its α -helicity. Finally, this study showcases the diversity in the structural response to allostery within the NR superfamily. We are drawn to hypothesize that such structural responses have been evolutionarily selected to optimize the specific behaviour of individual members of these NR transcription factors.

Acknowledgements

We thank Drs. Francisco Barrera, Chris Dealwis, Engin Serpersu and Edward Wright for reading the manuscript. William Alexander and B.-D. Kumar Putcha assisted with molecular cloning. Nina Martyris assisted with editing. CR is a PEER fellow and JA is a Chancellors Undergraduate Fellow at UTK. EJF is a recipient of funds from the Donald L. Akers foundation at UTK.

Author contributions

EJF designed and performed experiments, analysed data and wrote the manuscript. VG performed experiments and analysed data and CR and JA performed experiments.

References

- 1 Thompson CC, Weinberger C, Lebo R and Evans RM (1987) Identification of a novel thyroid hormone receptor expressed in the mammalian central nervous system. *Science* **237**, 1610–1614.
- 2 Lazar MA (1993) Thyroid hormone receptors: multiple forms, multiple possibilities. *Endocr Rev* **14**, 184–193.
- 3 Hodin RA, Lazar MA and Chin WW (1990) Differential and tissue-specific regulation of the multiple rat c-erbA messenger RNA species by thyroid hormone. *J Clin Invest* **85**, 101–105.
- 4 Cheng SY, Leonard JL and Davis PJ (2010) Molecular aspects of thyroid hormone actions. *Endocr Rev* **31**, 139–170.
- 5 Bradley DJ, Towle HC and Young WS 3rd (1992) Spatial and temporal expression of alpha- and beta-thyroid hormone receptor mRNAs, including the beta 2-subtype, in the developing mammalian nervous system. *J Neurosci* **12**, 2288–2302.
- 6 Wong J, Shi YB and Wolffe AP (1995) A role for nucleosome assembly in both silencing and activation of the *Xenopus* TR beta A gene by the thyroid hormone receptor. *Genes Dev* **9**, 2696–2711.
- 7 Grontved L., Waterfall J. J., Kim D. W., Baek S., Sung M.-H., Zhao L., Park J. W., Nielsen R., Walker R. L., Zhu Y. J. *et al.* (2015) Transcriptional activation by the thyroid hormone receptor through ligand-dependent receptor recruitment and chromatin remodelling. *Nat Commun* **6**, 7048–7058.
- 8 Chen JD and Evans RM (1995) A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* **377**, 454–457.
- 9 Horlein A. J., Naar A. M., Heinzel T., Torchia J., Gloss B., Kurokawa R., Ryan A., Kamei Y., Soderstrom M., Glass C. K. *et al.* (1995) Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor. *Nature* **377**, 397–404.
- 10 Tong GX, Jeyakumar M, Tanen MR and Bagchi MK (1996) Transcriptional silencing by unliganded thyroid hormone receptor beta requires a soluble corepressor that interacts with the ligand-binding domain of the receptor. *Mol Cell Biol* **16**, 1909–1920.
- 11 Tsai MJ and O’Malley BW (1994) Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu Rev Biochem* **63**, 451–486.
- 12 Lavery DN and McEwan IJ (2005) Structure and function of steroid receptor AF1 transactivation domains: induction of active conformations. *Biochem J* **391**, 449–464.
- 13 Suino K, Peng L, Reynolds R, Li Y, Cha JY, Repa JJ, Klier SA and Xu HE (2004) The nuclear xenobiotic receptor CAR; structural determinants of constitutive activation and heterodimerization. *Mol Cell* **16**, 893–905.

- 14 Xu RX, Lambert MH, Wisely BB, Warren EN, Weinert EE, Waitt GM, Williams JD, Collins JL, Moore LB, Willson TM *et al.* (2004) A structural basis for constitutive activity in the human CAR/RXRalpha heterodimer. *Mol Cell* **16**, 919–928.
- 15 Wright E, Vincent J and Fernandez EJ (2007) Thermodynamic characterization of the interaction between CAR-RXR and SRC-1 peptide by isothermal titration calorimetry. *Biochemistry* **46**, 862–870.
- 16 Wright E, Busby SA, Wisecarver S, Vincent J, Griffin PR and Fernandez EJ (2011) Helix 11 dynamics is critical for constitutive androstane receptor activity. *Structure* **19**, 37–44.
- 17 Putcha BD, Wright E, Brunzelle JS and Fernandez EJ (2012) Structural basis for negative cooperativity within agonist-bound TR:RXR heterodimers. *Proc Natl Acad Sci USA* **109**, 6084–6087.
- 18 Xu HE, Stanley TB, Montana VG, Lambert MH, Shearer BG, Cobb JE, McKee DD, Galardi CM, Plunket KD, Nolte RT *et al.* (2002) Structural basis for antagonist-mediated recruitment of nuclear co-repressors by PPARalpha. *Nature* **415**, 813–817.
- 19 Dussault I, Lin M, Hollister K, Fan M, Termini J, Sherman MA and Forman BM (2002) A structural model of the constitutive androstane receptor defines novel interactions that mediate ligand-independent activity. *Mol Cell Biol* **22**, 5270–5280.
- 20 Shan L, Vincent J, Brunzelle JS, Dussault I, Lin M, Ianculescu I, Sherman MA, Forman BM and Fernandez EJ (2004) Structure of the murine constitutive androstane receptor complexed to androstenol: a molecular basis for inverse agonism. *Mol Cell* **16**, 907–917.
- 21 Tomura H, Lazar J, Phyllaier M and Nikodem VM (1995) The N-terminal region (A/B) of rat thyroid hormone receptors alpha 1, beta 1, but not beta 2 contains a strong thyroid hormone-dependent transactivation function. *Proc Natl Acad Sci USA* **92**, 5600–5604.
- 22 Fondell JD, Roy AL and Roeder RG (1993) Unliganded thyroid hormone receptor inhibits formation of a functional preinitiation complex: implications for active repression. *Genes Dev* **7**, 1400–1410.
- 23 Baniahmad A, Ha I, Reinberg D, Tsai S, Tsai MJ and O'Malley BW (1993) Interaction of human thyroid hormone receptor beta with transcription factor TFIIB may mediate target gene derepression and activation by thyroid hormone. *Proc Natl Acad Sci USA* **90**, 8832–8836.
- 24 Hadzic E, Desai-Yajnik V, Helmer E, Guo S, Wu S, Koudinova N, Casanova J, Raaka BM and Samuels HH (1995) A 10-amino-acid sequence in the N-terminal A/B domain of thyroid hormone receptor alpha is essential for transcriptional activation and interaction with the general transcription factor TFIIB. *Mol Cell Biol* **15**, 4507–4517.
- 25 Fondell JD, Brunel F, Hisatake K and Roeder RG (1996) Unliganded thyroid hormone receptor alpha can target TATA-binding protein for transcriptional repression. *Mol Cell Biol* **16**, 281–287.
- 26 Baniahmad A, Tsai SY, O'Malley BW and Tsai MJ (1992) Kindred S thyroid hormone receptor is an active and constitutive silencer and a repressor for thyroid hormone and retinoic acid responses. *Proc Natl Acad Sci USA* **89**, 10633–10637.
- 27 Hilser VJ and Thompson EB (2011) Structural dynamics, intrinsic disorder, and allostery in nuclear receptors as transcription factors. *J Biol Chem* **286**, 39675–39682.
- 28 Hill KK, Roemer SC, Churchill ME and Edwards DP (2012) Structural and functional analysis of domains of the progesterone receptor. *Mol Cell Endocrinol* **348**, 418–429.
- 29 Wood JR, Greene GL and Nardulli AM (1998) Estrogen response elements function as allosteric modulators of estrogen receptor conformation. *Mol Cell Biol* **18**, 1927–1934.
- 30 Meijising SH, Pufall MA, So AY, Bates DL, Chen L and Yamamoto KR (2009) DNA binding site sequence directs glucocorticoid receptor structure and activity. *Science* **324**, 407–410.
- 31 Watson LC, Kuchenbecker KM, Schiller BJ, Gross JD, Pufall MA and Yamamoto KR (2013) The glucocorticoid receptor dimer interface allosterically transmits sequence-specific DNA signals. *Nat Struct Mol Biol* **20**, 876–883.
- 32 Putcha BD and Fernandez EJ (2009) Direct interdomain interactions can mediate allostery in the thyroid receptor. *J Biol Chem* **284**, 22517–22524.
- 33 Kumar R, Baskakov IV, Srinivasan G, Bolen DW, Lee JC and Thompson EB (1999) Interdomain signaling in a two-domain fragment of the human glucocorticoid receptor. *J Biol Chem* **274**, 24737–24741.
- 34 Bain DL, Franden MA, McManaman JL, Takimoto GS and Horwitz KB (2000) The N-terminal region of the human progesterone A-receptor. Structural analysis and the influence of the DNA binding domain. *J Biol Chem* **275**, 7313–7320.
- 35 Brodie J and McEwan IJ (2005) Intra-domain communication between the N-terminal and DNA-binding domains of the androgen receptor: modulation of androgen response element DNA binding. *J Mol Endocrinol* **34**, 603–615.
- 36 Kumar R and McEwan IJ (2012) Allosteric modulators of steroid hormone receptors: structural dynamics and gene regulation. *Endocr Rev* **33**, 271–299.
- 37 Velasco LF, Togashi M, Walfish PG, Pessanha RP, Moura FN, Barra GB, Nguyen P, Rebong R, Yuan C, Simeoni LA *et al.* (2007) Thyroid hormone response

- element organization dictates the composition of active receptor. *J Biol Chem* **282**, 12458–12466.
- 38 Forman BM, Umesono K, Chen J and Evans RM (1995) Unique response pathways are established by allosteric interactions among nuclear hormone receptors. *Cell* **81**, 541–550.
- 39 Shulman AI, Larson C, Mangelsdorf DJ and Ranganathan R (2004) Structural determinants of allosteric ligand activation in RXR heterodimers. *Cell* **116**, 417–429.
- 40 Pavlin MR, Brunzelle JS and Fernandez EJ (2014) Agonist Ligands Mediate the Transcriptional Response of Nuclear Receptor Heterodimers through Distinct Stoichiometric Assemblies with Coactivators. *J Biol Chem* **289**, 24771–24778.
- 41 Johnson Q. R., Lindsay R. J., Nellas R. B., Fernandez E. J. and Shen T. (2015) Mapping allostery through computational glycine scanning and correlation analysis of residue-residue contacts. *Biochemistry* **54**, 1534–1541.
- 42 Schulman IG, Li C, Schwabe JW and Evans RM (1997) The phantom ligand effect: allosteric control of transcription by the retinoid X receptor. *Genes Dev* **11**, 299–308.
- 43 Bain DL, Franden MA, McManaman JL, Takimoto GS and Horwitz KB (2001) The N-terminal region of human progesterone b-receptors: biophysical and biochemical comparison to a-receptors. *J Biol Chem* **276**, 23825–23831.
- 44 Kraus WL, McInerney EM and Katzenellenbogen BS (1995) Ligand-dependent, transcriptionally productive association of the amino- and carboxyl-terminal regions of a steroid hormone nuclear receptor. *Proc Natl Acad Sci USA* **92**, 12314–12318.
- 45 Rastinejad F, Perlmann T, Evans RM and Sigler PB (1995) Structural determinants of nuclear receptor assembly on DNA direct repeats. *Nature* **375**, 203–211.
- 46 Chen Y and Young MA (2010) Structure of a thyroid hormone receptor DNA-binding domain homodimer bound to an inverted palindrome DNA response element. *Mol Endocrinol* **24**, 1650–1664.
- 47 Greenfield NJ (2007) Using circular dichroism collected as a function of temperature to determine the thermodynamics of protein unfolding and binding interactions. *Nat Protoc* **1**, 2527–2535.
- 48 Wiedemann C, Bellstedt P and Grolach M (2013) CAPITO—a web server-based analysis and plotting tool for circular dichroism data. *Bioinformatics* **29**, 1750–1757.
- 49 Kumar R, Lee JC, Bolen DW and Thompson EB (2001) The conformation of the glucocorticoid receptor af1/tau1 domain induced by osmolyte binds co-regulatory proteins. *J Biol Chem* **276**, 18146–18152.
- 50 Reid J, Kelly SM, Watt K, Price NC and McEwan IJ (2002) Conformational analysis of the androgen receptor amino-terminal domain involved in transactivation. Influence of structure-stabilizing solutes and protein-protein interactions. *J Biol Chem* **277**, 20079–20086.
- 51 Knegtel RMA, Katahira M, Schilthuis JG, Bonvin AMJJ, Boelens R, Eib D, van der Saag PT and Kaptein R (1993) The solution structure of the human retinoic acid receptor- β DNA-binding domain. *J Biomol NMR* **3**, 1–17.
- 52 Holmbeck SM, Foster MP, Casimiro DR, Sem DS, Dyson HJ and Wright PE (1998) High-resolution solution structure of the retinoid X receptor DNA-binding domain. *J Mol Biol* **281**, 271–284.
- 53 Holmbeck SM, Dyson HJ and Wright PE (1998) DNA-induced conformational changes are the basis for cooperative dimerization by the DNA binding domain of the retinoid X receptor. *J Mol Biol* **284**, 533–539.
- 54 van Tilborg PJA, Mulder FAA, de Backer MME, Nair M, van Heerde EC, Folkers G, van der Saag PT, Karimi-Nejad Y, Boelens R and Kaptein R (1999) Millisecond to microsecond time scale dynamics of the retinoid X and retinoic acid receptor DNA-binding domains and dimeric complex formation. *Biochemistry* **38**, 1951–1956.
- 55 Rastinejad F, Wagner T, Zhao Q and Khorasanizadeh S (2000) Structure of the RXR-RAR DNA-binding complex on the retinoic acid response element DR1. *EMBO J* **19**, 1045–1054.
- 56 Baskakov IV, Kumar R, Srinivasan G, Ji YS, Bolen DW and Thompson EB (1999) Trimethylamine N-oxide-induced cooperative folding of an intrinsically unfolded transcription-activating fragment of human glucocorticoid receptor. *J Biol Chem* **274**, 10693–10696.
- 57 McEwan IJ, Lavery D, Fischer K and Watt K (2007) Natural disordered sequences in the amino terminal domain of nuclear receptors: lessons from the androgen and glucocorticoid receptors. *Nucl Recept Signal* **5**, e001.
- 58 Kumar R., Moure C. M., Khan S. H., Callaway C., Grimm S., Goswami D., Griffin P. R. and Edwards D. P. (2013) Regulation of the structurally dynamic amino-terminal domain of progesterone receptor by protein induced folding. *J Biol Chem* **288**, 30285–30299.
- 59 Simons SS Jr, Edwards DP and Kumar R (2014) Minireview: dynamic structures of nuclear hormone receptors: new promises and challenges. *Mol Endocrinol* **28**, 173–182.
- 60 Khorasanizadeh S and Rastinejad F (2016) Visualizing the architectures and interactions of nuclear receptors. *Endocrinology* **157**, 4212–4221.
- 61 Ford J, McEwan IJ, Wright AP and Gustafsson JA (1997) Involvement of the transcription factor IID protein complex in gene activation by the N-terminal transactivation domain of the glucocorticoid receptor *in vitro*. *Mol Endocrinol* **11**, 1467–1475.

- 62 Reid J, Murray I, Watt K, Betney R and McEwan IJ (2002) The androgen receptor interacts with multiple regions of the large subunit of general transcription factor TFIIF. *J Biol Chem* **277**, 41247–41253.
- 63 Ing NH, Beekman JM, Tsai SY, Tsai MJ and O'Malley BW (1992) Members of the steroid hormone receptor superfamily interact with TFIIB (S300-II). *J Biol Chem* **267**, 17617–17623.
- 64 Sabbah M, Kang KI, Tora L and Redeuilh G (1998) Oestrogen receptor facilitates the formation of preinitiation complex assembly: involvement of the general transcription factor TFIIB. *Biochem J* **336**, 639–646.
- 65 McEwan IJ, Wright AP and Gustafsson JA (1997) Mechanism of gene expression by the glucocorticoid receptor: role of protein-protein interactions. *BioEssays* **19**, 153–160.
- 66 Fischer K, Kelly SM, Watt K, Price NC and McEwan IJ (2010) Conformation of the mineralocorticoid receptor N-terminal domain: evidence for induced and stable structure. *Mol Endocrinol* **24**, 1935–1948.
- 67 Tsai FT and Sigler PB (2000) Structural basis of preinitiation complex assembly on human pol II promoters. *EMBO J* **19**, 25–36.
- 68 Petersen JM, Skalicky JJ, Donaldson LW, McIntosh LP, Alber T and Graves BJ (1995) Modulation of transcription factor Ets-1 DNA binding: DNA-induced unfolding of an alpha helix. *Science* **269**, 1866–1869.
- 69 Newman M, Strzelecka T, Dorner LF, Schildkraut I and Aggarwal AK (1995) Structure of Bam HI endonuclease bound to DNA: partial folding and unfolding on DNA binding. *Science* **269**, 656–663.
- 70 Helsen C, Dubois V, Verfaillie A, Young J, Trekels M, Vancraenenbroeck R, De Maeyer M and Claessens F (2012) Evidence for DNA-binding domain–ligand-binding domain communications in the androgen receptor. *Mol Cell Biol* **32**, 3033–3043.
- 71 Chandra V, Huang P, Hamuro Y, Raghuram S, Wang Y, Burris TP and Rastinejad F (2008) Structure of the intact PPAR-gamma-RXR- nuclear receptor complex on DNA. *Nature* **456**, 350–356.
- 72 Connaghan-Jones KD, Heneghan AF, Miura MT and Bain DL (2007) Thermodynamic analysis of progesterone receptor-promoter interactions reveals a molecular model for isoform-specific function. *Proc Natl Acad Sci USA* **104**, 2187–2192.
- 73 Wiseman T, Williston S, Brandts JF and Lin LN (1989) Rapid measurement of binding constants and heats of binding using a new titration calorimeter. *Anal Biochem* **179**, 131–137.