Open Access

Guillain-Barré Syndrome: Prevalence and Long-Term Factors Impacting Bladder Function in an Australian Community Cohort

Bhasker Amatya, a Fary Khan, a,b,c Michael Whishaw, Julie F. Pallante

http://dx.doi.org/10.3988/jcn.2013.9.3.144

Table 5 and Fig. 2 were inadvertently omitted from the original publication of this article. The errors took place during publication process. The editor-in-chief apologizes for the mistake. Table 5 and Fig. 2 are presented here:

Table 5. Descriptive statistics and Spearman correlations among bladder scales with other measures

	Descriptive statistics			Spearman correlations			
	Cronbach a	Mean	SD	AUA	UDI6	IIQ7	AUA QoL
AUA	0.87	5.7	7.5	1.0	0.79 [†]	0.56 [†]	0.86 [†]
UDI6	0.89	8.3	3.4	0.79 [†]	1.0	0.79 [†]	0.79†
IIQ7	0.89	1.3	2.7	0.56 [†]	0.79 [†]	1.0	0.63 [†]
AUA QoL	NA	1.6	1.6	0.85 [†]	0.78 [†]	0.63 [†]	1.0
FIM-self care		37.9	5.4	-0.17	-0.15	-0.09	-0.1
FIM-sphincter		12.8	1.3	-0.23	-0.25*	-0.26*	-0.23
FIM-mobility		18.9	2.6	-0.29*	-0.33 [†]	-0.25*	-0.26*
FIM-locomotion		12.0	1.8	-0.34 [†]	-0.39 [†]	-0.34 [†]	-0.31*
DASS-depression		5.5	8.1	0.35 [†]	0.47†	0.35 [†]	0.35 [†]
DASS-anxiety		5.6	7.1	0.39†	0.44 [†]	0.42 [†]	0.37 [†]
DASS-stress		8.6	10.7	0.29*	0.47†	0.41†	0.37 [†]
PIPP-psychological		10.0	6.0	0.24*	0.28*	0.27*	0.27*
PIPP-selfcare		4.7	2.7	0.32	0.19	0.02	0.12
PIPP-mobility		8.2	4.9	0.35 [†]	0.31*	0.24	0.26*
PIPP-participation		9.7	6.0	0.39†	0.37†	0.26*	0.3*
PIPP-relation		5.2	3.1	0.37 [†]	0.31*	0.21	0.33 [†]

^{*}Correlation significant at the 0.05 level (2-tailed), †Correlation significant at the 0.01 level (2-tailed).

AUA: American Urological Association Symptom Index, DASS: Depression Anxiety Stress Scale-21, FIM: functional Independence measure, IIQ7: Incontinence Impact Questionnaire, PIPP: perceived impact of problem profile, QoL: quality of life, SD: standard deviation, UD16: Urogenital Distress Inventory.

^aDepartment of Rehabilitation Medicine, Royal Melbourne Hospital, Victoria, Australia

^bDepartment of Medicine, Dentistry and Health Sciences, University of Melbourne, Victoria, Australia

^cSchool of Public Health and Preventive Medicine, Monash University, Victoria, Australia

dRoyal Park Continence Service, Royal Melbourne Hospital, Victoria, Australia

^eRural Health Academic Centre, University of Melbourne, Victoria, Australia

J Clin Neurol 2013;9:144-150

[®] This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/ licenses/by-nc/3.0) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Erratum




Fig. 2. Total American Urological Association Symptom Index (AUA) scores for participants scoring 0 to 2 versus 3 to 6 on the single quality of life (QoL) item.