

Hepatoprotective Effect of Gallotannin-enriched Extract Isolated from Gall on Hydrogen Peroxide-induced Cytotoxicity in HepG2 Cells

Jun Go, Ji Eun Kim, Eun Kyoung Koh, Sung Hwa Song, Hyun Gu Kang¹, Young Hee Lee², Han Do Kim², Jin Tae Hong³, Dae Youn Hwang

Department of Biomaterials Science, College of Natural Resources and Life Science/Life and Industry Convergence Research Institute, Pusan National University, Miryang 627-706, ¹Laboratory of Veterinary Theriogenology, College of Veterinary Medicine, Chungbuk National University, Cheongju 362-763, ²Department of Organic Material Science and Engineering, Pusan National University, Busan 609-735, ³Department of Pharmacy, Chungbuk National University, Cheongju 361-763, Korea

Submitted: 04-11-2015

Revised: 28-12-2015

Published: 11-07-2017


ABSTRACT

Background: Gall (*Galla Rhois* [GR]) is known to have antibacterial, anti-inflammatory, antimetastatic, and anti-invasion activities and exert hepatoprotective effects. However, the hepatoprotective effects of gallotannin-enriched GR (GEGR) and their mechanisms have not yet been investigated. **Objective:** The potential protective effect of GEGR against hepatotoxicity induced by hydrogen peroxide (H_2O_2) was investigated. **Materials and Methods:** Changes in cell viability, apoptosis protein expression, and reactive oxygen species (ROS) generation were determined in HepG2 cells that were pretreated with four different concentrations of GEGR (6.25–50 $\mu\text{g/ml}$) for 24 h before H_2O_2 exposure. **Results:** GEGR consisted of gallotannin (69.2%), gallic acid (26.6%), and methyl gallate (4.2%) and showed remarkable 2,2-diphenyl-1-picrylhydrazyl scavenging activity (inhibitory concentration 50% = 0.212 $\mu\text{g/ml}$). The lethal dose 50% and effective dose 50% values for the response of HepG2 cells to GEGR were determined to be 178 and 6.85 $\mu\text{g/ml}$, respectively. Significant reductions in the immunofluorescence intensity indicating apoptosis were also detected in the nuclei of HepG2 cells stained with 4',6-diamidino-2-phenylindole and Annexin V after GEGR treatment. The Bax/Bcl-2 ratio and active caspase-3 level were higher in H_2O_2 + vehicle-treated cells. However, these levels gradually decreased to those of the No-treated group in the GEGR pretreated group even though little or no decrease was observed in response to low GEGR concentrations. Furthermore, the GEGR pretreated group showed a reduced level of 2',7'-dichlorofluorescein diacetate stained cells, indicating ROS generation relative to the H_2O_2 + vehicle-treated group. **Conclusion:** The results of this study provide strong evidence that GEGR can prevent cell death induced by H_2O_2 in HepG2 cells through the induction of antioxidant conditions. **Key words:** Antioxidant, Caspase-3, cell death, hepatotoxicity, reactive oxygen species

SUMMARY

- The gallotannin (69.2%), gallic acid (26.6%), and methyl gallate (4.2%) are the main constituents of water extracts of GR
- GEGR was more potent in DPPH scavenging, and gallotannin contributes to this extract activity

- GEGR significantly reduced the increase of apoptosis, Bax/Bcl-2 ratio, and active caspase-3 level after H_2O_2 treatment
- GEGR pretreatment showed protection against H_2O_2 -induced ROS production in DCFH-DA staining analysis.



Abbreviations used: COX: Cyclooxygenase; DAPI: 4',6-diamidino-2-phenylindole; DMSO: Dimethyl sulfoxide; DPPH: 2,2-diphenyl-1-picrylhydrazyl; GEGR: Gallotannin-enriched Galla Rhois; GR: Galla Rhois; HPLC: High-performance liquid chromatography; H_2O_2 : Hydrogen peroxide; MMP: Metalloproteinase; MTT: 3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide; ROS: Reactive oxygen species; UV-Vis: Ultraviolet-visible.

Correspondence:

Prof. Dae Youn Hwang,
Department of Biomaterial Science, College of Natural Resources and Life Science, Pusan National University, 50 Cheonghak-ri, Samnangjin-eup Miryang-si, Gyeongsangnam-do 627-706, Korea.
E-mail: dyhwang@pusan.ac.kr
DOI: 10.4103/pm.pm_424_15

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

Gall (*Galla Rhois* [GR]) is the excrescence formed by parasitic aphids, primarily *Schlechtendalia chinensis* Bell, on the leaves of the nutgall sumac tree, *Rhus javanica* L. (*Anacardiaceae*).^[1,2] GR has long been used as a traditional Korean medicine for the treatment of diarrhea, seminal emissions, excessive sweating, bleeding, and chronic cough although there is little scientific evidence supporting these pharmacological effects.^[3-5] Recent studies have revealed the therapeutic effects of GR against various human diseases and their mechanisms. For example, several compounds and extracts purified from GR exhibited good antibacterial activity against many pathogenic bacteria strains including *Salmonella* spp., *Escherichia coli*, *Eimeria tenella*, *Brucella abortus*, *Staphylococcus aureus*,

and *Clostridium perfringens*.^[6-11] Methyl gallate and ethyl gallate isolated from GR were also found to exert significant anti-inflammatory

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Go J, Kim JE, Koh EK, Song SH, Kang HG, Lee YH, et al. Hepatoprotective effect of gallotannin-enriched extract isolated from gall on hydrogen peroxide-induced cytotoxicity in HepG2 cells. *Phcog Mag* 2017;13:S294-300.

activity in lipopolysaccharide-stimulated RAW264.7 macrophages through the induction of heme oxygenase-1 and the suppression of inducible nitric oxide synthase/cyclooxygenase-2 (COX-2).^[12,13] Moreover, galloylglucose (GG6-10) isolated from GR inhibited the invasion of metastatic HT-1080 cells into a reconstituted basement membrane through inhibition of gelatinolysis mediated by matrix metalloproteinase (MMP-2) and MMP-9, whereas the ellagic acid extracted from GR showed anticancer activity against nasopharyngeal carcinoma cells through downregulated expression of COX-2 and stathmin.^[14,15] Furthermore, oral administration of GR 85% methanol extract reduced brain infarct volume by 37.5% and lipid peroxidation in middle cerebral artery occlusion, while also improving sensory motor function in a transient focal cerebral ischemia rat model.^[16] Moreover, tacrine, nitrofurantoin, and *tert*-butyl hydroperoxide-induced hepatotoxicity in HepG2 cells were greatly alleviated by two hepatoprotective constituents of GR, an equilibrium mixture of 3-galloyl-gallic acid (3GGA), 4-galloyl-gallic acid isomers (4GGAI), and 1,2,3,4,6-penta-*O*-galloyl- β -*D*-glucose (PGG).^[17,18] However, more studies are needed to determine the novel functions and mechanisms of GR extract in hydrogen peroxide (H₂O₂)-induced hepatotoxicity.

In this study, we investigated the hepatoprotective effects of gallotannin-enriched GR (GEGR) on H₂O₂-induced hepatotoxicity in HepG2 cells. The results provide novel data indicating that GEGR may associate with protective hepatocytes during apoptosis.

MATERIALS AND METHODS

Preparation of GEGR

Samples of GR were collected from plantations in the Hongcheon area of Korea in October 2013 by Professor Young-Hee Lee, then dried in a hot-air drying machine (JSR, Seoul, Korea) for 24 h at 60°C. Voucher specimens of GR (WPC-14-001) were deposited in the functional materials bank of the PNU-Wellbeing RIS Center at Pusan National University. GEGR was prepared using the modified extract quoting method as previously described.^[19-22] First, dry samples of GR were reduced to powder using an electric blender. Water extract was then obtained by placing the powder in a fixed liquor ratio (solid GR powder/water ratio, 1:10) and heating at 90°C for 9 h using circulating extraction equipment (IKA Labortechnik, Staufen, Germany). The extracts were subsequently filtered through a 0.4 μ m filter, after which they were concentrated by vacuum evaporation and lyophilization using circulating extraction equipment (IKA Labortechnik, Staufen, Germany). Finally, the powder of GEGR was dissolved in distilled water (dH₂O) to 1 mg/ml, then further diluted with phosphate-buffered saline (PBS) to the required concentration.

Analysis of main components in GEGR

During analysis of the main components of GEGR, gallic acid monohydrate, methyl gallate, and gallotannin were purchased from Sigma-Aldrich Co., St. Louis, MO, USA. The wavelengths of the maximum absorption of pure gallic acid, pure methyl gallate, commercial gallotannin, and nutgall extract were 212/257, 214/268, 213/278, and 212/275 nm, respectively. The ultraviolet-visible (UV-Vis) spectra of pure gallic acid, pure methyl gallate, pure gallotannin, and the nutgall extract showed two bands at 212–214 and 257–278 nm, which were both assigned to the $\pi \rightarrow \pi^*$ transitions of the given aromatic units and C=O groups in the UV-Vis region.^[23] Finally, the UV-Vis spectra were analyzed using a curve-resolving technique based on linear least squares analysis to fit the combined Lorentzian and Gaussian curves.

The high-performance liquid chromatography (HPLC) analysis was conducted as previously described.^[24] The HPLC system used for these experiments consisted of a Summit Dual-Gradient HPLC

System (Dionex, USA) with a photodiode array UV-Vis detector at the Korea Bio-IT Foundry Busan Center. Separation was carried out on an YMC-Triart C18 column (S-5 mm/12 mm, 150 mm \times 4.6 mm I.D.) maintained at 40°C. The mobile phase consisted of solvent A (0.4% formic acid in water) and solvent B (acetonitrile). The gradient condition of the mobile phase was as follows: 0–5 min, 10% B; 5–6 min, 10–15% B; 6–40 min, 15% B; 40–41 min, 15–30% B; 41–50 min, 30% B; 50–55 min, 30–10% B; and 55–60 min, 10% B. The injection volume was 5 ml in full loop injection. The flow rate was 0.8 mL/min, and detection was performed at 280 nm.

Free radical scavenging activity

The scavenging activity of 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical was measured as previously described.^[24] Briefly, the powder of GEGR was dissolved in 50% ethyl alcohol (100 μ l) in 13 different concentrations (0.122–500 μ g/ml) and mixed with 100 μ l of 0.1 mM DPPH (Cat. No. D9132, Sigma-Aldrich Co.) in 95% ethanol solution or 100 μ l of 95% ethanol solution, then incubated for 30 min at room temperature. Next, the absorbance of the reaction mixture was measured at 517 nm using a VersaMax plate reader (Molecular Devices, Sunnyvale, CA, USA). The DPPH radical scavenging activity of the GEGR was expressed as the percent decrease in absorbance relative to the control. The inhibitory concentration 50% (IC₅₀) value is defined as the concentration of substrate that causes a 50% loss in DPPH activity.

Cell culture

The HepG2 cell line was purchased from the Korean Cell Line Bank (Seoul, Korea) and grown in a humidified incubator at 37°C under 5% CO₂ and 95% air in Eagle's minimal essential medium with Earle's balanced salt solution (MEM/EBSS, Cat. No. SH30024.01, Thermo Scientific, Waltham, MA, USA) containing 10% fetal bovine serum, 2 mM glutamine, 100 U/ml of penicillin, and 100 μ g/ml streptomycin.

Cell viability assay

HepG2 cells treated as described above were further treated with 5 mM H₂O₂, then incubated for another 12 h, after which the cell viability was determined using the tetrazolium compound 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) (Cat. No. M2128, Sigma-Aldrich Co.). To determine cell viability, HepG2 cells were seeded at a density of 5 \times 10⁵ cells/2 ml of MEM media and grown for 24 h in a 37°C incubator. When the cells attained 70%–80% confluence, they were either untreated (control group), treated with vehicle (dH₂O), or pretreated with 6.25, 12.5, 25, or 50 μ g/ml of GEGR dissolved in distilled H₂O added to each well. Following incubation for 24 h, the supernatants were discarded, after which 2 ml of fresh MEM media and 500 μ l of MTT solution (2 mg/ml in PBS) were added to each well. The cells were then incubated at 37°C for 4 h, after which the formazan precipitate was dissolved in dimethyl sulfoxide (DMSO, Cat. No. D1370.0100, Duchefa Biochemie, Haarlem, the Netherlands), and the absorbance was read at 570 nm directly in the wells using a Vmax plate reader (Molecular Devices).

Fluorescence-activated cell sorter analysis

Apoptotic cells were detected using a Muse™ Annexin V and Dead Cell Kit (Cat. No. MCH100105, Millipore Co., Billerica, MA, USA) according to the manufacturer's protocols. Briefly, cells of subset groups were suspended in MEM media (1 \times 10⁵ cells/ml), after which 100 μ l of the cell suspension (1 \times 10⁴ cells/ml) was incubated with Muse™ Annexin V and Dead Cell Reagent (Cat. No. 12-0563, Millipore Co.) for 20 min at room temperature. After the final incubation, the reaction mixture

was analyzed using Muse™ Cell Analyzer (Cat. No. PB4455ENEU, Millipore Co.).

4',6-diamidino-2-phenylindole (DAPI) staining

The morphological changes in apoptotic cells were assessed by fluorescent microscopy following DAPI staining. Briefly, HepG2 cells were seeded at a density of 5×10^5 cells/2 ml in 6-well plates, then grown with four different concentrations of GEGR for 24 h in a 37°C incubator. After washing once with $\times 1$ PBS, cells were incubated with 5 mM H_2O_2 for another 12 h, then fixed in 4% formaldehyde (Cat. No. 69360-0380, Junsei Chemical Co. Ltd., Tokyo, Japan) for 1 h and permeated with 0.1% Triton X-100 (Cat. No. T1020, Biosesang Inc., Seongnam, Korea) for 5 min. Next, the DNA-specific fluorochrome DAPI (100 μ M, Cat. No. D1306, Invitrogen) was applied to each well, after which samples were incubated for 10 min in the dark at room temperature. Finally, the cells were washed three times with $\times 1$ PBS and examined using a fluorescent microscope (Olympus IX71, Tokyo, Japan) at $\times 400$ magnification.

Analysis of intracellular reactive oxygen species level

Intracellular reactive oxygen species (ROS) levels were measured by staining with 2',7'-dichlorofluorescein diacetate (DCFH-DA) (Cat. No. D6883, Sigma-Aldrich Co.), which is a cell permeable and nonfluorescent agent that can be deacetylated by intracellular esterases to nonfluorescent DCFH. In the presence of ROS, DCFH was converted to highly fluorescent DCF intracellularly. Briefly, HepG2 cells were seeded at a density of 5×10^5 cells/2 ml in 6-well plates, then grown with four different concentrations of GEGR for 24 h in a 37°C incubator. After washing once with $\times 1$ PBS, the cells were incubated with 5 μ M H_2O_2 for another 12 h. Next, cells were incubated with 25 μ M DCFH-DA for 30 min at 37°C. The cells were then washed twice with PBS, after which the green fluorescence was observed using a fluorescent microscope (Eclipse TX100, Nikon, Tokyo, Japan).

Western blot

Proteins prepared from HepG2 cells were separated by 4%–20% sodium dodecyl sulfate-polyacrylamide gel electrophoresis for 2 h, after which they were transferred to nitrocellulose membranes for 2 h at 40 V. Each membrane was then incubated separately at 4°C with the following primary antibodies overnight: anti-Bax (Cat. No. ab7977, Abcam, Cambridge, UK), anti-Bcl-2 (Cat. No. ab7973, Abcam), anti-Caspase-3 (Cat. No. 9662, Cell Signaling, Danvers, MA, USA), and anti-actin antibody (Cat. No. A5316, Sigma-Aldrich). Next, the membranes were washed with washing buffer (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4 , and 0.05% Tween 20) and incubated with 1:1000 diluted horseradish peroxidase-conjugated goat anti-rabbit IgG (Cat. No. G21234, Invitrogen) at room temperature for 1 h. Finally, membrane blots were developed using Amersham ECL Select Western Blotting detection reagent (Cat. No. RPN2235, GE Healthcare, Little Chalfont, UK). The chemiluminescence signals that originated from specific bands were detected using FluorChemⁱFC2 (Alpha Innotech Co., San Leandro, CA, USA).

Statistical analysis

One-way ANOVA was used to identify significant differences between No- and H_2O_2 -treated groups (SPSS for Windows, Release 10.10, Standard Version, Chicago, IL, USA). Differences in the responses of the H_2O_2 + vehicle and H_2O_2 + GEGR treated groups were evaluated using a *post hoc* test (SPSS for Windows, Release 10.10, Standard Version). All values are reported as the mean \pm standard deviation and $P < 0.05$ was considered statistically significant.

RESULTS

Distribution of functional components in GEGR

As shown in Figure 1a, the fitting curves of pure gallic acid were detected at 257 nm, whereas pure methyl gallate and pure gallotannin were observed at 268 and 278 nm, respectively. GEGR contained high concentrations of three bioactive components related to its hepatoprotective effects, gallotannin (69.2%), gallic acid (26.6%), and methyl gallate (4.2%). HPLC curves of GEGR were found to have three characteristic peaks for gallic acid (3.58 min), methyl gallate (11.44 min), and gallotannin (48.26, 51.55, 52.46, and 53.23 min) [Figure 1b]. Furthermore, GEGR showed a very small peak near 11.4 min, indicating the existence of a small amount of methyl gallate, but a large amount of gallotannin and gallic acid. These results demonstrate that GEGR had a high concentration of gallotannin.




Figure 1: Analysis of the distribution of major components and free radical scavenging activity of GEGR. (a) Ultraviolet-visible spectra of GEGR. (b) High-performance liquid chromatography chromatograms of GEGR. (c) 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity of GEGR. The data shown represent the means \pm standard deviation of three replicates

Free radical scavenging activity of GEGR

To measure the antioxidant activity of GEGR, the free radical scavenging activity of DPPH was analyzed following treatment with various doses of GEGR. The inhibitory activity against DPPH radical was gradually increased by the addition of 0.12–500 µg/ml of GEGR. Based on these data, the IC₅₀ value of GEGR was determined to be 0.212 µg/ml [Figure 1c]. Taken together, these results indicate that GEGR has very strong DPPH radical scavenging activity and therefore has the potential for use as an antioxidant.

Protective effects of GEGR treatment against hydrogen peroxide-induced apoptosis

H₂O₂ treatment induces apoptosis through the production of ROS and oxidative stress in various cells.^[25,26] Therefore, we examined whether GEGR pretreatment can prevent hepatic cell death induced by H₂O₂ exposure. To accomplish this, cell viability was measured in HepG2 cells pretreated with different concentrations of GEGR by MTT assay, fluorescence-activated cell sorter staining, and DAPI staining analysis. Toxicity analysis revealed no significant alterations in cell viability in HepG2 cells pretreated with 6.25 µg/ml to 50 µg/ml of GEGR alone [Figure 2a], while the lethal dose 50% of GEGR was determined to be 178 µg/ml (data not shown). Overall, these findings indicated that GEGR exerted no toxicity at <50 µg/ml.

H₂O₂ + vehicle-treated cells showed low (53%) cell viability relative to the control group. However, their level was significantly higher in HepG2 cells treated with H₂O₂ and four concentrations of GEGR although the maximum level remained consistent in response to pretreatment with 12.5–50 µg/ml of GEGR [Figure 2b]. Based on the above data, the effective dose 50% was determined to be 6.85 µg/ml of GEGR. Similar results were observed upon DAPI staining. Specifically, more irregular nuclei were detected in H₂O₂ + vehicle-treated cells relative to the control group, whereas a lower number of irregular nuclei were observed in H₂O₂ + GEGR-treated cells than H₂O₂ + vehicle-treated cells [Figure 3]. The total number of apoptotic cells showed the reverse pattern. A significant increase in the number (347%–409%) of apoptotic cells was detected in H₂O₂ + vehicle-treated cells relative to the control group. Following pretreatment with GEGR, the number of apoptotic cells gradually decreased although a constant number was detected in HepG2 cells pretreated with 12.5, 25, or 50 µg/ml [Figure 4]. In conclusion, these findings indicate that GEGR pretreatment may prevent the cytotoxicity induced by H₂O₂ treatment in HepG2 cells.

Effects of GEGR on apoptosis-related protein expression

To determine if the increase in apoptotic cells was accompanied by altered expression of apoptosis-related proteins, the expression levels




Figure 2: Effects of GEGR pretreatments on hydrogen peroxide-induced cytotoxicity. (a) Cytotoxicity of GEGR. (b) Protective effects of GEGR on hydrogen peroxide-induced hepatotoxicity. The data shown represent the means ± standard deviation of three replicates. a: *P* < 0.05 relative to the no-treated group. b: *P* < 0.05 relative to the vehicle treated group




Figure 3: Effects of GEGR pretreatments on the increase of apoptotic cells induced by hydrogen peroxide exposure. Cells were stained with 4',6-diamidino-2-phenylindole to detect changes in the nuclei. Cellular morphology was observed under a fluorescent microscope (original magnification ×400)




Figure 4: Effects of GEGR on the number of Annexin V stained cells. Fluorescence-Activated cell sorter data represented four independent phases (live, dead, early apoptosis, and late apoptosis). The data shown represent the means ± standard deviation of three experiments. a: $P < 0.05$ relative to the control group. b: $P < 0.05$ relative to the hydrogen peroxide + vehicle-treated group




Figure 5: Effects of GEGR on the expression of Bax, Bcl-2, and caspase-3 in hydrogen peroxide-exposed HepG2 cells after GEGR treatment. The data represent the means ± standard deviation of three replicates. a: $P < 0.05$ compared to the control group. b: $P < 0.05$ compared to the hydrogen peroxide + vehicle-treated group

of Bax, Bcl-2, and caspase-3 were measured by Western blot analysis in HepG2 cells pretreated with GEGR before H₂O₂ exposure. The Bax/Bcl-2 ratio increased by 19.8% in the H₂O₂ + vehicle-treated group relative to the control group, whereas it decreased significantly in the GEGR pretreated group although cells pretreated with 6.25 µg/ml of GEGR showed a slight increase in this ratio. Furthermore, a dramatic change in the active form of caspase-3 was observed. Specifically, the intensity of the active form of caspase-3 in the H₂O₂ + vehicle-treated group was about two times higher than that in the control group, while their level was dramatically decreased in the GEGR pretreated group relative to the H₂O₂ + vehicle-treated group. HepG2 cells pretreated with 50 µg/ml of GEGR before exposure to 5 mM H₂O₂ showed a similar level of active caspase-3 as observed in the control group [Figure 5]. In conclusion, these results indicate that GEGR pretreatment may prevent H₂O₂-induced apoptosis through the regulation of Bax/Bcl-2 expression and the suppression of caspase-3 activation.

Suppression of reactive oxygen species by GEGR

To determine if the prevention of apoptosis was correlated with the suppression of ROS production, alterations in ROS level were measured by DCFH-DA assay. The number of DCFH-DA-stained cells was higher in the H₂O₂ + vehicle-treated group than the control group. Following GEGR pretreatment, this number decreased

greatly in all groups although concentration-dependent changes were only detected in the 6.25 and 12.5 µg/ml pretreated groups. A constant level was maintained in groups pretreated with 12.5, 25, and 50 µg/ml GEGR [Figure 6]. In conclusion, these results suggest that GEGR pretreatment inhibits ROS production induced by H₂O₂ treatment.

DISCUSSION

The liver is a greater target of human toxicants than other organs. Various toxicants induce hepatic damage, necrosis,^[27] and apoptosis,^[28] and their administration for long periods of time leads to fibrosis, cirrhosis, and hepatic carcinoma.^[29] Therefore, novel therapeutic drugs with hepatoprotective activity are of great interest. In this study, we investigated the hepatoprotective effects of GEGR against H₂O₂-induced cytotoxicity. The results demonstrated that pretreatment with GEGR may prevent apoptosis of HepG2 cells through suppression of caspase-3 activation and ROS production.

Previous studies have shown that several compounds in methanol and ethanol extracts of GR protect against tacrine- and nitrofurantoin-induced cytotoxicity in mammalian cells. Four main compounds, gallic acid methyl ester, gallic acid, an equilibrium mixture of 3GGA + 4GGAI, and PGG were identified in methanol extract of GR.^[17] The ethanol extract of GR contained syringic acid (18.5 mg), as well as the phenolics methyl gallate (20.7 mg) and gallic acid (19.5 mg). In addition, a study conducted by Lee *et al.*^[30] revealed that aqueous extract of GR contained gallocatechin (69.0%), gallic acid (25.7%), and methyl gallate (5.3%) and showed deodorizing function and antibacterial activity. The concentrations of these components were very similar to those observed in the present study although they did differ slightly, possibly due to the origin of the GR used.

Tannic acid and its derivatives have promising hepatoprotective activity.^[31,32] Specifically, phlorotannins from *Eisenia bicyclis* that showed high contents of total phenolics and strong antioxidant activity




Figure 6: Effects of GEGR on intracellular reactive oxygen species production. Cells in each square of a $\times 100$ magnification image (left column) were further examined under $\times 400$ magnification (right column). Arrows indicate cells stained with 2',7'-dichlorofluorescein diacetate

significantly protected HepG2 cells against *tert*-butyl hydroperoxide-induced cell death.^[33] Treatment with Punarnavashtak kwath (an Ayurvedic formulation), which contains alkaloids, tannins, flavonoids, and saponins, induced a significant increase in the viability of HepG2 cells relative to a carbon tetrachloride (CCl_4)-treated group.^[34] Moreover, HepG2 cells were effectively protected against *tert*-butyl hydroperoxide-induced cytotoxicity by treatment with ethanol extract of *Ecklonia stolonifera* containing several phlorotannins.^[35] Phenolic constituents *namely* PGG and mixture of 3GGA + 4GGAI also showed significant hepatoprotective effects against tacrine- and nitrofurantoin-induced cytotoxicity in HepG2 cells.^[17] In the present study, the gallotannin-enriched complex of GEGR exerted hepatoprotective effects against H_2O_2 -treated HepG2 cells. Therefore, the present results provide novel evidence that the hepatotoxicity induced by H_2O_2 treatment may be prevented by GEGR.

Caspase-3 is considered a key factor responsible for cleavage and inactivation of PPAR, as well as mitochondrial dysfunction (caspase-9 activation) and death receptor pathway activation (caspase-8 activation) in apoptosis of hepatocytes.^[36,37] Treatment with PGG-inhibited production of the activated form of caspase-3 in primary rat hepatocytes treated with 100 μM glycochendoxycholic acid.^[13] Moreover, the dose-dependent activation of caspase-3 in HepG2 cells by tacrine was inhibited by 2-phloroecol, one of four types of phlorotannins isolated from the ethyl acetate of *E. stolonifera*.^[38] Similar results were observed in the present study even though the concentration and composition of treatment compounds varied.

Excessive production of ROS can induce oxidative stress, cell dysfunction, and ultimately apoptosis and necrosis.^[39] Tannic acid and several related products have been reported to inhibit ROS production. Specifically, ROS production induced by *tert*-butyl hydroperoxide and tacrine treatment significantly inhibited the procyanidin fractions from defatted grape seeds, as well as 2-phloroecol from *E. stolonifera* in HepG2 cells.^[38,40] In the present study, the ROS level was decreased in the H_2O_2 + GEGR-treated group relative to the H_2O_2 + vehicle-treated group. These findings are similar to those of previous studies although the decrease ratio varied.

CONCLUSION

The results of the present study indicated that GEGR was closely associated with prevention of H_2O_2 -induced HepG2 cell death through regulation of apoptosis-related protein expression. These findings also indicate that GEGR plays a crucial role in the prevention of apoptosis through inhibition of ROS production. Furthermore, GEGR has the potential for use as a food supplement for the prevention of hepatotoxicity.

Financial support and sponsorship

This study was supported by grants to Professor Dae Youn Hwang from the Korea Institute of Planning & Evaluation for Technology In Food, Agriculture, Forestry & Fisheries (114034-03-1-HD030 and 116027-3-2- HD030).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Lee SM, Lee DW, Park JD, Kim JI. Study on formation and development of gall in *Rhus javanica*. Korean J Appl Entomol 1997;36:83-7.
- Ren Z, Zhu B, Wang D, Ma E, Su D, Zhong Y. Comparative population structure of Chinese sumac aphid *Schlechtendalia chinensis* and its primary host-plant *Rhus chinensis*. Genetica 2008;132:103-12.
- Shim YJ, Doo HK, Ahn SY, Kim YS, Seong JK, Park IS, *et al.* Inhibitory effect of aqueous extract from the gall of *Rhus chinensis* on alpha-glucosidase activity and postprandial blood glucose. J Ethnopharmacol 2003;85:283-7.
- Song GY, Park BJ, Kim SH. Antithrombotic effect of Galla Rhois. Korean J Pharmacogn 2002;33:120-3.
- Kim SH, Park HH, Lee S, Jun CD, Choi BJ, Kim SY, *et al.* The anti-anaphylactic effect of the gall of *Rhus javanica* is mediated through inhibition of histamine release and inflammatory cytokine secretion. Int Immunopharmacol 2005;5:1820-9.
- Ahn YJ, Lee CO, Kweon JH, Ahn JW, Park JH. Growth-inhibitory effects of Galla Rhois-derived tannins on intestinal bacteria. J Appl Microbiol 1998;84:439-43.
- Choi JG, Kang OH, Lee YS, Oh YC, Chae HS, Jang HJ, *et al.* *In vitro* activity of methyl gallate isolated from Galla Rhois alone and in combination with ciprofloxacin against clinical isolates of *Salmonella*. J Microbiol Biotechnol 2008;18:1848-52.
- Choi JG, Kang OH, Lee YS, Oh YC, Chae HS, Jang HJ, *et al.* Antibacterial activity of methyl gallate isolated from Galla Rhois or carvacrol combined with nalidixic acid against nalidixic acid resistant bacteria. Molecules 2009;14:1773-80.

9. Choi JG, Mun SH, Chahar HS, Bharaj P, Kang OH, Kim SG, *et al.* Methyl gallate from Galla Rhois successfully controls clinical isolates of *Salmonella* infection in both *in vitro* and *in vivo* systems. *PLoS One* 2014;9:e102697. Available from: <http://www.journals.plos.org/plosone/article?id=10.1371/journal.pone.0102697>. [Last cited on 2014 Jul 21].
10. Lee JJ, Bae JH, Kim DH, Lim JJ, Kim DG, Lee HJ, *et al.* Intracellular replication inhibitory effects of Galla Rhois ethanol extract for *Brucella abortus* infection. *J Ethnopharmacol* 2011;138:602-9.
11. Lee JJ, Kim DH, Lim JJ, Kim DG, Min W, Kim GS, *et al.* Anticoccidial effect of supplemental dietary Galla Rhois against infection with *Eimeria tenella* in chickens. *Avian Pathol* 2012;41:403-7.
12. Chae HS, Kang OH, Choi JG, Oh YC, Lee YS, Brice OO, *et al.* Methyl gallate inhibits the production of interleukin-6 and nitric oxide via down-regulation of extracellular-signal regulated protein kinase in RAW 264.7 cells. *Am J Chin Med* 2010;38:973-83.
13. Park PH, Hur J, Kim YC, An RB, Sohn DH. Involvement of heme oxygenase-1 induction in inhibitory effect of ethyl gallate isolated from Galla Rhois on nitric oxide production in RAW 264.7 macrophages. *Arch Pharm Res* 2011;34:1545-52.
14. Ata N, Oku T, Hattori M, Fujii H, Nakajima M, Saiki I. Inhibition by galloylglucose (GG6-10) of tumor invasion through extracellular matrix and gelatinase-mediated degradation of type IV collagens by metastatic tumor cells. *Oncol Res* 1996;8:503-11.
15. Xiang Q, Fan C, Xiao S, Pan H, Wang J, Zhao N, *et al.* Effect of gallnut extract on nasopharyngeal carcinoma 5-8F cells and its mechanism. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2012;37:871-5.
16. Lee K, Kim J, Lee BJ, Park JW, Leem KH, Bu Y. Protective effects of Galla Rhois, the excrescence produced by the sumac aphid, *Schlechtendalia chinensis*, on transient focal cerebral ischemia in the rat. *J Insect Sci* 2012;12:10.
17. An RB, Oh H, Kim YC. Phenolic constituents of Galla Rhois with hepatoprotective effects on tacrine- and nitrofurantoin-induced cytotoxicity in HepG2 cells. *Biol Pharm Bull* 2005;28:2155-7.
18. Park EJ, Zhao YZ, An RB, Kim YC, Sohn DH. 1,2,3,4,6-penta-O-galloyl-beta-D-glucose from Galla Rhois protects primary rat hepatocytes from necrosis and apoptosis. *Planta Med* 2008;74:1380-3.
19. Lee YH. Dyeing, fastness, and deodorizing properties of cotton, silk, and wool fabrics dyed with coffee sludge (*Coffea arabica* L.) extract. *J Appl Polym Sci* 2007;103:251-7.
20. Lee YH, Hwang EK, Jung YJ, Do SK, Kim HD. Dyeing and deodorizing properties of cotton, silk, wool fabrics dyed with amurcorktree, *Dryopteris crassirhizoma*, *Chrysanthemum boreale*, *Artemisia* extracts. *J Appl Polym Sci* 2010;115:2246-53.
21. Lee YH, Hwang EK, Baek YM, Lee MS, Lee DJ, Jung YJ, *et al.* Deodorizing and antibacterial performance of cotton, silk and wool fabrics dyed with *Punicagranatum* L. extracts. *Fibers Polym* 2013;14:1445-53.
22. Hwang EK, Lee YH, Kim HD. Dyeing and deodorizing properties of cotton, silk, and wool fabrics dyed with various natural colorants. *J Korean Soc Dyers Finishers* 2007;19:12-20.
23. Albu MG, Ghica MV, Giurginca M, Trandafir V, Popa L, Cotrut MC. Spectral characteristics and antioxidant properties of tannic acid immobilized on collagen drug-delivery systems. *Rev Chim (Bucur)* 2009;60:666-72.
24. Oh H, Ko EK, Kim DH, Jang KK, Park SE, Lee HS, *et al.* Secoiridoid glucosides with free radical scavenging activity from the leaves of *Syringa dilatata*. *Phytother Res* 2003;17:417-9.
25. Emanuele S, Calvaruso G, Lauricella M, Giuliano M, Bellavia G, D'Anneo A, *et al.* Apoptosis induced in hepatoblastoma HepG2 cells by the proteasome inhibitor MG132 is associated with hydrogen peroxide production, expression of Bcl-XS and activation of caspase-3. *Int J Oncol* 2002;21:857-65.
26. Lee JH, Lee JS, Kim YR, Jung WC, Lee KE, Lee SY, *et al.* Hispidin isolated from *Phellinus linteus* protects against hydrogen peroxide-induced oxidative stress in pancreatic MIN6N β -cells. *J Med Food* 2011;14:1431-8.
27. Siviková K, Piesová E, Dianovský J. The protection of Vitamin E and selenium against carbon tetrachloride-induced genotoxicity in ovine peripheral blood lymphocytes. *Mutat Res* 2001;494:135-42.
28. Shi J, Aisaki K, Ikawa Y, Wake K. Evidence of hepatocyte apoptosis in rat liver after the administration of carbon tetrachloride. *Am J Pathol* 1998;153:515-25.
29. Wernke MJ, Schell JD. Solvents and malignancy. *Clin Occup Environ Med* 2004;4:513-27, vii.
30. Lee YH, Hwang EK, Baek YM, Kim HD. Deodorizing function and antibacterial activity of fabrics dyed with gallnut (*Galla chinensis*) extract. *Text Res J* 2015;85:1045-54. Available from: <http://www.trj.sagepub.com/content/early/2014/11/27/0040517514559580>. [Last cited on 2015 Apr 07].
31. Tikoo K, Tanta A, Ali IY, Gupta J, Gaikwad AB. Tannic acid prevents azidothymidine (AZT) induced hepatotoxicity and genotoxicity along with change in expression of PARG and histone H3 acetylation. *Toxicol Lett* 2008;177:90-6.
32. Shimoda H, Tanaka J, Kikuchi M, Fukuda T, Ito H, Hatano T, *et al.* Walnut polyphenols prevent liver damage induced by carbon tetrachloride and d-galactosamine: hepatoprotective hydrolyzable tannins in the kernel pellicles of walnut. *J Agric Food Chem* 2008;56:4444-9.
33. Kim SM, Kang K, Jeon JS, Jho EH, Kim CY, Nho CW, *et al.* Isolation of phlorotannins from *Eisenia bicyclis* and their hepatoprotective effect against oxidative stress induced by tert-butyl hydroperoxide. *Appl Biochem Biotechnol* 2011;165:1296-307.
34. Shah VN, Shah MB, Bhatt PA. Hepatoprotective activity of punarnavashtak kwath, an Ayurvedic formulation, against CCl4-induced hepatotoxicity in rats and on the HepG2 cell line. *Pharm Biol* 2011;49:408-15.
35. Kim YC, An RB, Yoon NY, Nam TJ, Choi JS. Hepatoprotective constituents of the edible brown alga *Ecklonia stolonifera* on tacrine-induced cytotoxicity in HepG2 cells. *Arch Pharm Res* 2005;28:1376-80.
36. Allen RT, Hunter WJ 3rd, Agrawal DK. Morphological and biochemical characterization and analysis of apoptosis. *J Pharmacol Toxicol Methods* 1997;37:215-28.
37. Yoon JH, Gores GJ. Death receptor-mediated apoptosis and the liver. *J Hepatol* 2002;37:400-10.
38. Lee MS, Shin T, Utsuki T, Choi JS, Byun DS, Kim HR. Isolation and identification of phlorotannins from *Ecklonia stolonifera* with antioxidant and hepatoprotective properties in tacrine-treated HepG2 cells. *J Agric Food Chem* 2012;60:5340-9.
39. Halliwell B, Gutteridge JM, Cross CE. Free radicals, antioxidants, and human disease: Where are we now? *J Lab Clin Med* 1992;119:598-620.
40. Kim Y, Choi Y, Ham H, Jeong HS, Lee J. Protective effects of oligomeric and polymeric procyanidin fractions from defatted grape seeds on tert-butyl hydroperoxide-induced oxidative damage in HepG2 cells. *Food Chem* 2013;137:136-41.