



Minor migrants' age estimation: Comparison of two dental methods

Francesco Lupariello*, Alessandro Gabriele, Federica Mirri, Giuliana Mattioda, Emilio Nuzzolese, Giancarlo Di Vella

Dipartimento di Scienze Della Sanità Pubblica e Pediatriche, Sezione di Medicina Legale, "Università Degli Studi di Torino", Corso Galileo Galilei 22, 00126, Torino, Italy



ARTICLE INFO

Article history:

Received 6 October 2020
Received in revised form
7 January 2021
Accepted 8 January 2021
Available online 16 January 2021

Keywords:

Forensic sciences
Forensic anthropology
Child rights
Age estimation
Legal age

ABSTRACT

In case of minor migrants, absence of valid identification documents that clearly define age is a critical issue, because without these data the child cannot enjoy the rights provided by the Convention on the Rights of the Child. Differentiation between minors and adults is fundamental when age is disputed in human identification, asylum seeking, criminal liability, and child abuse fields. Few indications are available about qualitative/statistical agreement of different age estimation methods. Ages of 301 individuals were estimated through two dental methods in order to: determine quantitative and statistical agreements in legal age definition; identify practical recommendations. The study pointed out discrepancy between the two methods in 7/301 cases. From a statistical point of view, this finding corresponded to an almost perfect agreement. Thus, authors suggested that the two methods can be alternately used for legal age assessment, but operators should use both methods when the estimated age is 18.5 years.

© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The Convention on the Rights of the Child (1989) states that "States Parties undertake to respect the right of the child to preserve his or her identity, including nationality, name and family relations as recognized by law without unlawful interference. Where a child is illegally deprived of some or all of the elements of his or her identity, States Parties shall provide appropriate assistance and protection, with a view to re-establishing rapidly his or her identity" [1]. In case of minor migrants, absence of valid identification documents that clearly define age is a critical issue, because without these data the child cannot enjoy the rights provided by the Convention. In this scenario, child best interest matches with careful assessment of his/her age.

Several age estimation methods have been proposed [2–5]. They usually rely on the correlation between bone/dental maturation and chronological age [6–13]. In 2010 AlQahtani and colleagues described an atlas (The London Atlas of Human Tooth Development and Eruption) of tooth development in order to estimate age between 28 weeks intrauterine and 23 years [14].

Besides, this method was applied to several population samples, in order to evaluate its accuracy in people of different geographical areas [15–17]. Cameriere and colleagues described an innovative method (third molar maturity index – I_{3M}) that allows to distinguish adults ($I_{3M} < 0.08$) from minors ($I_{3M} \geq 0.08$) [18]. I_{3M} is calculated "as the sum of the widths of the inner margins of the two open apices than divided by tooth length" [19].

From a practical point of view, in the most part of European and non-European countries legal age is reached when an individual turns 18 years old [20,21]. In case of subjects who do not have proper documents, differentiation between minors (<18 years old) and adults (≥ 18 years old) is fundamental when age is disputed in human identification, asylum seeking, criminal liability, and child abuse fields. In the literature, there are several studies that evaluate accuracy and intra-observer/inter-observer agreement of each of the two abovementioned methods [15,17]. Nevertheless, comparative analyzes of results obtained through these methods are not available. In particular, there are no indications that allow: 1) to determine quantitative and statistical agreements between the AlQahtani atlas and the Cameriere method in legal age definition; 2) to identify recommendations that operators should take into account when they use these methods in order to minimize errors.

In the light of the above, the authors propose a comparison between the results of the London Atlas and the third molar

* Corresponding author. corso Galileo Galilei 22, 10126, Turin, Italy.
E-mail addresses: francesco.lupariello@unito.it, francesco.lupariello@gmail.com (F. Lupariello).

maturity index method on an unknown age population, in order to define quantitative and statistical agreements, and to identify practical recommendations when it comes to determine legal age using the aforementioned methods.

2. Materials and methods

The study included 301 orthopantomography (OPG) images corresponding to 301 individuals without valid identify documents; actual chronological ages of the study sample were unknown. These individuals had been consecutively evaluated from November 2014 to April 2018 by the operators of the "A.O.U. Città della Salute e della Scienza di Torino" (Italy) hospital, in order to produce identification/age estimation assessments. Individuals' nationality, sex, and OPG images were obtained from electronic medical records (Trakcare® platform). Medical record review excluded recurrence of systemic diseases or developmental problems in the study population. A numerical code was assigned to each image in order to anonymize them. In addition, these numerical codes were reported into two different Excel® files. Left and right mandibular third molars were respectively defined as 38 and 48, according to classification by Federation Dentaire Internationale [22]. In the first session, a forensic odontologist (experienced in age estimation; with ten year experience in age assessment through the two aforementioned methods) analyzed 38s and 48s of the 301 OPG images throughout the MPDicom Viewer® software. For each molar, the abovementioned operator/observer assigned a dental age (in the form of binary categorical variables: < 18 years old if $I_{3M} \geq 0.08$; ≥ 18 years old if $I_{3M} < 0.08$), using the third molar index method described by Cameriere and colleagues [18]. Then, the observer collected ages in one of the aforementioned Excel® file in association with the correspondent numerical code.

In the second session (performed one week later the first one), the same forensic odontologist analyzed 38s and 48s of the same 301 OPG images throughout the MPDicom Viewer® software. In this session, the operator/observer had not access to the results of the first Excel® file. For each molar, the observer assigned an estimated age, using the atlas described by AlQahtani and colleagues [14]. Then, the operator collected ages in the second Excel® file in association with the correspondent numerical code.

After these sessions, a second operator merged the two Excel® files in a third one in which numerical codes, nationality, sex, and estimated ages were matched. Assessed ages obtained using the AlQahtani atlas were transformed in binary categorical variables (<18 or ≥ 18 years old), in order to allow quantitative and statistical comparisons.

At first, quantitative comparison of the estimated ages was carried out. Then, statistical agreement in legal age assessment of the methods was separately calculated for 38s and 48s through the IBM SPSS Statistics 20 software. In particular, the Cohen's kappa coefficient (K) was used to evaluate statistical agreement of binary qualitative categories (<18 or ≥ 18 years old) [23]. Even if K is commonly performed to test observations of a single dataset between two observers, according to the scientific literature its use for the same rater evaluating the same data at two time points is considered as acceptable [24]. However, this use has limitations because it does not take into account the magnitude of differences (especially for ordinal data) [24].

3. Results

All results are summarized in Table 1. Among the 301 individuals, 44 were female and 257 were male (Table 1). They mostly came from the African continent; in particular from Nigeria (52/

301), Guinea (39/301), Senegal (32/301), Mali (22/301), Gambia (20/301), Ivory Coast (18/301), and Morocco (16/301). In both sessions: 38 and 48 were simultaneously absent in 10/301 individuals; 38 was absent in 16/301 subjects; 48 was absent in 14/301 cases. Quantitative analysis yielded the following results:

- for 38s, the Cameriere method defined 216 ages as ≥ 18 years and 59 as < 18 years;
- for 38s, the AlQahtani atlas defined 223 ages as ≥ 18 years and 52 as < 18 years;
- for 48s, the Cameriere method defined 212 ages as ≥ 18 years and 65 as < 18 years;
- for 48s, the AlQahtani atlas defined 219 ages as ≥ 18 years and 58 as < 18 years.

In 7 cases, there was discrepancy in legal age assessment of both molars, because the Cameriere method identified ages as < 18 years, while the AlQahtani atlas as correspondent to 18.5 years (≥ 18 years old).

In one case, the operator/observer reported different maturation stages for 38 and 48, defining (through both methods) 38 as correspondent to ≥ 18 years and 48 as correspondent to < 18 years in both sessions.

Statistical analysis of the agreement (K) of the aforementioned methods yielded the following results:

- $K = 0.921$ for left mandibular third molar (38);
- $K = 0.927$ for right mandibular third molar (48).

4. Discussion

The results of the present manuscript pointed out that quantitative agreement in the identification of legal age was correspondent to 97.59%. Indeed, lower third molars of the same 7 individuals were identified as correspondent to ≥ 18 years through the AlQahtani method and as < 18 years through the Cameriere one. This study does not allow to define the reasons of the discrepancy. However, it is important to highlight that in all 7 cases the operator/observer assessed through the London Atlas both 38 and 48 as correspondent to 18.5 years. Therefore, it is possible to suggest that the two methods are more likely to differ when the AlQahtani atlas defines a chronological age that is just above 18 years old. However, it is important to note that in the other 9 cases in which the same atlas identified a chronological age of 18.5 years, there was concordance with the third molar index method. These results demonstrate that the abovementioned discrepancy does not seem to be a constant finding.

From a statistical point of view, identification of binary categorical variables can differ or agree as a result of chance. For this reason, Cohen Kappa coefficient (K) evaluated agreement between the two methods by eliminating the so-called per chance agreement/disagreement [23]. The results of Kappa statistics yielded a value of >0.8 . Therefore, this result suggested that the London Atlas and the Cameriere method are characterized by an almost perfect agreement ($K > 0.8$) [22,24,25]. Indeed, according to Ranganathan and colleagues (2017) and Vierra and Garrett (2005) a value of K from 0.81 to 0.99 is defined as almost perfect agreement [24,25]; thus, the two methods are statistically comparable.

From a practical point of view, the abovementioned considerations are meaningful. Indeed, the almost perfect agreement allows to state that the AlQahtani method and the Cameriere third molar maturity index can be alternately used for legal age assessment [24,25]. Nevertheless, quantitative data suggest a careful approach when the London Atlas identifies a chronological age of 18.5 years.

Table 1
Caption: Summary of the results (in years).

Case number	Sex	Nationality	Cameriere 38	AlQahtani 38	Cameriere 48	AlQahtani 48
1	Female	Mali	≥18	≥18 (21.5)	≥18	≥18 (21.5)
2	Male	Nigeria	Absent	Absent	<18	<18 (16.5)
3	Male	Bangladesh	≥18	≥18 (20.5)	≥18	≥18 (20.5)
4	Male	Pakistan	≥18	≥18 (22.5)	≥18	≥18 (22.5)
5	Male	Mali	≥18	≥18 (23.5)	≥18	≥18 (23.5)
6	Male	Mali	≥18	≥18 (22.5)	≥18	≥18 (22.5)
7	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
8	Male	Nigeria	<18	<18 (15.5)	<18	<18 (15.5)
9	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
10	Male	Gambia	<18	<18 (16.5)	<18	<18 (16.5)
11	Male	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
12	Male	Ivory Coast	<18	<18 (16.5)	<18	<18 (16.5)
13	Male	Bangladesh	≥18	≥18 (22.5)	≥18	≥18 (22.5)
14	Male	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
15	Male	Nigeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
16	Male	Nigeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)
17	Female	Nigeria	<18	<18 (16.5)	<18	<18 (16.5)
18	Female	Nigeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)
19	Female	Nigeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)
20	Male	Cameroon	≥18	≥18 (21.5)	≥18	≥18 (21.5)
21	Male	Senegal	≥18	≥18 (20.5)	≥18	≥18 (20.5)
22	Female	Eritrea	<18	<18 (16.5)	<18	<18 (16.5)
23	Male	Ethiopia	≥18	≥18 (18.5)	≥18	≥18 (18.5)
24	Male	Cameroon	≥18	≥18 (20.5)	≥18	≥18 (20.5)
25	Male	Sudan	≥18	≥18 (18.5)	≥18	≥18 (18.5)
26	Male	Somalia	≥18	≥18 (20.5)	≥18	≥18 (20.5)
27	Female	Nigeria	<18	<18 (17.5)	<18	<18 (17.5)
28	Male	syria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
29	Female	Eritrea	<18	<18 (17.5)	<18	<18 (17.5)
30	Female	Eritrea	<18	<18 (16.5)	<18	<18 (16.5)
31	Female	Nigeria	<18	<18 (16.5)	<18	<18 (16.5)
32	Female	Nigeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)
33	Female	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
34	Female	Nigeria	≥18	≥18 (18.5)	≥18	≥18 (18.5)
35	Male	Ghana	≥18	≥18 (19.5)	≥18	≥18 (19.5)
36	Male	Eritrea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
37	Male	Eritrea	<18	<18 (15.5)	<18	<18 (16.5)
38	Male	Sudan	≥18	≥18 (22.5)	Absent	Absent
39	Male	Eritrea	<18	<18 (17.5)	<18	<18 (17.5)
40	Male	Somalia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
41	Male	Somalia	<18	<18 (17.5)	<18	<18 (17.5)
42	Female	Nigeria	≥18	≥18 (21.5)	≥18	≥18 (21.5)
43	Female	Nigeria	≥18	≥18 (18.5)	≥18	≥18 (18.5)
44	Female	Nigeria	≥18	≥18 (21.5)	≥18	≥18 (21.5)
45	Female	Nigeria	≥18	≥18 (23.5)	≥18	≥18 (23.5)
46	Male	Morocco	Absent	Absent	Absent	Absent
47	Male	Morocco	≥18	≥18 (20.5)	≥18	≥18 (20.5)
48	Male	Morocco	≥18	≥18 (21.5)	≥18	≥18 (21.5)
49	Male	Pakistan	<18	<18 (17.5)	<18	<18 (17.5)
50	Male	Gambia	≥18	≥18 (19.5)	≥18	≥18 (19.5)
51	Male	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
52	Male	Pakistan	≥18	≥18 (22.5)	≥18	≥18 (22.5)
53	Male	Algeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
54	Male	Mali	≥18	≥18 (21.5)	≥18	≥18 (21.5)
55	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
56	Female	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
57	Male	Gabon	≥18	≥18 (21.5)	≥18	≥18 (21.5)
58	Male	Guinea	≥18	≥18 (21.5)	≥18	≥18 (21.5)
59	Male	Ivory Coast	≥18	≥18 (22.5)	≥18	≥18 (22.5)
60	Female	Nigeria	<18	≥18 (18.5)	<18	≥18 (18.5)
61	Male	Morocco	≥18	≥18 (19.5)	≥18	≥18 (19.5)
62	Female	Nigeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)
63	Female	Nigeria	<18	<18 (16.5)	<18	<18 (16.5)
64	Female	Nigeria	≥18	≥18 (18.5)	≥18	≥18 (18.5)
65	Male	Iraq	≥18	≥18 (19.5)	≥18	≥18 (19.5)
66	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
67	Female	Nigeria	<18	<18 (15.5)	<18	<18 (15.5)
68	Male	Ghana	≥18	≥18 (23.5)	≥18	≥18 (23.5)
69	Male	Morocco	≥18	≥18 (21.5)	≥18	≥18 (21.5)
70	Male	Ivory Coast	≥18	≥18 (19.5)	≥18	≥18 (19.5)
71	Female	Nigeria	<18	<18 (16.5)	<18	<18 (16.5)
72	Male	Egypt	<18	<18 (16.5)	<18	<18 (16.5)
73	Female	Nigeria	<18	<18 (15.5)	<18	<18 (15.5)

(continued on next page)

Table 1 (continued)

Case number	Sex	Nationality	Cameriere 38	AlQahtani 38	Cameriere 48	AlQahtani 48
74	Male	Somalia	≥18	≥18 (18.5)	≥18	≥18 (18.5)
75	Male	Nigeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
76	Female	Ivory Coast	≥18	≥18 (20.5)	≥18	≥18 (20.5)
77	Male	Mali	≥18	≥18 (20.5)	≥18	≥18 (20.5)
78	Male	Gambia	<18	≥18 (18.5)	<18	≥18 (18.5)
79	Male	Bangladesh	Absent	Absent	Absent	Absent
80	Male	Bangladesh	≥18	≥18 (20.5)	Absent	Absent
81	Male	Ghana	≥18	≥18 (23.5)	≥18	≥18 (23.5)
82	Male	Mali	≥18	≥18 (22.5)	≥18	≥18 (22.5)
83	Male	Gabon	≥18	≥18 (22.5)	≥18	≥18 (22.5)
84	Male	Togo	<18	<18 (17.5)	<18	<18 (17.5)
85	Male	Ivory Coast	≥18	≥18 (21.5)	≥18	≥18 (21.5)
86	Male	Bangladesh	Absent	Absent	≥18	≥18 (22.5)
87	Male	Bangladesh	≥18	≥18 (20.5)	≥18	≥18 (20.5)
88	Female	Nigeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
89	Male	Senegal	≥18	≥18 (19.5)	≥18	≥18 (19.5)
90	Male	Ghana	≥18	≥18 (19.5)	≥18	≥18 (19.5)
91	Male	Nigeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
92	Male	Senegal	≥18	≥18 (21.5)	≥18	≥18 (21.5)
93	Male	Guinea	≥18	≥18 (21.5)	≥18	≥18 (21.5)
94	Male	Guinea	≥18	≥18 (20.5)	≥18	≥18 (20.5)
95	Male	Morocco	≥18	≥18 (22.5)	≥18	≥18 (22.5)
96	Male	Nigeria	<18	<18 (17.5)	<18	<18 (17.5)
97	Male	Egypt	<18	<18 (15.5)	<18	<18 (15.5)
98	Male	Benin	Absent	Absent	<18	<18 (15.5)
99	Female	Nigeria	<18	≥18 (18.5)	<18	≥18 (18.5)
100	Female	Nigeria	<18	≥18 (18.5)	<18	≥18 (18.5)
101	Male	Nigeria	<18	<18 (17.5)	<18	<18 (17.5)
102	Male	Burkina Faso	≥18	≥18 (19.5)	≥18	≥18 (19.5)
103	Female	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
104	Female	Nigeria	≥18	≥18 (21.5)	≥18	≥18 (21.5)
105	Male	Somalia	<18	<18 (17.5)	<18	<18 (17.5)
106	Male	Bangladesh	≥18	≥18 (19.5)	≥18	≥18 (19.5)
107	Female	Nigeria	<18	≥18 (18.5)	<18	≥18 (18.5)
108	Male	Bangladesh	<18	<18 (16.5)	<18	<18 (16.5)
109	Male	Somalia	≥18	≥18 (19.5)	≥18	≥18 (19.5)
110	Male	Ivory Coast	≥18	≥18 (20.5)	≥18	≥18 (20.5)
111	Male	Gambia	Absent	Absent	≥18	≥18 (21.5)
112	Male	Algeria	<18	<18 (17.5)	<18	<18 (17.5)
113	Male	Morocco	≥18	≥18 (22.5)	≥18	≥18 (22.5)
114	Male	Egypt	≥18	≥18 (19.5)	≥18	≥18 (19.5)
115	Male	Guinea	≥18	≥18 (23.4)	≥18	≥18 (23.5)
116	Male	Mali	≥18	≥18 (21.5)	≥18	≥18 (21.5)
117	Male	Mali	≥18	≥18 (20.5)	≥18	≥18 (20.5)
118	Male	Mali	≥18	≥18 (21.5)	≥18	≥18 (21.5)
119	Male	Gabon	≥18	≥18 (22.5)	Absent	Absent
120	Male	Ivory Coast	≥18	≥18 (20.5)	≥18	≥18 (20.5)
121	Male	Ivory Coast	≥18	≥18 (20.4)	≥18	≥18 (20.5)
122	Male	Pakistan	<18	<18 (15.5)	<18	<18 (15.5)
123	Male	Senegal	≥18	≥18 (23.5)	Absent	Absent
124	Male	Senegal	≥18	≥18 (23.5)	≥18	≥18 (23.5)
125	Female	Nigeria	<18	<18 (15.5)	<18	<18 (15.5)
126	Male	Nigeria	<18	<18 (15.5)	<18	<18 (15.5)
127	Male	Bangladesh	≥18	≥18 (21.5)	≥18	≥18 (21.5)
128	Male	Myanmar	<18	<18 (17.5)	<18	<18 (17.5)
129	Male	Afgahnistan	<18	<18 (17.5)	<18	<18 (17.5)
130	Male	Afgahnistan	Absent	Absent	Absent	Absent
131	Male	Guinea	≥18	≥18 (23.5)	≥18	≥18 (23.5)
132	Male	Senegal	≥18	≥18 (23.5)	Absent	Absent
133	Male	Mali	≥18	≥18 (18.5)	≥18	≥18 (18.5)
134	Male	Guinea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
135	Male	Guinea	≥18	≥18 (18.5)	≥18	≥18 (18.5)
136	Male	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
137	Male	Algeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
138	Male	Guinea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
139	Male	Senegal	≥18	≥18 (21.5)	≥18	≥18 (21.5)
140	Male	Senegal	≥18	≥18 (21.5)	≥18	≥18 (21.5)
141	Male	Morocco	≥18	≥18 (19.5)	≥18	≥18 (19.5)
142	Male	Mali	≥18	≥18 (20.5)	Absent	Absent
143	Male	Pakistan	Absent	Absent	<18	<18 (17.5)
144	Male	Gambia	≥18	≥18 (21.5)	≥18	≥18 (21.5)
145	Male	Guinea	≥18	≥18 (22.5)	Absent	Absent
146	Male	Gambia	≥18	≥18 (20.5)	≥18	≥18 (20.5)
147	Male	Gabon	Absent	Absent	Absent	Absent
148	Female	Nigeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)

Table 1 (continued)

Case number	Sex	Nationality	Cameriere 38	AlQahtani 38	Cameriere 48	AlQahtani 48
149	Female	Romania	<18	<18 (15.5)	<18	<18 (15.5)
150	Female	Nigeria	<18	≥18 (18.5)	<18	≥18 (18.5)
151	Male	Pakistan	<18	<18 (16.5)	<18	<18 (16.5)
152	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
153	Male	Senegal	≥18	≥18 (19.5)	≥18	≥18 (19.5)
154	Male	Senegal	<18	<18 (17.5)	<18	<18 (17.5)
155	Male	Senegal	≥18	≥18 (20.5)	Absent	Absent
156	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
157	Male	Guinea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
158	Male	Senegal	≥18	≥18 (20.5)	≥18	≥18 (20.5)
159	Male	Gambia	≥18	≥18 (21.5)	≥18	≥18 (21.5)
160	Male	Bangladesh	≥18	≥18 (20.5)	≥18	≥18 (20.5)
161	Male	Senegal	Absent	Absent	≥18	≥18 (22.5)
162	Male	Roania	<18	<18 (17.5)	<18	<18 (17.5)
163	Male	Ivory Coast	≥18	≥18 (19.5)	≥18	≥18 (19.5)
164	Male	Tunisia	<18	<18 (17.5)	<18	<18 (17.5)
165	Male	Guinea	≥18	≥18 (20.5)	≥18	≥18 (20.5)
166	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
167	Male	Guinea	≥18	≥18 (21.5)	≥18	≥18 (21.5)
168	Male	Morocco	<18	<18 (17.5)	<18	<18 (17.5)
169	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
170	Male	Gabon	≥18	≥18 (23.5)	≥18	≥18 (23.5)
171	Male	Guinea	≥18	≥18 (23.5)	≥18	≥18 (23.5)
172	Male	Gambia	≥18	≥18 (23.5)	≥18	≥18 (23.5)
173	Male	Mali	≥18	≥18 (22.5)	≥18	≥18 (22.5)
174	Female	Nigeria	≥18	≥18 (21.5)	≥18	≥18 (21.5)
175	Male	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
176	Male	Morocco	<18	<18 (15.5)	<18	<18 (15.5)
177	Male	Guinea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
178	Male	Guinea	≥18	≥18 (21.5)	≥18	≥18 (21.5)
179	Male	Cameroon	≥18	≥18 (23.5)	≥18	≥18 (23.5)
180	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
181	Male	Guinea	≥18	≥18 (21.5)	≥18	≥18 (21.5)
182	Male	Cameroon	≥18	≥18 (22.5)	≥18	≥18 (22.5)
183	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
184	Female	Serbia	<18	<18 (15.5)	<18	<18 (15.5)
185	Male	Senegal	Absent	Absent	Absent	Absent
186	Male	Gambia	≥18	≥18 (23.5)	≥18	≥18 (23.5)
187	Male	Nigeria	≥18	≥18 (23.5)	≥18	≥18 (23.5)
188	Male	Nigeria	≥18	≥18 (23.5)	≥18	≥18 (23.5)
189	Male	Guinea	<18	<18 (17.5)	<18	<18 (17.5)
190	Male	Senegal	≥18	≥18 (22.5)	≥18	≥18 (22.5)
191	Male	Guinea	≥18	≥18 (20.5)	≥18	≥18 (20.5)
192	Male	Guinea	<18	<18 (17.5)	<18	<18 (17.5)
193	Male	Guinea	≥18	≥18 (20.5)	≥18	≥18 (20.5)
194	Male	Mali	≥18	≥18 (19.5)	≥18	≥18 (19.5)
195	Male	Ivory Coast	≥18	≥18 (20.5)	≥18	≥18 (20.5)
196	Male	Mali	≥18	≥18 (21.5)	≥18	≥18 (21.5)
197	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
198	Male	Guinea	Absent	Absent	≥18	≥18 (22.5)
199	Male	Guinea	≥18	≥18 (23.5)	≥18	≥18 (23.5)
200	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
201	Male	Nigeria	≥18	≥18 (20.5)	≥18	≥18 (20.5)
202	Male	Ivory Coast	≥18	≥18 (21.5)	≥18	≥18 (21.5)
203	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
204	Female	Nigeria	<18	≥18 (18.5)	<18	≥18 (18.5)
205	Male	Libya	Absent	Absent	≥18	≥18 (22.5)
206	Male	Sudan	≥18	≥18 (22.5)	≥18	≥18 (22.5)
207	Male	Ivory Coast	≥18	≥18 (23.5)	≥18	≥18 (23.5)
208	Male	Eritrea	≥18	≥18 (21.5)	≥18	≥18 (21.5)
209	Male	Egypt	<18	<18 (17.5)	<18	<18 (17.5)
210	Male	Senegal	≥18	≥18 (21.5)	Absent	Absent
211	Male	Senegal	≥18	≥18 (23.5)	≥18	≥18 (23.5)
212	Male	Tunisia	≥18	≥18 (23.5)	≥18	≥18 (23.5)
213	Male	Zambia	≥18	≥18 (23.5)	≥18	≥18 (23.5)
214	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
215	Male	Gambia	≥18	≥18 (21.5)	≥18	≥18 (21.5)
216	Male	Gambia	≥18	≥18 (20.5)	≥18	≥18 (20.5)
217	Male	Guinea Bassau	≥18	≥18 (20.5)	≥18	≥18 (20.5)
218	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
219	Male	Senegal	≥18	≥18 (22.5)	≥18	≥18 (22.5)
220	Male	Senegal	≥18	≥18 (21.5)	≥18	≥18 (21.5)
221	Male	Gabon	≥18	≥18 (22.5)	Absent	Absent
222	Male	Libya	Absent	Absent	Absent	Absent

(continued on next page)

Table 1 (continued)

Case number	Sex	Nationality	Cameriere 38	AlQahtani 38	Cameriere 48	AlQahtani 48
223	Female	Morocco	<18	<18 (16.5)	<18	<18 (16.5)
224	Male	Senegal	≥18	≥18 (22.5)	≥18	≥18 (22.5)
225	Male	Guinea Bissau	≥18	≥18 (20.5)	Absent	Absent
226	Male	Mali	Absent	Absent	<18	<18 (17.5)
227	Female	Nigeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
228	Male	Senegal	≥18	≥18 (23.5)	≥18	≥18 (23.5)
229	Male	Gambia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
230	Male	Morocco	≥18	≥18 (19.5)	≥18	≥18 (19.5)
231	Female	Nigeria	≥18	≥18 (23.5)	≥18	≥18 (23.5)
232	Male	Bangladesh	Absent	Absent	<18	<18 (17.5)
233	Male	Mali	≥18	≥18 (19.5)	≥18	≥18 (19.5)
234	Male	Senegal	≥18	≥18 (19.5)	≥18	≥18 (19.5)
235	Male	Benin	≥18	≥18 (22.5)	≥18	≥18 (22.5)
236	Male	Senegal	≥18	≥18 (21.5)	≥18	≥18 (21.5)
237	Male	Nigeria	Absent	Absent	Absent	Absent
238	Male	Nigeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
239	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
240	Male	Ivory Coast	≥18	≥18 (23.5)	Absent	Absent
241	Male	Senegal	<18	<18 (14.5)	Absent	Absent
242	Male	Morocco	≥18	≥18 (20.5)	≥18	≥18 (20.5)
243	Male	Eritrea	<18	<18 (15.5)	<18	<18 (15.5)
244	Male	Eritrea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
245	Male	Egypt	<18	<18 (17.5)	<18	<18 (17.5)
246	Male	Ivory Coast	<18	<18 (17.5)	<18	<18 (17.5)
247	Male	Guinea Bissau	≥18	≥18 (20.5)	≥18	≥18 (20.5)
248	Male	Bangladesh	≥18	≥18 (22.5)	≥18	≥18 (22.5)
249	Male	Senegal	≥18	≥18 (22.5)	≥18	≥18 (22.5)
250	Male	Gabon	≥18	≥18 (23.5)	≥18	≥18 (23.5)
251	Male	Mali	Absent	Absent	≥18	≥18 (21.5)
252	Male	Senegal	Absent	Absent	≥18	≥18 (22.5)
253	Male	Mali	≥18	≥18 (23.5)	≥18	≥18 (23.5)
254	Male	Ivory Coast	≥18	≥18 (22.5)	≥18	≥18 (22.5)
255	Male	Guinea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
256	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
257	Male	Mali	≥18	≥18 (20.5)	≥18	≥18 (20.5)
258	Male	Ivory Coast	≥18	≥18 (23.5)	≥18	≥18 (23.5)
259	Male	Afghanistan	<18	<18 (15.5)	<18	<18 (15.5)
260	Male	Sudan	≥18	≥18 (20.5)	≥18	≥18 (20.5)
261	Male	Bosnia-Erzegovina	<18	<18 (14.5)	<18	<18 (14.5)
262	Male	Mali	≥18	≥18 (23.5)	≥18	≥18 (23.5)
263	Male	Morocco	Absent	Absent	≥18	≥18 (22.5)
264	Male	Gambia	Absent	Absent	Absent	Absent
265	Male	Tunisia	≥18	≥18 (19.5)	≥18	≥18 (19.5)
266	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
267	Male	Senegal	≥18	≥18 (20.5)	≥18	≥18 (20.5)
268	Male	Senegal	≥18	≥18 (23.5)	≥18	≥18 (23.5)
269	Male	Gabon	≥18	≥18 (23.5)	≥18	≥18 (23.5)
270	Female	Tunisia	<18	<18 (17.5)	<18	<18 (17.5)
271	Female	Mali	<18	<18 (16.5)	<18	<18 (16.5)
272	Male	Senegal	≥18	≥18 (19.5)	≥18	≥18 (19.5)
273	Male	Morocco	Absent	Absent	<18	<18 (17.5)
274	Male	Senegal	Absent	Absent	≥18	≥18 (18.5)
275	Male	Senegal	≥18	≥18 (22.5)	≥18	≥18 (22.5)
276	Male	Romania	Absent	Absent	Absent	Absent
277	Male	Mali	≥18	≥18 (19.5)	≥18	≥18 (19.5)
278	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
279	Male	Guinea	≥18	≥18 (22.5)	≥18	≥18 (22.5)
280	Male	Tunisia	≥18	≥18 (20.5)	≥18	≥18 (20.5)
281	Male	Tunisia	≥18	≥18 (21.5)	≥18	≥18 (21.5)
282	Male	Tunisia	≥18	≥18 (22.5)	≥18	≥18 (22.5)
283	Male	Chad	≥18	≥18 (22.5)	≥18	≥18 (22.5)
284	Male	Tunisia	≥18	≥18 (23.5)	≥18	≥18 (23.5)
285	Male	Guinea	≥18	≥18 (19.5)	≥18	≥18 (19.5)
286	Male	Tunisia	Absent	Absent	Absent	Absent
287	Male	Algeria	≥18	≥18 (19.5)	≥18	≥18 (19.5)
288	Male	Tunisia	≥18	≥18 (20.5)	≥18	≥18 (20.5)
289	Male	Mauritania	≥18	≥18 (22.5)	Absent	Absent
290	Male	Cameroon	≥18	≥18 (22.5)	≥18	≥18 (22.5)
291	Male	Guinea	≥18	≥18 (20.5)	≥18	≥18 (20.5)
292	Male	Tunisia	≥18	≥18 (19.5)	≥18	≥18 (19.5)
293	Male	Morocco	≥18	≥18 (21.5)	≥18	≥18 (21.5)
294	Female	Hungary	<18	<18 (14.5)	<18	<18 (14.5)
295	Male	Senegal	Absent	Absent	≥18	≥18 (21.5)
296	Male	Ivory Coast	≥18	≥18 (20.5)	≥18	≥18 (20.5)
297	Male	Tunisia	≥18	≥18 (18.5)	<18	<18 (17.5)

Table 1 (continued)

Case number	Sex	Nationality	Cameriere 38	AlQahtani 38	Cameriere 48	AlQahtani 48
298	Male	Afghanistan	<18	<18 (17.5)	<18	<18 (17.5)
299	Male	Algeria	≥18	≥18 (22.5)	≥18	≥18 (22.5)
300	Male	Ivory Coast	≥18	≥18 (23.5)	≥18	≥18 (23.5)
301	Male	Gambia	≥18	≥18 (23.5)	≥18	≥18 (23.5)

In these cases, the latter method should be used in association with the Cameriere one, in order to minimize the risk to define a minor as adult. The present manuscript does not allow to understand which of the two methods correctly defines legal age in the aforementioned 7 cases, because actual chronological ages of the study sample were unknown. However, it is important to note that in 2017 the Council of European Union (Child Right Division) recommended “Authorities to interpret inconclusive results in the applicant’s favour, *in dubio pro refugio* or *in dubio pro minore*” [26]. Thus, in such cases the use of both methods allows a proper implementation of the following fundamental principle: the so called benefit of the doubt.

The absence of both 38 and 48 in the same individual was registered in 10/301 cases (3.32%). The latter result shows that age estimation methods that rely on mandibular third molars can be widely useable for age assessment. In addition, the present study shows a high concordance in legal age definition, comparing results obtained evaluating 38 and 48 of the same person. Indeed, except for one individual, in all cases right and left mandibular third molars were characterized by the same developmental stage. Only in one case the operator/observer defined (through both methods) the left mandibular third molar as correspondent to ≥ 18 years and the right one as correspondent to < 18 years. These findings are meaningful because they highlight that both methods are applicable to age estimation of people who have only one of mandibular third molars.

Limitations of the present study rely on the lack of actual chronological ages of the study sample; for this reason, bias and inaccuracy could not be calculated [27]. In particular, sensitivity/specificity and positive/negative predictive values of K could not be obtained, because according to the scientific literature neither AlQahtani atlas nor Cameriere third molar index can be identified as “gold-standard” method [26,27]. However, these limitations are consistent with study design that was aimed to apply two dental ageing methods onto migrants without valid identify documents in a real forensic casework.

Further limitations rely on the study population. Indeed, it is well known that age estimation methods are strictly dependent on the underlying structure of datasets used to construct them. Therefore, results of their application in different study populations can be biased because of intrinsic differences between study samples. It is important to note that the London Atlas of Human Tooth Development and Eruption by AlQahtani and colleagues is based on a sample of half Caucasian and half Bangladeshi individuals [14]; whereas the Cameriere index was obtained by way of the study of an entirely Caucasian population [18]. In the present paper, the latter methods were used to analyze a population composed by a significant number of North African and sub-Saharan Africans. It is important to highlight this discrepancy because it can negatively influence study results. In the scientific literature, there are several manuscripts in which authors validate age estimation methods on populations of different geographical areas. For example, the third molar index was used to determine minor/major age of South African samples [28,29], stating that “I_{3M} is a valuable method to distinguish subjects who are around legal adult age in South Africa” [28]. However, even if an increasing number of population studies have been reported [28,29], the scientific literature is far behind the

validation of age estimation methods for several populations of different geographical areas. For these reasons, the above-mentioned considerations should be taken into account in order to better understand study results, and to promote the implementation of further researches in this field.

5. Conclusions

The study demonstrates that AlQahtani and Cameriere methods are characterized by a high statistical agreement. For this reason, they can be alternately used for legal age assessment. In case of the London Atlas identifies a chronological age of 18.5 years, operators should use both methods in order to minimize the risk to define a minor as adult.

Disclaimers

None.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors have not conflict of interest to disclose.

Acknowledgments

None.

References

- [1] Assembly Un General, Convention on the Rights of the Child. United Nations, Treaty Series vol. 1577, 1989, 3, <https://www.ohchr.org/en/professionalinterest/pages/crc.aspx>.
- [2] S.J. Alqahtani, M.P. Hector, H.M. Liversidge, Accuracy of dental age estimation charts: schour and massler, ubelaker and the London atlas, *Am. J. Phys. Anthropol.* 154 (1) (2014) 70–78.
- [3] R.W. Nahhas, R.J. Sherwood, W.C. Chumlea, D.L. Duren, An update of the statistical methods underlying the FELS method of skeletal maturity assessment, *Ann. Hum. Biol.* 40 (6) (2013) 505–514.
- [4] S. Serinelli, V. Panetta, P. Pasqualetti, D. Marchetti, Accuracy of three age determination X-ray methods on the left hand-wrist: a systematic review and meta-analysis, *Leg. Med.* 13 (3) (2011) 120–133.
- [5] A. Demirjian, H. Goldstein, J.M. Tanner, A new system of dental age assessment, *Hum. Biol.* 45 (2) (1973) 211–227.
- [6] P. Chariot, C. Denis, Commentary on: hackman L, black S. The reliability of the greulich and pyle atlas when applied to a modern scottish population, *J. Forensic Sci.* 58 (1) (2013) 114–119. *J. Forensic Sci.* 2013;58(4):1106.
- [7] A. Schmeling, C. Grundmann, A. Fuhrmann, et al., Criteria for age estimation in living individuals, *Int. J. Leg. Med.* 122 (6) (2008) 457–460.
- [8] O. Ekizoglu, E. Hocaoglu, E. Inci, et al., Forensic age estimation by the Schmeling method: computed tomography analysis of the medial clavicular epiphysis, *Int. J. Leg. Med.* 129 (1) (2015) 203–210.
- [9] A. Olze, W. Reisinger, G. Geserick, A. Schmeling, Age estimation of unaccompanied minors. Part II. Dental aspects, *Forensic Sci. Int.* 159 (Suppl 1) (2006) S65–S67.
- [10] A. Olze, P. Van niekerk, R. Schulz, S. Ribbecke, A. Schmeling, The influence of impaction on the rate of third molar mineralisation in male black Africans, *Int. J. Leg. Med.* 126 (6) (2012) 869–874.
- [11] A. Olze, B.R. Pynn, V. Kraul, et al., Studies on the chronology of third molar mineralization in First Nations people of Canada, *Int. J. Leg. Med.* 124 (5)

- (2010) 433–437.
- [12] A. Olze, P. Van niekerk, T. Ishikawa, et al., Comparative study on the effect of ethnicity on wisdom tooth eruption, *Int. J. Leg. Med.* 121 (6) (2007) 445–448.
- [13] T.Y. Marroquin, S. Karkhanis, S.I. Kvaal, S. Vasudavan, E. Kruger, M. Tennant, Age estimation in adults by dental imaging assessment systematic review, *Forensic Sci. Int.* 275 (2017) 203–211.
- [14] S.J. Alqahtani, M.P. Hector, H.M. Liversidge, Brief communication: the London atlas of human tooth development and eruption, *Am. J. Phys. Anthropol.* 142 (3) (2010) 481–490.
- [15] D.M. Alsudairi, S.J. Alqahtani, Testing and comparing the accuracy of two dental age estimation methods on Saudi children: measurements of open apices in teeth and the London Atlas of Tooth Development, *Forensic Sci. Int.* 295 (2019) 226.e1–226.e9.
- [16] S. Pavlović, C. Palmela Pereira, R.F. Vargas de Sousa Santos, Age estimation in Portuguese population: the application of the London atlas of tooth development and eruption, *Forensic Sci. Int.* 272 (2017) 97–103.
- [17] P. Sharma, V. Wadhwan, Comparison of accuracy of age estimation in Indian children by measurement of open apices in teeth with the London Atlas of tooth development, *J. Forensic Odontostomatol* 1 (38) (2020) 39–47.
- [18] R. Cameriere, L. Ferrante, D. De Angelis, F. Scarpino, F. Galli, The comparison between measurement of open apices of third molars and Demirjian stages to test chronological age of over 18 year olds in living subjects, *Int. J. Leg. Med.* 122 (6) (2008) 493–497.
- [19] L. Ribier, P. Saint-martin, M. Seignier, A. Paré, L. Brunereau, C. Rérolle, Cameriere's third molar maturity index in assessing age of majority: a study of a French sample, *Int. J. Leg. Med.* 134 (2) (2020) 783–792.
- [20] A. Schmeling, R. Dettmeyer, E. Rudolf, V. Vieth, G. Geserick, Forensic age estimation, *Dtsch Arztebl Int* 113 (4) (2016) 44–50.
- [21] J. Costa, J. Montero, S. Serrano, A. Albaladejo, A. López-valverde, I. Bica, Accuracy in the legal age estimation according to the third molars mineralization among Mexicans and Columbians, *Atención Primaria* 46 (Suppl 5) (2014) 165–175.
- [22] J.C. Turp, K.W. Alt, Designating teeth: the advantages of the FDI's two-digit system, *Quintessence Int.* 26 (7) (1995) 501–504.
- [23] P.F. Watson, A. Petrie, Method agreement analysis: a review of correct methodology, *Theriogenology* 73 (9) (2010) 1167–1179.
- [24] P. Ranganathan, C.S. Pramesh, R. Aggarwal, Common pitfalls in statistical analysis: measures of agreement, *Perspect Clin Res* 8 (4) (2017) 187–191.
- [25] A.J. Viera, J.M. Garrett, Understanding interobserver agreement: the kappa statistic, *Fam. Med.* 37 (5) (2005) 360–363.
- [26] D. Wenke, Age Assessment: Council of Europe Member States' Policies, Procedures and Practices Respectful of Children's Rights in the Context of Migration, Council of Europe, 2017, in: <https://rm.coe.int/age-assessment-council-of-europe-member-states-policies-procedures-and/168074b723>.
- [27] J. Peacock, P. Peacock, *Oxford Handbook of Medical Statistics*, Oxford University Press, 2011, pp. 340–345.
- [28] N. Angelakopoulos, S. De Luca, L.A. Velandia Palacio, E. Coccia, L. Ferrante, R. Cameriere, Third molar maturity index (I3m) for assessing age of majority: study of a black South African sample, *Int. J. Leg. Med.* 132 (5) (2018) 1457–1464.
- [29] J. Cavrić, I. Galić, M. Vodanović, et al., Third molar maturity index (I3m) for assessing age of majority in a black African population in Botswana, *Int. J. Leg. Med.* 130 (4) (2016) 1109–1120.