



# Complete Genome Sequence of *Cupriavidus necator* KK10, an Azaarene-Degrading and Polyhydroxyalkanoate-Producing Soil Bacterium

 Jiro F. Mori,<sup>a</sup> Miho Nagai,<sup>a</sup> Robert A. Kanaly<sup>a</sup>

<sup>a</sup>Department of Life and Environmental System Science, Graduate School of Nanobiosciences, Yokohama City University, Kanagawa, Yokohama, Japan

**ABSTRACT** *Cupriavidus necator* KK10 has been investigated in azaarene and diesel fuel biodegradation studies and is capable of polyhydroxyalkanoate (PHA) production. Its complete genome sequence revealed two closed circular sequences, the chromosome (6.68 Mb) and plasmid (1.67 Mb). The KK10 genome carries functional genes potentially responsible for azaarene biodegradation and polyhydroxyalkanoate biosynthesis.

*Cupriavidus necator* KK10, a member of the *Betaproteobacteria*, was isolated from a cattle pasture soil bacterial consortium by streaking on Noble agar with the heavy fraction of a diesel fuel as the carbon source (1–5). KK10 biotransforms azaarenes and single-ring aromatic compounds (4–6) and produces polyhydroxyalkanoates (PHA). Sequencing the genome of KK10 expands our understanding of this versatile and potentially useful bacterium.

Hybrid assembly was conducted by utilizing short-read (DNBSEQ-G400; MGI Tech, China) and long-read (GridION X5; Oxford Nanopore Technologies, UK) sequencing to close the chromosomal and plasmid sequences similarly to a method described previously (7). The strain was grown on 20 mM glycerol in Stanier's basal medium for 3 days at 30°C by rotary shaking (150 rpm in darkness), DNA was extracted with the NucleoBond high-molecular-weight (HMW) DNA kit (Macherey-Nagel, Germany), and a DNBseq library was prepared with the MGIEasy FS DNA library prep set (MGI Tech). A total of 6,210,182 reads were obtained with 200-bp paired-end read lengths that were trimmed and quality filtered using Cutadapt v. 2.7, SeqKit v. 0.11.0, and Sickle v. 1.33. Then, 1,000 ng of genomic DNA was bar-coded using native barcoding expansion for GridION analysis, and a library was created using a ligation sequencing kit (SQK-LSK109; Oxford Nanopore Technologies; after adapter ligation, >3-kb fragments were enriched). From the GridION platform, using R9.4.1 flow cells and Guppy v. 4.0.11 for live base-calling, a total of 130,465 reads with an average length of 6,801 bp were generated. Raw reads were trimmed and quality filtered using Porechop v. 0.2.3 and Filtlong v. 0.2.0 (minimum length, 1,000 bp). All software was used with default settings unless otherwise indicated. *De novo* hybrid assembly was conducted after trimming and quality filtering with Unicycler v. 0.4.7 (8) and was validated with Bandage v. 0.8.1.

The complete genome was determined (8,350,386 bp; 255× coverage) and consisted of a circular chromosome (6,679,877 bp) and plasmid (1,670,509 bp) with an overall G+C content of 65.6%. The NCBI Prokaryotic Genome Annotation Pipeline v. 5.1 was utilized to perform gene annotation, whereby 7,750 coding sequences (CDSs), 15 rRNAs, 65 tRNAs, and 2 transfer-messenger RNAs (tmRNAs) were identified; 6,142 (79.25%) and 1,608 (20.75%) CDSs were located on the chromosome and plasmid, respectively.

According to homologous gene identification using the Integrated Microbial Genome and Microbiomes (IMG/MER) system, a *phaCABR* gene cluster, which includes genes responsible for PHA biosynthesis (9) (PHA polymerase, beta-ketothiolase, acetoacetyl coenzyme A reductase, and PHA synthesis repressor, respectively) was found located on the chromosome and supported PHA synthesis in KK10. At the same time, homologues of the

**Citation** Mori JF, Nagai M, Kanaly RA. 2021. Complete genome sequence of *Cupriavidus necator* KK10, an azaarene-degrading and polyhydroxyalkanoate-producing soil bacterium. *Microbiol Resour Announc* 10: e00423-21. <https://doi.org/10.1128/MRA.00423-21>.

**Editor** Irene L. G. Newton, Indiana University, Bloomington

**Copyright** © 2021 Mori et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jiro F. Mori, [morij@yokohama-cu.ac.jp](mailto:morij@yokohama-cu.ac.jp).

**Received** 2 May 2021

**Accepted** 14 June 2021

**Published** 15 July 2021

*indAB* genes, which encode an indole monooxygenase (10), were found on the main chromosome, and their identification confirmed the capabilities of KK10 to biotransform azaarenes such as indole to produce indigoids (4, 5).

**Data availability.** The sequences were deposited in NCBI GenBank under the accession numbers [CP073677](#) and [CP073678](#) and in the IMG/MER database under accession number [2913661577](#). The raw sequences are available from SRA accession numbers [SRR14308055](#) and [SRR14308056](#) under BioProject number [PRJNA722091](#) and BioSample number [SAMN18744514](#).

## ACKNOWLEDGMENTS

This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI grant 17K00555 to R.A.K. and grant 19K15738 to J.F.M.

Hybrid assembly complete genome sequencing was conducted by the Bioengineering Lab Co., Ltd. (Sagamihara, Japan).

## REFERENCES

1. Fukuoka K, Ozeki Y, Kanaly RA. 2015. Aerobic biotransformation of 3-methylindole to ring cleavage products by *Cupriavidus* sp. strain KK10. *Biodegradation* 26:359–373. <https://doi.org/10.1007/s10532-015-9739-0>.
2. Fukuoka K, Tanaka K, Ozeki Y, Kanaly RA. 2015. Biotransformation of indole by *Cupriavidus* sp. strain KK10 proceeds through *N*-heterocyclic and carbocyclic-aromatic ring cleavage and production of indigoids. *Int Biodeterior Biodegradation* 97:13–24. <https://doi.org/10.1016/j.ibiod.2014.11.007>.
3. Kanaly RA, Bartha R. 1999. Cometabolic mineralization of benzo[*a*]pyrene caused by hydrocarbon additions to soil. *Environ Toxicol Chem* 18:2186–2190. <https://doi.org/10.1002/etc.5620181010>.
4. Kanaly RA, Bartha R, Watanabe K, Harayama S. 2001. Enhanced mineralization of benzo[*a*]pyrene in the presence of nonaqueous phase liquids. *Environ Toxicol Chem* 20:498–501. <https://doi.org/10.1002/etc.5620200307>.
5. Kanaly RA, Watanabe K. 2004. Multiple mechanisms contribute to the biodegradation of benzo[*a*]pyrene by petroleum-derived multicomponent nonaqueous-phase liquids. *Environ Toxicol Chem* 23:850–856. <https://doi.org/10.1897/03-191>.
6. Mori JF, Kanaly RA. 2020. Multispecies diesel fuel biodegradation and niche formation are ignited by pioneer hydrocarbon-utilizing proteobacteria in a soil bacterial consortium. *Appl Environ Microbiol* 87:e02268-20. <https://doi.org/10.1128/AEM.02268-20>.
7. Mori JF, Kanaly RA. 2021. Complete genome sequence of *Sphingobium barthaii* KK22, a high-molecular-weight polycyclic aromatic hydrocarbon-degrading soil bacterium. *Microbiol Resour Announc* 10:e01250-20. <https://doi.org/10.1128/MRA.01250-20>.
8. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
9. Fukui T, Ohsawa K, Mifune J, Orita I, Nakamura S. 2011. Evaluation of promoters for gene expression in polyhydroxyalkanoate-producing *Cupriavidus necator* H16. *Appl Microbiol Biotechnol* 89:1527–1536. <https://doi.org/10.1007/s00253-011-3100-2>.
10. Dai CX, Ma Q, Li Y, Zhou DD, Yang BY, Qu YY. 2019. Application of an efficient indole oxygenase system from *Cupriavidus* sp. SHE for indigo production. *Bio-process Biosyst Eng* 42:1963–1971. <https://doi.org/10.1007/s00449-019-02189-4>.