

CORRECTION

Correction: Association between Perfluorinated Compound Exposure and Miscarriage in Danish Pregnant Women

Tina Kold Jensen, Louise Bjørkholt Andersen, Henriette Boye Kyhl, Flemming Nielsen, Henrik Thybo Christesen, Philippe Grandjean

In [Table 2](#), there are errors in the “N Miscarriage/birth” column. Please see the corrected [Table 2](#) here.



 OPEN ACCESS

Citation: Jensen TK, Andersen LB, Kyhl HB, Nielsen F, Christesen HT, Grandjean P (2016) Correction: Association between Perfluorinated Compound Exposure and Miscarriage in Danish Pregnant Women. PLoS ONE 11(2): e0149366. doi:10.1371/journal.pone.0149366

Published: February 9, 2016

Copyright: © 2016 Jensen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 2. Unadjusted and adjusted odds ratio (OR) and 95% confidence interval (95% CI) for miscarriage according to PFAS exposure as a continuous variable and in tertiles among 392 pregnant women from Odense, Denmark 2010–2012.

PFAS exposure (ng/mL)	N Miscarriage/birth	Case-cohort design				Matched case-control design			
		Unadjusted OR for miscarriage		Adjusted ¹⁾ OR for miscarriage		Unadjusted OR for miscarriage		Adjusted ¹⁾ OR for miscarriage	
		OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
PFHxS									
1 st tertile	19/142	Reference		Reference		Reference		Reference	
2 nd tertile	19/99	1.43	0.72–2.85	1.46	0.72–3.00	1.46	0.68–3.16	1.30	0.58–2.89
3 rd tertile	18/95	1.42	0.71–2.84	1.46	0.68–3.13	1.73	0.80–3.78	1.53	0.67–3.51
Continuous ²⁾	56/336	1.47	0.98–2.21	1.53	0.99–2.38				
PFOS									
1 th tertile	19/128	Reference		Reference		Reference		Reference	
2 th tertile	19/114	1.12	0.57–2.23	1.15	0.57–2.35	1.12	0.52–2.40	1.03	0.47–2.29
3 th tertile	18/94	1.29	0.64–2.59	1.33	0.64–2.78	1.25	0.57–2.71	1.06	0.46–2.41
Continuous ²⁾	56/336	1.12	0.59–2.12	1.16	0.59–1.29				
PFOA									
1 st tertile	19/116	Reference		Reference		Reference		Reference	
2 nd tertile	19/98	1.18	0.59–2.36	1.23	0.60–2.51	1.15	0.53–2.51	1.21	0.53–2.73
3 rd tertile	18/122	0.90	0.45–1.80	0.93	0.42–2.03	0.69	0.29–1.62	0.68	0.28–1.67
Continuous ²⁾	56/336	0.67	0.41–1.10	0.64	0.36–1.18				
PFNA									
1 th tertile	18/274	Reference		Reference		Reference		Reference	
2 th tertile	19/34	8.51	4.07–17.77	10.88	4.76–24.84	17.55	5.73–53.75	18.97	5.49–65.58
3 th tertile	19/28	10.33	4.87–21.93	16.17	6.88–38.03	32.06	9.31–110.36	37.93	9.90–145.2
Continuous ²⁾	56/336	11.82	5.78–24.19	16.46	7.39–36.62				
PFDA									
1 th tertile	19/180	Reference		Reference		Reference		Reference	
2 th tertile	18/86	1.98	0.99–3.97	1.86	0.91–3.83	1.89	0.86–4.17	1.85	0.77–4.46
3 th tertile	19/70	2.57	1.29–5.14	2.67	1.31–5.44	3.08	1.45–6.57	3.71	1.60–8.60
Continuous ²⁾	56/336	2.29	1.20–4.39	2.30	1.18–4.47				

In addition, conditional logistic regression on 1:4 matched data (on parity and gestational week at inclusion) with 51 cases of miscarriage and 204 controls.

1. Adjusted for age, BMI, parity and gestational age at serum sampling.

2. Transformed by the use of the natural logarithm. *p<0.05

doi:10.1371/journal.pone.0149366.t001

Reference

1. Jensen TK, Andersen LB, Kyhl HB, Nielsen F, Christesen HT, Grandjean P (2015) Association between Perfluorinated Compound Exposure and Miscarriage in Danish Pregnant Women. PLoS ONE 10(4): e0123496. doi:10.1371/journal.pone.0123496 PMID: 25848775