

# Review of the Psychodinae from Mallorca, Spain, with description of *Pericoma unipennata*, sp. n. (Diptera, Psychodidae)

Gunnar Mikalsen Kvifte<sup>1,2</sup>, Morten Stokkan<sup>2,3</sup>, Rüdiger Wagner<sup>1</sup>

**1** Dept. of Zoology-Limnology, University of Kassel, Heinrich-Plett-Strasse 40, 34132 Kassel-Oberwehren, Germany **2** Dept. of Natural History, University Museum of Bergen, P.O. Box 7800, University of Bergen, 5040 Bergen, Norway **3** Department of Biodiversity and Conservation, Instituto Mediterraneo de Estudios Avanzados (IMEDEA, CSIC-UIB), Esporles, Spain

Corresponding author: Gunnar Mikalsen Kvifte ([Gunnar.Kvifte@uib.no](mailto:Gunnar.Kvifte@uib.no))

---

Academic editor: J. Salmela | Received 2 January 2016 | Accepted 6 March 2016 | Published 5 April 2016

---

<http://zoobank.org/99814AF8-365D-43BA-AEE7-521ECCF64238>

---

**Citation:** Kvifte GM, Stokkan M, Wagner R (2016) Review of the Psychodinae from Mallorca, Spain, with description of *Pericoma unipennata*, sp. n. (Diptera, Psychodidae). ZooKeys 577: 149–160. doi: 10.3897/zookeys.577.7679

---

## Abstract

We review the Psychodinae of Mallorca, recognising fifteen species based on recent collections and available literature. Previously unpublished data is presented for eleven species, of which *Neoariseumus ibericus* Wagner, 1978, *Mormia tenebricosa* (Vaillant, 1954), *Clogmia albipunctata* (Williston, 1893), *Lepiseodina rothschildi* (Eaton, 1913), *Paramormia ustulata* (Walker, 1856), *Philosepedon pyrenaicus* Vaillant, 1974 and *Psychoda (Psycha) griseascens* Tonnoir, 1922 are first records for Mallorca. An old record of *Pericoma trifasciata* (Meigen, 1804) is considered doubtful. *Pericoma unipennata* **sp. n.** is described and illustrated based on a male collected at Deía. Distributional data are reviewed for all newly recorded species. Based on the Psychodinae fauna, the zoogeographical affinities of Mallorca are briefly discussed.

## Keywords

Moth flies, Balearic Isles, new species, faunistics, distribution, check list

## Introduction

The Psychodidae (moth flies and sand flies) are a species-rich and widespread group of small insects mainly associated with humid habitats. The most thoroughly studied faunas of the group are found in Europe, from where more than 500 species have been described and new discoveries continue to be made (Wagner 2004, Wagner and Kvifte 2015). The most poorly studied moth fly faunas of Europe are those of the Iberian peninsula (Wagner 2001) and the Balearic islands of Spain.

The most thoroughly studied moth flies from the Balearic islands belong to the subfamily Phlebotominae, the sand flies. On Mallorca, the largest of the Balearic islands, phlebotomine sand flies have received much attention due to their significance in veterinary medicine as vectors of canine leishmaniasis. This disease is widespread and long established, to the point that some local dog breeds have evolved resistance to the parasite (Solano-Gallego et al. 2000). Four species of *Phlebotomus* and one of *Sergentomyia* have been recorded from the island, of which *Phlebotomus ariasi* Tonnoir, 1921 has been suggested as a doubtful record (Alcover et al. 2014).

Other subfamilies of Psychodidae have been studied far less. Wagner (1990) listed a single species, namely *Pericoma barbarica* Vaillant, 1955 based on Vaillant (1978). Wagner et al. (2002) further included a record of *Pericoma trifasciata* (Meigen, 1818), most likely based on a typographical error in Vaillant (1978). The *P. trifasciata* record was not mentioned by Wagner (2004), who instead listed *Psychoda minuta* Banks, 1894, *P. phalaenoides* (Linnaeus, 1758), *P. (Tineria) alternata* Say, 1824 and *P. (Tinearia) lativentris* Berdén, 1952 based on unpublished material (the latter two placed in *Tinearia* Schellenberg, 1803, treated by Wagner 2004 as a genus).

In the present study, we review existing records of Mallorcan Psychodinae and present new material for eleven species; seven of which are previously unknown from Mallorca. In addition, we describe *Pericoma unipennata* sp. n. as new to science.

## Material and methods

Specimens were collected mainly by sweep netting and with aspirators from vegetation near the presumed larval habitats and preserved in 70–100% alcohol. Male specimens were sorted, dissected and mounted on slides in euparal (material in coll. ZFMK and ZMUB) or Canada balsam (material in coll. RW). Morphological terminology is according to Quate and Brown (2004) and Kvifte (2014, 2015). The „median moveable appendage“ in Kvifte et al. (2013) is here recognised as a *parameral sclerite*. Measurements are given in  $\mu\text{m}$  with an accuracy of 3  $\mu\text{m}$ ; except wing length which is given in mm to an accuracy of 100  $\mu\text{m}$ .

Both literature records and new material that we have examined are included in our present checklist. Tribe-level classification is given according to Duckhouse (1987), while genus-level taxonomy is according to Wagner (2004) except where noted otherwise in the text. Species recorded as new to Mallorca are marked with an asterisk (\*).

The material is deposited in the following collections:

**RW** Private collection of Rüdiger Wagner, Kassel

**ZFMK** Alexander-König Zoologischer Forschungsmuseum, Bonn, Germany

**ZMUB** Entomology Collections, Dept. of Natural History, University Museum of Bergen, Bergen, Norway

## Species list

### Maruinini

\* *Neoariseinus ibericus* Wagner, 1978

**First record from Mallorca.** Puigpunyent, 39.6167°N, 2.5167°E, 6.x.1981, H. Malicky leg. 1♂ (RW).

**Remarks.** Previously only recorded from the type locality in northern Spain.

### Mormiini

\* *Mormia tenebricosa* (Vaillant, 1954)

**First records from Mallorca.** Calobra, 39.85°N, 2.8°E, 9.v.1978, 90 m a.s.l., H. Malicky leg. 1♂ (RW);

Deiá, town fountain, 39.748072°N, 2.643385°E, 8.ii.2015, G.Kvifte, M. Stokkan & C. Garcia leg. 1♂ (ZFMK);

Puigpunyent, 39.62°N, 2.85°E, 12.v.1978, 200m a.s.l., H. Malicky leg., 1♂; same but 1.X.1979, 6♂♂; same but 6.x.1979, 6♂♂ (all RW);

South slope of Puig Major, north of Soller, 39.783°N, 2.767°E, 7.–9.v.1978, 700 m a.s.l., H. Malicky leg. 2♂♂ (RW).

**Remarks.** Previously known from France, mainland Spain, Italy, Morocco and Algeria (Vaillant 1974). The generic and subgeneric classification of Mormiina is unstable and requires revision; therefore the placement in *Mormia* may change.

### Paramormiini

\* *Clogmia albipunctata* (Williston, 1893)

**First records from Mallorca.** Esporles, IMEDEA research center, 39.666438°N, 2.580863°E, 13.iv.2015, M. Stokkan leg. 1♀ (ZMFK);

E of Puigpunyent, 39.616667°N, 2.85°E, 1.x.1979, 200m a.s.l., H. Malicky leg. 1♀; same but 6.x.1979, 4♀♀ (all RW).

**Remarks.** A widespread near-cosmopolitan species, first recorded from Spain by Tonnoir (1920) under the synonym *Telmatoscopus meridionalis* (Eaton, 1894). Its biology is summarised in Boumans et al. (2009) and Ježek et al. (2012); for a review of its taxonomy see Ibañez-Bernal (2008).

\* *Lepiseodina rothschildi* (Eaton, 1912)

**First record from Mallorca.** SW Pollensa, 39.8833°N, 2.9833°E, 3–5.x.1981, H. Malicky leg. 1♂ (RW).

**Remarks.** Placed in *Lepiseodina* Enderlein, 1937 (type species *Psychoda tristis* Meigen, 1830) by Ježek (1990) because of its asymmetric genitalia. The placement in *Clogmia* by Ježek (1984), Vaillant (1989) and Wagner (2004) may be a valid option to consider for the future when further character systems are explored.

Previous records are from Austria, Belgium, Czech Republic, France, Germany, Great Britain, Ireland and Slovakia (Oboňa and Ježek 2012).

\* *Paramormia ustulata* (Walker, 1856)

**First records from Mallorca.** Banyalbufar, 39.691635°N, 2.514367°E, 25.x.2012, G.Kvifte leg. 1♂1♀ (ZMUB);

Pond west of Cala Figuera, 39.335635°N, 3.152597°E, 11.ii.2015, G.Kvifte leg. 1♂ (ZFMK).

**Remarks.** A widespread species or complex of species occurring in the Holarctic region. Ježek and Yağci (2005) list occurrences from the following countries: Afghanistan, Algeria, the Azores, Belgium, Bosnia-Herzegovina, Bulgaria, the Canary Islands, China, Corsica, Czech Republic, Denmark, France, Germany, Great Britain, Greece, Hungary, Iran, Ireland, Israel, Italy, Macedonia, Madeira, Mongolia, Morocco, the Netherlands, Olanda island, Poland, Romania, Sardinia, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Tunisia, Turkey and the USA.


## Pericomaini

*Pericoma unipennata* Kvifte, Stokkan & Wagner, sp. n.

<http://zoobank.org/14619D1C-5A7E-4197-B99C-2C004DE54E83>

**Type material.** Holotype male. Deiá, town fountain, 39.748072°N, 2.643385°E, 8.ii.2015, G.Kvifte, M. Stokkan & C. Garcia leg (ZMUB).

**Diagnosis.** *Pericoma unipennata* can be separated from all other *Pericoma* species on the presence of one feather-tipped and four spatulate tenacula on each surstylus, as well as the following combination of characters: parameral sheath with shallow U-sha-



**Figure 1.** *Pericoma unipennata* sp. n., holotype male. **A** Head **B** Wing **C** Aedeagus **D** Male genitalia **E** Tenacula. The following abbreviations are used: bp – basiphallus, cr – cercal region, dp lm – distiphallic lateral margin, dp s – distiphallic spatula, ft – feathery tenacula, gcx – gonocoxite, gst – gonostyle, hyp – hypandrium, pm – parameral sclerite, pm j – parameral joint, pms – parameral sheath, s t – spatuliform tenacula.

ped apical indentation 1/6th as deep as width of parameral appendage base, gonostyle with narrow distal part 1/4th as long as broad basal part, distiphallic spatula with sclerotized side margins not converging apically, parameral appendage concave at lateral sides with distal 1/8th protruding over distiphallic spatula.

**Description.** Male (n=1). Head (Fig. 1A) longer than wide; vertex rounded with posterior lobe pointed, about half length of head; eyebridge comprising four rows of facets, separated by five facet diameters; interocular suture broadly V-shaped with median swelling; 7 supraocular setae present, 3 on ventral side and 4 on dorsal side; frontal scar patch inversely T-shaped, reaching level of uppermost facet row of eyebridge; clypeus subrectangular, weakly concave at anterior margin, densely setose, not

protruding in front of level of eyes; palp with 4th segment fleshy, corrugated, length of palpus segments 93 : 106 : 129 : 231; labellum fleshy, micropilose with 11 larger setae; antennae with 14 fusiform flagellomeres; scape cylindrical, pedicel elongate globular to barrell-shaped; ascoids present on flagellomeres 4-8; flagellomere 14 with apiculus about as long as base of segment; length of antennal segments 69 : 63 : 48 : 45 : 45 : 42 : 42 : 42 : 42 : 42 : 36 : 36 : 33 : 27 : 24 : 36.

Thorax without accessory organs; mesonotum and scutellum covered in setae alveoli except on lateral margins; anepisternum and laterotergite covered with setae alveoli; coxae and trochanters with dorsal and ventral stripes of setae alveoli, mid coxa with apicoanterior setose projection; legs without special features;

Wing (Fig. 1B) 2.3 mm long, subovate, membrane without setation or infuscation; radial fork very slightly distad of medial fork, both slightly distad of  $CuA_2$ ; C with two breaks; Sc weakly curved towards C; apical section of  $R_2$  curved anteriorly; wing apex between  $R_4$  and  $R_5$ ; medial fork incomplete; apical section of  $CuA_2$  curved posteriorly; jugum obtuse;

Genitalia (Fig. 1C, D) with hypandrium of even width; aedeagus (Fig. 1D) with basiphallus compressed laterally; distiphallus consisting of two phallomeres forming a spatula with sclerotized lateral margins, lateral margins not converging distally; parameral sheath fused with gonocoxites basally, distally narrowing to nearly half its width at base; distally with narrow median incision; parameral sclerite meeting distal section of basiphallus, jointed to parameral sheath with slightly curved transverse band-like sclerite; parameral sclerite half length of distiphallus, consisting of two fused sclerites which are concave laterally, median suture complete; gonocoxites (Fig. 1C) stout, reniform, parabasal process absent but setose field present near connection with parameral sheath; gonocoxal condyles apparently fused with parameral sheath, gonostyle (Fig. 1C) with basal four 5ths liver-shaped, stout, distal 5th pointed, weakly sinous; epandrium slightly wider than long, expanding distally, number of apertures not discernable in specimen, carrying sparse hairs on ventral surface; surstyli about as long as epandrium, with lateral inner third more weakly sclerotized and densely micropilose; apex of surstylus with five aseriate tenacula of which one apical is feathery at the apex, four subapical tenacula spatulate; hypoproct obtusely isosceles trapezoid, micropilose; epiproct oval, densely pilose.

**Etymology.** *unipennata* = with one feather; refers to the presence of a single feather-tipped and four spatulate tenacula on its surstyli.

**Biology.** The specimen was collected at a spring with many bryophytes growing in a seepage stream. Bryophyte material was collected and extracted, however, no psychodid larvae or pupae were found.

### *Pericoma barbarica* Vaillant, 1955

**New records.** Calobra, 39.85°N, 2.8°E, 9.v.1978, 90m a.s.l., H. Malicky leg. 1 ♂ (RW);

Deiá, town fountain, 39.748072°N, 2.643385°E, 8.ii.2015, G.Kvifte, M. Stokkan & C. Garcia leg. 3♂♂ (ZMUB);

Esporles, Torrent de San Vic, 39.670459°N, 2.569193°E, 11.ii.2015, G.Kvifte leg. 1♂ (ZMFK);

Puigpunyent, 39.616667°N, 2.85°E, 15.v.1978, 200m a.s.l., H. Malicky leg. 12 ♂♂; same but 1.x.1979, 5 ♂♂; same but 6.x.1979, 2 ♂♂ (all RW);

South slope of Piug Major, north of Soller, 39.783333°N, 2.766667°E, 7.–9.v.1978, 700 m a.s.l., H. Malicky leg. 11♂♂ (RW).

**Literature record.** Vaillant (1978).

**Remarks.** This species was described from Algeria and subsequently recorded from Mallorca by Vaillant (1978). It also occurs in mainland Spain and in North Africa (Vaillant 1978).

### *Pericoma trifasciata* (Meigen, 1818)

**Literature record.** Wagner et al. (2002).

**Remarks.** Vaillant (1978) wrote “[*Pericoma*] *trifasciata* ist ebenfalls gemein auf der Mallorca-Insel [...]” (“*trifasciata* is also common on the island of Mallorca”) in the paragraph summarizing the geographical distribution of *Pericoma barbarica* Vaillant, 1955; this is probably a lapsus as Mallorca is not mentioned in the species account for *Pericoma trifasciata*. No other records are available in the literature prior to the listing by Wagner et al. (2002), which was based on Vaillant (1978). In Wagner (2004) and the present paper we deem the records from the Balears to be doubtful and in need of verification through examination of specimens.

## Psychodini

### \* *Philosepedon pyrenaicus* Vaillant, 1974

**First records from Mallorca.** Deiá, town fountain, 39.748072°N, 2.643385°E, 8.ii.2015, G.Kvifte leg. 1♂ (ZMFK);

Esporles, town fountain, 39.669519°N, 2.576953°E, 10.ii.2015, G.Kvifte leg. 1♂ (ZMFK);

Palma de Mallorca, 39.570725°N, 2.641432°E, 7.ii.2015, G.Kvifte leg. 3♂1♀ (2♂ ZMFK, 1♂1♀ ZMUB).

**Remarks.** The males recorded here were identified as *Philosepedon pyrenaicus* according to the key in Vaillant (1974) because of the eyes separated by three facet diameters, the wing “mittwinkel” 105° and the left phallomere of the aedeagus being longer than the right. However, the specimens differ in that the 1st flagellomere is longer than the combined length of the scape and pedicel, and the eyebridge is two facet rows wide at several points. We deem these differences not to be taxonomically meaningful, as the differences in proportions and eyebridge composition could be interpreted as interspecific variation.

***Psychoda (Psychodocha) cinerea* Banks, 1894****Literature record.** Wagner (2004)**New records.** Cala Figuera, 39.332245°N, 3.166440°E, 11.ii.2015, G. Kvifte leg. 2♂♂ (ZMFK);

Deiá, town fountain, 39.748072°N, 2.643385°E, 8.ii.2015, G.Kvifte, M. Stokkan &amp; C. Garcia leg. 1♂ (ZMUB);

Esporles, Torrent de San Vic, 39.670459°N, 2.569193°E, 10.ii.2015, G.Kvifte leg. 1♂ (ZMFK).

***Psychoda (Psychodocha) gemina* (Eaton, 1904)****Literature record.** Wagner (2004)***Psychoda (Psychoda) phalaenoides* (Linnaeus, 1758)****Literature record.** Wagner (2004).**\* *Psychoda (Psychoda) grisescens* Tonnoir, 1922****First records from Mallorca.** Esporles, Torrent de San Vic, 39.670459°N, 2.569193°E, 11.ii.2015, G.Kvifte leg. 1♂.**Remarks.** Recorded from Algeria, Austria, Belgium, Bosnia-Herzegovina, Czech Republic, Denmark, the Faroe Islands, France, Germany, Great Britain, Greece, Hungary, Ireland, Italy, Morocco, the Netherlands, Norway, Slovakia, Slovenia, Sweden, Tunisia and Turkey (Andersen 1999, Ježek 2004, Ježek and Yağci 2005).***Psychoda (Psychodula) minuta* Banks, 1894****Literature record.** Wagner (2004).**New record.** Esporles, Torrent de San Vic, 39.670459°N, 2.569193°E, 10.ii.2015, G.Kvifte leg. 1♂.***Psychoda (Tinearia) alternata* +Say, 1824****Literature record.** Wagner (2004).**New record.** Esporles, town fountain, 39.669519°N, 2.576953°E, 10.ii.2015, G.Kvifte leg. 1♀ (ZMFK).



**Remarks.** *Tinearia* Schellenberg, 1803 was treated as a genus by Wagner (2004) following Ježek (1977). This species group is undoubtedly monophyletic, however based on morphology and some analyses of molecular data it appears to be deeply nested within *Psychoda* s.l. (Vaillant 1990, Espindola et al. 2012, Kvitte and Andersen 2012, Kvitte unpubl.). Inclusion of *Tinearia* as a subgenus within *Psychoda* would thus allow ease of identification of monophyletic units, whereas recognising *Tinearia* as a separate taxon would render *Psychoda* paraphyletic. For a discussion of the monophyly criterion in supraspecific taxonomy, see Komarek and Beutel (2006).

### ***Psychoda (Tinearia) lativentris* Berdén, 1952**

**Literature record.** Wagner (2004)

**Remarks.** For genus taxonomy, see remarks under *Psychoda alternata* above.

### **Discussion**

The material examined in the present study was collected opportunistically and does not reflect the diversity of suitable habitats on Mallorca. Nevertheless, a few preliminary conclusions about the diversity and zoogeographic affinities of the fauna can be made.

Most of the species encountered are widespread European, Holarctic or even cosmopolitan species (*Clogmia albipunctata*, *Paramormia ustulata*, *Psychoda* spp. and arguably *Lepiseodina rothschildi*). Four species appear to have more limited distributions as local West Mediterranean elements, namely *Neoarisemus ibericus*, *Mormia tenebricosa*, *Pericoma barbarica* and *Philosepedon pyrenaicus*. A single species, *Pericoma unipennata* sp. n., has yet to be recorded outside of Mallorca but may have been overlooked elsewhere.

Our records of *Neoarisemus ibericus* and *Philosepedon pyrenaicus* are the first since the original descriptions, which in both cases were based on very few specimens collected in Northern Spain: Montes Universales and the Pyrenees respectively. The records from Mallorca represent a major range extension for both species, suggesting them to be more widespread than previously expected and that they may have been overlooked elsewhere.

Both *M. tenebricosa* and *P. barbarica* have similar distribution patterns; occurring on the north and south coasts of the West Mediterranean. Both species appear widespread in North Africa, having been recorded from Morocco, Algeria and Tunisia (Vaillant 1974, 1978, Wagner 1987). They differ in their European distributions, with the range of *P. barbarica* extending northeast to the Pyrenees and *M. tenebricosa* reaching the Western Alps in southern France and Italy.

Vaillant (1974) mentions consistent minor morphological differences between adult males of *M. tenebricosa* in the North African and European populations, but refrained from using them to delimit species. We consider that these differences warrant further study as they may be evidence of cryptic species or ongoing speciation; DNA sequences will be useful in illuminating this question.

*Pericoma unipennata* sp. n. appears to be a member of the Mediterranean *Pericoma modesta* Tonnoir, 1922 species group as defined by Vaillant (1978, p. 226). In this group, *Pericoma modesta* has a wide Mediterranean distribution whereas *P. albambra* Vaillant, 1978, *P. graecica* Vaillant, 1978 and *P. motasi* Vaillant, 1978 are localized endemics in Southern Spain, Southern Balkan and the Romanian Carpathian mountains, respectively. The Psychodidae of the Iberian peninsula are too incompletely known to tell whether *Pericoma unipennata* sp. n. is an island endemic of the Balearic islands or whether it is more widespread in the Mediterranean region.

## Acknowledgements

We are grateful to Hans Malicky, Lunz am See, for useful material, to Pelayo Secades, Bergen, for valuable help in arranging GMK's field stays in Mallorca, and to Cecilia Enriques García, Bergen, for assistance in the field. Michelle Vine, Gold Coast, kindly checked the English. We thank our colleagues Greg Curler, Sergio Ibañez-Bernal, Jan Ježek and Jukka Salmela for their helpful reviews of the manuscript. We are indebted to the University Library of Kassel's Open Access Publication Fund for covering publication costs.

## References

- Alcover MM, Ballart C, Martín-Sánchez J, Serra T, Castillejo S, Portús M, Gállego M (2014) Factors influencing the presence of sand flies in Majorca (Balearic Islands, Spain) with special reference to *Phlebotomus perniciosus*, vector of *Leishmania infantum*. *Parasites and vectors* 7: 421. doi: 10.1186/1756-3305-7-421
- Andersen T (1999) Moth flies (Diptera, Psychodidae) from the Faroes. *Norwegian Journal of Entomology* 46: 66.
- Boumans L, Zimmer J-Y, Verheggen F (2009) First record of the 'bathroom mothmidge' *Clogmia albipunctata*, a conspicuous element of the Belgian fauna that went unnoticed (Diptera: Psychodidae). *Phegea* 37: 153–160. <http://dare.uva.nl/document/2/71517>
- Duckhouse DA (1987) A revision of Afrotropical *Setomima*, elucidation of their genealogical relationships and descriptions of other Afrotropical Psychodinae (Diptera: Psychodidae). *Annals of the Natal Museum* 28: 231–281. [http://reference.sabinet.co.za/webx/access/journal\\_archive/03040798/404.pdf](http://reference.sabinet.co.za/webx/access/journal_archive/03040798/404.pdf)
- Espindola A, Buerki S, Jacquier A, Ježek J, Alvarez N (2012) Phylogenetic relationships in the subfamily Psychodinae (Diptera, Psychodidae). *Zoologica Scripta* 41: 489–498. doi: 10.1111/j.1463-6409.2012.00544.x
- Ibañez-Bernal S (2008) New records and descriptions of Mexican moth flies (Diptera: Psychodidae, Psychodinae). *Transactions of the American Entomological Society* 134: 87–131. doi: 10.3157/0002-8320(2008)134[87:NRADOM]2.0.CO;2
- Ježek J (1977) Reinstatement of the genus *Tinearia* Schellenberg (Diptera, Psychodidae). *Acta Entomologica Bohemoslovaca* 74: 232–241.

- Ježek J (1984) Nomenclatorial changes of some higher taxa of palaeartic Psychodinae (Diptera, Psychodidae). *Acta Faunistica Entomologica Musei Nationalis Pragae* 17: 155–170.
- Ježek J (1990) Contribution to the taxonomy of some genera of Paramormiine moth flies (Diptera, Psychodidae) with description of a new genus *Karakovounimerus*. *Acta Entomologica Musei Nationalis Pragae* 43: 129–157.
- Ježek J (2004) New faunistic data of non-phlebotomine moth flies (Diptera: Psychodidae) from the Palearctic Region. *Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis, Biologia* 109: 141–151.
- Ježek J, Yağci Ş (2005) Common non-biting moth flies (Insecta, Diptera, Psychodidae) new to the fauna of Turkey. *Acta Parasitologica Turcica* 29: 188–192. [http://www.turkiyeparazitolog.org/sayilar/19/buyuk/pdf\\_TPD\\_351.pdf](http://www.turkiyeparazitolog.org/sayilar/19/buyuk/pdf_TPD_351.pdf)
- Ježek J, Lukáš J, Kvitte GM, Oboňa J (2012) New faunistic records of non-biting moth flies (Diptera: Psychodidae) from the Czech Republic and Slovakia. *Klapalekiana* 48: 121–126.
- Komarek A, Beutel RG (2006) Problems in taxonomy and suggestions for a standardized description of new taxa. *Entomological problems* 36(2): 55–70.
- Kvitte GM (2014) Nomenclature and taxonomy of *Telmatoscopus* Eaton and *Seoda* Enderlein; with a discussion of parameral evolution in Paramormiini and Pericomaini (Diptera: Psychodidae, Psychodinae). *Zootaxa* 3878: 390–400. doi: 10.11646/zootaxa.3878.4.5
- Kvitte GM (2015) Review of the genus *Psychomasina* (Diptera: Psychodidae: Psychodinae), with descriptions of two new species. *Acta Entomologica Musei Nationalis Pragae* 55: 495–503. [http://aemnp.eu/PDF/55\\_2/55\\_2\\_495.pdf](http://aemnp.eu/PDF/55_2/55_2_495.pdf)
- Kvitte GM, Andersen T (2012) Moth flies (Diptera, Psychodidae) from Finnmark, northern Norway. *Norwegian Journal of Entomology* 59: 108–119. <http://www.entomologi.no/journals/nje/2012-2/pdf/nje-vol59-no2-108-119-kvitte.pdf>
- Kvitte GM, Ivković M, Klarić A (2013) New records of moth flies (Diptera: Psychodidae) from Croatia, with the description of *Berdeniella keroveci* sp.nov. *Zootaxa* 3737: 57–67. doi: 10.11646/zootaxa.3737.1.4
- Oboňa J, Ježek J (2012) First records of dendrolimnetic moth flies (Diptera: Psychodidae) from Slovakia. *Klapalekiana* 48: 279–287.
- Quate LW, Brown BV (2004) Revision of Neotropical Setomimini (Diptera: Psychodidae: Psychodinae). *Contributions in Science* 500: 1–117. [http://www.nhm.org/site/sites/default/files/pdf/contrib\\_science/CS500.pdf](http://www.nhm.org/site/sites/default/files/pdf/contrib_science/CS500.pdf)
- Solano-Gallego L, Llull J, Ramos G, Riera C, Arboix M, Alberola J, Ferrer L (2000) The Ibiza hound presents a predominantly cellular immune response against natural *Leishmania* infection. *Veterinary Parasitology* 90: 37–45. doi: 10.1016/S0304-4017(00)00223-5
- Tonnoir AL (1920) Notes sur quelques Psychodidae Africains. *Revue Zoologique Africaine* 8: 127–147. doi: 10.5962/bhl.part.22391
- Vaillant F (1974) Psychodidae–Psychodinae. In: Lindner E (Ed.) *Die Fliegen Der Palearktischen Region. Lieferung 305*. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 109–142.
- Vaillant F (1978) Psychodidae–Psychodinae. In: Lindner E (Ed.) *Die Fliegen der Palearktischen Region. Lieferung 317*. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 207–238.
- Vaillant F (1989) Les Psychodidae dendrolimnophiles et dendrolimnobiontes paléarctiques et néarctiques (Insecta, Diptera, Nematocera, Psychodidae). *Spixiana* 12: 193–208.
- Vaillant F (1990) Propositions pour une révision de la classification des Diptères Psychodidae Psychodinae. *Bulletin de la Société vaudoise des sciences naturelles* 80: 141–163.

- Wagner R (1987) Tunesische Psychodiden (Diptera, Psychodidae). Entomofauna 8: 9–25. [http://www.landesmuseum.at/pdf\\_frei\\_remote/ENT\\_0008\\_0009-0025.pdf](http://www.landesmuseum.at/pdf_frei_remote/ENT_0008_0009-0025.pdf)
- Wagner R (1990) Family Psychodidae. In: Soós Á, Papp L (Eds) Catalogue of Palearctic Diptera: Psychodidae–Chironomidae. Akadémiai Kiadó, Budapest, 11–65.
- Wagner R (2001) Contribution to the knowledge of Spanish Psychodidae (Diptera) with description of two new species. Zoologica Baetica 12: 83–99. [http://wdb.ugr.es/~zool\\_bae/articulos/v12a7.pdf](http://wdb.ugr.es/~zool_bae/articulos/v12a7.pdf)
- Wagner R (2004) Fauna Europaea: Psychodidae. In: Beuk P, Pape T (Eds) Fauna Europaea Web Service version 2.6.2. [www.faunaeur.org](http://www.faunaeur.org)
- Wagner R, Kvifte GM (2015) Description of *Jezekiella patera* gen. et sp. nov. from Europe (Diptera: Psychodidae). Acta Entomologica Musei Nationalis Pragae 55: 505–511. [http://aemnp.eu/PDF/55\\_2/55\\_2\\_505.pdf](http://aemnp.eu/PDF/55_2/55_2_505.pdf)
- Wagner R, Lucientes J, Báez M (2002) Psychodidae. In: Carles-Tolrá Hjort-Andersen M (Ed.) Catálogo de los Diptera de España, Portugal y Andorra (Insecta). Monografías S.E.A., Zaragoza, 66–68. <http://www.sea-entomologia.org/PDF/MSEA08.pdf>