

# Seroprevalence and Associated Risk Factors of Lumpy Skin Disease of Cattle in Selected Districts of Afar Region, Ethiopia

Teshager Dubie , Fentaw Hussen Abegaz, Beyene Dereje, Wossene Negash , Muhammed Hamid 

Department of Veterinary Medicine, College of Veterinary Medicine, Samara University, Semera, Afar, Ethiopia

Correspondence: Teshager Dubie, Tel +251-910344667, Email teshager.dubie@yahoo.com

**Background:** Lumpy skin disease (LSD) is one of Ethiopia's most economically significant transboundary livestock illnesses. The disease has a significant economic impact on pastoral household livestock owners, who rely significantly on their cattle as a source of income.

**Methods:** A cross-sectional study was undertaken in selected districts of Afar region from November 2018 to May 2019 primarily intended to estimate the prevalence of lumpy skin disease serologically in local Afar cattle as well as identify potential associated factors. A multistage sampling method was employed to select study districts, peasant association, herd size and study units. A total of 384 sera were processed using serum neutralization test (SNT) method to detect antibodies against lumpy skin disease virus. Relevant data were refined and further analyzed using stata version 14.

**Results:** In the study districts, the overall animal level seroprevalence was found to be 7.6% (N = 29/384; 95% confidence interval: 4.90–10.20) and the overall herd level prevalence was found to be 20.8% (n = 15/72; 95% confidence interval: 11.42–30.18). Only district was shown to be statistically significant (P = 0.004) in terms of LSD occurrence among the relevant factors studied. Cattle in Chifra district were 20.18 times more likely to contract LSD infection than cattle in Dubti district, when Asayita district was used as the reference group.

**Conclusion:** The present study finding confirmed the presence of the disease in the study districts of afar region and coordinated intervention set to be in place.

**Keywords:** afar, cattle, LSD, associated factors, seroprevalence, serum neutralization test

## Introduction

Ethiopia is the first in Africa with huge livestock population, with an estimated 59.5 million cattle, 31.3 million sheep, 32.74 million goats, and 54.5 million chickens.<sup>1</sup> In Ethiopia, the livestock subsector is a significant contributor to the majority of the population's livelihood as a source of meat, milk, drought power, and income, particularly in pastoral and agro-pastoral areas.<sup>2</sup> Despite having the highest livestock figure in Ethiopia, production and productivity has remained low, owing to livestock diseases and other constraints.<sup>1</sup> In Ethiopia in general and the study areas in particular, livestock diseases are the leading cause of significant economic losses to pastoral livestock owners. Various fragmented study reports show that livestock diseases are widespread across the country, with yearly mortality rates of 8–10% for cattle herds, 15% for sheep and 12% for goat flocks respectively. Furthermore, diseases were estimated to reduce livestock productivity by 50–60% per year.<sup>3,4</sup> Transboundary livestock diseases have continued to cause significant devastating losses in Ethiopia's livestock sector, possibly by imposing trade restrictions, aggravating poverty. Lumpy skin disease (LSD), foot and mouth disease (FMD), and contagious bovine pleuropneumonia (CBPP) are three of the most economically important livestock diseases in developing countries including Ethiopia.<sup>5,6</sup>

Lumpy skin disease (LSD) is one of the most economically important transboundary viral infections affecting particularly of cattle. Lumpy skin disease (LSD) is a highly infectious disease caused by lumpy skin disease virus


(LSDV) that belongs to the *Capripoxvirus* genus of Poxviridae family.<sup>7,8</sup> The genus *Capripoxvirus* is comprised of three genetically closely related important virus species, such as sheep pox virus (SPV), goat pox virus (GPV) and LSDV, affecting sheep, goats and cattle respectively.<sup>9</sup> Arthropod vectors, mechanical means and tick species of the family Ixodidae mainly play an important role in the transmission of LSDV. More importantly, *Stomoxys calcitrans* (Stable fly) has been reported as the most probable vector for LSDV due to its abundance and being associated with outbreaks.<sup>10,11</sup> As a result, LSD episodes mainly occurred during the rainy season, when bug activity is at its peak<sup>12</sup> and the severity of the disease is also depends on virulence nature of the viral strain, susceptibility of the host, immune status of the cattle, animal breed and other related factors.<sup>13</sup> Clinically, the disease is characterized by elevated fever, enlarged lymph nodes and bordered nodules on the skin, emaciation, edema of the skin, nasal discharge, and death.<sup>8,14</sup> Since the disease was first discovered in Zambia in 1929, LSD has gradually and widely expanded over Africa, the Middle East, Southeastern Europe, Central Asia, and more recently South Asia and China. Several countries in Africa, states of the Middle East, and Turkey currently have endemic to cases of the disease. Recently, LSD was introduced into China, Bangladesh and India, beginning from July 2019. In 2020, the disease then spread to other parts of China and India as well as Nepal and Bhutan, indicating the continued and widespread presence of the disease.<sup>15</sup> In case of Ethiopia, the disease was first discovered in 1981 in northwestern regions,<sup>16</sup> and it has since spread to practically all of the country's territory. According to epidemiological studies conducted in various regions of Ethiopia, LSD seroprevalence ranges from 6.4% to 31% at the animal level and up to 64% at the herd level.<sup>17,18</sup> Due to its rapid spread and significant economic losses, the World Organization for Animal Health (OIE) categorized LSD as a notifiable disease under "Cattle diseases and infections".<sup>19-21</sup> Lumpy skin disease virus (LSDV) mostly affects ruminants, imposing financial pressure on pastoral livestock owners whose main source of income is livestock production and rearing.<sup>22</sup> Its substantial economic losses of the disease in livestock industry results from a sharp decline in milk yield and meat production, permanent damage to skin and hide, body weight loss, abortion, retarded growth and infertility. Trade limitations and national and international livestock movement are to blame for the LSD's indirect economic losses.<sup>23-25</sup> Morbidity rates may vary significantly during LSD outbreaks and reach up to 100%, whereas mortality due to LSD varies between 1 and 3%, but up to 40% have been reported in severe outbreak case.<sup>26</sup> Control and prevention of LSDV rely on application of vaccination, restricted animal movement and vector control.<sup>8</sup> Vaccination is reported to be the most effective among the methods used for controlling LSD in both disease endemic and non-endemic areas.<sup>27,28</sup> Currently, the strain KSGP-O180 is being used for vaccine to control the LSD outbreak in Ethiopia.<sup>29</sup> Ethiopia has been striving to control LSD using mass vaccination at a specified season as well as following a report of cases. In the study areas, the annual vaccination coverage seems extremely below the immunity threshold of the population that is necessary to control LSD outbreak occurrence.<sup>30</sup> Moreover, unpublished reports have been coming out from animals' owner and experts about a suspected vaccine failure.<sup>31</sup>

Despite the fact that LSD has a significant economic impact, especially in pastoral areas of afar region, there is lack of studies on LSD disease and associated factors that contribute for occurrence the disease. Therefore, this study was initiated to estimate the seroprevalence of LSD at the animal and herd level as well as its associated factors in the study areas.

## Materials and Methods

### Description of the Study Areas

The study was conducted from November, 2018 to May, 2019 in three districts namely (Asayita, Dubti and Chifra), which are located in the administrative zone one of Afar Region, Ethiopia. The Afar Pastoral Region is located in northeast of Ethiopia between 39°34' to 42°28'E longitude and 8°49' to 14°30'N latitude (Figure 1). The region shares common international boundaries with Eritrea in the northeast and Djibouti in the east and it is characterized by an arid and semi-arid climate with low and erratic rainfall. Rainfall is bi-modal throughout the region, with a mean annual rainfall below 500 mm in the semi-arid western escarpments and decreasing to 150 mm in the arid zones to the east. The region's elevation varies from 120 meters below sea level in the Danakil depression to 1500 meters above sea level elsewhere. Temperatures range from 20°C. at higher elevations to 48°C. at lower levels. Pastoralists, who rely heavily on livestock production for their livelihood, make up the majority of the human population in the afar region. In the afar



**Figure 1** Map of the study areas.

region, there are approximately 1.9 million Afar breed cattle, 90% of which are handled under a pastoral production system and the other 10% in an agro-pastoral production system.<sup>1</sup>

## Study Population

The study populations were non-vaccinated cattle for the last six months and aged above six months. Basically, the neutralizing antibodies developed against LSD vaccine are supposed to be short lived and get cleared before six months. However, cattle vaccinated before six months and young animals aged six months and above months were included in the study for assurance of antibody absence. The age groups of the study animals were categorized as Young ( $\leq 3.5$  years), Adult ( $3.5 \text{ years} < x \leq 5.5 \text{ years}$ ) and Old ( $> 5.5 \text{ years}$ ).<sup>32</sup> In addition, herd size was grouped as small ( $< 40$  cattle), medium ( $40\text{--}75$  cattle) and large ( $> 75$  cattle) according to.<sup>33</sup> Some studies revealed that LSD infection level varied among the study population. Hence, the aim of categorizing the study population in this study was to check whether LSDV infection is affected by herd size or not from highly infectious nature of this virus.

## Study Design and Period

A cross sectional study design was carried out from November 2018 to May 2019 aimed to estimate the seroprevalence of LSD and to assess associated factors selected districts of Afar region.

## Sampling Technique and Sample Size Determination

The sampling method employed in this study was simple random sampling to select the study population since the study districts were purposively selected based on higher study population, access to transportation, history of no vaccination for the last six months, absence of outbreak cases and willingness of pastoralists to participate in this research work. Moreover, the study kebeles within each district were selected purposively. Lastly, the study units were randomly selected in each kebele. As no previous study was found on LSD in cattle in the study areas, 50% expected prevalence, 95% confidence level and 5% absolute precision or marginal error were used respectively. Taking these variables into equation, the total number of animals to be included in the study was determined applying the formula.<sup>34</sup>

$$n = \frac{Z^2 \times P_{exp}(1 - P_{exp})}{d^2}$$

Where n = required sample size; d = desired absolute precision (0.05); Z = Multiplier from normal distribution at 95% Confidence interval (1.96);  $P_{exp}$  = expected prevalence (50%);  $(1 - P_{exp})$  = Probability of having no disease 50% (0.5). Based on the equation, a total of 384 study units were determined to be sampled of all the study districts. A total of 147, 97, and 140 sera were collected from Asayita, Dubti and Chifra, respectively using proportional sample allocation strategy.

## Sample Collection and Transportation

With precaution and gloved hands, approximately 8–10mL blood was drawn from jugular vein of cattle using disposable needles and 10mL non-heparinized vacutainer tube and 21 Gauge needles of 384 animal units. Following sample collection, vacutainer tubes were labeled, packed and transported to Samara University, college of veterinary medicine (SU-CVM) laboratory and kept overnight at room temperature to allow the blood to clot at a slant position. Then, the serum samples were transferred into identically labeled sterile cryovials and stored at  $-20^{\circ}\text{C}$  refrigerator prior to laboratory analysis. Finally, the sera were transported to National Veterinary Institute (NVI), Bishoftu and the samples were processed using serum neutralization test (SNT) for detection of LSD antibody presence in the blood of the sampled cattle.

## Administration of Questionnaire Survey

Open and closed ended questionnaires were prepared and administered to herd owners to assess potential risk factors of the disease alongside with sample collection. Respondents from each district were randomly selected and interviewed to assess associated risk factors of the disease. The questionnaires were interpreted into Afaraf language. Herd owner having cattle were the sampling units for questionnaire survey. Using herd owner as study unit, a total of 72 herd owners were selected proportionally allocating 27 to Asayita, 20 to Dubti and 25 to Chifra. All needed epidemiological data were collected, tabulated, coded and analyzed using stata software version 14 (Stata Corp, College Station, TX, USA).

## Serological Analysis

### Serum Neutralization Test

The serum neutralization test (SNT) protocol was employed by OIE.<sup>8</sup> The SNT was performed using a constant-virus /varying-serum method as described by.<sup>35</sup> Briefly, each serum was tested in duplicate wells at serial dilutions of 1/5, 1/25, 1/125, 1/625 and 1/ 3125. In 96 well flat-bottomed tissue culture microtiter plates, Kenyan sheep pox virus (KS1) was employed at 100 TCID<sub>50</sub> per well. To acquire consistent results, the Vero cell was utilized as the culture host for the assay.<sup>36</sup> The plates were incubated at 37 °C in an atmosphere containing 5% carbon dioxide for 9 days. The presence of cytopathic effect (CPE) was examined using an inverted phase-contrast microscope at day 4 and final reading was made at day 9. In this study, both positive and negative control samples were used during this laboratory work. Kenyan sheep pox (KS1 O-180) Capripox virus strain was used as positive control. The serum samples were considered LSD positive, when CPE was inhibited either in both or in one of the duplicate wells at 1/25 or higher dilutions.

## Data Management and Analysis

Both the laboratory analytical output and the necessary questionnaire response data were recorded, coded, and filtered in Microsoft Excel before being uploaded to Stata 14.0 (Stata Corp, College Station, TX, USA) for data analysis. To determine frequencies, diagrams, and tables, descriptive statistics were used. Association between risk factors and the disease positivity was assessed using Chi-square ( $\chi^2$ ). Bivariate logistic regression was computed to estimate the magnitude association between risk factors and the disease. In all the analyses, confidence level at 95% was calculated and the  $P < 0.05$  was used for statistical significance level between the potential confounders and the outcome variable.

### Limitation of the Study

Limitation of the present study supposed to be sampling bias in selecting study populations and false positive or negative result by SNT during laboratory work.

## Results

### Animal Level Seroprevalence

Out of 384 serum samples tested using SNT, the overall seroprevalence at animal level was found to be 7.6% (95% CI= 4.90–10.20;  $n=29/384$ ) and herd level seroprevalence was found to be 20.8% (95% CI= 11.42–30.18;  $n=15/72$ ) seropositive for the presence of antibodies against LSDV infection. Descriptive computations are summarized (Table 1). The majority of the animals studied,  $n = 312$  (81.25%), were female animals, while the rest,  $n = 72$  (18.75%), were male animals. In comparison to Dubti and Asayita districts, Chifra had more seropositive animals ( $N = 140$ , or 12.14%) than the other two districts.

### Herd Level Seroprevalence

In an attempt to assess the attributes of herd over the disease occurrence, a total of 72 herds were examined over the three study districts of which only 15 herds (20.8%;  $n=15/72$ ; 95% CI= 11.42–30.18) have been confirmed at least one positive cattle by SNT test for antibodies against LSDV.

### Associated Factors of Lumpy Skin Disease

In the current study, four factors were considered presumably supposed to have influence the infection and establishment of LSD disease among the study animals applying Chi-square ( $\chi^2$ ) statistics. As a result, only one study district ( $P = 0.004$ ) was

**Table 1** Descriptive Statistics Result Showing Variables with Animal Level Seroprevalence

| Variable  | Category | Frequency (No) | Positive | Prevalence in % (95% CI) |
|-----------|----------|----------------|----------|--------------------------|
| Sex       | Female   | 312            | 25       | 8.00 (5.25–11.60)        |
|           | Male     | 72             | 4        | 5.556 (1.53–13.62)       |
| Age       | Young    | 77             | 4        | 5.19 (0.29–10.90)        |
|           | Adult    | 210            | 21       | 10 (5.90–14.10)          |
|           | Old      | 97             | 4        | 4.12 (0.16–8.10)         |
| Herd Size | Small    | 124            | 8        | 6.45 (2.15–10.77)        |
|           | Medium   | 142            | 10       | 7.04 (2.84–11.25)        |
|           | Large    | 118            | 11       | 9.32 (4.08–14.56)        |
| District  | Asayita  | 147            | 1        | 0.68 (–0.65–2.00)        |
|           | Dubti    | 97             | 11       | 11.34 (5.15–17.56)       |
|           | Chifra   | 140            | 17       | 12.14 (6.74–17.69)       |

**Table 2** Logistic Regression Analysis of Risk Factors for LSD Infection

| Variable  | Category | Animals Sampled | Positive | $\chi^2$ | P-value |
|-----------|----------|-----------------|----------|----------|---------|
| Age       | Young    | 77              | 4        | 4.048    | 0.132   |
|           | Adult    | 210             | 21       |          |         |
|           | Old      | 97              | 4        |          |         |
| Sex       | Female   | 312             | 25       | 0.506    | 0.477   |
|           | Male     | 72              | 4        |          |         |
| Herd size | Small    | 124             | 8        | 0.797    | 0.671   |
|           | Medium   | 142             | 10       |          |         |
|           | Large    | 118             | 11       |          |         |
| District  | Asayita  | 147             | 1        | 16.162   | 0.000   |
|           | Dubti    | 97              | 11       |          |         |
|           | Chifra   | 140             | 17       |          |         |

**Table 3** Bivariate Logistic Regression Output by Districts

| District | No of Animals Sampled | Positive | COR   | P-value | 95% CI        |
|----------|-----------------------|----------|-------|---------|---------------|
| Asayita  | 147                   | 1        | -     | -       | -             |
| Dubti    | 97                    | 11       | 18.67 | 0.005   | 2.369–147.162 |
| Chifra   | 140                   | 17       | 20.18 | 0.004   | 2.648–153.798 |
| Total    | 384                   | 29       |       |         |               |

found to be correlated to LSD occurrence. Seropositivity variations were observed among age groups; however, they were statistically insignificant. Adult cattle showed higher seropositivity than young and old animals, however, the difference was statistically insignificant (Table 2).

Bivariate logistic regression analysis was computed using 95% CI and  $P < 0.05$  to assess the strength of association between the disease and risk factors (Crude Odds Ratio=COR) with the occurrence of disease as depicted (Table 3). In comparison to study animals identified in Asayita district, study animals found in Chifra district were 20.18 times more likely to be at risk of acquiring LSD.

## Discussion

Lumpy skin disease seroprevalence reports of have been reported with various serological results in different agro-ecological parts. There was no even a single scientific study finding on the seroprevalence of LSD and its associated factors in the study districts of afar region except unpublished reports of woreda animal health bureau. That is why the current study is the first of its kind to estimate the seroprevalence of LSD in cattle using serum neutralization test (SNT) and assesses associated risk factors in afar region, Ethiopia. In the present study animal level seroprevalence of LSD infection was found to be 7.6% which is comparable with previous reports of 6.43% at West Wollega Zone of Oromia region,<sup>37</sup> 7.4% in north-eastern Ethiopia<sup>38</sup> and 7% at around Nekemt.<sup>39</sup> In contrast to this study, recent studies have found that the Southern region has a higher LSD seroprevalence of 11.6%,<sup>40</sup> 27.9% at Woliso town<sup>41</sup> and 28% in three districts in Amhara's eastern region.<sup>42</sup> This seropositivity variation could be due to differences in individual animal breed, immune status, interaction of cattle with other animals, production system, variation in geographical area,

sampling period and testing methods employed for the studies, sample size, seasons, agro-ecological conditions and introduction of new animals without screening which could affect the incidence of LSD.<sup>42,43</sup>

In this study the herd level seroprevalence was compared with other previous studies and showed lower finding than previous herd level prevalence in lowland (50%), midland (26%) and highland agro-climate zones (64%) in Ethiopia.<sup>17</sup> Seroprevalence at the herd level was reported to be 52.6% in central and northwestern Ethiopia, and 5.95% in western Ethiopia<sup>44</sup> and 44% in eastern part of Ethiopia.<sup>45</sup> This herd level LSD seroprevalence variation could be due to vectors population availability in the study areas, density of livestock at the grazing and watering points, husbandry methods, rainy seasons, geographical situations and introducing new animals from outside without screening that would be affect the occurrence of LSD.<sup>46</sup> Among the associated factors assessed, only the study area was found to be statistically significant with regard to LSD serostatus. LSD occurrence and age groups were found to be statistically insignificant. Adult and old cattle, on the other hand, have a higher risk of developing the disease than young animals. The results of this seroprevalence study are consistent with previous findings.<sup>44</sup> This difference could be due to fatigue from lactation and intense effort for milk and draft animals as well as stress.<sup>42,44,46</sup> The lower prevalence of LSD in young animals, on the other hand, could be associated to traditional young animal care practices that separate young animals from the herd, possibly reducing vector exposure.<sup>38,47</sup>

Furthermore, this study found no significant association between sex and LSD serostatus, which is consistent with the previous study results, despite females having a relatively high seroprevalence.<sup>44</sup> In contrast, statistically significant association between sex and serostatus of LSD was reported by previous seroprevalence studies.<sup>39,46</sup> These seroprevalence results variations might be due to female animals are usually kept longer by farmers while males are sold off at a younger age and thus the effect of sex may be an artifact of duration of exposure.<sup>18</sup> This study found a significantly variation in LSD seroprevalence across the study districts, which is consistent with previous seroprevalence reports.<sup>37,44,45</sup> This seroprevalence discrepancy could be because of differences in herd management system, the presence of flooding and irrigation in the study areas that may facilitate multiplication of potential mechanical vectors to enhance disease transmission, herd sizes, sharing common boundary with neighboring states, and introduction of new animals without screening could contribute for occurrence of LSDV.<sup>17,40,45</sup> In addition, in the present study, even though no statistical association between herd size and LSD serostatus was observed, it has been found a relative high seroprevalence of LSD in medium and large herd size as compared to small herd size. Previous LSD serostatus studies back up this assertion.<sup>40</sup> Furthermore, cattle with larger herd sizes were found to be more affected than cattle with smaller herd sizes. Stressful conditions, frequency of transmission, intensity of LSDV exposure and other related factors could all influence LSD serostatus variation in herd size.<sup>41</sup>

## Conclusions

The current study finding indicated that the overall LSD seroprevalence of cattle was 7.6% at individual animal and 20.8% at herd level. Hence, the current study revealed that LSD was prevalent in the study areas. Unlike other risk factors, the study areas were found to be associated factors for occurrence of the disease. Therefore, further studies should be conducted to estimate its region wise seroprevalence and assess the economic impact of the disease. From transboundary nature and economic impact point of view, the disease requires technically sound and coordinated efforts for its prevention and control. Therefore, the study findings strongly urge a prompted and coordinated interventional to be set in place.

## Data Sharing Statement

The data sets used and/or analyzed during the current study could be available from the corresponding author on reasonable request.

## Ethics Approval and Consent to Participate

Written ethical approval and written informed consent for this study was obtained from Samara University, College of Veterinary Medicine of Animal Research Ethics and Review committee (reference AREC019/2019). Written informed consent was also obtained from the herd owners to take samples from their cattle and for further research use of the

samples. The reason for this written informed consent is that, participants were required for interview and the individual participants were not subjected to any harm as much as their privacy is kept confidential. Confidentiality of collected data and the scientific honesty during write up was considered. These written informed consents were documented.

## Acknowledgments

The authors would like to express their appreciation to Samara University's Research and Community Service office for funding this study. We would like to express our sincere gratitude to the National Veterinary Institute of Ethiopia for offering laboratory facilities and space for this research, as well as to livestock owners and district animal health staff for their cooperation and technical support during field sample collection.

## Funding

This research work was financially supported by Samara University, Research and Community Service vice president office. However, the funder had no role in the design and execution of this study, sample collection, data analysis and interpretation and manuscript writing.

## Disclosure

The authors declare that they have no competing interests in the publication of this paper.

## References

1. CSA FDRoE. Central Statistical Agency: agricultural sample survey 2018, CSA, Addis Ababa, Ethiopia; 2017.
2. Ayelet G, Soressa M, Sisay T, et al. FMD virus isolates: the candidate strains for polyvalent vaccine development in Ethiopia. *Acta Trop*. 2013;126(3):244–248. doi:10.1016/j.actatropica.2013.02.005
3. FAO: Food and Agriculture Organization of the United Nations. Livestock sector brief: Ethiopia. Livestock information, sector analysis and policy branch. Rome: FAO; 2004.
4. Ganeshkumar B. Economic impact of foot-and-mouth disease in India, scientific developments and technical challenges in the progressive control of foot-and-mouth disease in South Asia, New Delhi, India. *Glob Res Alli*. 2012;13:5.
5. AU-IBAR. African union-inter-African bureau for animal resources, Pan African; 2011.
6. OIE. *Foot-and-Mouth Disease, Manual of Standard for Diagnostic Tests and Vaccine for Terrestrial Animals*. 6th ed. Paris: OIE; 2009:156–212.
7. Sprygin A, Pestova Y, Bjadovskaya O, et al. Evidence of recombination of vaccine strains of lumpy skin disease virus with field strains, causing disease. *PLoS One*. 2020;15(5):1–8. doi:10.1371/journal.pone.0232584
8. OIE. Manual of diagnostic tests and vaccines for terrestrial animals. *Lumpy Skin Dis*. 2010;2(4):768–778.
9. Babiuk S, Bowden T, Boyle D, Wallace D, Kitching RP. Capripoxviruses: an emerging worldwide threat to sheep, goats and cattle. *Transbound Emerg Dis*. 2008;55(7):263–272. doi:10.1111/j.1865-1682.2008.01043.x
10. Tuppurainen E, Alexandrov T, Beltrán-Alcruado D. Lumpy skin disease field manual – a manual for veterinarians. In: *FAO Animal Production and Health Manual No. 20 (P. 60)*. Food and Agriculture Organization of the United Nations(FAO); 2017.
11. Sprygin A, Pestova Y, Wallace DB, Tuppurainen E, Kononov AV. Transmission of lumpy skin disease virus. *Virus Res*. 2019;269:197637. doi:10.1016/j.virusres.2019.05.015
12. Ding YZ, Chen J, Zhang JH, Zhou LN, Ma L, Zhang Y. An overview of control strategy and diagnostic technology for foot-and-mouth disease in China. *Viro J*. 2013;10:78. doi:10.1186/1743-422X-10-78
13. CFSPH. The center for food security and public health, Iowa State University, College of veterinary medicine and institution of international cooperation in animal biologics an OIE Collaborating Center; 2008:1–4. Available from: <http://www.cfsph.iastate.edu/Factsheets/pdfs/lumpyskindisease>. Accessed August 12, 2022.
14. Radostits OM, Gay CC, Hinchcliff KW, Constable P. *Veterinary Medicine: A Textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses*. Saunders Elsevier; 2007.
15. FAO. *EMPRES-i Global Animal Disease Information System. LSD Disease Events Reported by Iran, Iraq, Israel, Jordan, Kuwait, Lebanon, Saudi Arabia, Turkey, and the West Bank (2012–2016)*. Rome: FAO Animal Production and Health Division; 2020. Available from <http://empres-i.fao.org/eipws3g/>. Accessed August 12, 2022.
16. Mebratu GY, Kassa B, Fikre Y, Berhanu B. Observation on the outbreak of lumpy skin disease in Ethiopia. *Rev Elev Méd Vét Pays Trop*. 1984;37:395–399.
17. Gari G, Grosbois V, Waret-Szkuta A, Babiuk S, Jacquet P, Roger F. Lumpy skin disease in Ethiopia: seroprevalence study across different agroclimate zones. *Acta Trop*. 2012;123:101–106. doi:10.1016/j.actatropica.2012.04.009
18. Gari G, Waret-Szkuta A, Grosbois V, Jacquet P, Roger F. Risk factors associated with observed clinical lumpy skin disease in Ethiopia. *Epidemiol Infect*. 2010;138:1657–1666. doi:10.1017/S0950268810000506
19. Tuppurainen E, Stoltz W, Troskie M, Wallace D, Oura A, Mellor P. A potential role for ixodid (Hard) tick vectors in the transmission of lumpy skin disease virus in cattle. *Transbound. Emerg Dis*. 2010;58:90–104.
20. Tuppurainen E, Klaas D, Janika W, Hannes B. Vaccines and Vaccination against Lumpy Skin Disease. *Vaccines*. 2022;9:1136. doi:10.3390/vaccines9101136
21. OIE. Lumpy skin disease. In: *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*. Paris: OIE; 2019.



22. Buller RM, Arif BM, Black D, Dumbell KR, Esposito J. *Virus Taxonomy: Classification and Nomenclature of Viruses. Eighth Report of the International Committee on Taxonomy of Viruses*. San Diego: Elsevier Academic Press; 2005:117–133.
23. Yacob H, Nesanet B, Dinka A. Prevalence of major skin diseases in cattle, sheep and goats at Adama Veterinary Clinic, Oromia regional state, Ethiopia. *Revue Méd Vet*. 2008;159(8–9):455–461.
24. Merk Veterinary Manual. Integumentary system: pox diseases: lumpy skin disease, economic impact of lumpy skin disease; 2011.
25. Tuppurainen E, Alexandrov T, Beltrán-Alcruo D. Lumpy skin disease field manual – a manual for veterinarians. FAO Animal Production and Health Manual No. 20 (pp. 1–60). Rome: Food and Agriculture Organization of the United Nations (FAO); 2017.
26. Coetzer J. Lumpy skin disease. In: Coetzer JAW, Tustin RC, editors. *Infectious Diseases of Livestock*. Cape Town: Oxford University Press Southern Africa; 2004:1268–1276.
27. EFSA: European Food Safety Authority. Lumpy skin disease: vaccination is most effective control method; 2019. Available from: <https://www.efsa.europa.eu/en/press/news/160809>. Accessed August 12, 2022.
28. OIE. Manual of diagnostic tests and vaccines for terrestrial manual. In: *Version Adopted by the World Assembly of Delegates of the OIE*. 7th ed. Vol. I. Paris, France: OIE; 2012:1401–1405.
29. Klement E, Broglia A, Antoniou S, Tsiamadis V, Plevraki E, Petrović T. Neethling vaccine proved highly effective in controlling lumpy skin disease epidemics in the Balkans. *Prev Vet Med*. 2018;181:104595.
30. Mo A. Livestock disease outbreak database. Addis Ababa, Ethiopia; 2016.
31. Afar Pastoral, Agricultural, Development B. Baseline survey made on the potential, constraints, and opportunity on the production system of 29 districts of Afar National Regional State; 2006.
32. Bertram MR, Delgado A, Pauszek SJ, et al. Effect of vaccination on cattle subclinically infected with foot-and-mouth disease virus in Cameroon. *Prev Vet Med*. 2018;155:1–10. doi:10.1016/j.prevetmed.2018.04.003
33. Zemedu L. Contribution of livestock sector in Ethiopian economy. *Rev Adv Life Sci Technol*. 2015;29:1–14.
34. Thrusfield MVM. *Veterinary Epidemiology*. 3rd ed. Singapore: Blackwell Science; 2007:233.
35. Gari G, Biteau-Corolle F, LeGoff C, Caufour P, Roger F. Evaluation of indirect fluorescent antibody test (IFAT) for the diagnosis and screening of lumpy skin disease using Bayesian method. *Vet Microbiol*. 2008;129(3–4):269–280. doi:10.1016/j.vetmic.2007.12.005
36. OIE. Manual of diagnostic tests and vaccines for terrestrial animals, Chapter 2. 4.14. In: *Lumpy Skin Disease*. Paris: OIE; 2017.
37. Gari G, Biteau-Corolle F, LeGoff C, Caufour P, Roger F. Evaluation of indirect fluorescent antibody test (IFAT) for the diagnosis and screening of lumpy skin disease using Bayesian method. *Vet Microbiol*. 2008;128:269–280.
38. Paulos A, Eshetu Y, Bethelhem N, Abebe B, Badeg Z. A study on the prevalence of animal rabies in Addis Ababa during 1999–2002. *Ethiop Vet J*. 2003;7:69–77.
39. Sylvester O, Kimberly V, Anna M, et al. Seroprevalence and risk factors for lumpy skin disease virus seropositivity in cattle in Uganda. *BMC Vet Res*. 2019;15:236. doi:10.1186/s12917-019-1983-9
40. Bossche P, Coetzer J. Climate change and animal health in Africa. *Rev Sci Tech Int Epiz*. 2008;27(2):551–562.
41. Zelalem A, Hailu D, Getachew G, Menbere K. Sero-prevalence of lumpy skin disease in selected districts of West Wollega zone, Ethiopia. *BMC Vet Res*. 2015;11:7–9. doi:10.1186/s12917-015-0317-9
42. Hunter P, Wallace D. Lumpy skin disease in Southern Africa: a review of the disease and aspects of control. *J S Afr Vet Assoc*. 2001;72(2):68–71. doi:10.4102/jsava.v72i2.619
43. Gezahegne M, Fekadu A, Yalelet W, et al. Bovine tuberculosis and its associated risk factors in pastoral and agro-pastoral cattle herds of Afar Region, Northeast Ethiopia. *J Vet Med Anim Health*. 2013;5(6):171–179.
44. Abera Z, Degefu H, Gari G, Kidane M. Sero-prevalence of lumpy skin disease in selected districts of west Wollega zone. *BMC Vet Res*. 2015;11:1–9. doi:10.1186/s12917-015-0432-7
45. Hailu B, Tolosa T, Gari G, Teklue T, Beyene B. Estimated prevalence and risk factors associated with clinical lumpy skin disease in North-Eastern Ethiopia. *Prev Vet Med*. 2014;115(2):64–68. doi:10.1016/j.prevetmed.2014.03.013
46. Tuppurainen ES, Oura C. Review: lumpy skin disease: an emerging threat to Europe, the Middle East and Asia. *Transbound Emerg Dis*. 2012;59:40–48. doi:10.1111/j.1865-1682.2011.01242.x
47. Gari G, Bonnet P, Roger F, Waret-Szkuta A. Epidemiological aspects and financial impact of lumpy skin disease in Ethiopia. *Prev Vet Med*. 2011;102:274–283. doi:10.1016/j.prevetmed.2011.07.003

## Veterinary Medicine: Research and Reports

Dovepress

### Publish your work in this journal

Veterinary Medicine: Research and Reports is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of veterinary medicine. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/veterinary-medicine-research-and-reports-journal>