

BMJ Open Living through the psychological consequences of COVID-19 pandemic: a systematic review of effective mitigating interventions

Angkana Lekagul ¹, Peeraya Piancharoen,¹ Anamika Chattong,¹ Chawisa Suradom,² Viroj Tangcharoensathien¹

To cite: Lekagul A, Piancharoen P, Chattong A, *et al*. Living through the psychological consequences of COVID-19 pandemic: a systematic review of effective mitigating interventions. *BMJ Open* 2022;**12**:e060804. doi:10.1136/bmjopen-2022-060804

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-060804>).

Received 11 January 2022

Accepted 14 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹International Health Policy Program, Amphur Muang, Nonthaburi, Thailand

²Psychiatry, Chiang Mai University Faculty of Medicine, Chiang Mai, Thailand

Correspondence to

Dr Angkana Lekagul; angkana@ihpp.thaigov.net

ABSTRACT

Objective This review assesses interventions and their effectiveness in mitigating psychological consequences from pandemic.

Method Published English literatures were searched from four databases (Medline, PubMed, Embase and PsycINFO) from January 2020 and September 2021. A total of 27 papers with 29 studies (one paper reported three studies) met inclusion criteria. Cochrane risk-of-bias tool is applied to assess the quality of all randomised controlled trials (RCT).

Results All studies were recently conducted in 2020. Publications were from high-income (13, 44.8%), upper middle-income (12, 41.4%) and lower middle-income countries (3, 10.3%) and global (1, 3.5%). Half of the studies conducted for general population (51.7%). One-third of studies (8, 27.6%) provided interventions to patients with COVID-19 and 20.7% to healthcare workers. Of the 29 studies, 14 (48.3%) were RCT. All RCTs were assessed for risk of biases; five studies (15, 35.7%) had low risk as measured against all six dimensions reflecting high-quality study.

Of these 29 studies, 26 diagnostic or screening measures were applied; 8 (30.9%) for anxiety, 7 (26.9%) for depression, 5 (19.2%) for stress, 5 (19.2%) for insomnia and 1 (3.8%) for suicide. Measures used to assess the baseline and outcomes of interventions were standardised and widely applied by other studies with high level of reliability and validity. Of 11 RCT studies, 10 (90.9%) showed that anxiety interventions significantly lowered anxiety in intervention groups. Five of the six RCT studies (83.3%) had significantly reduced the level of depression. Most interventions for anxiety and stress were mindfulness and meditation based.

Conclusions Results from RCT studies (11%, 78.6%) were effective in mitigating psychological consequences from COVID-19 pandemic when applied to healthcare workers, patients with COVID-19 and general population. These effective interventions can be applied and scaled up in other country settings through adaptation of modes of delivery suitable to country resources, pandemic and health system context.

INTRODUCTION

COVID-19 was first reported in December 2019.¹ By March 2020, the WHO declared the COVID-19 a pandemic and as of November 2021, there have been over 259 million

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Effective interventions which targeted patients, healthcare workers and general population can be applied and scaled up by other countries.
- ⇒ The heterogeneity of included studies such as the mode of delivery of interventions, participant characteristics and measurement tools prevents efforts to conduct meta-analysis.
- ⇒ There is a lack of assessment of the effects of cointerventions, provided to the participants in the intervention and control groups, which influence the outcomes in both groups.
- ⇒ Measurement of outcome at short duration after interventions does not offer understanding of long-term outcome or its sustainability.

cumulative confirmed cases and 5 million deaths from COVID-19 worldwide.² The rapid global spread of disease has had physical health impacts and psychological consequences on the population.

The COVID-19 outbreak has had direct and indirect impacts on mental health. Many experienced fear and anxiety due to uncertainty about the pandemic's evolution, feelings of hopelessness, despair and grief in the face of uncontrollable events.^{3 4} The impact of public health measures such as quarantine measures, social restrictions and isolation has also resulted in mental health consequences including anxiety, depression and feelings of neglect; anxiety was highly prevalent in people undergoing quarantine or social isolation.⁵⁻⁷ The pandemic's indirect effects, such as unemployment due to economic downturn and the death of family members from disease, have led to depression and suicide.⁸⁻¹⁰ Decreased population mobility and high rates of COVID-19 infection were found to be significantly related to an increased prevalence of major depression disorder, which compared with prepandemic,

there were additional 53.2 million cases of major depressive disorder and 76.2 million cases of anxiety disorders in 2020.¹¹

Psychological consequences from the pandemic also developed more significantly in certain groups of people. Long hours of COVID-19 clinical services caused significant pandemic-related burnout for frontline healthcare workers worldwide.^{12–14} Fear of transmitting the infection to family members and increased demand for health services also caused stress and sleep problems among medical professionals.^{15–17}

Multiple mental health interventions have been implemented during previous epidemics such as Ebola and Middle East respiratory syndrome, particularly for healthcare workers experiencing higher levels of burnout and psychological consequences than others.¹⁸ Effective interventions like psychosocial and administrative support in workplaces have been shown to reduce the rates of emotional exhaustion in healthcare workers.¹⁸ Interventions in other population groups during Ebola outbreaks reduced depression, stress and anxiety.^{19–21} Implementing interventions to alleviate mental health problems during the COVID-19 pandemic has proven to be difficult due to various social restrictions; face-to-face interventions seem not feasible where online interventions including telehealth and multimedia application were used as alternatives.^{22–24}

Though various interventions to mitigate psychosocial consequences from the COVID-19 pandemic have been launched, the measures used, their delivery channels and effectiveness on population groups are not well understood. This systematic review assesses interventions and measures used, and the effectiveness of interventions in mitigating psychological consequences on different populations so that effective interventions can be upscaled accordingly. The psychological consequences affect patients who suffered from the illnesses and stress from quarantine and isolation, healthcare workers who suffered from stressful long hours of work and general population facing indirect effects of government measures such as lockdown and limited social activities.

We propose three review questions. First, what interventions are used to mitigate psychological consequences in the population and in healthcare workers? Second, what measures are used for assessing intervention outcome? Third, what is the effectiveness of different interventions? Findings from this review can inform health professionals and governments of effective interventions to prevent or mitigate mental health problems from the COVID-19 pandemic.

METHODS

Search strategy

We used PICO framework, a common tool used to develop literature search strategies, to ensure the comprehensive searches recommended by the Cochrane Collaboration.²⁵ It is used in evidence-based practice to formulate

healthcare-related question. PICO elements include Population, Interventions, Comparisons and Outcomes. In this review, population are individuals in countries or areas affected by COVID-19; interventions are exposure to COVID-19 pandemic and the clinical or non-clinical interventions which mitigate the psychological impact from the pandemic; comparison of outcomes between case and control groups or before and after interventions; and outcome are mental consequences including suicide, depression, anxiety, stress and insomnia/reduced sleep quality. See the full search strategy in online supplemental annex 1.

Published literatures were electronically retrieved from four databases including Medline, PubMed, Embase and PsycINFO. The eligibility criteria are English-language studies between January 2020 and September 2021 where interventions were provided with measurable outcomes. [Table 1](#) provides inclusion and exclusion criteria.

Selection of publications

The search terms were identified based on search strategy and applied to search in title, abstract, keyword and full text. After searching, the duplicated studies or data were removed then the abstracts were reviewed; those were excluded for the following reasons with agreement in inclusion and exclusion criteria.

[Figure 1](#) shows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow of this process.

In the process of abstract screening, three researchers (AL, PP and AC) independently reviewed the abstracts. Consensus was reached if two researchers agreed based on eligibility criteria. In case of disagreement, the third researcher reviewed and decided.

Assessment of the risk of bias

The Cochrane risk-of-bias tool²⁵ was applied to assess the methodological quality of 14 randomised controlled trials (RCT) included by this systematic review. There are six domains of risk of bias: (1) random sequence generation (selection bias), (2) allocation concealment (selection bias), (3) blinding of participants, personnel and outcome assessors, (4) incomplete outcome data (attrition bias), (5) selective reporting (reporting bias), and (6) other bias.²⁵ Three authors (AL, PP and AC) separately evaluated each domain and categorised the risk of bias into 'low risk', 'high risk' or 'unclear risk'. The results of assessment were shown in terms of the number of 'low risk' of bias, which is 6 in total. Different criteria were used for making risk of bias judgements for each domain. Insufficient information reported in studies resulted in 'unclear risk' of bias judgement. When there were conflicting views, discussion among the three authors was convened to reach consensus.

Data extraction and synthesis

Relevant contents in all included articles were extracted and synthesised into three variables in an Excel spreadsheet: (a) characteristics of study: author, year of

Table 1 Search terms, inclusion and exclusion criteria

	Inclusion and exclusion criteria
Population	<p>Inclusion: Individuals in countries or areas affected by COVID-19 including patients, healthcare workers, women and minority groups.</p> <p>Exclusion:</p> <ul style="list-style-type: none"> ▶ Individuals in countries or areas not affected by COVID-19. ▶ Individuals having prior psychological symptoms.
Intervention/exposure	<p>Inclusion:</p> <ul style="list-style-type: none"> ▶ Exposure to COVID-19 pandemic. ▶ Both clinical and non-clinical interventions, provided by therapists, psychotherapists/counsellors, psychologists and qualified mental health staff including music and physical activity. <p>Exclusion: Not any.</p>
Comparison	<p>Inclusion:</p> <ul style="list-style-type: none"> ▶ Comparing prevalence or degree of psychological consequences before and after interventions in the same group. ▶ Comparing prevalence or degree of psychological consequences in control and intervention groups. <p>Exclusion: Not any.</p>
Outcome	<p>Inclusion: Mental health consequences including suicide, depression, anxiety, stress and insomnia/reduced sleep quality. Outcomes must be quantified as proportions/prevalence rate or defined categorically.</p> <p>Exclusion: Not any.</p>
Study design	<p>Inclusion:</p> <ul style="list-style-type: none"> ▶ Randomised controlled trials (RCTs). ▶ Controlled before-and-after studies (CBAS), with baseline and postintervention measurement for both groups. ▶ Experimental research designs. <p>Exclusion:</p> <ul style="list-style-type: none"> ▶ Type of publication: editorials, commentary, letters to the editor, reviews. ▶ Studies that did not have the outcome measures specified in the protocol.

publication, objective, country, study design; (b) interventions: provider, participant; and (c) outcomes: different types of psychological consequence, and measures used for assessment of psychological consequence.

Patient and public involvement

No patients were involved in this study.

RESULTS

Our search strategies identified a total of 5623 records, of which 5619 were from the four databases and four from hand-searching from references of key articles. After abstract screening, 705 duplicated records were removed while 4602 records were not relevant. The full texts of the remaining 316 records were retrieved and reviewed for eligibility. We found that 289 publications were not relevant to the review questions. Finally, 27 articles comprising 29 studies were included for systematic review.

Study characteristics

Characteristics of 27 included articles are described in table 2. One paper reported three studies,²⁶ totalling 29 studies, all of which were conducted in 2020. Of the 29 studies, one was conducted globally (3.5%)²⁷ and the rest were categorised according to WHO geographical area. Three studies (10.3%) were conducted in the Eastern

Mediterranean region,^{28–30} 10 (34.5%) in the European region,^{31–40} 4 (13.8%) in the Americas^{41–44} and 11 (37.9%) in the Western Pacific.^{26 45–52} Applying the World Bank's income group categories, 13 (44.8%) studies were in high-income group,^{31–33 35–41 43–45} 12 (41.4%) in upper middle-income group^{26 34 42 46–52} and 3 (10.3%) in lower middle-income group^{28–30}; there were no studies from the low-income group.

On population, out of the total 29 studies, 8 (27.6%) targeted patients with COVID-19,^{28–30 45 48 49 51 52} 6 (20.7%) among healthcare workers^{31 34 35 44 46 47} and more than half (51.7%) were general population.^{26 27 32 33 36–43 50} Out of the 29 studies, 14 (48.3%) were RCTs.^{27–29 33 34 38–42 44} (2B,3)^{48 50} The remaining 15 (51.7%) were non-RCT studies.

The interventions reported by the studies can be classified into three types: 19 online-based interventions (65.5%),^{26 27 29–31 33 35 36 38 40–45 47 48} 6 on-site-based interventions (20.7%)^{28 34 39 49–51} and 4 combined online and on-site interventions (13.8%).^{32 37 46 52}

On outcome measurement, out of the total 29 studies, only 1 (3.5%) assessed suicide ideation,⁴⁵ while 14 (48.3%) measured depression outcome,^{32 35 38–41 43–47 50–52} 24 (82.8%) measured anxiety,^{26 28 29 32 34–36 38–52} 9 (31.0%) measured psychological stress,^{30 35–40 43 50} 9 (31.0%) measured sleep quality^{27 33 35 43 45 47–49 52} and 6 (20.7%) measured other outcomes such as loneliness and




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of the review process. *Some studies measure multiple outcomes.

self-compassion.^{26 31 36 41 44} Table 2 shows the characteristics of the studies.

Measures used for the assessment of psychological consequences

Out of 29 studies, 20 measures were used to assess psychological outcomes. We categorised the measures based on the outcome they measured including suicidal ideation depression, anxiety, stress and insomnia (online supplemental annex 2).

Suicidal ideation was assessed by using one measure, the 9-item Beck Depression Inventory (BDI), whereas depression and anxiety were assessed by using seven and eight measures, respectively. The Hospital Anxiety and Depression Scale (HADS) was most used for the assessment of depression, while the Spielberger State-Trait Anxiety Inventory (STAI) was used in six studies (25.0%) on anxiety.^{26 28 29 34 36 48} Two different measures were used to assess both depression and anxiety, including the Patient Health Questionnaire-4 and HADS. The Brief Symptom Inventory-18 and Depression Anxiety Stress Scale-21 were used to assess depression, anxiety and psychological distress.

Stress was evaluated primarily using five measures (55.6%) in nine studies, though the 10-item Perceived Stress Scale (PSS-10) was most commonly used. Among the other outcomes, insomnia assessments were distinctive in that they consisted of five tests, each of which measured the patients' sleep quality. The Insomnia Severity Index was widely used to determine sleep quality

in four studies.^{27 33 35 45} There were several measures assessing outcomes beyond the focus of this study, one test for self-compassion and two tests for loneliness.^{26 36 41 44}

Interventions, measurements and effectiveness

Table 3 (RCT studies) and table 4 (non-RCT studies) describe study design, providers of intervention, target population, intervention, measurement and psychological outcome (see full description in online supplemental annex 3).

Of 24 anxiety interventions from 18 publications, 21 (87.5%) significantly lowered anxiety in intervention groups; 11 were RCT^{26 28 29 34 39 41-43 48 50} and 10 quasi-experimental studies.^{26 32 36 44-47 49 51 52} Among the 11 RCT studies, five interventions (45.5%) were mindfulness and meditation based.^{26 39 42 43 50} Two studies (18.2%) applied specific techniques such as emotional freedom techniques³⁴ and progressive muscle relaxation and deep breathing techniques.⁴⁸

Fourteen studies provided different interventions to reduce depression, of which 11 (78.6%) significantly reduced depression.^{32 38-41 43 44 46 50-52} Of 14 studies, 5 (35.7%) of them were RCT; 4 (28.6%) related to mindfulness and meditation-based stress reduction,^{43 50} focusing on mind-body exercises including yoga⁴⁰ and tai chi,³⁹ and 1 (7.1%) provided empathetic conversations through phone calls.⁴¹

Of nine studies on insomnia, seven (77.8%) found that the interventions significantly reduced insomnia or improved sleep quality.^{27 33 43 47-49 52} Of these nine

Table 2 Characteristics of the 29 included studies*

	Characteristics (%)	Reference
Geographical area (WHO regions)		
Eastern Mediterranean region	3 (10.3)	28–30
European region	10 (34.5)	31–40
Region of the Americas	4 (13.8)	41–44
Western Pacific region	11 (37.9)	26 45–52
Global	1 (3.5)	27
Income groups (World Bank)		
High-income economies	13 (44.8)	31–33 35–41 43–45
Upper middle-income economies	12 (41.4)	26 34 42 46–52
Lower middle-income economies	3 (10.3)	28–30
Global	1 (3.5)	27
Participants		
Patients with COVID-19 (confirmed/suspected)	8 (27.6)	28–30 45 48 49 51 52
Healthcare workers	6 (20.7)	31 34 35 44 46 47
General population (included student and teacher)	15 (51.7)	26 27 32 33 36–43 50
Study design		
Randomised controlled trial	14 (48.3)	26 28–30 34 35 39–43(2B,3) 48 50
Non-randomised controlled trial	15 (51.7)	26 27 31–33 36–38 4445–47 49 51 52
Intervention		
Online	19 (65.5)	26 27 29–31 33 35 36 38 40–45 47 48
On-site	6 (20.7)	28 34 39 49–51
Combined online and on-site	4 (13.8)	32 37 46 52
Psychological outcome		
Suicidal idea	1 (3.5)	45
Depression	14 (48.3)	32 35 38–41 43–47 50–52
Anxiety	24 (82.8)	26 28 29 32 34–36 38–52
Stress	9 (31.0)	30 35–40 43 50
Insomnia/sleep quality	9 (31.0)	27 33 35 43 45 47–49 52

*There are 29 studies out of 27 papers.

studies, only two (22.2%) were RCT, with one (11.1%) using a mobile meditation application,⁴³ and the other using progressive muscle relaxation and deep breathing techniques.⁴⁸

Of the total nine interventions against stress, seven (77.8%) had shown significant positive outcomes in stress reduction.^{30 36–39 43 50} One study (11.1%) shows no association of interventions and outcome,⁴⁰ while one (11.1%) study showed positive and negative results due to timing of assessment.⁴⁵ Of the total 29 studies, three (10.3%) reported the effectiveness of interventions on self-compassion; participants reported significantly higher perception of self-compassion than those in the control group.^{26 36 44} One of 29 studies (3.4%) using preassessment and postassessment shows improvement of loneliness.⁴¹ Only one study (3.4%) examined suicide, 30 min telephone consultation was provided which was effective after 1 week.⁴⁵

Risk of bias

The quality of reporting for 14 RCT studies was assessed using the Cochrane risk-of-bias tool. Table 5 shows the risk of bias by six domains. These RCT studies had no risk of attrition bias (incomplete outcome data) or reporting bias (selective reporting). Of 14 studies, five (35.7%) had low risk of bias for all six dimensions reflecting high-quality RCT. Most studies had unclear reporting on the risk of allocation concealment, random sequence generation and blinding participants and personnel.^{26 28 39 42 48} Two studies (14.3%) reported high risk of bias on blinding of participants and personnel.^{40 50}

DISCUSSION

Study characteristics

All 29 studies in this review were carried out in 2020 in all geographical continents except Africa and Australia. The majority of studies (25, 86.2%) were from high-income

Table 3 Characteristics of the reviewed randomised controlled trial (RCT) studies, intervention, measurement and psychological outcome

Author	Provider (channel)	Target population	Intervention	Significance					Suicidal idea	
				N=total sample Nc: control Ni: intervention	Anxiety	Stress	Depression	Insomnia/ sleep quality		
Dincer and Inangil ³⁴ (Turkey)	Certified emotional freedom technique	Nurses	Activating and deactivating signals to brain by stimulating points on skin	N=72 Nc=37 Ni=35	*					
Fiol-DeRoque et al ³⁵ (Spain)	Self-conducted (online application)	Healthcare workers	PsyCovidApp, targeting emotional skills, healthy lifestyle behaviour, burnout and social support	N=436 Nc=234 Ni=248	NoAssoc	NoAssoc	NoAssoc	NoAssoc	NoAssoc	
Guan et al ²⁶ (study 2B) (China)	Self-conducted (online)	General population	Self-compassion writing task	N=79 Nc=39 Ni=40	*					
Guan et al ²⁶ (study 3) (China)	Self-conducted (online)	College students	Self-compassion meditation and writing tasks	N=95 Nc=45 Ni=50	*					
Kahlon et al ⁴¹ (USA)	Trained callers (telephone based)	Homebound older adults and people with disabilities	Phone call with empathetic conversations	N=240 Nc=120 Ni=120	*			*		
Khademi et al ²⁸ (Iran)	Self-conducted	Patients with COVID-19	Performed mandala colouring	N=70 Nc=35 Ni=35	*					
Liu et al ⁴⁸ (China)	Trainer	Patients with COVID-19	Jacobson's relaxation techniques	Intervention=51 Nc=26 Ni=25	*				*	
Malboeuf-Hurtubise et al ⁴² (Canada)	Trained students in psychology (video)	School students	Mindfulness-based interventions and philosophical discussions on COVID-19	N=37 (pre-post intervention assessment)	*					

Continued

Table 3 Continued

Author	Provider (channel)	Target population	Intervention	N=total sample Nc: control Ni: intervention	Significance					
					Anxiety	Stress	Depression	Insomnia/ sleep quality	Suicidal idea	
Parizad <i>et al</i> ²⁹ (Iran)	Nurses	Patients with COVID-19	Guided imagery session with five audio tracks	N=110 Nc=55 Ni=55	*					
Shaygan <i>et al</i> ³⁰ (Iran)	Self-conducted (online multimedia)	Patients with COVID-19	Cognitive-behavioural techniques, stress management, mindfulness and positive psychotherapy	N=48 Nc=22 Ni=26		*				
Smith <i>et al</i> ⁴³ (USA)	Self-conducted (online application)	Obstetric and gynaecology patients	Mobile meditation app: encouragement, mindfulness meditation, sleep stories, nature sounds	N=101 Nc=51 Ni=50	*	*	*	*	*	
Solianik <i>et al</i> ³⁹ (Lithuania)	Tai chi instructor (online)	General population >60 years	Tai chi practice	N=30 Nc=15 Ni=15	*	*	*	*	*	
Wadhen and Cartwright ⁴⁰ (UK)	Yoga teacher (online)	General population	Yoga class via Zoom meetings	N=34 Nc=17 Ni=17	NoAssoc	NoAssoc	NoAssoc	*		
Zhang <i>et al</i> ⁵⁰ (China)	Self-conducted (online)	Residents	Mindfulness-based stress reduction programme	N=51 Nc=28 Ni=29	*	*	*	*	*	

NoAssoc denotes no association between measure and postintervention symptoms.
 Grey shade denotes not applicable.
 *Denotes statistically significant association with postintervention mental health improvements.

Table 4 Characteristics of the reviewed non-randomised controlled studies (NRS), intervention, measurement and psychological outcome

Author	Provider (channel)	Target population	Intervention	N=total sample Nc: control Ni: intervention	Anxiety	Stress	Depression	Insomnia/ sleep quality	Suicidal idea
Guan <i>et al</i> ²⁶ (China)	Self-conducted (online)	Young adults	A live session of self-compassion meditation	N=64 (pre-post intervention assessment)	*				
Giordano <i>et al</i> ³¹ (Italy)	Trained music therapist	Clinical staff	Music therapy supplemented by guided imagery and music	N=34 (pre-post intervention assessment)					*
González-García <i>et al</i> ³⁶ (Spain)	Mindfulness teacher and psychologist (online)	University students	Mini lectures, mindfulness and compassion practices and self-reflection exercises	N=66 (pre-post intervention assessment)	*	*	*	*	
Gorbeña <i>et al</i> ³⁷ (Spain)	Facilitators (interview and music playlists)	University students	Well-being and personal development programme	N=151 Nc=39 Ni=112				*	
Kim <i>et al</i> ⁴⁵ (South Korea)	Psychiatrists (telephone based)	Patients with COVID-19	30 min telephone consultation providing information and support for COVID-19 infection and hospital isolation	Week 1, n=33 Week 2, n=19 (pre-post intervention assessment)	†		†	†	†
Li <i>et al</i> ⁴⁶ (China)	Teacher (video)	Nurses	Simulation training about using personal protective devices	N=60 (pre-post intervention assessment)	*		*		
Liu <i>et al</i> ⁴⁷ (China)	Self-conducted	Nurses	Diaphragmatic breathing relaxation training	N=140 (pre-post intervention assessment)	*		NoAssoc	*	
Luo <i>et al</i> ⁴⁹ (China)	Nurses	Patients with COVID-19	Auricular point pressure	N=84 Nc=18 Ni=66	*			*	
Matiz <i>et al</i> ³² (Italy)	Socio-health educator and psychologist	Schoolteachers	Mindfulness-oriented meditation training programme	N=67 (pre-post intervention assessment)	*		*		

Continued

Table 4 Continued

Author	Provider (channel)	Target population	Intervention	N: total sample Nc: control Ni: intervention	Anxiety	Stress	Depression	Insomnia/ sleep quality	Suicidal idea
Niu <i>et al</i> ⁵¹ (China)	Nurses	(Suspected) COVID-19	Providing information about isolation, treatment and social-emotional support	N=137 (pre-post intervention assessment)	*		*		
Philip <i>et al</i> ³³ (France)	Self-conducted (online application)	General population	KANOPEE app with virtual agent interaction	N=47 (pre-post intervention assessment)				*	
Riva <i>et al</i> ³⁸ (Italy)	Self-conducted	General population	360° virtual reality (VR) video entitled 'The Secret Garden' and a series of social exercises	N=38 (pre-post intervention assessment)	NoAssoc	*	*		
Sylvia <i>et al</i> ⁴⁴ (USA)	Psychologists, physicians, social workers, nurses (online)	Healthcare workers	Stress management and resilience training, relaxation response programme	N=102 (pre-post intervention assessment)	*		*		
Tunuguntia <i>et al</i> ²⁷ (global level)	Self-conducted (online)	General population	Yoga and meditation with sound therapy	N=820 (pre-post intervention assessment)				*	
Yang <i>et al</i> ⁵² (China)	Psychotherapist and nurse (in person and online)	Patients with COVID-19	Psychotherapy, empathy, muscle and breath relaxation and cognitive-behavioural therapy	N=35 (pre-post intervention assessment)	*		*	*	

NoAssoc denotes no association between measure and postintervention symptoms.
 Grey shade denotes not applicable.
 *Denotes statistically significant association with postintervention mental health improvements.
 †Denotes mixed results of association and no association due to difference in timing of outcome measurement.

Table 5 Risk of bias assessments for all 14 RCT studies

	Random sequence generation	Allocation concealment	Blinding of participants and personnel	Incomplete outcome data	Selective reporting	Other bias	Number of low risks out of six criteria
Dincer and Inangil ³⁴	Low	Low	Low	Low	Low	Low	6
Fiol-DeRoque <i>et al</i> ³⁵	Low	Low	Low	Low	Low	Low	6
Guan <i>et al</i> ²⁶ (study 2B)	Unclear	Unclear	Unclear	Low	Low	Low	3
Guan <i>et al</i> ²⁶ (study 3)	Unclear	Unclear	Unclear	Low	Low	Low	3
Kahlon <i>et al</i> ⁴¹	Low	Low	Low	Low	Low	Low	6
Khademi <i>et al</i> ²⁸	Low	Unclear	Unclear	Low	Low	Low	4
Liu <i>et al</i> ⁴⁸	Unclear	Unclear	Unclear	Low	Low	Low	3
Malboeuf-Hurtubise <i>et al</i> ⁴²	Unclear	Unclear	Low	Low	Low	Unclear	3
Parizad <i>et al</i> ²⁹	Low	Low	Low	Low	Low	Low	6
Shaygan <i>et al</i> ³⁰	Low	Low	Low	Low	Low	Low	6
Smith <i>et al</i> ⁴³	Low	Low	Unclear	Low	Low	Low	5
Solianik <i>et al</i> ³⁹	Unclear	Unclear	Low	Low	Low	Low	4
Wadhen and Cartwright ⁴⁰	Low	Unclear	High	Low	Low	Low	4
Zhang <i>et al</i> ⁵⁰	Low	Unclear	High	Low	Low	Low	4

RCT, randomised controlled trial.

and upper middle-income countries. Therefore, the varieties of interventions in this review can be useful for adaptive application in different country contexts, although there is no study from low-income countries. Psychological impacts are caused by direct consequences of the pandemic,^{53–55} and by public health and social containment measures used by governments, notably movement restrictions and lockdowns.⁵⁶ Over a half (19, 65.5%) of the psychological interventions reviewed in this study were delivered to patients with COVID-19, healthcare workers and general population via an online platform, as on-site intervention was not possible in the pandemic context. The psychological consequences from this pandemic are diverse, requiring a wide range of interventions and tools for measuring subsequent psychological outcomes, including suicidal ideation, depression, anxiety, stress and insomnia.

Each outcome was evaluated using a variety of tools. The BDI was the most commonly used to assess suicidal ideation, followed by the HADS for screening depression, the STAI for screening anxiety, the PSS for screening stress and the Insomnia Severity Index for screening insomnia. The measures were all used for screening rather than diagnosis. Several measures were validated, with Cronbach's alpha values greater or equal to 0.7 indicating high reliability. The majority of the tools were self-assessment questionnaires that participants in the study could complete using the sample scoring system. However, no articles reported on the sensitivity and specificity of each measure.

Interventions and psychological outcomes

Overall, most interventions have positive impact in mitigating psychological consequences in patients with COVID-19, healthcare workers and general population. A single intervention can have combined effects on multiple psychological outcomes; for example, one RCT using mobile meditation application, including verbal and written encouragement, mindfulness meditation, sleep stories and nature sounds, has significant positive effects on reduction of depression, anxiety, stress and insomnia.⁴³ However, another study using an online application targeting emotional skills, healthy lifestyle behaviour, burnout and social support reported no significant difference in the same measured outcomes.³⁵

Almost all 14 RCT studies that included mindfulness and meditation interventions to reduce depression, anxiety and stress showed significantly improved outcomes (depression,^{39–41 43 50} anxiety,^{26 28 29 34 39–43 50} stress^{39 43 50}) with the exception of one study,⁴⁰ which did not improve anxiety. Stress, anxiety and depression are caused by sympathetic nervous system activation due to widespread depolarisation throughout the brain and body; on the other hand, meditation and deep breathing lead to parasympathetic nervous system activation due to widespread inhibition and hyperpolarisation.⁵⁷ Recently, mindfulness and meditation-based interventions have increasingly been integrated into mental health interventions⁵⁸; meditation practices facilitate training of the self-regulation of attention and awareness, thereby enhancing control of mental processes, and consequently

increasing well-being. Meditation and breathing techniques such as yoga and tai chi, targeting body and mind, are also applied and are effective in reducing stress, anxiety and depression.^{59 60} Yoga, for example, helps with intellectual and mental exercise, improves the psychological management and monitoring of stress and negative emotions and supports mental balance.^{60–62} Mindfulness and meditation-based interventions are readily available, and do not pose threats or risks. It can be applied as a first-line or complementary intervention for stress, anxiety, depression and some emotional disorders during COVID-19.

Moreover, combining mindfulness intervention with online application technology can benefit people who are quarantined or in isolation wards with no direct access to psychotherapy. In this review, online multimedia psychoeducational interventions were demonstrated to significantly reduce stress³⁰; mindfulness-based stress reduction, cognitive-behavioural techniques, stress management techniques and positive psychotherapy have been delivered via WhatsApp. Nowadays, internet-based, self-administrated multimedia and education can promote access to mental health interventions, particularly amidst COVID-19. Online-based interventions enable people to social distance and be time efficient and cost-effective.

The pandemic does not allow in-person on-site intervention, causing a major shift to remote care^{63–65} and telemedicine; the use of electronic communication to deliver healthcare services at a distance becomes a key supporting measure for health service delivery systems during this pandemic.^{66 67} Several channels of telemedicine are accessible by patients and health professionals including telephone calls, video calls, interactive mobile health applications (m-Health), short message service, email, secure remote patient monitoring and videoconferencing. Video call via social media is the most common channel of communication, but special communication software (eg, Zoom) or current messenger applications (eg, WhatsApp and WeChat) are popular and convenient options as well.⁶⁸

Furthermore, several countries have developed digital health technology to facilitate healthcare delivery through an online platform.^{69 70} Several challenges remain; online interventions cannot replace in-person therapy, notably when there is high risk of suicide which necessitates the presence of a psychotherapist or human interaction.^{71 72} Furthermore, several online counsellors⁶⁵ raised concerns from their experiences, such as a lack of non-verbal language for interpreting patients who struggle to verbally communicate their feelings⁷³ or significantly shorter counselling periods through the telephone due to 'more superficial responses to questions'.⁷⁴ However, the pandemic has raised awareness for the necessity of the online interventions where limitations can be resolved, as several studies have reported benefits of online services.^{75–78}

Bias assessment

The most common reason across items of bias assessment was related to incomplete or unclear reporting in the RCT.⁷⁹ Between 2011 and 2014, 24.7% of the RCTs included in the Cochrane Review demonstrated an unclear risk of bias in the domain of incomplete outcome data.⁸⁰ But all 14 RCTs in the review presented complete outcome data and reported loss to follow-up for risk of attrition bias. In this review, each article was assessed by at least two authors and we compared the risk of bias assessment for each item in terms of 'high', 'low' or 'unclear' risk of bias between the two reviews.

More than half of the studies had unclear risk on allocation concealment because the method of concealment was not described, or due to inadequate detail to allow a clear judgement. Similarly, 50.1% of the reported April 2011 issue of the Cochrane Database of Systematic Reviews had unclear risk of bias on allocation concealment.⁸¹ Concealment of randomised allocation prevents an influence of patient characteristics on allocation to intervention and control groups. One study found that lack of adequate allocation concealment was associated with overestimation of treatment effect.⁸²

Two studies reported high risk of bias on blinding of participants and personnel.^{40 50} Blinding of participants and personnel prevents differences in patient management between groups and blinding of outcome assessors also prevents knowledge of the assigned intervention group influencing outcome measurement. However, both studies were conducted through online platform and the group memberships were not blind.

Limitations

Causal interpretations are limited by 15 non-randomised controlled study designs. In addition to variation in sample size across studies, there were also variations in the mode of delivery of interventions, participant characteristics and measurement tools across different study sites. Furthermore, a majority of the trials included small sample sizes which is commonplace for studies on psychological interventions. High heterogeneity of selected studies led to inability to conduct meta-analysis. There was also a lack of assessment of cointerventions, which provide external sources of support to the participants in the intervention and control groups and may have influenced the results of the outcomes of an intervention. Lastly, postintervention outcomes from the studies were measured from short follow-up durations, so it is not possible to conclude the long-term psychological effects of those interventions.

CONCLUSION AND POLICY IMPLICATIONS

Of the total reviewed 29 studies, 26 diagnostic or screening measures were applied; 8 for anxiety (30.9%), 7 for depression (26.9%), 5 for stress (19.2%), 5 for insomnia (19.2%) and 1 for suicide (3.8%). Most studies conducted in high-income and upper middle-income



countries. Half of studies targeted general population and the remaining in patients with COVID-19 and healthcare workers. Among 14 RCT studies, most interventions were effective in reducing psychological consequences.

Despite study limitations, the use of measurement tools and the strength of evidence from this review suggest an overall positive impact of the different interventions in terms of reducing psychological consequences in the population and healthcare workers. These effective interventions can be replicated and applied in other country settings, for which modes of delivery can be adjusted in line with country resources, epidemic situation and social and health system contexts. Moreover, it is necessary to improve the use of evidence in psychological interventions in mental health services.

We recommend that these effective interventions can be replicated and applied in other country settings, for which modes of delivery can be adjusted in line with country resources, epidemic situation and social and health system contexts. In the first quarter of 2022, the pandemic is still far from over due to low immunisation coverage in African countries and continued emergence of variants of concerns in the context of pressure to resuming the economic activities and opening the country borders.⁸³ In ending the acute phase of the pandemic, rapid scale-up and promoting vaccine acceptance are key policy actions.⁸⁴

Acknowledgements The authors gratefully acknowledge the funding support from the Thailand Science Research and Innovation (TSRI) under the Senior Research Scholar on Health Policy and System Research (contract number: RTA6280007) and the National Research Council of Thailand. The authors are grateful for the partial support from Chiang Mai University and wish to thank Saranya Sachdev for her support on proofreading.

Contributors Conceptualisation: AL, PP, AC and VT. Data extraction: AL, PP and AC. Formal analysis: AL, PP and AC. Writing—original draft preparation: AL, PP and AC. Writing—review and editing: CS and VT. All authors have read and agreed to the published version of the manuscript. The scientific guarantor of this publication is AL.

Funding This study is funded by the National Research Council of Thailand (NRCT) and the Senior Research Scholar on Health Policy and System Research (grant number: RTA6280007).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Angkana Lekagul <http://orcid.org/0000-0002-7335-9028>

REFERENCES

- Harapan H, Wagner AL, Yufika A, *et al*. Acceptance of a COVID-19 vaccine in Southeast Asia: a cross-sectional study in Indonesia. *Front Public Health* 2020;8:381.
- World Health Organization. *Coronavirus disease (COVID-19) dashboard*, 2021.
- Levin J. Mental health care for survivors and healthcare workers in the aftermath of an outbreak. In: *Psychiatry of pandemics*, 2019: 127–41.
- Rubin GJ, Wessely S. The psychological effects of quarantining a City. *BMJ* 2020;368:m313.
- Abad C, Fearday A, Safdar N. Adverse effects of isolation in hospitalised patients: a systematic review. *J Hosp Infect* 2010;76:97–102.
- Purssell E, Gould D, Chudleigh J. Impact of isolation on hospitalised patients who are infectious: systematic review with meta-analysis. *BMJ Open* 2020;10:e030371.
- Sharma A, Pillai DR, Lu M, *et al*. Impact of isolation precautions on quality of life: a meta-analysis. *J Hosp Infect* 2020;105:35–42.
- Adhanom Ghebreyesus T. Addressing mental health needs: an integral part of COVID-19 response. *World Psychiatry* 2020;19:129–30.
- Kawohl W, Nordt C. COVID-19, unemployment, and suicide. *Lancet Psychiatry* 2020;7:389–90.
- Posel D, Oyenubi A, Kollamparambil U. Job loss and mental health during the COVID-19 lockdown: evidence from South Africa. *PLoS One* 2021;16:e0249352.
- COVID-19 Mental Disorders Collaborators. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet* 2021;398:1700–12.
- Khasne RW, Dhakulkar BS, Mahajan HC, *et al*. Burnout among healthcare workers during COVID-19 pandemic in India: results of a questionnaire-based survey. *Indian J Crit Care Med* 2020;24:664–71.
- Jalili M, Niroomand M, Hadavand F, *et al*. Burnout among healthcare professionals during COVID-19 pandemic: a cross-sectional study. *Int Arch Occup Environ Health* 2021;94:1345–52.
- Nishimura Y, Miyoshi T, Hagiya H, *et al*. Burnout of healthcare workers amid the COVID-19 pandemic: a Japanese cross-sectional survey. *Int J Environ Res Public Health* 2021;18:2434.
- Preti E, Di Mattei V, Perego G, *et al*. The psychological impact of epidemic and pandemic outbreaks on healthcare workers: rapid review of the evidence. *Curr Psychiatry Rep* 2020;22:43.
- Xiao X, Zhu X, Fu S, *et al*. Psychological impact of healthcare workers in China during COVID-19 pneumonia epidemic: a multi-center cross-sectional survey investigation. *J Affect Disord* 2020;274:405–10.
- Marvaldi M, Mallet J, Dubertret C, *et al*. Anxiety, depression, trauma-related, and sleep disorders among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Neurosci Biobehav Rev* 2021;126:252–64.
- Magnavita N, Chirico F, Garbarino S, *et al*. Sars/mers/sars-cov-2 outbreaks and burnout syndrome among healthcare workers. An umbrella systematic review. *Int J Environ Res Public Health* 2021;18:4361.
- Waterman S, Hunter ECM, Cole CL, *et al*. Training Peers to treat Ebola centre workers with anxiety and depression in Sierra Leone. *Int J Soc Psychiatry* 2018;64:156–65.
- Decosimo CA, Hanson J, Quinn M, *et al*. Playing to live: outcome evaluation of a community-based psychosocial expressive arts program for children during the Liberian Ebola epidemic. *Glob Ment Health* 2019;6:e3.
- Cénat JM, Mukunzi JN, Noorishad P-G, *et al*. A systematic review of mental health programs among populations affected by the Ebola virus disease. *J Psychosom Res* 2020;131:109966.
- Figuerola CA, Aguilera A. The need for a mental health technology revolution in the COVID-19 pandemic. *Front Psychiatry* 2020;11:523.
- Madigan S, Racine N, Cooke JE, *et al*. COVID-19 and telemental health: benefits, challenges, and future directions. *Canadian Psychology / Psychologie canadienne* 2020;62:5–11.
- Zhu S, Zhuang Y, Lee P, *et al*. The changes of suicidal ideation status among young people in Hong Kong during COVID-19: a longitudinal survey. *J Affect Disord* 2021;294:151–8.

- 25 Higgins JPT, Thomas J, Chandler J, *et al*, eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.2. Handbook*, 2021.
- 26 Guan F, Wu Y, Ren W, *et al*. Self-compassion and the mitigation of negative affect in the era of social distancing. *Mindfulness* 2021;12:2184–95.
- 27 Tunuguntla R, Tunuguntla HSGR, Kathuria H, *et al*. Effectiveness of App-Based yoga of Immortals (YOI) intervention for insomnia in Asian population during pandemic restrictions. *Int J Environ Res Public Health* 2021;18:5706.
- 28 Khademi F, Rassouli M, Rafiei F, *et al*. The effect of mandala colouring on anxiety in hospitalized COVID-19 patients: A randomized controlled clinical trial. *Int J Ment Health Nurs* 2021;30 Suppl 1:1437–44.
- 29 Parizad N, Goli R, Faraji N, *et al*. Effect of guided imagery on anxiety, muscle pain, and vital signs in patients with COVID-19: a randomized controlled trial. *Complement Ther Clin Pract* 2021;43:101335.
- 30 Shaygan M, Yazdani Z, Valibeygi A. The effect of online multimedia psychoeducational interventions on the resilience and perceived stress of hospitalized patients with COVID-19: a pilot cluster randomized parallel-controlled trial. *BMC Psychiatry* 2021;21:93.
- 31 Giordano F, Scarlata E, Baroni M, *et al*. Receptive music therapy to reduce stress and improve wellbeing in Italian clinical staff involved in COVID-19 pandemic: a preliminary study. *Arts Psychother* 2020;70:101688.
- 32 Matiz A, Fabbro F, Paschetto A, *et al*. Positive impact of mindfulness meditation on mental health of female teachers during the COVID-19 outbreak in Italy. *Int J Environ Res Public Health* 2020;17:6450.
- 33 Philip P, Dupuy L, Morin CM, *et al*. Smartphone-Based virtual agents to help individuals with sleep concerns during COVID-19 confinement: feasibility study. *J Med Internet Res* 2020;22:e24268.
- 34 Dincer B, Inangil D. The effect of emotional freedom techniques on nurses' stress, anxiety, and burnout levels during the COVID-19 pandemic: a randomized controlled trial. *Explore* 2021;17:109–14.
- 35 Fiol-DeRoque MA, Serrano-Ripoll MJ, Jiménez R, *et al*. A mobile phone-based intervention to reduce mental health problems in health care workers during the COVID-19 pandemic (PsyCovidApp): randomized controlled trial. *JMIR Mhealth Uhealth* 2021;9:e27039.
- 36 González-García M, Álvarez JC, Pérez EZ, *et al*. Feasibility of a brief online mindfulness and Compassion-Based intervention to promote mental health among university students during the COVID-19 pandemic. *Mindfulness* 2021;12:1685–95.
- 37 Gorbeña S, Gómez I, Govillard L, *et al*. The effects of an intervention to improve mental health during the COVID-19 quarantine: comparison with a COVID control group, and a pre-COVID intervention group. *Psychol Health* 2022;37:178–93.
- 38 Riva G, Bernardelli L, Castelnovo G, *et al*. A virtual Reality-Based self-help intervention for dealing with the psychological distress associated with the COVID-19 Lockdown: an effectiveness study with a two-week follow-up. *Int J Environ Res Public Health* 2021;18:8188.
- 39 Solianik R, Mickevičienė D, Žibinaitė L, *et al*. Tai chi improves psychoemotional state, cognition, and motor learning in older adults during the COVID-19 pandemic. *Exp Gerontol* 2021;150:111363.
- 40 Wadhen V, Cartwright T. Feasibility and outcome of an online streamed yoga intervention on stress and wellbeing of people working from home during COVID-19. *Work* 2021;69:331–49.
- 41 Kahlon MK, Aksan N, Aubrey R, *et al*. Effect of Layperson-Delivered, Empathy-Focused program of telephone calls on loneliness, depression, and anxiety among adults during the COVID-19 pandemic: a randomized clinical trial. *JAMA Psychiatry* 2021;78:616–22.
- 42 Malboeuf-Hurtubise C, Léger-Goodes T, Mageau GA, *et al*. Philosophy for children and mindfulness during COVID-19: results from a randomized cluster trial and impact on mental health in elementary school students. *Prog Neuropsychopharmacol Biol Psychiatry* 2021;107:110260.
- 43 Smith RB, Mahner ND, Foote J, *et al*. Mindfulness effects in obstetric and gynecology patients during the coronavirus disease 2019 (COVID-19) pandemic: a randomized controlled trial. *Obstet Gynecol* 2021;137:1032–40.
- 44 Sylvia LG, George N, Rabideau DJ, *et al*. Moderators of a resiliency group intervention for frontline clinicians during the COVID-19 pandemic. *J Affect Disord* 2021;293:373–8.
- 45 Kim J-W, Stewart R, Kang S-J, *et al*. Telephone based interventions for psychological problems in hospital isolated patients with COVID-19. *Clin Psychopharmacol Neurosci* 2020;18:616–20.
- 46 Li D-F, Shi C-X, Shi F-Z, *et al*. Effects of simulation training on COVID-19 control ability and psychological states of nurses in a children's Hospital. *Eur Rev Med Pharmacol Sci* 2020;24:11381–5.
- 47 Liu Y, Jiang T-T, Shi T-Y, *et al*. The effectiveness of diaphragmatic breathing relaxation training for improving sleep quality among nursing staff during the COVID-19 outbreak: a before and after study. *Sleep Med* 2021;78:8–14.
- 48 Liu K, Chen Y, Wu D, *et al*. Effects of progressive muscle relaxation on anxiety and sleep quality in patients with COVID-19. *Complement Ther Clin Pract* 2020;39:101132.
- 49 Luo Y, Ling C, Liu Y, *et al*. The beneficial role of auricular point pressure in insomnia and anxiety in isolated COVID-19 patients. *Evid Based Complement Alternat Med* 2021;2021:6611942.
- 50 Zhang H, Zhang A, Liu C, *et al*. A brief online Mindfulness-Based group intervention for psychological distress among Chinese residents during COVID-19: a pilot randomized controlled trial. *Mindfulness* 2021;12:1502–12.
- 51 Niu W, Ma X, Zhang Y, *et al*. The psychological effects of nursing interventions on patients with suspected COVID-19 during isolation. *Ann Palliat Med* 2021;10:6344–50.
- 52 Yang X, Yang X, Kumar P, *et al*. Social support and clinical improvement in COVID-19 positive patients in China. *Nurs Outlook* 2020;68:830–7.
- 53 Bekele F, Hajure M. Magnitude and determinants of the psychological impact of COVID-19 among health care workers: a systematic review. *SAGE Open Med* 2021;9:20503121211012510.
- 54 Della Monica A, Ferrara P, Dal Mas F, *et al*. The impact of Covid-19 healthcare emergency on the psychological well-being of health professionals: a review of literature. *Ann Ig* 2022;34:27–44.
- 55 Nochaiwong S, Ruengorn C, Thavorn K, *et al*. Global prevalence of mental health issues among the general population during the coronavirus disease-2019 pandemic: a systematic review and meta-analysis. *Sci Rep* 2021;11:10173.
- 56 Brooks SK, Webster RK, Smith LE, *et al*. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet* 2020;395:912–20.
- 57 Jerath R, Crawford MW, Barnes VA, *et al*. Self-Regulation of breathing as a primary treatment for anxiety. *Appl Psychophysiol Biofeedback* 2015;40:107–15.
- 58 Wielgosz J, Goldberg SB, Kral TRA, *et al*. Mindfulness meditation and psychopathology. *Annu Rev Clin Psychol* 2019;15:285–316.
- 59 Wang C, Bannuru R, Ramel J, *et al*. Tai chi on psychological well-being: systematic review and meta-analysis. *BMC Complement Altern Med* 2010;10:23.
- 60 Dhruva A, Miaskowski C, Abrams D, *et al*. Yoga breathing for cancer chemotherapy-associated symptoms and quality of life: results of a pilot randomized controlled trial. *J Altern Complement Med* 2012;18:473–9.
- 61 Streeter CC, Whitfield TH, Owen L, *et al*. Effects of yoga versus walking on mood, anxiety, and brain GABA levels: a randomized controlled MRS study. *J Altern Complement Med* 2010;16:1145–52.
- 62 Dumbala S, Bhargav H, Satyanarayana V, *et al*. Effect of yoga on psychological distress among women receiving treatment for infertility. *Int J Yoga* 2020;13:115.
- 63 European Parliament. The rise of digital health technologies during the pandemic | think tank |. Available: [https://www.europarl.europa.eu/thinktank/en/document/EPRS_BRI\(2021\)690548](https://www.europarl.europa.eu/thinktank/en/document/EPRS_BRI(2021)690548) [Accessed 07 Dec 2021].
- 64 Renn BN, Hoeft TJ, Lee HS, *et al*. Preference for in-person psychotherapy versus digital psychotherapy options for depression: survey of adults in the U.S. *NPJ Digit Med* 2019;2:6.
- 65 Feijt M, de Kort Y, Bongers I, *et al*. Mental Health Care Goes Online: Practitioners' Experiences of Providing Mental Health Care During the COVID-19 Pandemic. *Cyberpsychol Behav Soc Netw* 2020;23:860–4.
- 66 Zhang MWB, Ho RCM. Moodle: the cost effective solution for Internet cognitive behavioral therapy (I-CBT) interventions. *Technol Health Care* 2017;25:163–5.
- 67 Tran BX, Hoang MT, Vo LH, *et al*. Telemedicine in the COVID-19 pandemic: motivations for integrated, interconnected, and community-based health delivery in Resource-Scarce settings? *Front Psychiatry* 2020;11:564452.
- 68 Shanbehzadeh M, Kazemi-Arpanahi H, Kalkhajah SG, *et al*. Systematic review on telemedicine platforms in lockdown periods: lessons learned from the COVID-19 pandemic. *J Educ Health Promot* 2021;10:211.
- 69 Gunasekeran DV, Tseng RMWW, Tham Y-C, *et al*. Applications of digital health for public health responses to COVID-19: a systematic scoping review of artificial intelligence, telehealth and related technologies. *NPJ Digit Med* 2021;4:40.
- 70 World Health Organization. The impact of COVID-19 on mental, neurological and substance use services: results of a rapid assessment on JSTOR [Internet]. 2020. Available: https://www.jstor.org/stable/resrep27862?turn_away=true [Accessed 07 Dec 2021].



- 71 Gilmore AK, Ward-Ciesielski EF. Perceived risks and use of psychotherapy via telemedicine for patients at risk for suicide. *J Telemed Telecare* 2019;25:59–63.
- 72 Jobs DA, Crumlish JA, Evans AD. The COVID-19 pandemic and treating suicidal risk: the telepsychotherapy use of CAMS. *J Psychother Integr* 2020;30:226–37.
- 73 Liberati E, Richards N, Parker J. Remote care for mental health: qualitative study with service users carers and staff during the COVID-19 pandemic. *medRxiv* 2021.
- 74 Irvine A, Drew P, Bower P, *et al.* Are there interactional differences between telephone and face-to-face psychological therapy? A systematic review of comparative studies. *J Affect Disord* 2020;265:120–31.
- 75 Fu Z, Burger H, Arjadi R, *et al.* Effectiveness of digital psychological interventions for mental health problems in low-income and middle-income countries: a systematic review and meta-analysis. *Lancet Psychiatry* 2020;7:851–64.
- 76 Castro A, Gili M, Ricci-Cabello I, *et al.* Effectiveness and adherence of telephone-administered psychotherapy for depression: a systematic review and meta-analysis. *J Affect Disord* 2020;260:514–26.
- 77 Andersson G, Bergström J, Holländare F, *et al.* Internet-Based self-help for depression: randomised controlled trial. *Br J Psychiatry* 2005;187:456–61.
- 78 Zhou X, Edirippulige S, Bai X, *et al.* Are online mental health interventions for youth effective? A systematic review. *J Telemed Telecare* 2021;27:638–66.
- 79 Bertizzolo L, Bossuyt P, Atal I, *et al.* Disagreements in risk of bias assessment for randomised controlled trials included in more than one Cochrane systematic reviews: a research on research study using cross-sectional design. *BMJ Open* 2019;9:e028382.
- 80 Dechartres A, Trinquart L, Atal I, *et al.* Evolution of poor reporting and inadequate methods over time in 20920 randomised controlled trials included in Cochrane reviews: research on research study. *BMJ* 2017;357:j2490.
- 81 Savović J, Turner RM, Mawdsley D, *et al.* Association between risk-of-bias assessments and results of randomized trials in Cochrane reviews: the ROBES meta-epidemiologic study. *Am J Epidemiol* 2018;187:1113–22.
- 82 Herbison P, Hay-Smith J, Gillespie WJ. Adjustment of meta-analyses on the basis of quality scores should be abandoned. *J Clin Epidemiol* 2006;59:1249–e1.
- 83 Tangcharoensathien V, Carroll D, Lekagul A. Resilient and equitable recovery from the covid-19 pandemic. *BMJ* 2022;376:o311.
- 84 Gallè F, Sabella EA, Roma P, *et al.* Acceptance of COVID-19 vaccination in the elderly: a cross-sectional study in southern Italy. *Vaccines* 2021;9:1222.