

# Hearing Loss in the Elderly: Is the Hearing Handicap Inventory for the Elderly - Screening Version Effective in Diagnosis When Compared to the Audiometric Test?

Alexandre Barbosa Servidoni<sup>1</sup> Lucieni de Oliveira Conterno<sup>2</sup>

<sup>1</sup>Otorhinolaryngologist, Faculdade de Medicina de Marília (Famema), SP, Brazil

<sup>2</sup>Infectologist & Clinical Epidemiologist, Faculdade de Medicina de Marília (Famema), SP, Brazil

Address for correspondence Alexandre Barbosa Servidoni, MD, MSc, Rua José Bonifácio, 1185, 64-D, Jardim Cristo Rei, Marília-SP, 17513-230, Brazil (e-mail: absmed@hotmail.com; alexandreservidoni@gmail.com).

Int Arch Otorhinolaryngol 2018;22:1–8.

## Abstract

**Introduction** Hearing losses inherent to the natural process of aging represent today a major public health issue, despite the little attention that their adequate care still receives. Early recognition and proper management of these shortcomings can significantly improve hearing, as well as the patient's general quality of life, reducing the overall impact of this important and prevalent condition of the aging process.

**Objective** The aim of this research was to evaluate the accuracy of the Hearing Handicap Inventory for the Elderly - Screening Version (HHIE-S) in the diagnosis of hearing loss in the elderly when compared with the audiometric test.

**Methods** Through a cross-sectional study, our target population was composed of 138 individuals, aged over 60 and with any otorhinolaryngological complaints, recruited at the Clinic of Otorhinolaryngology and Speech Therapy of the Faculdade de Medicina de Marília (Famema), in the city of Marília, SP, Brazil. Patients already in the process of auditory rehabilitation were excluded, as well as those who did not demonstrate the minimum level of oral understanding necessary to allow the interview.

**Results** The prevalence of hearing loss according to the questionnaire was of 76.1%, while audiometry showed 79.7%. We found the diagnostic accuracy of the instrument to be of 86.2%, with a sensitivity of 89.1% and a specificity of 75.0%, regardless of gender.

**Conclusion** Thereby, we conclude that the standardized questionnaire under rating is suitable for the screening of hearing loss in the elderly, given its high accuracy and user-friendly quality.

## Keywords

- ▶ hearing loss
- ▶ presbycusis
- ▶ surveys and questionnaires
- ▶ audiometry
- ▶ aged
- ▶ aging

## Introduction

The hearing losses inherent to the natural process of aging represent a major public health issue, despite the little attention that they receive regarding their proper management.<sup>1–10</sup>

The test considered the gold standard for evaluating the auditory function is the pure tone audiometry (PTA), which does not consider the loss qualitatively, but classifies the extent of the loss at different frequencies tested.<sup>11–17</sup> There is still no consensus on the most appropriate classification of

received  
September 21, 2016  
accepted after revision  
January 2, 2017  
published online  
March 31, 2017

DOI <https://doi.org/10.1055/s-0037-1601427>.  
ISSN 1809-9777.

Copyright © 2018 by Thieme Revinter Publicações Ltda, Rio de Janeiro, Brazil

License terms



hearing loss regarding the degree of intensity, which is obtained from the arithmetic average of the hearing thresholds found in the different frequencies tested.<sup>18–26</sup>

Hearing impairment refers to any restriction or lack of ability to perceive sound and to perform tasks due to hearing problems experienced by those affected by hypoacusis. On the other hand, hearing handicap refers to the emotional and social aspects resulting from hearing impairment, which can sometimes restrict the individual's participation in everyday activities, as well as impact family relationships and social interactions.<sup>27–29</sup>

In 1982, Ventry and Weinstein<sup>30</sup> developed in the United States the Hearing Handicap Inventory for the Elderly (HHIE), a self-assessment tool containing 25 questions aimed to assess the impact of hearing loss in the emotional and social-situational adjustments of elderly patients who are not institutionalized.<sup>31–36</sup> In 1983, the same authors published a reduced version of the HHIE containing only 10 questions, but also divided between social and emotional aspects: the Hearing Handicap Inventory for the Elderly - Screening Version (HHIE-S), proposed as a screening tool to detect the degree of hearing complaints.<sup>34,37–39</sup>

In 1997, Wieselberg<sup>40</sup> released the pioneering work of translation and adaptation of the HHIE into the Portuguese language. And according to Paiva et al,<sup>41</sup> the translation into Portuguese maintained the validity of the English version.

In view of the relevance of the subject due to the high prevalence of hearing disorders and their harmful consequences;<sup>42</sup> and considering that there are more simple and inexpensive tools for tracking them, we aimed to assess the accuracy of the HHIE-S in the diagnosis of hearing losses in the elderly when compared with PTA; besides, the sensitivity and specificity of the instrument were defined, and the prevalence of hearing disorder in the target population was determined both by the questionnaire and by PTA.

## Methods

A cross-sectional study with a quantitative approach was conducted to evaluate the accuracy of the HHIE-S questionnaire as a diagnostic test when compared with PTA, which is considered the gold standard for detecting hearing loss.

The target population of this study consisted of elderly individuals aged over 60 years, who were patients at the Clinic of Otorhinolaryngology and Speech Pathology of Faculdade de Medicina de Marília (Famema), in the city of Marília, SP, Brazil. The individuals included in the study were previously scheduled or referenced in for elective outpatient consultations in the morning, Monday to Friday, thus fulfilling the requirement of being able to present an otorhinolaryngologic complaint. As the location is not a specific hearing loss clinic, individuals scheduled randomly throughout the week did not necessarily have hearing complaints.

Users of hearing aids or patients already in the process of audiological rehabilitation were excluded, as well as those who did not demonstrate a minimum level of listening to allow the questionnaire;<sup>43–46</sup> individuals with active infec-

tious diseases or tumors diagnosed during the otoscopy were also excluded and referred to proper treatment.

The sample size (N) was calculated taking into account previous literature data evaluating the performance of the HHIE-S questionnaire in a population without known hearing disorders.<sup>23,25,47</sup> Thereby, the final N = 138 was achieved.

Data collection only started after the patients read and signed the consent form, thus allowing the use of the data obtained in accordance with the standards set by the Ethics Committee of our Academic Institute. Therefore, the execution of this research was duly approved by the Ethics Committee of the institution, through protocol number 950,987, released on February 10, 2015.

After routine otoscopy, all participants were initially submitted to the HHIE-S questionnaire, aimed at analyzing the perception of hearing handicap.<sup>34,37,48–51</sup> This questionnaire was always applied by the same researcher, and the items read orally by the interviewer and immediately answered by the interviewee, a technique known as face to face. Individuals responded to questions related to hearing skills, indicating when a situation represented a problem.<sup>32,52–54</sup> If requested by the patient, a brief explanation or a new reading was provided. The average time for the whole questionnaire ranged between 5 and 10 minutes, depending on the ability of each interviewee.

After the interview with the HHIE-S questionnaire, the individual was then referred for further examination by PTA, performed by the speech therapists from our clinic, preferably and mostly on the same day as the initial interview; when this was not possible due to logistical or time constraint reasons, the PTA examination was scheduled as a routine service, to be performed within a maximum period of two months. On the day set for the audiometric examination, a new otoscopy was performed initially, but repetition of the HHIE-S was not necessary.<sup>34,37,55–58</sup> For this research, we used the audiometer 229E Interacoustics model (Middelfart, Denmark), set to the proper calibrations, as well as an appropriate acoustic booth.

To include a wider range of frequencies that encompass both the energy of speech sounds and their intelligibility,<sup>59</sup> we adopted the classification proposed by the World Health Organization (WHO),<sup>60</sup> which uses the arithmetic mean of the hearing thresholds at 500 Hz, 1,000 Hz, 2,000 Hz and 4,000 Hz, as well as 25 dB HL as normality threshold. It is important to note that the frequency of 4,000 Hz is among the first to demonstrate an age-related decline, and is, therefore, fundamental in understanding speech, especially in noisy environments.<sup>19,57,61</sup> The classification adopted was made as shown in ►Table 1:<sup>60</sup>

The accuracy of the HHIE-S questionnaire was determined by calculating the sensitivity and specificity, comparing the results that were true positive, false positive, true negative and false negative with the PTA. The confidence intervals (CIs) used to estimate the sensitivity, specificity, positive predictive value and negative predictive value were calculated using the exact method of Clopper-Pearson.<sup>62,63</sup> Analyses were performed using the IBM statistics software SPSS 17.0 (North Castle, NY, USA).

**Table 1** Audiological classification adopted (WHO)

Normal hearing	0–25 dB HL
Mild hearing loss	26–40 dB HL
Moderate hearing loss	41–60 dB HL
Severe hearing loss	61–80 dB HL
Profound hearing loss	≥ 81 dB HL

Abbreviations: dB HL, decibels hearing level; WHO, World Health Organization.

## Results

The demographic and epidemiological data obtained in this study are summarized and presented in ►Table 2.

The evaluation by the HHIE-S showed that 33 elderly patients (23.9%) had no perception of hearing handicap; 65 (47.1%) had mild/moderate perception of handicap; and 40 (29.0%) had a significant perception of handicap. Thus, the prevalence of hearing impairment by the HHIE-S questionnaire, considering the absence or presence of any perception of hearing handicap, was of 76.1% (105/138). The prevalence data as classified by the HHIE-S are described in ►Table 3 (irrespective of sex) and ►Table 4 (comparing sexes).

**Table 2** Clinical and epidemiological characteristics of 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

	Number of patients	Percentage
	<b>Total</b>	<b>%</b>
	<b>138</b>	<b>100%</b>
Male	71	51.4%
Female	67	48.6%
<b>Average age ± SD</b>	<b>71.6 ± 8.1</b>	–
60–70 years old	71	51.4%
71–80 years old	45	32.6%
> 80 years old	22	15.9%
<b>Schooling</b>		
Elementary School	34	24.6%
Middle School	93	67.4%
High School	06	4.3%
Higher Education	05	3.6%
<b>Diseases*</b>		
Diabetes mellitus	30	21.7%
Hypertension	89	64.5%
Dyslipidemia	23	16.7%
Thyroid disease	17	12.3%
Otologic disease	15	10.9%
Other diseases	68	49.3%

Abbreviation: SD, standard deviation.

Note: \* Individuals may have more than one associated disease.

**Table 3** Hearing disorder prevalence according to the HHIE-S, irrespective of sex, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

Handicap	Elderly population studied	
	Number of patients	Percentage
	<b>TOTAL (138)</b>	<b>% (100%)</b>
Absence	33	23.9%
Mild/Moderate	65	47.1%
Severe	40	29.0%
Any handicap	105	76.1%

Abbreviation: HHIE-S, Hearing Handicap Inventory For The Elderly – Screening Version.

The audiometric evaluation of these patients, considering the overall PTA, showed the following proportions: normal hearing in 28 subjects (20.3%); mild hearing loss in 39 (28.3%); moderate hearing loss in 53 (38.4%); severe hearing loss in 13 (9.4%); and profound hearing loss in 5 subjects (3.6%). The prevalence of hearing impairment by audiometric testing, considering the overall result and including all its degrees of intensity, was of 79.7% (110/138).

►Table 5 shows the prevalence data of hearing impairment according to PTA and its intensity levels, separated by right and left ear and globally.

Comparing the results obtained from the HHIE-S questionnaire and the overall PTA, we observed that among the 28 individuals with normal hearing by audiometric examination, 21 (75.0%) revealed no handicap, that is, had no perception of hearing handicap. At the other end of the results, among the five individuals with profound hearing loss according to the audiometric examination, four (80.0%) showed a severe handicap, that is, had a significant perception of hearing handicap.

In relation to the moderate hearing losses, we note that among the 53 individuals with such intensity level of hearing loss by audiometric examination, 26 (49.1%) showed a severe

**Table 4** Hearing disorder prevalence according to the HHIE-S, comparing sexes, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

Handicap	Elderly population studied (N = 138)			
	Number of patients		Percentage	
	♀	♂	♀	♂
Absence	18	15	26.9%	21.1%
Mild/Moderate	31	34	46.3%	47.9%
Severe	18	22	26.9%	31.0%
Any handicap	49	56	73.1%	78.9%

Abbreviations: HHIE-S, Hearing Handicap Inventory for the Elderly – Screening Version; N, number of patients.

Notes: ♀, female; ♂, male.

**Table 5** Hearing disorder prevalence according to PTA and its intensity levels, divided by ear (right and left) and overall, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

Hearing level (WHO)	Right ear		Left ear		Overall	
	N	%	N	%	N	%
Normal (0–25 dB)	24	17.4%	19	13.8%	28	20.3%
Mild hearing loss (26–40 dB)	36	26.1%	37	26.8%	39	28.3%
Moderate hearing loss (41–60 dB)	51	37.0%	45	32.6%	53	38.4%
Severe hearing loss (61–80 dB)	20	14.5%	24	17.4%	13	9.4%
Profound hearing loss ( $\geq 81$ dB)	07	5.1%	13	9.4%	05	3.6%

Abbreviations: dB, decibels; N, number of patients; PTA, pure tone audiometry; WHO, World Health Organization.

handicap, another 26 (49.1%) showed a mild to moderate handicap, and only 1 (1.9%) revealed no handicap.

The results found in the comparison between both diagnostic methods (HHIE-S and PTA) are shown in **Table 6**.

When comparing the overall results of HHIE-S and PTA, irrespective of the intensity levels, that is, simply considering the handicap as absent or present and the audiometric test as normal or abnormal, we found the following results: among the 28 individuals with normal audiometric examinations, 21 (75.0%) showed no handicap, and only 7 (25.0%) showed some perception of hearing handicap. Among the 110 individuals with abnormal audiometric examinations, 98 (89.1%) revealed the presence of some degree of handicap, while only 12 (10.9%) had no perception of any hearing handicap.

**Table 7** shows the comparison of the overall results of the two diagnostic methods (HHIE-S and PTA) without taking into account the degrees of intensity.

From the previous table (**Table 7**), we calculated the sensitivity, specificity, and positive and negative predictive values, with their CIs, and the accuracy of the HHIE-S questionnaire compared with the gold standard PTA, considering the overall assessment.

**Table 8** describes the statistical data of the HHIE-S, showing their calculations and the final results.

When comparing sexes, we calculated the prevalence of hearing impairment according to three criteria: the PTA, the HHIE-S questionnaire, and both methods together.

Thus, according to the overall audiometric examination and considering all the intensity levels, the prevalence of hearing impairment was of 73.1% (49/67) among women, and of 85.9% (61/71) among men.

According to the HHIE-S, and considering any level of handicap, the prevalence of hearing impairment was also of 73.1% (49/67) among women, and of 78.9% (56/71) among men.

And considering both methods together (PTA and HHIE-S), we found a prevalence of 68.7% (46/67) among women, and of 73.2% (52/71) among men.

When we evaluated the sensitivity, specificity and positive and negative predictive values, and their CIs, again considering the two methods together, but now comparing the sexes, we observed that all results were higher for females.

Thus, women showed a sensitivity of 93.9% (46/49), a specificity of 83.3% (15/18), a positive predictive value of 93.9% (46/49) and a negative predictive value of 83.3% (15/18). Among men, we found a sensitivity of 85.3% (52/61), a specificity of 60.0% (6/10), a positive predictive value of 92.9% (52/56) and a negative predictive value of 40.0% (6/15).

**Table 9** displays the statistical results obtained and described before.

**Table 6** Relationship between the results of the HHIE-S and the PTA, according to intensity levels, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

	PTA					
	Normal hearing	Mild hearing loss	Moderate hearing loss	Severe hearing loss	Profound hearing loss	Total
	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
<b>HHIE-S</b>						
Absence of handicap	21 (75.0%)	11 (28.2%)	01 (1.9%)	0 (0.0%)	0 (0.0%)	33 (23.9%)
Mild/Moderate handicap	06 (21.4%)	26 (66.7%)	26 (49.1%)	06 (46.2%)	01 (20.0%)	65 (47.1%)
Severe handicap	01 (3.6%)	02 (5.1%)	26 (49.1%)	07 (53.8%)	04 (80.0%)	40 (29.0%)
<b>Total</b>	28 (100%)	39 (100%)	53 (100%)	13 (100%)	05 (100%)	138 (100%)

Abbreviation: HHIE-S, Hearing Handicap Inventory For The Elderly – Screening Version; N, number of patients; PTA, pure tone audiometry.

Note: %, percentage (regarding the PTA).

**Table 7** Comparison between the overall results of the HHIE-S and the PTA, irrespective of the intensity levels, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

	PTA		Total
	Normal	Abnormal	
<b>Handicap</b>			
Absent	21 (75.0%)	12 (10.9%)	33 (23.9%)
Present	07 (25.0%)	98 (89.1%)	105 (76.1%)
<b>Total</b>	<b>28 (100.0%)</b>	<b>110 (100.0%)</b>	<b>138 (100.0%)</b>

Abbreviations: HHIE-S, Hearing Handicap Inventory For The Elderly – Screening Version; PTA, pure tone audiometry.

## Discussion

Considering only the HHIE-S questionnaire, we observed that 23.9% of the subjects did not perceive a handicap, while 76.1% reported the perception of some level of handicap (mild/moderate or significant); comparing sexes, we observed perception in 73.1% of women and 78.9% of men. Cruickshanks et al<sup>16</sup> found 33.7% with hearing handicap, considering both sexes; Wieselberg<sup>40</sup> observed 91.0% among women and 86.0% among men; Sestrem<sup>64</sup> found 58.0% among women and 43.0% among men. After reviewing the literature, we found no other studies reporting the prevalence of hearing impairment determined by HHIE-S alone, so that we could make a broader comparative analysis.

It is difficult to compare data on the prevalence of hearing loss among the various published studies, due to differences in sampling, populations and scenarios, as well as due to lack of consensus on audiometric criteria and/or different questionnaires used.<sup>65–68</sup>

**Table 8** Statistics of the HHIE-S, compared with the overall PTA, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

	Calculus	Results	95% CI
Accuracy	(21 + 98)/138	86.2%	79.4–91.0
Sensitivity	98/110	89.1%	81.7–94.2
Specificity	21/28	75.0%	55.1–89.3
Positive predictive value	98/105	93.3%	86.8–97.3
Negative predictive value	21/33	63.6%	45.1–79.6
Positive LR	Sen/1 – Spe	3.56	1.87–6.79
Negative LR	1 – Sen/Spe	0.15	0.08–0.26

Abbreviations: CI, confidence interval; HHIE-S, Hearing Handicap Inventory For The Elderly – Screening Version; LR, likelihood ratio; PTA, pure tone audiometry; Sen, sensitivity; Spe, specificity.

**Table 9** Statistics of the HHIE-S, compared with the overall PTA and comparing sexes, for 138 elderly patients evaluated at the Clinic of Otorhinolaryngology and Speech Therapy of Famema

	Results	95% CI (♀)	Results	95% CI (♂)
Accuracy	91.0%	84.2–97.9	81.7%	72.7–90.7
Sensitivity	93.9%	82.1–98.4	85.2%	73.3–92.6
Specificity	83.3%	57.7–95.6	60.0%	27.4–86.3
Positive predictive value	93.9%	82.1–98.4	92.9%	81.9–97.7
Negative predictive value	83.3%	57.7–95.6	40.0%	17.5–67.1
Positive LR	5.63	1.99–15.86	2.13	0.99–4.59
Negative LR	0.07	0.02–0.22	0.25	0.12–0.51

Abbreviations: CI, confidence interval; HHIE-S, Hearing Handicap Inventory For The Elderly – Screening Version; LR, likelihood ratio; PTA, pure tone audiometry.

Notes: ♀, female; ♂, male.

Considering the overall result of the PTA and including all its degrees of intensity, according to the audiometric classification of the WHO,<sup>19,57,60</sup> we observed a high prevalence of hearing loss: 79.7%. The less serious hearing losses were the most frequently observed, with 28.3% of the subjects with mild hearing loss and 38.4% with moderate hearing loss; the more disabling hearing losses were observed to a lesser extent, with 9.4% of the subjects showing severe hearing loss, and 3.6% a profound hearing loss.

Despite the difficulty in comparing prevalence data, we note singular differences between our research and the studies reviewed. Cruickshanks et al,<sup>16</sup> in a study published in 1998, found an overall prevalence of hearing impairment of 45.9%; Calais,<sup>15</sup> in 2005, found a prevalence of 12% in the age group of 45–64 years, rising to 24% in the 65–74 years age group, and reaching 39% at ages over 75 years; and finally, Sousa et al,<sup>13</sup> in 2009, observed an overall prevalence of 36.1%.

In our study, we found high values in the HHIE-S accuracy (86.2%), sensitivity (89.1%), specificity (75.0%) and positive predictive value (93.3%), whereas only the negative predictive value (63.6%) showed a lower percentage. Rosis et al<sup>25</sup> published in 2009 a study on the accuracy of the HHIE-S in two different scenarios at Universidade Federal de São Paulo: at the Audiology Clinic, with 55 elderly subjects ( $\geq 60$  years) presenting hearing complaints, and at the Geriatrics Clinic, with 23 elderly subjects ( $\geq 60$  years) with some complaints. In the first scenario, they found an accuracy of 58.2%, a sensitivity of 23.5%, a specificity of 73.7%, a positive predictive value of 28.6%, and a negative predictive value of 68.3%. In the second scenario, they found an accuracy of 91.3%, a sensitivity of 94.7%, a specificity of 75.0%, a positive predictive value of 94.7%, and a negative predictive value of 75.0%. Thereby, in the group without mandatory hearing complaints (Geriatrics Clinic) there was a statistical association between the results

obtained from the HHIE-S and the possibility of a hearing disorder. Therefore, only the Geriatrics Clinic has provided results similar to ours. It is important to emphasize that the participants in our research, despite being cared for at a specialized clinic (the Clinic of Otorhinolaryngology and Speech Therapy of Famema), might have otorhinolaryngologic complaints that are not just hearing-related. The previous data also seem to agree with the study of Menegotto et al,<sup>20</sup> published in 2011. This study concluded that, at least at audiology clinics, a scenario in which all patients necessarily have hearing complaints, the questionnaires have not proved to be effective tools for hearing screening. In a sample of 51 adults and elderly subjects, they obtained a low sensitivity (47%) and a high specificity (75%); for this, they used the HHIE-S questionnaire for the elderly, as well as its equivalent for adults, the Hearing Handicap Inventory for the Adults - Screening Version (HHIA-S),<sup>69,70</sup> both later compared with the audiometric examination. Thus, we believe that the questionnaires could be valid in health services not specific to audiology, such as a general clinic of otorhinolaryngology, geriatrics or even primary care. However, further studies are needed to confirm this hypothesis.

Evaluating the results found in our current research, we noticed a greater agreement on the extremes, that is, between individuals with normal hearing and absent handicap (75.0%), and between those with profound hearing loss and severe handicap (80.0%). For this reason, our data suggest that the HHIE-S is more suitable to identify more disabling hearing losses (severe and profound), just as Deepthi and Kasthuri<sup>71</sup> observed in 2012, and to identify with good reliability the absence of hearing disorder.

Among the 110 subjects with abnormal audiometry, 98 (89.1%) showed some degree of perceived handicap, while only 12 (10.9%) showed no perception of hearing handicap. And among 28 individuals with normal hearing, 21 (75.0%) had not perceived handicap, while only 7 (25.0%) showed some degree of perception of hearing handicap.

From the above discussion, we can see that the PTA alone is insufficient to describe the reaction of the individual before a hearing disorder, that is, it does not detect the hearing handicap. At the same time, the HHIE-S alone is not always able to accurately detect auditory sensitivity.

According to several authors,<sup>31,65-68</sup> despite the high prevalence of hearing disorders in the elderly, such an in-depth investigation is still generally restricted to medical evaluations. For this reason, they suggest the hearing screening as a useful tool in primary care, using a standardized questionnaire, such as the HHIE-S.<sup>72,73</sup>

## Conclusion

From the results obtained in this study, we conclude that the instrument under evaluation, the HHIE-S questionnaire, is suitable in the screening for hearing loss in the elderly, given its high accuracy and user-friendly quality.

Considering the methods used and the results obtained, we also conclude that there is 10.9% of probability that the elderly with no handicap show a hearing loss, as well as 89.1%

of probability that some degree of hearing loss exists in the elderly with a hearing handicap.

Moreover, it is a simple and inexpensive tool, requiring little time for its application, 10 minutes at most, and thus, it can be incorporated into traditional clinical care.

We must keep in mind that no patient should be in a process of rehabilitation without both instruments, the PTA and the HHIE-S, complementing each other.

Besides extending this research to other settings, we believe that the future association of a qualitative research might clarify some yet unknown relevant aspects, such as the possible attitudes and meanings that the elderly take on when undergoing a hearing care interview, which are likely to influence the final results.

We thus hope that such issues may be useful for all health professionals who are interested in furthering such an important subject, which concerns all of us who seek a healthy and humanized aging process.

## References

- 1 Bogardus ST Jr, Yueh B, Shekelle PG. Screening and management of adult hearing loss in primary care: clinical applications. *JAMA* 2003;289(15):1986-1990
- 2 Weinstein BE. Evaluation and management of the hearing-impaired elderly. *Geriatrics* 1990;45(02):75-83, 79-80, 83
- 3 IBGE. Censo 2010 [Internet]. Rio de Janeiro(RJ): IBGE; 2016 [citado em 5 maio 2016]. Disponível em: <http://censo2010.ibge.gov.br/resultados>
- 4 Pinzan-Faria VM, Iorio MCM. Sensibilidade auditiva e autopercepção do *handicap*: um estudo em idosos. *Distúrb Comun* 2004; 16(03):289-299
- 5 Gates GA, Murphy M, Rees TS, Fraher A. Screening for handicap hearing loss in the elderly. *J Fam Pract* 2003;52(01):56-62
- 6 do Carmo LC, Médicis da Silveira JA, Marone SA, D'Ottaviano FG, Zagati LL, Dias von Söhsten Lins E. Audiological study of an elderly Brazilian population. *Rev Bras Otorrinolaringol* 2008;74(03): 342-349
- 7 Alvarenga KF, Bevilacqua MC, Martinez MANS, Melo TM, Blasca WQ, Taga MFL. Proposta para capacitação de agentes comunitários de saúde em saúde auditiva. *Pró-Fono R Atual Cient* 2008; 20(03):171-6
- 8 Bramatti L, Morata TC, Marques JM. Ações educativas com enfoque positivo em programa de conservação auditiva e sua avaliação. *Rev CEFAC* 2008;10(03):398-408
- 9 Karlsmose B, Lauritzen T, Engberg M, Parving A. A randomised controlled trial of screening for adult hearing loss during preventive health checks. *Br J Gen Pract* 2001;51(466):351-355
- 10 Gândara MER, Alves FRA. Presbiacusia. Neto SC, Mello JF Jr, Martins RHG, Costa SS. *Tratado de Otorrinolaringologia e Cirurgia Cervicofacial*. Volume II Otologia-Otoneurologia. 2ª ed. São Paulo (SP): Roca; 2011:302-13
- 11 Veras RP, Mattos LC. Audiology and Aging: literature review and current horizons. *Rev Bras Otorrinolaringol* 2007;73(01):122-128
- 12 Samelli AG, Negretti CA, Ueda KS, Moreira RR, Schochat E. Comparação entre avaliação audiológica e screening: um estudo sobre presbiacusia. *Rev Bras Otorrinolaringol* 2011;77(01):70-76
- 13 Sousa CS, Castro Júnior Nd, Larsson EJ, Ching TH. Risk factors for presbycusis in a socio-economic middle-class sample. *Rev Bras Otorrinolaringol (Engl Ed)* 2009;75(04):530-536
- 14 Paiva KM, Cesar CLG, Alves MCGP, Barros MBA, Carandina L, Goldbaum M. Aging and self-reported hearing loss: a population-based study. *Cad Saude Publica* 2011;27(07):1292-1300

- 15 Calais LL. Acuidade e sensibilidade auditiva no idoso [CD-ROM]. In: Anais do 20º Encontro Internacional de Audiologia. 2005 São Paulo, Brasil. São Paulo(SP): Academia Brasileira de Audiologia; 2005
- 16 Cruickshanks KJ, Wiley TL, Tweed TS, et al; The Epidemiology of Hearing Loss Study. Prevalence of hearing loss in older adults in Beaver Dam, Wisconsin. *Am J Epidemiol* 1998;148(09):879–886
- 17 Griz SMS, Menezes DC, Menezes PL. Audiometria Tonal e Logoaudiometria. Neto SC, Mello JF Jr, Martins RHG, Costa SS. Tratado de Otorrinolaringologia e Cirurgia Cervicofacial. Volume I Fundamentos. 2ª ed. São Paulo(SP): Roca; 2011:389–411
- 18 Yueh B, Shapiro N, MacLean CH, Shekelle PG. Screening and management of adult hearing loss in primary care: scientific review. *JAMA* 2003;289(15):1976–1985
- 19 Lopes AC. Audiometria Tonal Liminar. Bevilacqua MC, Martinez MAN, Balen SA, Pupo AC, Reis ACMB, Frota S. Tratado de Audiologia. 1ª ed. São Paulo(SP): Ed. Santos; 2012:63–80
- 20 Menegotto IH, Soldera CLC, Anderle P, Anhaia TC. Correlation between hearing loss and the results of the following questionnaires: Hearing Handicap Inventory for the Adults – Screening Version HHIA-S and Hearing Handicap Inventory for the Elderly – Screening Version - HHIE-S. *Int Arch Otorhinolaryngol* 2011; 15(03):319–326
- 21 Calviti KCFK, Pereira LD. Sensibilidade, especificidade e valores preditivos da queixa auditiva comparados com diferentes médias audiométricas. *Braz J Otorhinolaryngol* 2009;75(06):794–800
- 22 Valet-Rosalino CM, Rozenfeld S. Triagem auditiva em idosos: comparação entre auto-relato e audiometria. *Rev Bras Otorrinolaryngol* 2005;71(02):193–200
- 23 Lopes AC, Nelli MP, Lauris JRP, Amorim RB, Melo ADP. Conditions of auditory health at work: inquiry of the auditory effects in workers exposed to occupational noise. *Int Arch Otorhinolaryngol* 2009;13(01):49–54
- 24 Davis A, Smith P, Ferguson M, Stephens D, Gianopoulos I. Acceptability, benefit and costs of early screening for hearing disability: a study of potential screening tests and models. *Health Technol Assess* 2007;11(42):1–294
- 25 Rosis ACA, Souza MRF, Iório MCM. Questionário *Hearing Handicap Inventory for the Elderly – Screening version* (HHIE-S): estudo da sensibilidade e especificidade. *Rev Soc Bras Fonoaudiol*. 2009; 14(03):339–345
- 26 Weinstein BE, Richards AM, Montano J. Handicap versus impairment: an important distinction. *J Am Acad Audiol* 1995;6(03):250–255
- 27 Romero J. Questionários de auto-avaliação de dificuldades auditivas: questões psicométricas na utilização clínica e em pesquisa [dissertação]. São Paulo (SP): Pontifícia Universidade Católica de São Paulo; 2003:71
- 28 Costa TRS. Distúrbios de audição e a classificação internacional de funcionalidade, incapacidade e saúde [tese]. São Paulo (SP): Pontifícia Universidade Católica de São Paulo; 2013:111
- 29 Corrêa GF, Russo ICP. Autopercepção do *handicap* em deficientes auditivos adultos e idosos. *Rev CEFAC* 1999;10(01):54–63 Available at: <http://www.cefac.br/revista/revista11/Artigo%207.pdf>
- 30 Ventry IM, Weinstein BE. The hearing handicap inventory for the elderly: a new tool. *Ear Hear* 1982;3(03):128–134
- 31 Santiago LM, Novaes CO. Auto-avaliação da audição em idosos. *Rev CEFAC* 2009;11(Suppl 1):98–105
- 32 Marques ACDO, Kozłowski L, Marques JM. Reabilitação auditiva no idoso. *Rev Bras Otorrinolaryngol*. 2004;70(06):806–811
- 33 Weinstein BE. Validity of a screening protocol for identifying elderly people with hearing problems. *ASHA* 1986;28(05):41–45
- 34 Weinstein BE, Ventry IM. Audiometric correlates of the Hearing Handicap Inventory for the elderly. *J Speech Hear Disord* 1983; 48(04):379–384
- 35 Lichtenstein MJ, Bess FH, Logan SA. Diagnostic performance of the hearing handicap inventory for the elderly (screening version) against differing definitions of hearing loss. *Ear Hear* 1988;9(04): 208–211
- 36 United States. Guidelines for the identification of hearing impairment/handicap in adult/elderly persons. *ASHA* 1989;31(08):59–63
- 37 Ventry IM, Weinstein BE. Identification of elderly people with hearing problems. *ASHA* 1983;25(07):37–42
- 38 Weinstein BE, Spitzer JB, Ventry IM. Test-retest reliability of the Hearing Handicap Inventory for the Elderly. *Ear Hear* 1986;7(05): 295–299
- 39 Newman CW, Weinstein BE. Test-retest reliability of the Hearing Handicap Inventory for the Elderly using two administration approaches. *Ear Hear* 1989;10(03):190–191
- 40 Wieselberg MB. A auto-avaliação do *handicap* em idosos portadores de deficiência auditiva: o uso do H.H.I.E. [dissertação]. São Paulo (SP): Pontifícia Universidade Católica de São Paulo; 1997:108
- 41 de Paiva SM, Simões J, Paiva A, Newman C, Castro E Sousa F, Bébér JP. Validity and Reliability of the Hearing Handicap Inventory for Elderly: Version Adapted for Use on the Portuguese Population. *J Am Acad Audiol*. 2016 Sep; 27(08):677–682
- 42 Cassol M, Ferreira MIDC, Pogliana DMA. A utilização do questionário HHIE-S associado à avaliação audiológica e vocal num grupo de idosos. *Estud Interdiscipl Envelhec*. 2007;12(01):81–95
- 43 Sindhusake D, Mitchell P, Smith W, et al. Validation of self-reported hearing loss. The Blue Mountains Hearing Study. *Int J Epidemiol* 2001;30(06):1371–1378
- 44 Gopinath B, Hickson L, Schneider J, et al. Hearing-impaired adults are at increased risk of experiencing emotional distress and social engagement restrictions five years later. *Age Ageing* 2012;41(05): 618–623
- 45 Dalton DS, Cruickshanks KJ, Klein BEK, Klein R, Wiley TL, Nondahl DM. The impact of hearing loss on quality of life in older adults. *Gerontologist* 2003;43(05):661–668
- 46 Pugh KC, Crandell CC. Hearing loss, hearing handicap, and functional health status between African American and Caucasian American seniors. *J Am Acad Audiol* 2002;13(09):493–502
- 47 Mattos LC, Veras RP. A prevalência da perda auditiva em uma população de idosos da cidade do Rio de Janeiro: um estudo seccional. *Rev Bras Otorrinolaryngol* 2007;73(05):654–659
- 48 Yamamoto CH, Ferrari DV. Relação entre limiares audiométricos, handicap e tempo para procura de tratamento da deficiência auditiva. *Rev Soc Bras Fonoaudiol*. 2012;17(02):135–141
- 49 Jupiter T. Screening for hearing loss in the elderly using distortion product otoacoustic emissions, pure tones, and a self-assessment tool. *Am J Audiol* 2009;18(02):99–107
- 50 Gopinath B, Schneider J, Hickson L, et al. Hearing handicap, rather than measured hearing impairment, predicts poorer quality of life over 10 years in older adults. *Maturitas* 2012;72(02):146–151
- 51 Saito H, Nishiwaki Y, Michikawa T, et al. Hearing handicap predicts the development of depressive symptoms after 3 years in older community-dwelling Japanese. *J Am Geriatr Soc* 2010; 58(01):93–97
- 52 John AB, Kreisman BM, Pallett S. Validity of hearing impairment calculation methods for prediction of self-reported hearing handicap. *Noise Health* 2012;14(56):13–20
- 53 Wu HY, Chin JJ, Tong HMM. Screening for hearing impairment in a cohort of elderly patients attending a hospital geriatric medicine service. *Singapore Med J* 2004;45(02):79–84
- 54 Tomioka K, Ikeda H, Hanaie K, et al. The Hearing Handicap Inventory for Elderly-Screening (HHIE-S) versus a single question: reliability, validity, and relations with quality of life measures in the elderly community, Japan. *Qual Life Res* 2013;22(05):1151–1159
- 55 Chou R, Dana T, Bougatsos C, Fleming C, Beil T. Screening adults aged 50 years or older for hearing loss: a review of the evidence for the U.S. preventive services task force. *Ann Intern Med* 2011; 154(05):347–355
- 56 Chang HP, Ho CY, Chou P. The factors associated with a self-perceived hearing handicap in elderly people with hearing impairment—results from a community-based study. *Ear Hear* 2009; 30(05):576–583

- 57 Russo ICP. Reabilitação Auditiva de Idosos. Bevilacqua MC, Martinez MAN, Balen SA, Pupo AC, Reis ACMB, Frota S. Tratado de Audiologia. 1ª ed. São Paulo(SP): Ed. Santos; 2012:775-87
- 58 Freitas CDD, Costa MJ. Processo de adaptação de próteses auditivas em usuários atendidos em uma instituição pública federal - parte II: resultados dos questionários de auto-avaliação. Rev Bras Otorrinolaringol 2007;73(05):660-670
- 59 Russo ICP, Pereira LD, Carvalho RMM, Anastásio ART. Encaminhamentos sobre a classificação do grau de perda auditiva em nossa realidade. Rev Soc Bras Fonoaudiol. 2009;14(02):287-288
- 60 World Health Organization (WHO). Prevention of blindness and deafness. Grades of hearing impairment [Internet]. Geneva: World Health Organization (WHO); 2016 [cited 2016 May 5]. Available from: [http://www.who.int/pbd/deafness/hearing\\_impairment\\_grades/en/](http://www.who.int/pbd/deafness/hearing_impairment_grades/en/)
- 61 Lichtenstein MJ, Bess FH, Logan SA. Validation of screening tools for identifying hearing-impaired elderly in primary care. JAMA 1988;259(19):2875-2878
- 62 Pereira MG. Epidemiologia: Teoria e Prática. 1ª ed. Rio de Janeiro (RJ): Guanabara Koogan; 1995
- 63 Fletcher RM, Fletcher SW, Wagner EH. Clinical Epidemiology, the essentials. 1st edition. Philadelphia: Lippincott Williams & Wilkins; 1983
- 64 Sestrem E. Avaliação da auto-percepção do *handicap* auditivo em idosos e percepção da fala: um estudo comparativo [dissertação]. Curitiba (PR): Universidade Tuiuti do Paraná; 2000:127
- 65 Salonen J, Johansson R, Karjalainen S, Vahlberg T, Isoaho R. Relationship between self-reported hearing and measured hearing impairment in an elderly population in Finland. Int J Audiol 2011;50(05):297-302
- 66 Wiley TL, Cruickshanks KJ, Nondahl DM, Tweed TS. Self-reported hearing handicap and audiometric measures in older adults. J Am Acad Audiol 2000;11(02):67-75
- 67 Chew HS, Yeak S. Quality of life in patients with untreated age-related hearing loss. J Laryngol Otol 2010;124(08):835-841
- 68 Hands S. Hearing loss in over-65s: is routine questionnaire screening worthwhile? J Laryngol Otol 2000;114(09):661-666
- 69 Macedo LS, Pupo AC, Balieiro CR. Aplicabilidade dos Questionários de Auto-Avaliação em Adultos e Idosos com Deficiência Auditiva. Distúrb Comun 2006;18(01):19-25
- 70 Aiello CP, Lima II, Ferrari DV. Validity and reliability of the hearing handicap inventory for adults. Rev Bras Otorrinolaringol (Engl Ed) 2011;77(04):432-438
- 71 Deepthi R, Kasthuri A. Validation of the use of self-reported hearing loss and the Hearing Handicap Inventory for elderly among rural Indian elderly population. Arch Gerontol Geriatr 2012;55(03):762-767
- 72 Camargo MK. Percepção da desvantagem auditiva em idosos - estudo da sensibilidade e especificidade do HHIE-S [monografia de conclusão do Curso de Especialização em Audiologia]. Porto Alegre (RS): Universidade Federal do Rio Grande do Sul; 2013:42
- 73 Tavares PF. Perda auditiva no idoso - suas interferências na vida psicossocial [monografia de conclusão do Curso de Especialização em Audiologia Clínica]. Itajaí (SC): Centro de Especialização em Fonoaudiologia Clínica; 2001:30