

Uncoupling neurogenic gene networks in the *Drosophila* embryo

William A. Rogers,^{1,3} Yogesh Goyal,^{2,3}
Kei Yamaya,¹ Stanislav Y. Shvartsman,²
and Michael S. Levine¹

¹Department of Molecular Biology, Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, New Jersey 08544, USA; ²Department of Chemical and Biological Engineering, Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, New Jersey 08544, USA

The EGF signaling pathway specifies neuronal identities in the *Drosophila* embryo by regulating developmental patterning genes such as *intermediate neuroblasts defective* (*ind*). EGFR is activated in the ventral midline and neurogenic ectoderm by the Spitz ligand, which is processed by the Rhomboid protease. CRISPR/Cas9 was used to delete defined *rhomboid* enhancers mediating expression at each site of Spitz processing. Surprisingly, the neurogenic ectoderm, not the ventral midline, was found to be the dominant source of EGF patterning activity. We suggest that *Drosophila* is undergoing an evolutionary transition in central nervous system (CNS)-organizing activity from the ventral midline to the neurogenic ectoderm.

Supplemental material is available for this article.

Received February 5, 2017; revised version accepted March 20, 2017.

Transcriptional control of ligand production is important for the spatiotemporal patterning of cell fate during embryogenesis (Marinić et al. 2013). Limiting components of signaling pathways, such as a ligand or ligand-processing enzyme, are often controlled by multiple enhancers mediating distinct patterns of expression. There are many examples of complex gene control, including the FGF ligands *Thisbe* and *Branchless* and the BMP ligands *Dpp* and *Gbb* (Housden and Perrimon 2014). However, the specific contributions of individual enhancers to any given signaling process remain unclear. Here we address this question during the patterning of the neurogenic ectoderm in the *Drosophila* embryo, which is controlled by two EGF ligands: *Vein* and *Spitz*.

Vein is secreted from its site of synthesis, whereas *Spitz* requires processing by the membrane-bound *Rhomboid* protease. *Vein* and *Spitz* induce the expression of regulatory genes required for the specification of ventral and lateral neurons (Mayer and Nüsslein-Volhard 1988; Rutledge et al. 1992; Raz and Shilo 1993; Schweitzer et al. 1995; Schnepf et al. 1996; Golembo et al. 1996). High levels of

EGF/ERK signaling trigger the expression of the *ventral nervous system defective* (*vnd*) determinant in ventral regions of the neurogenic ectoderm, while low levels activate *intermediate neuroblasts defective* (*ind*) in lateral regions (Ohlen and Doe 2000; Stathopoulos and Levine 2005; Ajuria et al. 2011; Lim et al. 2013, 2015).

Most previous studies on the establishment of this EGF signaling gradient have emphasized the paracrine effects of EGF ligands emanating from the ventral midline (Golembo et al. 1996; Chang et al. 2001; Rouso et al. 2010). However, both *vein* (*vn*) and *rhomboid* (*rho*) are also expressed in ventral regions of the presumptive neurogenic ectoderm (“lateral stripes”) prior to the onset of gastrulation (Kosman et al. 1991; Ip et al. 1992; González-Crespo and Levine 1993; Markstein et al. 2004). It has not been possible to assess the relative contributions of the two sites of EGF synthesis, the ventral midline and lateral stripes, in the patterning of the central nervous system (CNS).

Past studies identified separate enhancers for the regulation of *rho* expression in lateral stripes and the ventral midline. Midline expression depends on two separate enhancers, the midline enhancer (MLE) and a putative shadow enhancer (SHA) (Kvon et al. 2014; Pearson and Crews 2014), which is consistent with the view that the midline source of the *Vein* and *Spitz* ligands is particularly important for the patterning of the CNS. The role of the lateral stripe pattern, which is controlled by a well-characterized neurogenic ectoderm enhancer (NEE), remains uncertain (Supplemental Table 1).

Here, we used newly developed CRISPR/Cas9 methods to uncouple EGF synthesis in lateral stripes and the ventral midline (Barrangou et al. 2007; Jinek et al. 2012; Gratz et al. 2013). Mutant embryos harboring deletions of defined *rho* enhancers were combined with *vn* mutants, since past studies have shown that *Spitz* and *Vein* function in a somewhat redundant fashion to activate *vnd* and *ind* expression in ventral and lateral regions of the developing ventral nerve cord (Skeath 1998; Lim et al. 2015). *Spitz* activity is fully abolished by the loss of *rho* activity, and, consequently, *rho;vn* double mutants exhibit a complete loss of *ind* expression due to the absence of both *Vein* and activated *Spitz* (Hong et al. 2008).

We were therefore able to obtain mutant embryos containing a midline-only source of EGF signaling by the targeted deletion of the *rho* NEE in *vn⁻vn⁻* mutant embryos. These embryos exhibit incomplete patterns of *ind* expression, suggesting that the midline is not sufficient to pattern the CNS. Surprisingly, the reciprocal experiment, restricted processing of *Spitz* in lateral stripes, produces an essentially normal *ind* expression pattern. These findings underscore the importance of *rho* and *vn* lateral stripes in the formation of the EGF signaling gradient and patterning of the CNS. Genome-editing methods provide a powerful means for deleting defined enhancers within endogenous loci to uncouple the complex

[Keywords: *Drosophila* embryo; EGFR regulatory networks; *rhomboid*; *intermediate neuroblasts defective*; central nervous system]

³These authors contributed equally to this work.

Corresponding authors: msl2@princeton.edu, stas@princeton.edu

Article published online ahead of print. Article and publication date are online at <http://www.genesdev.org/cgi/doi/10.1101/gad.297150.117>.

© 2017 Rogers et al. This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genesdev.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

regulatory control of critical developmental patterning genes such as *rho*.

Results and Discussion

Differential activation of *vnd* and *ind* in ventral and lateral regions of the neurogenic ectoderm is controlled by Spitz and Vein EGF ligands (Ohlen and Doe 2000; Stathopoulos and Levine 2005; Ajuria et al. 2011; Lim et al. 2013, 2015) emanating from ventral regions of the neurogenic ectoderm (summarized in Fig. 1A). We used CRISPR/Cas9 genome-editing methods to determine the relative contributions of EGF signaling in lateral stripes (ventral neurogenic ectoderm) and the ventral midline by deleting defined *rho* enhancers: MLE, NEE, and SHA (summarized in Fig. 1B; Barrangou et al. 2007; Jinek et al. 2012; Gratz et al. 2013). The SHA and MLE mediate expression in the midline in response to the Sim activator, whereas the NEE activates *rho* in the presumptive neurogenic ectoderm (lateral stripes) in response to the maternal dorsal gradient.

Midline-specific EGF signaling was obtained by deleting the *rho* NEE. Mutant embryos exhibit a slight delay in the onset of *ind* expression but nonetheless grow to adulthood (Supplemental Fig. S1A,B''). Similar patterns of expression were also observed in *rho*⁻/*rho*⁻ mutant embryos (Skeath 1998; Hong et al. 2008; Lim et al. 2013) as well as mutants harboring different combinations of *rho* enhancer deletions (Supplemental Fig. S1). These observations are consistent with previous findings that Vein alone is sufficient for the normal induction of *ind* expression (Supplemental Fig. S1). In order to assess the individual contributions of *rho*⁺ gene activity in the midline and neurogenic ectoderm, it was necessary to introduce *rho* NEE deficiency homozygotes into a *vn*⁻/*vn*⁻ mutant background.

The resulting double mutants, *rho* ΔNEE;*vn*⁻/*rho* ΔNEE;*vn*⁻, completely lack both *rho* and *vn* lateral stripes (e.g., Fig. 1F). The only source of EGF signaling arises from the midline-specific expression of *rho* and localized pro-

cessing of Spitz (Fig. 1E,H). Double-labeling methods using *rho* and *ind* RNA probes indicate a significant delay in the activation of *ind* (Fig. 1, cf. D and G). Moreover, the late *ind* expression pattern is incomplete and displays considerable variation between embryos (Fig. 1H). These observations suggest that the midline processing of the Spitz ligand is not sufficient for proper induction of *ind* expression and patterning of the nerve cord.

Most previous studies of EGF-mediated patterning of the CNS emphasized the role of the ventral midline as the source of ligand production (Golembo et al. 1996; Chang et al. 2001; Rousso et al. 2010). However, embryos with individual deletions of the MLE and SHA exhibit normal *rho* and *ind* expression patterns (Supplemental Fig. S1A–A',C–D''). To more accurately assess the role of the midline, we produced embryos that selectively produce only lateral stripes of Spitz activity. This was achieved by creating *sim*;*vn* double-mutant embryos, which have transient lateral stripes of *rho* expression in the absence of midline activity (Fig. 2). *Sim* encodes a basic helix–loop–helix (bHLH)–PAS activator that functions as a “master regulator” of ventral midline differentiation. It is expressed in the presumptive midline prior to the onset of gastrulation, where it coordinates the expression of >50 different target genes, including *rho* and *vein*. *sim*⁻/*sim*⁻ mutant embryos display a severe loss of *rho* and *vn* expression in the ventral midline (Crews et al. 1988; Thomas et al. 1988; Nambu et al. 1990, 1991; Chang et al. 2001; Kearney et al. 2004; Hong et al. 2013).

sim;*vn* double mutants produce active Spitz ligands in lateral stripes due solely to the transient activity of the *rho* NEE. The initial induction of *ind* expression is virtually indistinguishable from that seen in normal embryos (e.g., Fig. 2A,D) despite the loss of midline targets of EGF signaling such as *orthodenticle* (Supplemental Fig. S2). We expected premature loss of *ind* expression due to the absence of sustained *rho* expression in the midline. However, *ind* expression was found to persist in advanced-stage embryos long after the loss of the transient *rho* lateral stripes (Fig. 2E,F). This pattern is significantly less variable than that seen when Spitz is processed solely in the mid-

line (Fig. 1H). We therefore conclude that transient lateral stripes of EGF signaling, rather than stable midline processing, provide the dominant source of EGF activity in the patterning of the *Drosophila* CNS (see below).

The maintenance of the *ind* expression pattern in *sim*;*vn* double mutants raises the possibility that autoregulatory mechanisms might be responsible for sustained expression at later stages of embryogenesis (Von Ohlen et al. 2007). For example, positive autofeedback is responsible for maintaining *sim* expression in the ventral midline of advanced-stage embryos (Kasai et al. 1992). To explore whether a similar mechanism might be used to regulate *ind*, we induced ectopic EGF signaling by placing the *rho* coding sequence under the control of *ind* regulatory sequences (Fig. 3).

Normally, *rho* lateral stripes straddle the ventral border of the *ind* expression pattern (e.g., Fig. 1C). The use of *ind* regulatory sequences causes a significant expansion of the *rho* expression pattern into more dorsal




Figure 1. Late *rho* enhancers insufficient for normal *ind* expression. (A) The EGFR pathway with arrows showing activation. (B) The locations of known *rho* enhancers in relation to the genes. Black bars are the enhancers for the *rho* gene (purple bar). The dotted line indicates intergenic region. (C–H) The *rho* ΔNEE;*vn* double mutant is orientated with anterior to the left and posterior to the right. Images from early NC14 (C,F), gastrulation (D,G), and germ band elongation (E,H). (C–E) Heterozygous mutants. (F–H) Homozygous mutants. (Magenta lines) *rho* expression; (anterior magenta) *hb-lacZ* was used to visualize heterozygous mutants; (green lines) *ind* expression; (gray dots) DAPI. Bar, 50 μm.




Figure 2. Early *rho* expression leads to stable late *ind* expression. The *sim;vn* heterozygous mutants (A–C) and homozygous mutants (D–F) are orientated with anterior to the left and posterior to the right. Images are from early NC14 (A,D), gastrulation (B,E), and germ band elongation (C,F). (Magenta lines) *rho* expression. (A–C) (Anterior magenta) *hb-lacZ* was used to visualize heterozygous mutants; (green lines) *ind* expression; (gray dots) DAPI. Bar, 50 μ m.

regions of the neurogenic ectoderm. This expanded pattern leads to ectopic EGF signaling and dpERK activity in lateral regions of the developing nerve cord (Fig. 3, cf. A and E). The initial *ind* expression pattern is considerably broader than the wild-type pattern (Fig. 3, cf. C and G) but nonetheless refines to a nearly normal pattern following gastrulation (Fig. 3, cf. D and H). This refinement occurs in spite of expanded EGF signaling, suggesting the occurrence of compensatory mechanisms of activation and repression of *ind* expression during development.

CRISPR/Cas9 genome-editing methods were used for the selective removal of defined developmental enhancers controlling the complex *rho* expression pattern during *Drosophila* embryogenesis (summarized in Supplemental Table 2). This approach has the potential to reveal new mechanisms of gene control. For example, the larval cuticles of *rho* Δ NEE, *vn* and *sim;vn* show no morphological defects (Supplemental Fig. S3), yet the removal of the 5' *rho* SHA and NEE appears to cause a delay in the onset of *rho* MLE activity in the ventral midline (Supplemental Fig. S4). The specific deletion of the *rho* NEE led to an incomplete and variable *ind* expression pattern, suggesting that the ventral midline may not be the central organizer of CNS patterning suggested by earlier studies (Golemboski et al. 1996; Chang et al. 2001; Rouso et al. 2010). Instead, we showed that transient lateral stripes of EGF signaling are sufficient to induce *ind* expression, while subsequent compensatory mechanisms maintain this expression during embryogenesis.

The ventral midline is thought to be the ancestral mode of EGF signaling in the developing CNS of insects and other arthropods. For example, the ventral midline of the crustacean *Parhyale* plays a dominant role in the patterning of the CNS (Vargas-Vila et al. 2010). Disruption of *sim*⁺ gene activity or ablation of midline cells causes severe patterning defects, whereas *sim* mutants in *Drosophila* exhibit essentially normal patterning of the nerve cord (except for the differentiation of mesectoderm cells arising from the midline) (Vargas-Vila et al. 2010; Lynch and Roth 2011). We propose that *Drosophila* represents a transitional patterning system in which the ancestral midline mode of EGF signaling is replaced by lateral

stripes (Fig. 4). Rhomboid does not exhibit obvious lateral stripes of expression in other insects such as flour beetles and mosquitoes. This novel pattern of expression appears to coincide with the duplication of *rhomboid* genes in the Drosophilids (Crocker et al. 2010; Rouso et al. 2010).

Drosophila appears to use two distinct gene regulatory networks for deploying EGF signaling during embryogenesis. The expression of EGF signaling components in the ventral midline depends on the *Sim* activator, which in turn is induced by Notch signaling (Zinzen et al. 2006). In contrast, lateral stripes of *rho* and *vn* expression depend on dorsal and bHLH activators, including proneural determinants such as Daughterless and Achaete/Scute (González-Crespo and Levine 1993). The latter mode of EGF signaling is dominant in *Drosophila*, but the retention of midline patterning activity might render the system robust to genetic and environmental variation. This view of network evolution, the co-option of novelty while retaining ancestral mechanisms, might be a common property of animal development.

Materials and methods

CRISPR–Cas9 (Supplemental Tables 3–5)

Guide RNA (gRNA) The protocol was followed based on the publication by the O'Connor-Giles laboratory, and the unique PAM recognition sites were designed with the CRISPR optimal target finder (<http://tools.flycrispr.molbio.wisc.edu/targetFinder>; Gratz et al. 2014). The two unique PAM recognition sites were inserted into individual pU6b-gRNA vectors by BbsI. PAM sites were verified in the injection line. The pU6b-gRNA was obtained from the Perrimon laboratory (Supplemental Material; Ren et al. 2013).

Donor vectors The p2xattP-dsRed donor vector was modified from the O'Connor-Giles laboratory pHD-DsRed-attP vector by the addition of an inverted attP site 3' of the dsRed cassette (Supplemental Material). An ~1-kb 5' homology arm was inserted with NheI and SacII. An ~1-kb 3' homology arm was inserted with SbfI and AscI (modified MCS). The GFP donor vector (gift from Thomas Gregor's laboratory) used XhoI/AscI and AgeI/NotI for the 5' and 3' homology arms, respectively. CRISPR vectors (donor vector and two unique guide vectors) were injected into nos-Cas9 (*y sc v;nos-Cas9[attP40/CyO]* [Bestgene: TH00788.N]). The double enhancer deletions were produced by injecting previous dsRed CRISPR fly lines with the GFP donor vector (Supplemental Material), two unique




Figure 3. Early ectopic *ind* expression compensates for normal late expression. (A–D) Wild type (WT). (E–H) Homozygous *rho* overexpression. Images were taken at gastrulation (A,E,G) and germ band extension (B,F,D,H). All images are anterior to posterior. (Red lines) dpERK expression; (purple lines) *rho* expression; (green lines) *ind* expression; (gray dots) DAPI. Bar, 50 μ m.




Figure 4. The ancestral *rho* MLE shifts its role to NEE for midline *ind* expression. (A) Mosquitos (*Anopheles gambiae*) rely primarily on the midline expression of *rho* for late *ind* expression. (B) Fruit fly (*Drosophila melanogaster*) *ind* regulation has a unique lateral expression followed by the ancestral late stage mediated mostly by *rho*. (C) *A. gambiae* has no lateral regulation of *ind* (X) and roles of *rho* SHA, and *ind* autoregulation are unknown (?). *D. melanogaster* seems to have acquired a major role for the *rho* NEE (large arrow) for regulating *ind* (arrow) while minimizing the role of the midline *rho* MLE and SHA (small arrows). (Purple) *rho*; (orange) *vn*; (green) *ind*.

guide vectors, and the nos-Cas9 plasmid. All lines were PCR-validated and imaged. All injections were performed by Bestgene. Primers were made by IDT.

Immunostaining and fluorescent in situ hybridization (FISH)

Immunostaining and FISH protocols were performed as described elsewhere (Lim et al. 2015). Sheep anti-digoxigenin (DIG; 1:125; Roche), mouse anti-biotin (1:125; Jackson ImmunoResearch), and rabbit anti-dpERK (1:100; CST4370) were used as primary antibodies. DAPI (1:10,000; Molecular Probes) was used to stain for nuclei, and Alexa fluor conjugates (1:500; Invitrogen) were used as secondary antibodies.

Fly recombination

;;vn, sim The *vn*^{L6}/TM3, *hb-lacZ* was crossed to *sp*⁺/*sim*²/TM3, *hb-lacZ*. The virgin female *vn/sim* flies were crossed to *dsRed*/TM3, *hb-lacZ*. Recombination was assayed by crossing individual recombinant males to virgins of each mutant line. Validated lines exhibited TM3 from the assay (*n* = 137).

;;vn, rho ΔNEE The *vn*^{L6}/TM3, *hb-lacZ* line was crossed to *rho* ΔNEE/TM3, *hb-lacZ*. The female virgin *vn/rho* ΔNEE flies were crossed to TM3, *hb-lacZ*. Individual males were tested for recombination by crossing to virgin *vn* mutants. Validated lines exhibited both TM3 and *dsRed* from the assays (*n* = 63).

Microscopy and image processing

Fluorescent imaging was performed on Nikon A1-RS scanning confocal (Fig. 1, 2) and Zeiss 880 (Fig. 3) microscopes with a 20× objective. For pairwise comparisons of wild-type and mutant backgrounds, embryos were collected, stained, and imaged together under the same experimental conditions. Broken embryos, embryos with intact vitelline membrane, or embryos undergoing mitosis were excluded from the analysis. The contrast and brightness for images were adjusted in ImageJ relative to the wild-type images for the final figure panels, thus allowing quantitative comparisons between two sets of images.

Acknowledgments

We thank members of the Levine and Shvartsman laboratories for their invaluable comments and advice. We also thank Hongtao Chen and Thomas Gregor for providing many of the CRISPR/Cas9 plasmids used in our experiments, and Gary Laevsky and Evangelos G. Gatzogiannis for their expert help with imaging. W.A.R. was supported by a National Institutes of Health (NIH) grant to M.S.L. (R35 GM118147). Y.G. and K.Y. were supported by an NIH grant to S.Y.S. (R01 GM086537). W.A.R., Y.G., S.Y.S., and M.S.L. conceived and designed the project; W.A.R. designed the CRISPR/Cas9 mutant flies; Y.G., W.A.R., and K.Y. performed the imaging experiments; W.A.R. and Y.G. analyzed the results; and W.A.R., Y.G., S.Y.S., and M.S.L. wrote the manuscript.

References

- Ajuria L, Nieva C, Winkler C, Kuo D, Samper N, Andreu MJ, Helman A, González-Crespo S, Paroush Z, Courey AJ, et al. 2011. Capicua DNA-binding sites are general response elements for RTK signaling in *Drosophila*. *Development* **924**: 915–924.
- Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, Romero DA, Horvath P. 2007. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **315**: 1709–1712.
- Chang J, Kim IO, Ahn JS, Kim SH. 2001. The CNS midline cells control the spitz class and Egr signaling genes to establish the proper cell fate of the *Drosophila* ventral neuroectoderm. *Int J Dev Biol* **45**: 715–724.
- Crews ST, Thomas JB, Goodman CS. 1988. The *Drosophila* single-minded gene encodes a nuclear protein with sequence similarity to the per gene product. *Cell* **52**: 143–151.
- Crocker J, Potter N, Erives A. 2010. Dynamic evolution of precise regulatory encodings creates the clustered site signature of enhancers. *Nat Commun* **1**: 99.
- Golembo M, Raz E, Shilo BZ. 1996. The *Drosophila* embryonic midline is the site of Spitz processing, and induces activation of the EGF receptor in the ventral ectoderm. *Development* **122**: 3363–3370.
- González-Crespo S, Levine M. 1993. Interactions between dorsal and helix-loop-helix proteins initiate the differentiation of the embryonic mesoderm and neuroectoderm in *Drosophila*. *Genes Dev* **7**: 1703–1713.
- Gratz SJ, Cummings AM, Nguyen JN, Hamm DC, Donohue LK, Harrison MM, Wildonger J, O'Connor-Giles KM. 2013. Genome engineering of *Drosophila* with the CRISPR RNA-guided Cas9 nuclease. *Genetics* **194**: 1029–1035.
- Gratz SJ, Ukken FP, Rubinstein CD, Thiede G, Donohue LK, Cummings AM, O'Connor-Giles KM. 2014. Highly specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in *Drosophila*. *Genetics* **196**: 961–971.
- Hong J-W, Hendrix DA, Papatzenko D, Levine MS. 2008. How the dorsal gradient works: insights from postgenome technologies. *Proc Natl Acad Sci* **105**: 20072–20076.
- Hong J-W, Park KW, Levine MS. 2013. Temporal regulation of single-minded target genes in the ventral midline of the *Drosophila* central nervous system. *Dev Biol* **380**: 335–343.
- Housden BE, Perrimon N. 2014. Spatial and temporal organization of signaling pathways. *Trends Biochem Sci* **39**: 457–464.
- Ip YT, Park RE, Kosman D, Bier E, Levine M. 1992. The dorsal gradient morphogen regulates stripes of rhomboid expression in the presumptive neuroectoderm of the *Drosophila* embryo. *Genes Dev* **6**: 1728–1739.
- Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. 2012. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**: 816–821.
- Kasai Y, Nambu JR, Lieberman PM, Crews ST. 1992. Dorsal-ventral patterning in *Drosophila*: DNA binding of snail protein to the single-minded gene. *Proc Natl Acad Sci* **89**: 3414–3418.
- Kearney JB, Wheeler SR, Estes P, Parente B, Crews ST. 2004. Gene expression profiling of the developing *Drosophila* CNS midline cells. *Dev Biol* **275**: 473–492.
- Kosman D, Ip YT, Levine M, Arora K. 1991. Establishment of the mesoderm-neuroectoderm boundary in the *Drosophila* embryo. *Science* **254**: 118–122.

- Kvon EZ, Kazmar T, Stampfel G, Yáñez-Cuna JO, Pagani M, Schernhuber K, Dickson BJ, Stark A. 2014. Genome-scale functional characterization of *Drosophila* developmental enhancers in vivo. *Nature* **517**: 91–95.
- Lim B, Samper N, Lu H, Rushlow C, Jiménez G, Shvartsman SY. 2013. Kinetics of gene derepression by ERK signaling. *Proc Natl Acad Sci* **110**: 10330–10335.
- Lim B, Dsilva CJ, Levario TJ, Lu H, Schüpbach T, Kevrekidis IG, Shvartsman SY. 2015. Dynamics of inductive ERK signaling in the *Drosophila* embryo. *Curr Biol* **25**: 1784–1790.
- Lynch JA, Roth S. 2011. The evolution of dorsal–ventral patterning mechanisms in insects. *Genes Dev* **25**: 107–118.
- Miranić M, Aktas T, Ruf S, Spitz F. 2013. An integrated holo-enhancer unit defines tissue and gene specificity of the Fgf8 regulatory landscape. *Dev Cell* **24**: 530–542.
- Markstein M, Zinzen R, Markstein P, Yee K, Erives A, Stathopoulos A, Levine M. 2004. A regulatory code for neurogenic gene expression in the *Drosophila* embryo. *Development* **131**: 2387–2394.
- Mayer U, Nüsslein-Volhard C. 1988. A group of genes required for pattern formation in the ventral ectoderm of the *Drosophila* embryo. *Genes Dev* **2**: 1496–1511.
- Nambu JR, Franks RG, Hu S, Crews ST. 1990. The single-minded gene of *Drosophila* is required for the expression of genes important for the development of CNS midline cells. *Cell* **63**: 63–75.
- Nambu JR, Lewis JO, Wharton KA, Crews ST. 1991. The *Drosophila* single-minded gene encodes a helix–loop–helix protein that acts as a master regulator of CNS midline development. *Cell* **67**: 1157–1167.
- Pearson JC, Crews ST. 2014. Enhancer diversity and the control of a simple pattern of *Drosophila* CNS midline cell expression. *Dev Biol* **392**: 466–482.
- Raz E, Shilo BZ. 1993. Establishment of ventral cell fates in the *Drosophila* embryonic ectoderm requires DER, the EGF receptor homolog. *Genes Dev* **7**: 1937–1948.
- Ren X, Sun J, Housden BE, Hu Y, Roesel C, Lin S, Liu L-P, Yang Z, Mao D, Sun L, et al. 2013. Optimized gene editing technology for *Drosophila melanogaster* using germ line-specific Cas9. *Proc Natl Acad Sci* **110**: 19012–19017.
- Rouso T, Lynch J, Yogev S, Roth S, Schejter ED, Shilo B. 2010. Generation of distinct signaling modes via diversification of the Egr ligand-processing cassette. *Development* **137**: 3427–3437.
- Rutledge BJ, Zhang K, Bier E, Jan YN, Perrimon N. 1992. The *Drosophila* spitz gene encodes a putative EGF-like growth factor involved in dorsal–ventral axis formation and neurogenesis. *Genes Dev* **6**: 1503–1517.
- Schnepp B, Grumbling G, Donaldson T, Simcox A. 1996. Vein is a novel component in the *Drosophila* epidermal growth factor receptor pathway with similarity to the neuregulins. *Genes Dev* **10**: 2302–2313.
- Schweitzer R, Shaharabany M, Seger R. 1995. Secreted Spitz triggers the DER signaling pathway and is a limiting component in embryonic ventral ectoderm determination. *Genes Dev* **9**: 1518–1529.
- Sketh JB. 1998. The *Drosophila* EGF receptor controls the formation and specification of neuroblasts along the dorsal–ventral axis of the *Drosophila* embryo. *Development* **125**: 3301–3312.
- Stathopoulos A, Levine M. 2005. Localized repressors delineate the neurogenic ectoderm in the early *Drosophila* embryo. *Dev Biol* **280**: 482–493.
- Thomas JB, Crews ST, Goodman CS. 1988. Molecular genetics of the single-minded locus: a gene involved in the development of the *Drosophila* nervous system. *Cell* **52**: 133–141.
- Vargas-Vila MA, Hannibal RL, Parchem RJ, Liu PZ, Patel NH. 2010. A prominent requirement for single-minded and the ventral midline in patterning the dorsoventral axis of the crustacean *Parhyale hawaiiensis*. *Development* **137**: 3469–3476.
- Von Ohlen T, Doe CQ. 2000. Convergence of dorsal, Dpp, and Egr signaling pathways subdivides the *Drosophila* neuroectoderm into three dorsal–ventral columns. *Dev Biol* **224**: 362–372.
- Von Ohlen TL, Harvey C, Panda M. 2007. Identification of an upstream regulatory element reveals a novel requirement for Ind activity in maintaining ind expression. *Mech Dev* **124**: 230–236.
- Zinzen RP, Cande J, Ronshaugen M, Papatsenko D, Levine M. 2006. Evolution of the ventral midline in insect embryos. *Dev Cell* **11**: 895–902.