

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Di- μ -chlorido-bis(chlorido{ N' -[phenyl-(pyridin-2-yl- κN)methylidene]pyridine-2-carbohydrazide- $\kappa^2 N', O$ }cadmium)

Mehmet Akkurt,^{a*} Ali Akbar Khandar,^b
Muhammad Nawaz Tahir,^c Farhad Akbari Afkhami^b and
Seyed Abolfazl Hosseini Yazdi^b

^aDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bDepartment of Inorganic Chemistry, Faculty of Chemistry, University of Tabriz, Tabriz, Iran, and ^cDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: akkurt@erciyes.edu.tr


Received 5 May 2014; accepted 9 May 2014

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.044; wR factor = 0.083; data-to-parameter ratio = 20.0.

The title compound, $[Cd_2Cl_4(C_{18}H_{14}N_4O)_2]$, was obtained from the reaction of $Cd(NO_3)_2 \cdot 4H_2O$ with 2-phenylpyridine-keton picolinoyl hydrazone and sodium chloride. Each Cd^{2+} cation is coordinated by two N atoms and one O atom of the tridentate ligand and three chloride anions, forming a distorted $CdNOCl_3$ octahedron. Each pair of adjacent metal cations is linked by two bridging chloride ligands, resulting in a dinuclear complex unit. The molecular conformation is stabilized by intramolecular $N-H \cdots N$ and $C-H \cdots O$ hydrogen bonds. In the crystal, molecules are linked by nonclassical $C-H \cdots O$ and $C-H \cdots Cl$ hydrogen bonds into a three-dimensional network. In addition, $\pi-\pi$ stacking interactions [centroid-centroid distances = 3.777 (2) and 3.631 (2) Å] contribute to the stabilization of the crystal packing.

Related literature

For related complexes with similar tridentate ligands, see: Akkurt *et al.* (2012); Chen *et al.* (2005); Datta *et al.* (2011).



Experimental

Crystal data

$[Cd_2Cl_4(C_{18}H_{14}N_4O)_2]$
 $M_r = 971.28$
Monoclinic, $P2_1/n$
 $a = 15.3215$ (4) Å
 $b = 14.4905$ (4) Å
 $c = 17.5457$ (5) Å
 $\beta = 104.667$ (1)°

$V = 3768.49$ (18) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.46$ mm⁻¹
 $T = 296$ K
0.23 × 0.20 × 0.18 mm

Data collection

Bruker Kappa APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)
 $T_{min} = 0.722$, $T_{max} = 0.769$

36565 measured reflections
9369 independent reflections
5625 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.066$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.083$
 $S = 0.98$
9369 reflections

469 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.47$ e Å⁻³
 $\Delta\rho_{min} = -0.43$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N3—H3A \cdots N4	0.86	2.29	2.646 (4)	105
N7—H7 \cdots N8	0.86	2.16	2.569 (4)	108
C1—H1 \cdots O2	0.93	2.53	3.303 (6)	140
C10—H10 \cdots O1 ⁱ	0.93	2.50	3.261 (6)	139
C36—H36 \cdots Cl3 ⁱⁱ	0.93	2.77	3.482 (5)	135

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *S SAINT* (Bruker, 2009); data reduction: *S SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON*.

This work was supported by the University of Tabriz Research Council. The authors acknowledge the provision of funds for the purchase of a diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supporting information for this paper is available from the IUCr electronic archives (Reference: RZ5124).

References

- Akkurt, M., Khandar, A. A., Tahir, M. N., Yazdi, S. A. H. & Afkhami, F. A. (2012). *Acta Cryst.* **E68**, m842.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, L., Zhang, W., Huang, S., Jin, X. & Sun, W.-H. (2005). *Inorg. Chem. Commun.* **8**, 41–43.
- Datta, A., Das, K., Jhou, Y.-M., Huang, J.-H. & Lee, H. M. (2011). *Acta Cryst.* **E67**, m123.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2014). E70, m213–m214 [doi:10.1107/S1600536814010630]

Di- μ -chlorido-bis(chlorido{ N' -[phenyl(pyridin-2-yl- κ N)methylidene]pyridine-2-carbohydrazide- $\kappa^2 N', O$ }cadmium)

Mehmet Akkurt, Ali Akbar Khandar, Muhammad Nawaz Tahir, Farhad Akbari Afkhami and Seyed Abolfazl Hosseini Yazdi

1. Comment


Schiff base complexes have attracted much attention due to their interesting structures and wide potential applications. Recently, the relative unsymmetrical tri- and tetradentate Schiff base ligands and their hydrogenated derivatives have been introduced in coordination chemistry to assemble polymers with beautiful molecular structures. Some organic N-donor ligands are often chosen to fabricate these complexes. In this connection, some complexes with similar tridentate ligands have been studied (Chen *et al.*, 2005; Datta *et al.*, 2011; Akkurt *et al.*, 2012). Herein, we report the structure of a new cadmium complex based on a pyridine based chelating Schiff base ligand. The title compound shows chloride-bridged dinuclear Cd(II) units (Fig. 1). The geometry around each Cd(II) ion is distorted octahedral, in which three positions are occupied by two N atoms and one O atom from the Schiff base ligand, two positions by two bridging chloride anions and the sixth position by one terminal chloride anion. Intramolecular non-classical hydrogen bonds of the type C—H \cdots O and N—H \cdots N are present (Table 1). Non-classical intermolecular hydrogen bonds of type C—H \cdots O and C—H \cdots Cl link complexes into a three-dimensional network (Table 1). In the crystal, π - π stacking interactions also contribute to the stabilization: $Cg6\cdots Cg9$ ($1 - x, 1 - y, 1 - z$) = 3.777 (2) Å, $Cg7\cdots Cg9(-1/2 + x, 1/2 - y, 1/2 + z)$ = 3.631 (2) Å; where $Cg6$, $Cg7$ and $Cg9$ are the centroids of the N1/C1–C5, N4/C14–C18 and N8/C32–C36 pyridine rings, respectively.

2. Experimental

The potentially tridentate ligand 2-phenylpyridineketon picolinoyl hydrazone was obtained by condensation of an equimolar mixture of 2-phenylpyridineketon and picolinic acid hydrazide in methanol. The title compound was synthesized by the reaction of a methanolic solution of the ligand and $Cd(NO_3)_2 \cdot 4H_2O$ in the presence of excess amount of NaCl. The ligand (1 mmol, 0.302 g) and cadmium nitrate (1 mmol, 0.308 g) were placed in the main arm of the branched tube; sodium chloride (2 mmol, 0.117 g) was added to the mixture too. Methanol was carefully added to fill the arms. The tube was sealed and the ligand-containing arm was immersed in an oil bath at 333 K while the branched arm was kept at ambient temperature. After three days, suitable single crystals had deposited in the cooler arm which were isolated, filtered off, washed with acetone and ether and air dried.

3. Refinement

The structure was solved by the Patterson method. All H atoms were positioned geometrically with C—H = 0.93, N—H = 0.86 Å, and refined using a riding model with $U_{iso}(H) = 1.2U_{eq}(C, N)$.


Figure 1

The molecular structure of the title compound with displacement ellipsoids drawn at the 40% probability level.

Di- μ -chlorido-bis(chlorido{ N' -[phenyl(pyridin-2-yl- κ N)methylidene]pyridine-2-carbohydrazide- κ^2N',O }cadmium)

Crystal data

$[\text{Cd}_2\text{Cl}_4(\text{C}_{18}\text{H}_{14}\text{N}_4\text{O})_2]$

$M_r = 971.28$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 15.3215(4)\ \text{\AA}$

$b = 14.4905(4)\ \text{\AA}$

$c = 17.5457(5)\ \text{\AA}$

$\beta = 104.667(1)^\circ$

$V = 3768.49(18)\ \text{\AA}^3$

$Z = 4$

$F(000) = 1920$

$D_x = 1.712\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 310 reflections

$\theta = 3.5\text{--}18.2^\circ$

$\mu = 1.46\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Prism, orange

$0.23 \times 0.20 \times 0.18\ \text{mm}$

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.722$, $T_{\max} = 0.769$

36565 measured reflections

9369 independent reflections

5625 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.066$

$\theta_{\max} = 28.4^\circ$, $\theta_{\min} = 1.9^\circ$

$h = -20 \rightarrow 17$

$k = -19 \rightarrow 19$

$l = -23 \rightarrow 23$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.083$	$w = 1/[\sigma^2(F_o^2) + (0.0251P)^2]$
$S = 0.98$	where $P = (F_o^2 + 2F_c^2)/3$
9369 reflections	$(\Delta/\sigma)_{\max} = 0.002$
469 parameters	$\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.21922 (2)	0.30193 (2)	0.59332 (2)	0.0425 (1)
Cd2	0.21468 (2)	0.45128 (2)	0.41438 (2)	0.0418 (1)
C11	0.23181 (8)	0.27779 (7)	0.44471 (6)	0.0563 (4)
C12	0.15115 (7)	0.45550 (7)	0.53781 (6)	0.0542 (4)
C13	0.20787 (7)	0.13663 (7)	0.62103 (6)	0.0505 (4)
C14	0.24456 (7)	0.61788 (7)	0.40603 (7)	0.0599 (4)
O1	0.09558 (16)	0.32324 (19)	0.66423 (15)	0.0490 (10)
O2	0.38776 (16)	0.43618 (19)	0.42115 (15)	0.0518 (10)
N1	0.3800 (2)	0.3236 (2)	0.63954 (18)	0.0444 (11)
N2	0.2716 (2)	0.3262 (2)	0.73505 (17)	0.0411 (11)
N3	0.20746 (19)	0.3262 (2)	0.77667 (17)	0.0445 (11)
N4	0.0893 (2)	0.3191 (2)	0.8634 (2)	0.0569 (14)
N5	0.0819 (2)	0.4427 (2)	0.30909 (18)	0.0416 (11)
N6	0.2485 (2)	0.4326 (2)	0.29130 (17)	0.0393 (11)
N7	0.3364 (2)	0.4316 (2)	0.28770 (18)	0.0474 (13)
N8	0.4953 (2)	0.4259 (2)	0.26459 (18)	0.0482 (12)
C1	0.4346 (3)	0.3258 (3)	0.5914 (3)	0.0551 (17)
C2	0.5277 (3)	0.3262 (3)	0.6176 (3)	0.0616 (17)
C3	0.5663 (3)	0.3206 (3)	0.6964 (3)	0.0613 (18)
C4	0.5109 (3)	0.3168 (3)	0.7475 (2)	0.0544 (16)
C5	0.4181 (2)	0.3203 (3)	0.7178 (2)	0.0428 (14)
C6	0.3553 (2)	0.3207 (2)	0.7699 (2)	0.0387 (12)
C7	0.3922 (2)	0.3131 (3)	0.8569 (2)	0.0414 (14)
C8	0.4468 (3)	0.3800 (3)	0.8997 (2)	0.0660 (19)
C9	0.4863 (3)	0.3687 (4)	0.9794 (3)	0.080 (2)
C10	0.4692 (3)	0.2909 (4)	1.0169 (3)	0.074 (2)

C11	0.4167 (3)	0.2230 (3)	0.9757 (3)	0.0662 (17)
C12	0.3767 (3)	0.2335 (3)	0.8955 (2)	0.0543 (17)
C13	0.1197 (2)	0.3233 (3)	0.7360 (2)	0.0397 (12)
C14	0.0537 (3)	0.3196 (3)	0.7857 (2)	0.0431 (14)
C15	-0.0371 (3)	0.3151 (3)	0.7504 (3)	0.0533 (17)
C16	-0.0959 (3)	0.3084 (3)	0.7988 (3)	0.0675 (19)
C17	-0.0605 (4)	0.3073 (3)	0.8781 (3)	0.079 (2)
C18	0.0301 (4)	0.3131 (3)	0.9079 (3)	0.075 (2)
C19	-0.0018 (3)	0.4465 (3)	0.3178 (3)	0.0534 (17)
C20	-0.0766 (3)	0.4436 (3)	0.2548 (3)	0.0621 (18)
C21	-0.0667 (3)	0.4379 (3)	0.1802 (3)	0.0638 (17)
C22	0.0197 (3)	0.4354 (3)	0.1698 (2)	0.0513 (17)
C23	0.0922 (2)	0.4377 (2)	0.2349 (2)	0.0381 (12)
C24	0.1858 (2)	0.4351 (2)	0.2271 (2)	0.0375 (12)
C25	0.2074 (2)	0.4374 (3)	0.1484 (2)	0.0399 (14)
C26	0.2287 (3)	0.5202 (3)	0.1180 (3)	0.0651 (19)
C27	0.2538 (4)	0.5219 (3)	0.0479 (3)	0.081 (2)
C28	0.2607 (3)	0.4422 (3)	0.0094 (3)	0.071 (2)
C29	0.2422 (3)	0.3600 (3)	0.0393 (3)	0.0686 (19)
C30	0.2160 (3)	0.3567 (3)	0.1094 (2)	0.0570 (17)
C31	0.4021 (2)	0.4331 (2)	0.3562 (2)	0.0403 (12)
C32	0.4946 (2)	0.4288 (2)	0.3411 (2)	0.0405 (12)
C33	0.5696 (3)	0.4262 (3)	0.4027 (2)	0.0503 (17)
C34	0.6528 (3)	0.4181 (3)	0.3849 (3)	0.0640 (19)
C35	0.6549 (3)	0.4157 (3)	0.3071 (3)	0.0621 (19)
C36	0.5760 (3)	0.4195 (3)	0.2494 (3)	0.0567 (17)
H1	0.40890	0.32720	0.53740	0.0660*
H2	0.56330	0.33020	0.58200	0.0740*
H3	0.62870	0.31940	0.71550	0.0740*
H3A	0.22240	0.32810	0.82730	0.0530*
H4	0.53580	0.31200	0.80150	0.0650*
H7	0.35000	0.43000	0.24300	0.0570*
H8	0.45750	0.43390	0.87470	0.0790*
H9	0.52440	0.41400	1.00720	0.0960*
H10	0.49350	0.28440	1.07090	0.0890*
H11	0.40730	0.16900	1.00120	0.0790*
H12	0.33970	0.18730	0.86800	0.0650*
H15	-0.05860	0.31640	0.69590	0.0640*
H16	-0.15790	0.30490	0.77750	0.0810*
H17	-0.09840	0.30250	0.91180	0.0950*
H18	0.05250	0.31300	0.96240	0.0900*
H19	-0.01010	0.45130	0.36830	0.0640*
H20	-0.13400	0.44560	0.26330	0.0750*
H21	-0.11680	0.43570	0.13720	0.0760*
H22	0.02880	0.43220	0.11950	0.0620*
H26	0.22610	0.57490	0.14510	0.0780*
H27	0.26600	0.57800	0.02690	0.0980*
H28	0.27820	0.44370	-0.03770	0.0860*
H29	0.24710	0.30560	0.01260	0.0830*

H30	0.20430	0.30020	0.13000	0.0680*
H33	0.56520	0.42970	0.45450	0.0610*
H34	0.70580	0.41440	0.42490	0.0770*
H35	0.70980	0.41160	0.29370	0.0740*
H36	0.57880	0.41760	0.19710	0.0680*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.0405 (2)	0.0555 (2)	0.0326 (2)	0.0014 (1)	0.0111 (1)	0.0028 (1)
Cd2	0.0438 (2)	0.0499 (2)	0.0342 (2)	-0.0006 (1)	0.0147 (1)	0.0042 (1)
Cl1	0.0807 (8)	0.0498 (7)	0.0450 (6)	0.0084 (5)	0.0281 (6)	0.0044 (5)
Cl2	0.0621 (7)	0.0590 (7)	0.0498 (6)	0.0107 (5)	0.0298 (5)	0.0078 (5)
Cl3	0.0498 (6)	0.0571 (7)	0.0486 (6)	0.0046 (5)	0.0201 (5)	0.0068 (5)
Cl4	0.0628 (7)	0.0495 (7)	0.0708 (8)	-0.0062 (5)	0.0234 (6)	0.0015 (6)
O1	0.0378 (15)	0.074 (2)	0.0358 (16)	0.0041 (14)	0.0105 (13)	-0.0022 (14)
O2	0.0431 (16)	0.079 (2)	0.0346 (16)	0.0045 (14)	0.0123 (13)	0.0039 (14)
N1	0.0399 (19)	0.054 (2)	0.043 (2)	0.0016 (16)	0.0176 (17)	0.0056 (16)
N2	0.0347 (18)	0.060 (2)	0.0305 (17)	-0.0030 (15)	0.0120 (15)	-0.0048 (15)
N3	0.0341 (18)	0.071 (2)	0.0291 (17)	-0.0005 (16)	0.0094 (15)	-0.0035 (16)
N4	0.054 (2)	0.076 (3)	0.046 (2)	0.0109 (19)	0.0227 (19)	0.0145 (19)
N5	0.0349 (18)	0.046 (2)	0.046 (2)	-0.0049 (15)	0.0144 (16)	0.0038 (15)
N6	0.0342 (18)	0.050 (2)	0.0356 (18)	0.0038 (15)	0.0121 (15)	0.0008 (15)
N7	0.0330 (18)	0.074 (3)	0.039 (2)	0.0101 (17)	0.0161 (16)	0.0026 (17)
N8	0.045 (2)	0.063 (2)	0.042 (2)	0.0088 (17)	0.0210 (17)	0.0026 (17)
C1	0.052 (3)	0.067 (3)	0.052 (3)	0.005 (2)	0.024 (2)	0.011 (2)
C2	0.052 (3)	0.075 (3)	0.067 (3)	0.004 (2)	0.032 (3)	0.008 (3)
C3	0.031 (2)	0.073 (3)	0.083 (4)	-0.005 (2)	0.020 (3)	0.002 (3)
C4	0.041 (2)	0.067 (3)	0.052 (3)	-0.003 (2)	0.006 (2)	-0.006 (2)
C5	0.036 (2)	0.050 (3)	0.043 (2)	0.0003 (18)	0.0111 (19)	-0.0006 (19)
C6	0.034 (2)	0.046 (2)	0.034 (2)	-0.0006 (18)	0.0050 (18)	-0.0017 (17)
C7	0.035 (2)	0.047 (3)	0.039 (2)	0.0000 (18)	0.0035 (18)	-0.0024 (19)
C8	0.085 (4)	0.050 (3)	0.050 (3)	-0.012 (2)	-0.007 (3)	0.001 (2)
C9	0.096 (4)	0.084 (4)	0.045 (3)	-0.014 (3)	-0.009 (3)	-0.005 (3)
C10	0.076 (4)	0.097 (4)	0.041 (3)	0.006 (3)	-0.002 (3)	0.008 (3)
C11	0.073 (3)	0.069 (3)	0.058 (3)	0.011 (3)	0.019 (3)	0.024 (3)
C12	0.052 (3)	0.062 (3)	0.049 (3)	-0.006 (2)	0.013 (2)	-0.004 (2)
C13	0.036 (2)	0.045 (2)	0.040 (2)	-0.0004 (18)	0.013 (2)	-0.0019 (18)
C14	0.043 (2)	0.043 (2)	0.049 (3)	0.0041 (19)	0.022 (2)	0.0009 (19)
C15	0.045 (3)	0.058 (3)	0.060 (3)	0.004 (2)	0.019 (2)	-0.006 (2)
C16	0.046 (3)	0.073 (3)	0.093 (4)	0.004 (2)	0.035 (3)	0.005 (3)
C17	0.078 (4)	0.085 (4)	0.095 (4)	0.020 (3)	0.059 (4)	0.027 (3)
C18	0.085 (4)	0.097 (4)	0.056 (3)	0.020 (3)	0.042 (3)	0.019 (3)
C19	0.041 (3)	0.063 (3)	0.061 (3)	-0.006 (2)	0.022 (2)	0.010 (2)
C20	0.034 (2)	0.069 (3)	0.085 (4)	-0.005 (2)	0.018 (3)	0.015 (3)
C21	0.041 (3)	0.075 (3)	0.068 (3)	-0.012 (2)	0.000 (2)	0.007 (3)
C22	0.044 (3)	0.061 (3)	0.046 (3)	-0.005 (2)	0.006 (2)	0.000 (2)
C23	0.038 (2)	0.035 (2)	0.042 (2)	-0.0034 (17)	0.0114 (19)	0.0005 (18)
C24	0.040 (2)	0.037 (2)	0.035 (2)	0.0057 (17)	0.0087 (19)	0.0013 (17)
C25	0.040 (2)	0.046 (3)	0.034 (2)	0.0056 (18)	0.0102 (18)	0.0000 (18)

C26	0.101 (4)	0.045 (3)	0.059 (3)	0.010 (2)	0.038 (3)	0.000 (2)
C27	0.138 (5)	0.058 (3)	0.064 (3)	0.006 (3)	0.055 (3)	0.015 (3)
C28	0.101 (4)	0.077 (4)	0.049 (3)	0.001 (3)	0.043 (3)	-0.003 (3)
C29	0.108 (4)	0.055 (3)	0.053 (3)	0.006 (3)	0.039 (3)	-0.012 (2)
C30	0.081 (3)	0.044 (3)	0.051 (3)	-0.001 (2)	0.026 (2)	0.001 (2)
C31	0.034 (2)	0.043 (2)	0.044 (2)	0.0101 (17)	0.010 (2)	0.0051 (19)
C32	0.035 (2)	0.041 (2)	0.047 (2)	0.0006 (17)	0.013 (2)	0.0002 (18)
C33	0.042 (3)	0.060 (3)	0.049 (3)	-0.003 (2)	0.012 (2)	-0.005 (2)
C34	0.037 (3)	0.081 (4)	0.070 (3)	-0.004 (2)	0.006 (2)	-0.005 (3)
C35	0.043 (3)	0.070 (3)	0.080 (4)	0.005 (2)	0.028 (3)	0.006 (3)
C36	0.053 (3)	0.070 (3)	0.056 (3)	0.010 (2)	0.030 (2)	0.011 (2)

Geometric parameters (Å, °)

Cd1—C11	2.6857 (11)	C16—C17	1.360 (7)
Cd1—C12	2.5444 (11)	C17—C18	1.356 (9)
Cd1—C13	2.4590 (11)	C19—C20	1.377 (7)
Cd1—O1	2.535 (3)	C20—C21	1.358 (7)
Cd1—N1	2.411 (3)	C21—C22	1.382 (7)
Cd1—N2	2.437 (3)	C22—C23	1.377 (5)
Cd2—C11	2.5693 (11)	C23—C24	1.475 (5)
Cd2—C12	2.5902 (11)	C24—C25	1.500 (5)
Cd2—C14	2.4684 (11)	C25—C26	1.385 (6)
Cd2—O2	2.633 (3)	C25—C30	1.378 (6)
Cd2—N5	2.380 (3)	C26—C27	1.379 (7)
Cd2—N6	2.360 (3)	C27—C28	1.356 (6)
O1—C13	1.219 (4)	C28—C29	1.360 (6)
O2—C31	1.215 (4)	C29—C30	1.388 (6)
N1—C1	1.331 (6)	C31—C32	1.508 (5)
N1—C5	1.350 (5)	C32—C33	1.364 (5)
N2—N3	1.365 (4)	C33—C34	1.392 (7)
N2—C6	1.276 (4)	C34—C35	1.374 (7)
N3—C13	1.354 (4)	C35—C36	1.367 (7)
N4—C14	1.334 (5)	C1—H1	0.9300
N4—C18	1.341 (7)	C2—H2	0.9300
N5—C19	1.331 (6)	C3—H3	0.9300
N5—C23	1.352 (5)	C4—H4	0.9300
N6—N7	1.364 (4)	C8—H8	0.9300
N6—C24	1.283 (4)	C9—H9	0.9300
N7—C31	1.359 (5)	C10—H10	0.9300
N8—C32	1.346 (5)	C11—H11	0.9300
N8—C36	1.332 (6)	C12—H12	0.9300
N3—H3A	0.8600	C15—H15	0.9300
N7—H7	0.8600	C16—H16	0.9300
C1—C2	1.384 (7)	C17—H17	0.9300
C2—C3	1.362 (7)	C18—H18	0.9300
C3—C4	1.383 (6)	C19—H19	0.9300
C4—C5	1.386 (6)	C20—H20	0.9300
C5—C6	1.486 (5)	C21—H21	0.9300
C6—C7	1.492 (5)	C22—H22	0.9300

C7—C12	1.388 (6)	C26—H26	0.9300
C7—C8	1.373 (6)	C27—H27	0.9300
C8—C9	1.386 (6)	C28—H28	0.9300
C9—C10	1.364 (8)	C29—H29	0.9300
C10—C11	1.358 (7)	C30—H30	0.9300
C11—C12	1.394 (6)	C33—H33	0.9300
C13—C14	1.494 (5)	C34—H34	0.9300
C14—C15	1.374 (7)	C35—H35	0.9300
C15—C16	1.389 (7)	C36—H36	0.9300
Cl1—Cd1—Cl2	82.27 (3)	N4—C18—C17	123.8 (5)
Cl1—Cd1—Cl3	95.23 (3)	N5—C19—C20	122.5 (4)
Cl1—Cd1—O1	137.62 (6)	C19—C20—C21	120.2 (5)
Cl1—Cd1—N1	91.44 (8)	C20—C21—C22	118.3 (4)
Cl1—Cd1—N2	157.41 (8)	C21—C22—C23	119.2 (4)
Cl2—Cd1—Cl3	152.25 (4)	N5—C23—C22	122.2 (3)
Cl2—Cd1—O1	78.26 (7)	N5—C23—C24	116.4 (3)
Cl2—Cd1—N1	107.44 (7)	C22—C23—C24	121.4 (3)
Cl2—Cd1—N2	105.16 (7)	N6—C24—C23	116.7 (3)
Cl3—Cd1—O1	85.56 (7)	N6—C24—C25	121.2 (3)
Cl3—Cd1—N1	100.24 (7)	C23—C24—C25	122.1 (3)
Cl3—Cd1—N2	87.73 (7)	C24—C25—C30	120.6 (3)
O1—Cd1—N1	130.24 (9)	C26—C25—C30	118.8 (4)
O1—Cd1—N2	64.89 (9)	C24—C25—C26	120.2 (4)
N1—Cd1—N2	66.02 (10)	C25—C26—C27	120.3 (4)
Cl1—Cd2—Cl2	83.70 (3)	C26—C27—C28	120.3 (4)
Cl1—Cd2—Cl4	163.43 (4)	C27—C28—C29	120.3 (5)
Cl1—Cd2—O2	81.98 (7)	C28—C29—C30	120.5 (4)
Cl1—Cd2—N5	97.59 (7)	C25—C30—C29	119.8 (4)
Cl1—Cd2—N6	92.45 (7)	O2—C31—C32	124.6 (3)
Cl2—Cd2—Cl4	98.11 (4)	N7—C31—C32	111.3 (3)
Cl2—Cd2—O2	123.47 (6)	O2—C31—N7	124.1 (3)
Cl2—Cd2—N5	102.82 (8)	N8—C32—C31	115.0 (3)
Cl2—Cd2—N6	169.63 (8)	N8—C32—C33	124.9 (3)
Cl4—Cd2—O2	83.35 (7)	C31—C32—C33	120.2 (3)
Cl4—Cd2—N5	98.05 (8)	C32—C33—C34	117.4 (4)
Cl4—Cd2—N6	88.33 (8)	C33—C34—C35	118.6 (4)
O2—Cd2—N5	133.15 (9)	C34—C35—C36	119.7 (5)
O2—Cd2—N6	65.16 (9)	N8—C36—C35	123.1 (5)
N5—Cd2—N6	68.06 (11)	N1—C1—H1	118.00
Cd1—Cl1—Cd2	92.58 (3)	C2—C1—H1	118.00
Cd1—Cl2—Cd2	95.45 (4)	C1—C2—H2	121.00
Cd1—O1—C13	116.0 (2)	C3—C2—H2	121.00
Cd2—O2—C31	112.3 (2)	C2—C3—H3	121.00
Cd1—N1—C1	122.9 (3)	C4—C3—H3	121.00
Cd1—N1—C5	118.6 (2)	C3—C4—H4	120.00
C1—N1—C5	117.8 (3)	C5—C4—H4	120.00
Cd1—N2—N3	116.8 (2)	C7—C8—H8	119.00
Cd1—N2—C6	120.8 (2)	C9—C8—H8	119.00

N3—N2—C6	121.1 (3)	C8—C9—H9	120.00
N2—N3—C13	118.1 (3)	C10—C9—H9	120.00
C14—N4—C18	115.7 (4)	C9—C10—H10	120.00
Cd2—N5—C19	124.6 (3)	C11—C10—H10	120.00
Cd2—N5—C23	117.7 (2)	C10—C11—H11	120.00
C19—N5—C23	117.6 (3)	C12—C11—H11	120.00
Cd2—N6—N7	119.4 (2)	C7—C12—H12	120.00
Cd2—N6—C24	120.7 (2)	C11—C12—H12	120.00
N7—N6—C24	119.2 (3)	C14—C15—H15	121.00
N6—N7—C31	118.6 (3)	C16—C15—H15	121.00
C32—N8—C36	116.4 (3)	C15—C16—H16	121.00
N2—N3—H3A	121.00	C17—C16—H16	121.00
C13—N3—H3A	121.00	C16—C17—H17	120.00
N6—N7—H7	121.00	C18—C17—H17	120.00
C31—N7—H7	121.00	N4—C18—H18	118.00
N1—C1—C2	123.4 (5)	C17—C18—H18	118.00
C1—C2—C3	118.9 (5)	N5—C19—H19	119.00
C2—C3—C4	118.7 (4)	C20—C19—H19	119.00
C3—C4—C5	119.6 (3)	C19—C20—H20	120.00
N1—C5—C4	121.5 (3)	C21—C20—H20	120.00
N1—C5—C6	116.4 (3)	C20—C21—H21	121.00
C4—C5—C6	122.1 (3)	C22—C21—H21	121.00
N2—C6—C7	124.7 (3)	C21—C22—H22	120.00
C5—C6—C7	119.5 (3)	C23—C22—H22	120.00
N2—C6—C5	115.8 (3)	C25—C26—H26	120.00
C8—C7—C12	118.4 (3)	C27—C26—H26	120.00
C6—C7—C8	121.9 (3)	C26—C27—H27	120.00
C6—C7—C12	119.6 (3)	C28—C27—H27	120.00
C7—C8—C9	121.2 (4)	C27—C28—H28	120.00
C8—C9—C10	119.8 (5)	C29—C28—H28	120.00
C9—C10—C11	120.2 (5)	C28—C29—H29	120.00
C10—C11—C12	120.6 (4)	C30—C29—H29	120.00
C7—C12—C11	119.8 (4)	C25—C30—H30	120.00
O1—C13—N3	123.0 (3)	C29—C30—H30	120.00
O1—C13—C14	122.0 (3)	C32—C33—H33	121.00
N3—C13—C14	115.0 (3)	C34—C33—H33	121.00
N4—C14—C15	124.5 (4)	C33—C34—H34	121.00
C13—C14—C15	119.8 (3)	C35—C34—H34	121.00
N4—C14—C13	115.8 (4)	C34—C35—H35	120.00
C14—C15—C16	117.9 (4)	C36—C35—H35	120.00
C15—C16—C17	118.3 (5)	N8—C36—H36	118.00
C16—C17—C18	119.9 (5)	C35—C36—H36	118.00
C12—Cd1—C11—Cd2	18.37 (4)	N2—N3—C13—O1	-2.1 (6)
C13—Cd1—C11—Cd2	170.55 (4)	N2—N3—C13—C14	177.6 (3)
O1—Cd1—C11—Cd2	81.41 (10)	C18—N4—C14—C15	0.5 (6)
N1—Cd1—C11—Cd2	-89.04 (8)	C14—N4—C18—C17	0.3 (6)
N2—Cd1—C11—Cd2	-92.78 (19)	C18—N4—C14—C13	-178.2 (4)
O1—Cd1—N1—C5	22.3 (3)	Cd2—N5—C19—C20	178.3 (3)

N2—Cd1—N1—C5	12.2 (3)	Cd2—N5—C23—C22	-177.9 (3)
Cl1—Cd1—N2—N3	-176.23 (15)	Cd2—N5—C23—C24	1.9 (3)
Cl2—Cd1—N2—N3	77.0 (2)	C23—N5—C19—C20	1.4 (6)
Cl1—Cd1—Cl2—Cd2	-18.28 (4)	C19—N5—C23—C24	179.0 (3)
Cl3—Cd1—Cl2—Cd2	-104.82 (7)	C19—N5—C23—C22	-0.8 (5)
O1—Cd1—Cl2—Cd2	-160.43 (7)	N7—N6—C24—C25	-0.3 (5)
N1—Cd1—Cl2—Cd2	70.79 (8)	C24—N6—N7—C31	177.1 (3)
N2—Cd1—Cl2—Cd2	139.93 (8)	Cd2—N6—C24—C23	-7.8 (4)
Cl3—Cd1—N2—C6	88.5 (2)	Cd2—N6—C24—C25	170.9 (2)
O1—Cd1—N2—C6	174.7 (3)	Cd2—N6—N7—C31	5.9 (4)
N1—Cd1—N2—C6	-13.8 (2)	N7—N6—C24—C23	-179.0 (3)
N2—Cd1—N1—C1	-177.4 (3)	N6—N7—C31—O2	-0.7 (5)
Cl1—Cd1—O1—C13	173.2 (3)	N6—N7—C31—C32	178.3 (3)
Cl2—Cd1—O1—C13	-122.4 (3)	C36—N8—C32—C33	0.4 (5)
Cl3—Cd1—O1—C13	80.3 (3)	C32—N8—C36—C35	0.2 (6)
N1—Cd1—O1—C13	-19.4 (3)	C36—N8—C32—C31	-178.2 (3)
N2—Cd1—O1—C13	-9.3 (3)	N1—C1—C2—C3	2.6 (7)
Cl3—Cd1—N1—C1	99.7 (3)	C1—C2—C3—C4	-1.4 (6)
O1—Cd1—N1—C1	-167.4 (3)	C2—C3—C4—C5	-1.1 (6)
Cl3—Cd1—N2—N3	-78.1 (2)	C3—C4—C5—N1	2.7 (7)
O1—Cd1—N2—N3	8.1 (2)	C3—C4—C5—C6	-177.2 (4)
N1—Cd1—N2—N3	179.7 (3)	N1—C5—C6—C7	177.4 (3)
Cl1—Cd1—N1—C1	4.2 (3)	C4—C5—C6—N2	178.5 (4)
Cl2—Cd1—N1—C1	-78.2 (3)	N1—C5—C6—N2	-1.5 (5)
Cl2—Cd1—N2—C6	-116.4 (2)	C4—C5—C6—C7	-2.7 (6)
Cl2—Cd1—N1—C5	111.5 (3)	C5—C6—C7—C8	64.3 (5)
Cl3—Cd1—N1—C5	-70.6 (3)	C5—C6—C7—C12	-111.0 (4)
Cl1—Cd1—N1—C5	-166.2 (3)	N2—C6—C7—C8	-117.0 (4)
Cl1—Cd1—N2—C6	-9.7 (4)	N2—C6—C7—C12	67.8 (5)
Cl2—Cd2—O2—C31	178.2 (2)	C6—C7—C8—C9	-174.9 (4)
N5—Cd2—Cl2—Cd1	115.46 (8)	C6—C7—C12—C11	175.1 (4)
Cl1—Cd2—O2—C31	101.2 (2)	C12—C7—C8—C9	0.5 (6)
N5—Cd2—Cl1—Cd1	-120.13 (8)	C8—C7—C12—C11	-0.4 (6)
N6—Cd2—Cl1—Cd1	171.68 (8)	C7—C8—C9—C10	-1.6 (7)
Cl4—Cd2—O2—C31	-86.5 (2)	C8—C9—C10—C11	2.7 (8)
N5—Cd2—O2—C31	8.3 (3)	C9—C10—C11—C12	-2.7 (7)
N6—Cd2—O2—C31	4.7 (2)	C10—C11—C12—C7	1.5 (7)
Cl2—Cd2—Cl1—Cd1	-17.98 (4)	O1—C13—C14—N4	179.3 (4)
O2—Cd2—Cl1—Cd1	107.19 (6)	N3—C13—C14—C15	-179.1 (4)
Cl1—Cd2—Cl2—Cd1	19.08 (4)	N3—C13—C14—N4	-0.4 (6)
Cl4—Cd2—Cl2—Cd1	-144.32 (4)	O1—C13—C14—C15	0.6 (7)
O2—Cd2—Cl2—Cd1	-56.95 (8)	C13—C14—C15—C16	177.7 (4)
N5—Cd2—N6—N7	177.6 (3)	N4—C14—C15—C16	-0.9 (7)
Cl1—Cd2—N6—C24	103.6 (2)	C14—C15—C16—C17	0.4 (6)
Cl1—Cd2—N5—C19	89.5 (3)	C15—C16—C17—C18	0.3 (7)
Cl2—Cd2—N5—C19	4.3 (3)	C16—C17—C18—N4	-0.7 (7)
Cl4—Cd2—N5—C19	-96.0 (3)	N5—C19—C20—C21	-0.9 (7)
O2—Cd2—N5—C19	175.6 (3)	C19—C20—C21—C22	-0.2 (7)
N6—Cd2—N5—C19	179.1 (3)	C20—C21—C22—C23	0.7 (6)

C11—Cd2—N5—C23	-93.6 (2)	C21—C22—C23—C24	179.9 (3)
C12—Cd2—N5—C23	-178.8 (2)	C21—C22—C23—N5	-0.3 (6)
C14—Cd2—N5—C23	80.9 (2)	C22—C23—C24—C25	4.9 (5)
O2—Cd2—N5—C23	-7.5 (3)	N5—C23—C24—C25	-174.9 (3)
N6—Cd2—N5—C23	-4.0 (2)	N5—C23—C24—N6	3.8 (4)
C11—Cd2—N6—N7	-85.3 (2)	C22—C23—C24—N6	-176.4 (3)
C14—Cd2—N6—N7	78.2 (2)	N6—C24—C25—C30	90.3 (4)
O2—Cd2—N6—N7	-5.3 (2)	C23—C24—C25—C26	96.0 (4)
O2—Cd2—N6—C24	-176.4 (3)	C23—C24—C25—C30	-91.1 (4)
N5—Cd2—N6—C24	6.4 (2)	N6—C24—C25—C26	-82.6 (5)
C14—Cd2—N6—C24	-92.9 (2)	C24—C25—C26—C27	176.1 (4)
Cd1—O1—C13—N3	9.9 (5)	C24—C25—C30—C29	-175.5 (4)
Cd1—O1—C13—C14	-169.8 (3)	C26—C25—C30—C29	-2.5 (6)
Cd2—O2—C31—N7	-4.0 (4)	C30—C25—C26—C27	3.1 (7)
Cd2—O2—C31—C32	177.1 (2)	C25—C26—C27—C28	-2.2 (8)
Cd1—N1—C5—C6	-10.9 (4)	C26—C27—C28—C29	0.5 (8)
Cd1—N1—C5—C4	169.1 (3)	C27—C28—C29—C30	0.1 (8)
C1—N1—C5—C6	178.3 (3)	C28—C29—C30—C25	1.0 (7)
Cd1—N1—C1—C2	-171.4 (3)	O2—C31—C32—N8	-179.9 (3)
C5—N1—C1—C2	-1.0 (6)	N7—C31—C32—C33	-177.6 (3)
C1—N1—C5—C4	-1.7 (6)	O2—C31—C32—C33	1.4 (5)
N3—N2—C6—C5	179.3 (3)	N7—C31—C32—N8	1.0 (4)
Cd1—N2—C6—C5	13.3 (4)	N8—C32—C33—C34	-1.3 (6)
N3—N2—C6—C7	0.5 (5)	C31—C32—C33—C34	177.2 (3)
C6—N2—N3—C13	-173.8 (3)	C32—C33—C34—C35	1.7 (6)
Cd1—N2—N3—C13	-7.3 (4)	C33—C34—C35—C36	-1.2 (6)
Cd1—N2—C6—C7	-165.5 (3)	C34—C35—C36—N8	0.3 (7)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3 <i>A</i> ...N4	0.86	2.29	2.646 (4)	105
N7—H7...N8	0.86	2.16	2.569 (4)	108
C1—H1...O2	0.93	2.53	3.303 (6)	140
C10—H10...O1 ⁱ	0.93	2.50	3.261 (6)	139
C36—H36...C13 ⁱⁱ	0.93	2.77	3.482 (5)	135

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x+1/2, -y+1/2, z-1/2$.