



Crystal structure of (*Z*)-7,8-dichloro-4-(2-oxopropylidene)-4,5-dihydro-1*H*-1,5-benzodiazepin-2(3*H*)-one

Sanae Lahmidi,^{a*} Abdelhanine Essaghouni,^a El Mokhtar Essassi,^a Mohamed Saadi^b and Lahcen El Ammari^b

^aLaboratoire de Chimie Organique Hétérocyclique URAC 21, Pôle de Compétence Pharmacochimie, Av. Ibn Battouta, BP 1014, Faculté des Sciences, Université Mohammed V, Rabat, Morocco, and ^bLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V de Rabat, Avenue Ibn Battouta, BP 1014, Rabat, Morocco. *Correspondence e-mail: lahmidi_s@yahoo.fr

Received 8 December 2015; accepted 10 December 2015

Edited by E. R. T. Tiekink, University of Malaya, Malaysia


In the title compound, C₁₂H₁₀Cl₂N₂O₂, the seven-membered heterocycle displays a half-chair conformation. The mean plane through the oxopropylidene group makes a dihedral angle of 36.44 (9)° with the fused benzene ring. An intramolecular N—H···O hydrogen bond to close an *S*(6) loop is noted. An important feature of the molecular packing are N—H···O hydrogen bonds that lead to the formation of helical supramolecular chains along the *b* axis.

Keywords: crystal structure; 2-oxopropylidene; 1,5-benzodiazepinone; hydrogen bonding.

CCDC reference: 1441702

1. Related literature

For the pharmaceutical and biochemical properties of 1,5-benzodiazepine and their derivatives, see: El Azzaoui *et al.* (1999); Gringauz (1999); Swamy *et al.* (2008). For related structures, see: El Abbassi *et al.* (1997); Akkurt *et al.* (2006).



2. Experimental

2.1. Crystal data

C ₁₂ H ₁₀ Cl ₂ N ₂ O ₂	<i>V</i> = 1229.93 (11) Å ³
<i>M_r</i> = 285.12	<i>Z</i> = 4
Monoclinic, <i>P</i> 2 ₁ / <i>n</i>	Mo <i>K</i> α radiation
<i>a</i> = 7.6789 (4) Å	<i>μ</i> = 0.52 mm ⁻¹
<i>b</i> = 13.2199 (6) Å	<i>T</i> = 296 K
<i>c</i> = 12.4129 (7) Å	0.36 × 0.33 × 0.24 mm
<i>β</i> = 102.561 (3)°	

2.2. Data collection

Bruker X8 APEX diffractometer	25693 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2009)	3299 independent reflections
<i>T</i> _{min} = 0.672, <i>T</i> _{max} = 0.746	2692 reflections with <i>I</i> > 2σ(<i>I</i>)
	<i>R</i> _{int} = 0.031

2.3. Refinement

<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)] = 0.038	163 parameters
<i>wR</i> (<i>F</i> ²) = 0.112	H-atom parameters constrained
<i>S</i> = 1.02	Δρ _{max} = 0.36 e Å ⁻³
3298 reflections	Δρ _{min} = -0.33 e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2···O2	0.86	1.96	2.6410 (18)	135
N1—H1···O2 ⁱ	0.86	1.95	2.8010 (19)	173

Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT-Plus* (Bruker, 2009); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *PLATON* (Spek, 2009) and *pubCIF* (Westrip, 2010).

Acknowledgements

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements and the University Mohammed V, Rabat, Morocco, for financial support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: TK5415).

References

- Akkurt, M., Karaca, S., Bouhfid, R., Essassi, E. M. & Büyükgüngör, O. (2006). *Anal. Sci. X-ray Struct. Anal. Online*, **22**, x147–x148.
- Bruker (2009). *APEX2*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- El Abbassi, M., Essassi, E. M. & Fifi, J. (1997). *Bull. Soc. Chim. Belg.* **106**, 205–210.
- El Azzaoui, B., Fifi, J., Tjiou, E. M., Essassi, E. M., Jaud, J., Lopez, L. & Bellan, J. (1999). *Tetrahedron Lett.* **40**, 4677–4680.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.

Gringauz, A. (1999). *Med. Chem.* pp. 578–580 New York: Wiley-VCH.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

Swamy, G. Y. S. K., Sridhar, B., Ravikumar, K., Reddy, K. S. & Reddy, V. V. N.
(2008). *J. Struct. Chem.* **49**, 775–779.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, o1059–o1060 [doi:10.1107/S2056989015023750]

Crystal structure of (Z)-7,8-dichloro-4-(2-oxopropylidene)-4,5-dihydro-1H-1,5-benzodiazepin-2(3H)-one

Sanae Lahmidi, Abdelhanine Essaghouani, El Mokhtar Essassi, Mohamed Saadi and Lahcen El Ammari

S1. Comment

1,5-Benzodiazepines are used as starting materials in the synthesis of several heterocyclic compounds studied for potential biological activities (El Azzaoui *et al.* 1999). They are used for the purpose of hypnotic effects, owing to their less toxic and less severe withdrawal effects when compared with barbiturates (Gringauz, 1999). Some benzodiazepine derivatives have been widely used as anti-bacterial, anti-fungal, analgesic and anti-convulsant agents (Swamy *et al.*, 2008). In our laboratory we were interested in the synthesis of new 1,5-benzodiazepine derivatives (El Abbassi *et al.*, 1997; Akkurt *et al.*, 2006). The purpose of this work is to synthesize (Z)-7,8-dichloro-4,5-dihydro-4-(2-oxopropylidene)-1H-benzo[*b*][1,4] diazepin-2(3H)-one by condensation of 4,5-dichloro-*o*-phenylenediamine with 4-hydroxy-6-methyl-2H-pyran-2-one.

The molecule of the title compound, Fig. 1, is build up from two fused six- and seven-membered rings linked to a 2-oxopropylidene group. The seven-membered ring displays a half-chair conformation as indicated by the puckering amplitude $QT = 0.811(2) \text{ \AA}$ and spherical polar angle $\theta_2 = 73.9(2)^\circ$, $\varphi_2 = 129.07(12)^\circ$ and $\varphi_3 = -76.3(4)^\circ$. Moreover, the dihedral angle between the mean plane through the oxopropylidene group and the dichlorobenzene ring is of $36.44(9)^\circ$.

In the crystal, the molecules are linked by hydrogen bonds in the way to build an helical chain along the *b* axis as shown in Fig. 2 and Table 1. An intramolecular hydrogen bond $N2-H2\cdots O2$ is also observed in this structure.

S2. Experimental

A mixture of 4,5-dichloro-*o*-phenylenediamine (1.13 g) and of 4-hydroxy-6-methyl-2H-pyran-2-one (0.25 g) in xylene (30 ml) was heated at reflux for 4 h with azeotropic distillation. The completion of the reaction was confirmed by TLC. The solid obtained upon cooling the mixture was recrystallized from ethanol to afford colourless crystals in 75% yield.

S3. Refinement

The H atoms were located in a difference map and treated as riding with $C-H = 0.93-0.97 \text{ \AA}$ and $N-H = 0.86 \text{ \AA}$, and with $U_{iso}(H) = 1.2-1.5U_{eq}(C, N)$.




Figure 1

Molecular structure of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small circles.




Figure 2

Structure of the title compound, showing molecules linked through N1—H1...O1 hydrogen bonds and the intramolecular hydrogen bond N2—H2...O2 (dashed lines).

(Z)-7,8-Dichloro-4-(2-oxopropylidene)-4,5-dihydro-1*H*-1,5-benzodiazepin-2(3*H*)-one

Crystal data

$C_{12}H_{10}Cl_2N_2O_2$
 $M_r = 285.12$
 Monoclinic, $P2_1/n$
 $a = 7.6789$ (4) Å
 $b = 13.2199$ (6) Å
 $c = 12.4129$ (7) Å
 $\beta = 102.561$ (3)°
 $V = 1229.93$ (11) Å³
 $Z = 4$

$F(000) = 584$
 $D_x = 1.540$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 3299 reflections
 $\theta = 2.3$ – 29.1 °
 $\mu = 0.52$ mm⁻¹
 $T = 296$ K
 Block, colourless
 $0.36 \times 0.33 \times 0.24$ mm

Data collection

Bruker X8 APEX
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
 $T_{\min} = 0.672$, $T_{\max} = 0.746$

25693 measured reflections
3299 independent reflections
2692 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\max} = 29.1^\circ$, $\theta_{\min} = 2.3^\circ$
 $h = -10 \rightarrow 10$
 $k = -18 \rightarrow 18$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.112$
 $S = 1.02$
3298 reflections
163 parameters
0 restraints

Hydrogen site location: difference Fourier map
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0553P)^2 + 0.6111P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5232 (2)	0.21313 (13)	0.49879 (13)	0.0338 (3)
C2	0.5740 (2)	0.30850 (13)	0.54157 (13)	0.0355 (3)
C3	0.6157 (2)	0.38307 (12)	0.47350 (13)	0.0346 (3)
H3	0.6479	0.4470	0.5024	0.042*
C4	0.6104 (2)	0.36438 (11)	0.36221 (13)	0.0303 (3)
C5	0.5633 (2)	0.26785 (11)	0.31938 (12)	0.0291 (3)
C6	0.5189 (2)	0.19345 (12)	0.38896 (13)	0.0330 (3)
H6	0.4858	0.1294	0.3607	0.040*
C7	0.6022 (2)	0.46740 (11)	0.19419 (14)	0.0349 (3)
C8	0.4552 (2)	0.40089 (12)	0.13165 (15)	0.0375 (4)
H8A	0.3586	0.3984	0.1707	0.045*
H8B	0.4087	0.4298	0.0593	0.045*
C9	0.5204 (2)	0.29534 (11)	0.11872 (13)	0.0310 (3)
C10	0.5298 (2)	0.25845 (12)	0.01690 (13)	0.0333 (3)
H10	0.4958	0.3007	-0.0440	0.040*
C11	0.5889 (2)	0.15901 (12)	-0.00056 (13)	0.0330 (3)
C12	0.6037 (3)	0.12898 (15)	-0.11491 (14)	0.0424 (4)
H12A	0.5680	0.1847	-0.1645	0.064*
H12B	0.7249	0.1111	-0.1145	0.064*
H12C	0.5275	0.0720	-0.1388	0.064*
N1	0.6678 (2)	0.44232 (10)	0.30134 (12)	0.0352 (3)

H1	0.7364	0.4859	0.3410	0.042*
N2	0.5688 (2)	0.23921 (10)	0.21168 (11)	0.0339 (3)
H2	0.6012	0.1781	0.2028	0.041*
C11	0.46370 (7)	0.11838 (4)	0.57991 (4)	0.04762 (15)
C12	0.58703 (8)	0.33611 (4)	0.67921 (4)	0.05696 (17)
O1	0.6604 (2)	0.54018 (9)	0.15280 (11)	0.0496 (3)
O2	0.6297 (2)	0.09619 (9)	0.07595 (10)	0.0451 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0353 (8)	0.0360 (8)	0.0318 (7)	0.0068 (6)	0.0109 (6)	0.0083 (6)
C2	0.0383 (9)	0.0408 (8)	0.0278 (7)	0.0116 (7)	0.0080 (6)	0.0008 (6)
C3	0.0383 (9)	0.0313 (7)	0.0330 (8)	0.0052 (6)	0.0053 (6)	-0.0031 (6)
C4	0.0316 (8)	0.0283 (7)	0.0309 (7)	0.0032 (6)	0.0066 (6)	0.0017 (6)
C5	0.0317 (8)	0.0281 (7)	0.0278 (7)	0.0031 (6)	0.0069 (6)	0.0018 (5)
C6	0.0389 (9)	0.0276 (7)	0.0330 (8)	0.0009 (6)	0.0089 (6)	0.0022 (6)
C7	0.0438 (9)	0.0245 (7)	0.0383 (8)	0.0055 (6)	0.0136 (7)	0.0015 (6)
C8	0.0410 (9)	0.0325 (8)	0.0370 (8)	0.0075 (7)	0.0043 (7)	0.0044 (6)
C9	0.0328 (8)	0.0278 (7)	0.0309 (7)	-0.0020 (6)	0.0039 (6)	0.0031 (6)
C10	0.0407 (9)	0.0315 (7)	0.0258 (7)	-0.0023 (6)	0.0029 (6)	0.0055 (6)
C11	0.0353 (8)	0.0346 (8)	0.0273 (7)	-0.0037 (6)	0.0028 (6)	0.0009 (6)
C12	0.0495 (10)	0.0468 (10)	0.0306 (8)	-0.0034 (8)	0.0083 (7)	-0.0031 (7)
N1	0.0416 (8)	0.0275 (6)	0.0359 (7)	-0.0056 (5)	0.0073 (6)	-0.0001 (5)
N2	0.0487 (8)	0.0253 (6)	0.0287 (6)	0.0033 (5)	0.0104 (6)	0.0020 (5)
C11	0.0591 (3)	0.0464 (3)	0.0420 (2)	0.0054 (2)	0.0213 (2)	0.01498 (18)
C12	0.0855 (4)	0.0567 (3)	0.0300 (2)	0.0141 (3)	0.0155 (2)	-0.00200 (18)
O1	0.0731 (10)	0.0323 (6)	0.0465 (7)	-0.0059 (6)	0.0199 (7)	0.0064 (5)
O2	0.0680 (9)	0.0334 (6)	0.0318 (6)	0.0095 (6)	0.0061 (6)	0.0038 (5)

Geometric parameters (Å, °)

C1—C6	1.381 (2)	C8—C9	1.503 (2)
C1—C2	1.391 (2)	C8—H8A	0.9700
C1—C11	1.7297 (16)	C8—H8B	0.9700
C2—C3	1.380 (2)	C9—N2	1.3541 (19)
C2—C12	1.7286 (17)	C9—C10	1.371 (2)
C3—C4	1.395 (2)	C10—C11	1.422 (2)
C3—H3	0.9300	C10—H10	0.9300
C4—C5	1.399 (2)	C11—O2	1.2491 (19)
C4—N1	1.404 (2)	C11—C12	1.502 (2)
C5—N2	1.3989 (19)	C12—H12A	0.9600
C5—C6	1.399 (2)	C12—H12B	0.9600
C6—H6	0.9300	C12—H12C	0.9600
C7—O1	1.220 (2)	N1—H1	0.8599
C7—N1	1.357 (2)	N2—H2	0.8600
C7—C8	1.506 (2)		

C6—C1—C2	119.49 (15)	C9—C8—H8B	109.3
C6—C1—C11	119.04 (13)	C7—C8—H8B	109.3
C2—C1—C11	121.46 (13)	H8A—C8—H8B	108.0
C3—C2—C1	119.79 (15)	N2—C9—C10	122.05 (14)
C3—C2—C12	118.88 (13)	N2—C9—C8	116.99 (14)
C1—C2—C12	121.34 (13)	C10—C9—C8	120.95 (14)
C2—C3—C4	121.26 (15)	C9—C10—C11	123.48 (14)
C2—C3—H3	119.4	C9—C10—H10	118.3
C4—C3—H3	119.4	C11—C10—H10	118.3
C3—C4—C5	119.15 (14)	O2—C11—C10	122.29 (15)
C3—C4—N1	117.25 (14)	O2—C11—C12	119.00 (15)
C5—C4—N1	123.37 (14)	C10—C11—C12	118.71 (15)
N2—C5—C4	123.42 (14)	C11—C12—H12A	109.5
N2—C5—C6	117.50 (14)	C11—C12—H12B	109.5
C4—C5—C6	118.95 (14)	H12A—C12—H12B	109.5
C1—C6—C5	121.34 (15)	C11—C12—H12C	109.5
C1—C6—H6	119.3	H12A—C12—H12C	109.5
C5—C6—H6	119.3	H12B—C12—H12C	109.5
O1—C7—N1	120.91 (17)	C7—N1—C4	127.87 (14)
O1—C7—C8	123.03 (16)	C7—N1—H1	116.6
N1—C7—C8	116.06 (14)	C4—N1—H1	114.1
C9—C8—C7	111.53 (14)	C9—N2—C5	127.32 (13)
C9—C8—H8A	109.3	C9—N2—H2	116.1
C7—C8—H8A	109.3	C5—N2—H2	116.5

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O2	0.86	1.96	2.6410 (18)	135
N1—H1 \cdots O2 ⁱ	0.86	1.95	2.8010 (19)	173

Symmetry code: (i) $-x+3/2, y+1/2, -z+1/2$.