






Complete Genome Sequence of *Fusobacterium necrophorum* subsp. *necrophorum* ATCC 25286

 Ariana Umaña,^a  Justin A. Lemkul,^a  Daniel J. Slade^a

^aDepartment of Biochemistry, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, USA

ABSTRACT *Fusobacterium necrophorum* is a pathogenic Gram-negative, anaerobic bacterium. In this study, we present the first complete genome sequence of *Fusobacterium necrophorum* subsp. *necrophorum* ATCC 25286. These data provide a critical advancement in our understanding of virulence factors that could contribute to *F. necrophorum* pathogenesis in both human and livestock infections.

Fusobacterium is a genus of anaerobic bacteria that cause serious and even fatal infections of cattle, sheep, and humans. Recent research has uncovered that *Fusobacterium nucleatum* is overrepresented in colonic tumors, and this bacterium can drive the progression and severity of colorectal cancer (1, 2). In humans, *Fusobacterium necrophorum* can cause the rare and life-threatening Lemierre syndrome, which begins as bacterial pharyngitis and rapidly progresses to septic thrombophlebitis of the jugular vein (3). Lemierre syndrome most frequently manifests in patients age 15 to 30 and can lead to further complications, including pharyngeal and lung abscesses (4). *F. necrophorum* have been classified into *F. necrophorum* subsp. *necrophorum* (biotype A) and *F. necrophorum* subsp. *funduliforme* (biotype B) (5). *Fusobacterium necrophorum* subsp. *funduliforme* is most commonly the cause of human infection, with *Fusobacterium necrophorum* subsp. *necrophorum* being the predominant animal pathogen. In cattle and sheep, biotype A is primarily responsible for liver abscesses and footrot (necrotic pododermatitis), for which antibiotics are an effective treatment (6).

In this study, we present the first complete genome sequence of *Fusobacterium necrophorum* subsp. *necrophorum* ATCC 25286 (5). The genome consists of one 2,678,415-bp circular chromosome (G+C content of 34%) and no plasmids. Annotation detected 2,343 protein open reading frames, 18 rRNAs, and 51 tRNAs. In addition, a 66-kb prophage region (75 protein-encoding genes; genomic position 477708 to 543489) was discovered, with the greatest similarity to a phage from the genus *Clostridium* identified. We have identified the *lktBAC* leukotoxin gene cluster and also report an increase in the number of type 5a-secreted autotransporter virulence factors compared with an annotation from the previous draft genome (87 contigs; NCBI BioProject accession number [PRJNA257880](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA257880)).

F. necrophorum subsp. *necrophorum* ATCC 25286 was purchased through the ATCC. Bacteria from a single colony were grown in Columbia Broth, supplemented with hemin (5 μ g/mL) and menadione (0.5 μ g/mL) (CBHK) to stationary phase at 37°C in an anaerobic chamber (90% N₂, 5% CO₂, and 5% H₂). Genomic DNA was isolated in deionized water (diH₂O) using a Wizard isolation kit (Promega) and was quantitated using a Qubit fluorimeter (Life Technologies, Inc.). Long-read sequences were acquired by preparing 500 ng of DNA with a rapid sequencing kit (catalog number SQK-RAD004; Oxford Nanopore Technologies) and sequencing using a SpotON flow cell (R9.4) and a MinION sequencer (Oxford Nanopore Technologies). MinKnow 3.1.8 software was used to acquire raw data and basecall DNA sequences. Sequencing adapters from Oxford Nanopore reads were removed with Porechop (7), and a total of 71,386 reads (mean

Citation Umaña A, Lemkul JA, Slade DJ. 2019. Complete genome sequence of *Fusobacterium necrophorum* subsp. *necrophorum* ATCC 25286. *Microbiol Resour Announc* 8:e00025-19. <https://doi.org/10.1128/MRA.00025-19>.

Editor David A. Baltrus, University of Arizona

Copyright © 2019 Umaña et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Daniel J. Slade, dslade@vt.edu.

Received 11 January 2019

Accepted 25 January 2019

Published 21 February 2019

read length, 1,731 bp; maximum read length, 91,983 bp) were used for genome assembly. The same DNA preparation used for MinION sequencing was used for short-read DNA sequencing on a MiSeq nano instrument at the Genomic Sequence Center at the Virginia Tech Biocomplexity Institute. Briefly, 150 ng of genomic DNA was fragmented to 400 bp using a Covaris M220 focused ultrasonicator. The ends were repaired, and an A base was added to the 3' end for ligation to the adapters, which have a single T base overhang at their 3' end. Following ligation, the libraries were amplified by 7 cycles of PCR and barcoded. The library generated was validated by the use of an Agilent TapeStation and quantitated using a Quant-iT double-stranded (dsDNA) high sensitivity (HS) kit (Invitrogen) and quantitative PCR (qPCR). The libraries were then pooled and sequenced using a MiSeq nano instrument. In summary, 999,276 reads at 150 bp each (150 mb total; 56× genome coverage) were used for genome assembly. The *F. necrophorum* ATCC 25286 genome was assembled using the open-source software package Unicycler under default parameters, version 0.4.3 (8), using our previously described pipeline (9). For Illumina and MinION data, the mean depth of coverage for the genome was 56× and 53×, respectively. Prokka 1.13 under default parameters was used to annotate genes (10). Potential integrated phage were detected using PHASTER (11). Annotation in GenBank was carried out by the NCBI Prokaryotic Genome Annotation Pipeline (12).

Data availability. The complete sequence and all raw data for the *Fusobacterium necrophorum* subsp. *necrophorum* ATCC 25286 genome have been deposited in NCBI under the accession numbers CP034842 (GenBank), PRJNA513186 (BioProject), SAMN10697414 (BioSample), and SRR8400822 and SRR8400823 (SRA raw sequencing reads). In addition, all data for this genome and additional bioinformatic analysis will be freely available on the FusoPortal (13) data repository of *Fusobacterium* genomes (<http://fusoportal.org>) and our Open Science Framework site (<https://osf.io/2c8pv/>). The deposited data are the first version of this genome.

ACKNOWLEDGMENTS

This work was supported by funding from the Commonwealth Health Research Board of Virginia and the USDA National Institute of Food and Agriculture to D.J.S. We thank the Open Access Subvention Fund of Virginia Tech for publication funding.

We thank Austin Compton (Virginia Tech) for MinION sequencing optimization.

REFERENCES

- Brennan CA, Garrett WS. 2018. *Fusobacterium nucleatum*—symbiont, opportunist and oncobacterium. *Nat Rev Microbiol*. <https://doi.org/10.1038/s41579-018-0129-6>.
- Han YW. 2015. *Fusobacterium nucleatum*: a commensal-turned pathogen. *Curr Opin Microbiol* 23:141–147. <https://doi.org/10.1016/j.mib.2014.11.013>.
- Riordan T. 2007. Human infection with *Fusobacterium necrophorum* (Necrobacillosis), with a focus on Lemierre's syndrome. *Clin Microbiol Rev* 20:622–659. <https://doi.org/10.1128/CMR.00011-07>.
- Faraone A, Fortini A, Nenci G, Boccadori C, Mangani V, Oggioni R. 2016. *Fusobacterium necrophorum* pharyngitis complicated by Lemierre's syndrome. *Case Rep Med* 2016:3608346. <https://doi.org/10.1155/2016/3608346>.
- Shinjo T, Fujisawa T, Mitsuoka T. 1991. Proposal of two subspecies of *Fusobacterium necrophorum* (Flügge) Moore and Holdeman: *Fusobacterium necrophorum* subsp. *necrophorum* subsp. nov., nom. rev. (ex Flügge 1886), and *Fusobacterium necrophorum* subsp. *funduliforme* subsp. nov., nom. rev. (ex Hallé 1898). *Int J Syst Bacteriol* 41:395–397. <https://doi.org/10.1099/00207713-41-3-395>.
- Zhou H, Bennett G, Hickford JGH. 2009. Variation in *Fusobacterium necrophorum* strains present on the hooves of footrot infected sheep, goats and cattle. *Vet Microbiol* 135:363–367. <https://doi.org/10.1016/j.vetmic.2008.09.084>.
- Wick RR. 2018. PoreChop. GitHub repository. <https://github.com/rwick/Porechop>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Todd SM, Settlege RE, Lahmers KK, Slade DJ. 2018. *Fusobacterium* genomics using MinION and Illumina sequencing enables genome completion and correction. *mSphere* 3:e00269-18. <https://doi.org/10.1128/mSphere.00269-18>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Sanders BE, Umaña A, Lemkul JA, Slade DJ. 2018. FusoPortal: an interactive repository of hybrid MinION-sequenced *Fusobacterium* genomes improves gene identification and characterization. *mSphere* 3:e00228-18. <https://doi.org/10.1128/mSphere.00228-18>.