




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

Intensive & Critical Care Nursing

journal homepage: www.elsevier.com/icc

Editorial

An outbreak of restrictive intensive care unit visiting policies



Intensive care units (ICU) are high-stress and fast-paced environments, where a specialized multidisciplinary team cares of critically ill patients (South and Adair, 2014). The discussion about patient- and family-centred care has become particularly important in the quest to understand the needs of the individuals involved as well as to improve satisfaction with the care provided in the ICUs (Ning and Cope, 2020). Among patient and family-centred care interventions, liberal ICU visiting policies have been associated with decreased rates of patients' anxiety symptoms and increased rates of family's satisfaction (Nassar Junior et al., 2018, Ning and Cope, 2020, Rosa et al., 2019).

The COVID-19 pandemic, with its high case and fatality rates, forced the world into lockdowns and social isolation that extended to critical care units. Almost every ICU has instituted a "no visitation" or, at least, a very restrictive visiting policy. These restriction visiting policies aimed mainly at reducing transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) among relatives, patients and healthcare providers. Secondly, they aimed at sparing personnel protection equipment (PPE) that could have run short and decrease healthcare team workload who already had to deal with an increasing number of severely ill patients. More than a year after all these severe restrictions were imposed, we can see all three concerns might be unfounded. First, only a minority of patients with COVID-19 had a hospital-acquired infection (Rhee et al., 2020). Second, there was no substantial shortage of PPE, at least in high and upper-middle income countries and visiting policies could have been relaxed in the second wave of the pandemic. Safe visitation could have been accomplished if visitors accurately reporting their symptoms and healthcare team helped them with the compliance with PPE (Munshi et al., 2021). Third, restrictive visiting policies ultimately led healthcare providers to regret about these restrictions and to witness hasty end-of-life decisions (Azoulay et al., 2020). For patients, an additional burden of restrictive visiting policies was delirium. In a multicentre multinational study, more than 50% of ICU patients with COVID-19 had developed delirium during their ICU stay. Family visitation (in person or virtual) was the only identifiable factor associated with a lower risk of delirium (Pun et al., 2021).

In this issue of *Intensive and Critical Care Nursing*, Jensen et al. (2022) present a study carried out in three Scandinavian countries on the challenges imposed by ICU visiting restrictions due to COVID-19 pandemic. Despite selection bias (Scandinavian ICUs only) and response bias (low return rate of questionnaires) this study sheds light on the impact of the pandemic on ICU routine and on nurses' care. All ICUs had visiting restrictions. In Denmark and Norway, most ICUs reported a restricted access. Swedish ICUs

mainly adopted a "no visiting" policy, only allowing access for relatives of dying patients. Surveyed nurses described several adversities imposed by the pandemic to the ICU routine: higher workload, a compromise on their usual professional standards of family care, an increased level of moral and emotional strain (as "they sought to balance the restrictions and the needs of patients and relatives"), among others. We can infer from this study that there were many causes for the high prevalence of psychological distress, anxiety and depression symptoms that comprised healthcare providers during the present pandemic. ICU restrictive visitation policies was one of them. Additionally, restricted visiting policies may have been impacting on patients' outcomes, such as ICU length of stay, delirium occurrence and wellbeing, and on relatives' satisfaction (Eltaybani and Ahmed, 2021, McLennan and Aggar, 2020), especially on those who could not be able to say goodbye to their loved ones, since facilitating a "good death" and humanized care at the end of life impact minimizing stress and processing grief (Otani et al., 2017).

Open visitation policies have advantages and disadvantages for patients, relatives, and healthcare providers (Table 1). Many relatives find visiting a critically ill patient in the ICU as a stressful event, fearing they will not be recognized by the patient, as well as finding them under mechanical ventilation and unable to interact (Schneeberger et al., 2020). Liberal visitation policies previously also seemed to be associated with increased burnout rates among healthcare providers (Nassar Junior et al., 2018), but this has not been confirmed in a recent randomized controlled trial (Rosa et al., 2019). As mentioned before, Jansen et al. suggested that restrictive visitation ICU policies during the current pandemic may have increased psychological distress among nurses. Therefore, evidence from clinical studies point towards more benefits than harms for patients and relatives. Jensen et al. study adds evidence this may be the case even for healthcare providers.

The COVID-19 pandemic has enormously burdened healthcare providers. However, the study of Jansen et al. has also suggested that creativity, common sense, and empathy of healthcare providers in supporting family members and facilitating communication/contact were essential for the trying to maintain humanized care within ICU. The study also stresses the importance of a qualitative data analysis to identify the subtleties of the impact of visitation restrictions on those involved. The evidence emerging from studies on ICU visitation policies during the COVID-19 pandemic has been shedding light on what practices should we modify, improve, or routinely adopt to improve our ability to provide family-centered and to preserve healthcare providers wellbeing.

Table 1
Potential advantages and disadvantages of an open visiting policy.

| | Advantages | Disadvantages | Scandinavian Experience during COVID-19 pandemic |
|--------------------------|---|--|---|
| For patients | More patient satisfaction; Promoting patient recovery by reducing stress; Positive psychological effect; Positive effects on vital signs, decreased ICU length of stay, less delirium | Patient not getting enough rest; nurses may have less time for patients; | Individual considerations must be considered; |
| For families | Less stress; less anxiety; improved communication with the healthcare team, engagement in patients' care | Harmful physiologic consequences Family members become exhausted and feel obligated to stay | Restricted access to loved ones; worse grieving (impossibility to say 'Good-bye') |
| For healthcare providers | Higher job satisfaction due to family members' positive feedback; improved communication with families, improvement in nursing care due to better knowledge of patients' values and preferences | Burnout; less time on direct patients' care; More conflicts | Additional workload to supply family absence, less knowledge of patients' values, increased stress, regret. |

References

Azoulay, E., Cariou, A., Bruneel, F., Demoule, A., Kouatchet, A., Reuter, D., et al, 2020. Symptoms of anxiety, depression, and peritraumatic dissociation in critical care clinicians managing patients with COVID-19. A cross-sectional study. *Am. J. Respir. Crit. Care Med.* 202, 1388–1398.

Eltaybani, S., Ahmed, F.R., 2021. Family satisfaction in Egyptian adult intensive care units: A mixed-method study. *Intens. Crit. Care Nurs.* 66, 103060.

Jensen, H.I., Akeman, E., Alfhelm, H.B., Frivold, G., Fridh, I., Agard, A.S., 2022. Conditions and strategies to meet the challenges imposed by the COVID-19-related visiting restrictions in the ICU: a Scandinavian cross-sectional study. *Intens. Crit. Care Nurs.* 68, 103116.

McLennan, M., Aggar, C., 2020. Family satisfaction with care in the intensive care unit: a regional Australian perspective. *Aust. Crit. Care* 33 (6), 518–525.

Munshi, L., Evans, G., Razak, F., 2021. The case for relaxing no-visitor policies in hospitals during the ongoing COVID-19 pandemic. *Can. Med. Assoc. J.* 193, E135–E137.

Nassar Junior, A.P., Besen, B., Robinson, C.C., Falavigna, M., Teixeira, C., Rosa, R.G., 2018. Flexible versus restrictive visiting policies in ICUs: a systematic review and meta-analysis. *Crit. Care Med.* 46 (7), 1175–1180.

Ning, J., Cope, V., 2020. Open visiting in adult intensive care units – A structured literature review. *Intens. Crit. Care Nurs.* 56, 102763.

Otani, H., Yoshida, S., Morita, T., Aoyama, M., Kizawa, Y., Shima, Y., et al, 2017. Meaningful communication before death, but not present at the time of death itself, is associated with better outcomes on measures of depression and complicated grief among bereaved family members of cancer patients. *J. Pain Symptom Manage.* 54 (3), 273–279.

Pun, B.T., Badenes, R., Heras La Calle, G., Orun, O.M., Chen, W., Raman, R., et al, 2021. Prevalence and risk factors for delirium in critically ill patients with COVID-19 (COVID-D): a multicentre cohort study. *Lancet Respir. Med.* 9 (3), 239–250.

Rhee, C., Baker, M., Vaidya, V., Tucker, R., Resnick, A., Morris, C.A., et al, 2020. Incidence of nosocomial COVID-19 in patients hospitalized at a large US Academic Medical Center. *JAMA Netw. Open.* 3, e2020498.

Rosa, R.G., Falavigna, M., da Silva, D.B., Sganzerla, D., Santos, M.M.S., Kochhann, R., et al, 2019. Effect of flexible family visitation on delirium among patients in the Intensive Care Unit: the ICU visits randomized clinical trial. *JAMA* 322, 216–218.

Schneeberger, A., Brandstetter, S., Bein, T., Blecha, S., Apfelbacher, C., 2020. Stressors and strains of next of kin of patients with ARDS in intensive care: A qualitative interview study using a stress-strain approach. *Intens. Crit. Care Nurs.* 57, 102783. <https://doi.org/10.1016/j.iccn.2019.102783>.

South, T., Adair, B., 2014. Open access in the critical care environment. *Crit. Care Nurs. Clin. North Am.* 26 (4), 525–532.

Silvana Soares dos Santos
Antonio Paulo Nassar Junior
Intensive Care Unit, A.C. Camargo Cancer Center, Brazil
E-mail addresses: silvana.santos@accamargo.org.br (S.S. dos Santos),
paulo.nassar@accamargo.org.br (A.P. Nassar Junior)