

Nutrition therapy for type 2 diabetes: confirmed efficacy on individualized management

Feng-Xian Wei^{1,2}

Xue-Ping Qi¹

¹Lanzhou University Second Clinical Medical College, Lanzhou University Second Hospital, Lanzhou University, Lanzhou 730030, China; ²Department of General Surgery, Lanzhou University Second Hospital, Lanzhou University Second Clinical Medical College, Lanzhou University, Lanzhou 730030, China

Dear editor

We read with great interest the article titled “Success of nutrition-therapy interventions in persons with type 2 diabetes: challenges and future directions” by Franz and Macleod,¹ recently published in your journal *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. We would like to comment on the article by explaining the key recommendations reported by the Academy of Nutrition and Dietetics Nutrition Practice Guideline for Type 1 and Type 2 Diabetes in Adults.^{2,3} Compared with their valuable review in 2014,⁴ Franz and Macleod have summarized the intervention recommendations for nutrition therapy in detail in this article, including energy intake, macronutrient composition, carbohydrate management strategies, fiber intake, glycemic index and glycemic load, nutritive sweeteners and nonnutritive sweeteners, protein intake and protein intake for diabetic kidney disease, cardioprotective eating patterns, vitamin, mineral, and/or herbal supplementation, alcohol consumption, physical activity, and glucose monitoring. It is clear that the authors have done a lot of work to make diabetes management and self-management easier for professionals and patients, respectively, and they also provide a practical alternative for use of insulin and glucose-lowering medications and surgery in clinic. A crucial question is how nutrition therapy can be effectively implemented in the individual patient. The authors provide a promising solution involving “eHealth” and “e-patient”. It will be interesting as well as innovative to apply modern digital health technology to solve this problem in the future.

We also reviewed few other studies regarding nutrition therapy for type 2 diabetes, and would like to focus on three recommendations based on literature review and also our single-center experience, which does not seem to be fully addressed in the above study but is very beneficial for applying a proper and successful nutrition therapy. First, combined efficacy monitoring is important. Nutrition therapy would be helpful to patients with type 2 diabetes when it is applied alone or combined with other therapies for different severity of diabetes. A meta-analysis showed that a Mediterranean diet improves hemoglobin A1c (HbA1c) but not fasting blood glucose.⁵ Thus, examining two indexes at least for a short-term period among fasting blood glucose, fasting insulin, and HbA1c was considered to be useful and necessary. Second, individualized nutrition therapy guided by a dietitian is important. Dietitian or trained professionals can provide more comprehensive and specific recommen-

Correspondence: Xue-Ping Qi
Lanzhou University Second Clinical Medical College, Lanzhou University Second Hospital, Lanzhou University, Cuiyingmen 82, Chengguan District, Lanzhou 730030, Gansu Province, China
Email weifx08@126.com

dations than just dietary advice given by other doctors or nurses. A meta-analysis demonstrated that a dietitian-guided nutrition therapy led to a greater improvement in HbA1c, weight, and low-density lipoprotein cholesterol.⁶ Third, the level of weight loss is important. Weight loss is an important recommendation for overweight and obese adults with type 2 diabetes; however, a weight loss of <5% was shown to not result in beneficial metabolic outcomes.⁷ Therefore, a weight loss of >5% achieved by proper nutrition therapy especially physical activity is necessary to improve HbA1c, lipid level, and blood pressure.

Clearly it is important that readers further understand nutrition therapy, and type 2 diabetes patients can benefit from its application in individualized management in clinical practice.

Disclosure

The authors report no conflicts of interest in this communication.

Reference

1. Franz MJ, Macleod J. Success of nutrition-therapy interventions in persons with type 2 diabetes: challenges and future directions. *Diabetes Metab Syndr Obes*. 2018;11:265–270.
2. Franz MJ, Macleod J, Evert A, et al. Academy of Nutrition and Dietetics Nutrition Practice Guideline for Type 1 and Type 2 Diabetes in Adults: Systematic Review of Evidence for Medical Nutrition Therapy Effectiveness and Recommendations for Integration into the Nutrition Care Process. *J Acad Nutr Diet*. 2017;117(10):1659–1679.
3. Macleod J, Franz MJ, Handu D, et al. Academy of Nutrition and Dietetics Nutrition Practice Guideline for Type 1 and Type 2 Diabetes in Adults: Nutrition Intervention Evidence Reviews and Recommendations. *J Acad Nutr Diet*. 2017;117(10):1637–1658.
4. Franz MJ, Boucher JL, Evert AB. Evidence-based diabetes nutrition therapy recommendations are effective: the key is individualization. *Diabetes Metab Syndr Obes*. 2014;7:65.
5. Carter P, Achana F, Troughton J, et al. A Mediterranean diet improves HbA1c but not fasting blood glucose compared to alternative dietary strategies: a network meta-analysis. *J Hum Nutr Diet*. 2014;27(3):280–297.
6. Møller G, Andersen HK, Snorgaard O. A systematic review and meta-analysis of nutrition therapy compared with dietary advice in patients with type 2 diabetes. *Am J Clin Nutr*. 2017;106(6):1394–1400.
7. Franz MJ, Boucher JL, Rutten-Ramos S, Vanwormer JJ. Lifestyle weight-loss intervention outcomes in overweight and obese adults with type 2 diabetes: a systematic review and meta-analysis of randomized clinical trials. *J Acad Nutr Diet*. 2015;115(9):1447–1463.

Authors' reply

Marion J Franz¹

Janice MacLeod²

¹Nutrition Concepts by Franz, Minneapolis, MN, USA; ²Clinical Innovation, WellDoc, Columbia, MD, USA

Correspondence: Marion J Franz
Nutrition Concepts by Franz, 6635, Limerick Drive, Minneapolis, MN 55439, USA
Tel +1 952 996 0434
Fax +1 952 941 6734
Email MarionFranz@aol.com

Dear editor

We thank the writers of the letter for their comments on our article regarding nutrition therapy for type 2 diabetes. Of importance are their review of important considerations for successful diabetes nutrition therapy. We would enthusiastically agree that strategic glucose monitoring is critical to assessing the efficacy of the treatment plan for type 2 diabetes and determining how to adjust the therapy plan as was well illustrated in the citation Carter et al referenced by the writers.¹ We also agree that ideally diabetes medical nutrition therapy be provided by a dietitian or similarly trained professional for all persons with diabetes. However, it is essential that all members of the health care team providing care be familiar with the basic of diabetes nutrition therapy in order to best support the individual's holistic treatment plan.

However, we would raise concerns regarding the emphasis on weight loss instead of a reduced energy (calorie) eating plan. From the review of the effectiveness of nutrition therapy for type 2 diabetes, it is clear what emerges in successfully improving glycemia is "how much" individuals are consuming.² It should be remembered that weight loss is the outcome and a reduced energy intake is the nutrition therapy intervention. For some individuals, likely those who have not implemented this strategy before or for some who have gained weight after weight loss, this will lead to weight loss. For some who have lost weight, it may prevent weight regain. The

glycemic outcome of a reduced energy intake is also dependent on how long an individual has had type 2 diabetes and whether the individual is hyperinsulinemic or insulin deficient. If the individual has progressed to an insulin-deficient state, weight loss may or may not improve glycemia. It should also be remembered that our bodies have hormonal adaptations that encourage weight gain after diet-induced weight loss and these hormonal changes likely continue for an extended period of time.³ Weight loss also results in adaptive thermogenesis (decreased resting metabolic rate) which is also maintained for long term. It is clear that after a reduced energy intake leading to weight loss our bodies adapt to prevent future weight loss and thus prevent starvation. Emphasizing the "scale" rather than food eaten can be a very frustrating experience for individuals when they maintain a low calorie eating plan and do not continue to lose weight. Therefore, the focus of nutrition therapy for individuals with type 2 diabetes is on a reduced energy intake with an emphasis on nutrient-dense, fiber-rich foods, and, of course, not be forgotten is the importance of regular physical activity.

We also strongly support the importance of health professionals and persons with type 2 diabetes understanding nutrition therapy and the importance of individualized diabetes management. Again, we thank the letter writers for their excellent letter.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Carter P, Achana F, Troughton J, et al. A Mediterranean diet improves HbA1c but not fasting blood glucose compared to alternative dietary strategies: a network meta-analysis. *J Hum Nutr Diet*. 2014;27(3):280–297.
2. Franz MJ, MacLeod J, Evert A, et al. Academy of Nutrition and Dietetics nutrition practice guideline for type 1 and type 2 diabetes in adults: systematic review of evidence for medical nutrition therapy effectiveness and recommendations for integration into the nutrition care process. *J Acad Nutr Diet*. 2017;117(10):1659–1679.
3. Evert AB, Franz MJ. Why weight loss maintenance is difficult. *Diabetes Spectr*. 2017;30(3):153–156.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>

opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.