

ARTICLE

Open Access

Inhibition of IKK β /NF- κ B signaling pathway to improve Dasatinib efficacy in suppression of cisplatin-resistant head and neck squamous cell carcinoma

Zeijia Yang¹, Jipei Liao¹, Kevin J. Cullen¹ and Hancai Dan^{1,2}

Abstract

Proto-oncogene tyrosine-protein kinase Src plays an important role in Head and Neck Squamous Cell Carcinoma (HNSCC). However, the FDA-approved SRC inhibitor Dasatinib shows very limited efficacy in HNSCC clinical trials, even though Dasatinib can completely inhibit SRC in the laboratory setting. These results suggest that SRC inhibition can cause compensatory up-regulation and/or activation of other survival pathways, which suggests that co-targeting of SRC and the potential signaling pathways may improve the Dasatinib efficacy. In this study, we investigated the role of IKK β /NF- κ B in regulation of the sensitivity of cisplatin-resistant HNSCC to Dasatinib. Additionally, we wished to determine whether inhibition of the IKK β /NF- κ B signaling pathway could enhance Dasatinib efficacy to inhibit cisplatin-resistant HNSCC without the use of cisplatin. Previous studies have shown that ETS-1 is a crucial SRC effector protein that regulates cancer cell proliferation, anti-apoptosis, and metastasis. We found that SRC kinase inhibition by Dasatinib decreased ETS-1 expression but caused elevation of IKK β /NF- κ B signaling in multiple cisplatin-resistant HNSCC. Interestingly, inhibition of IKK β /NF- κ B by CmpdA (Bay65-1942), a recently identified IKK β inhibitor, also led to a decrease in ETS-1 levels. Moreover, the knockdown of IKK, but not NF- κ B, dramatically decreased ETS-1 expression. In addition, IKK β and ETS-1 interacted in cisplatin-resistant HNSCC. These data demonstrated cross-talk between SRC and IKK to regulate NF- κ B and ETS-1. Furthermore, we found that simultaneous inhibition of SRC and IKK β through a Dasatinib and CmpdA combination synergistically inhibited NF- κ B activation and ETS-1 expression, suppressed cell proliferation, and induced apoptosis. Taken together, our data indicate that SRC and IKK β play crucial roles in cisplatin-resistant HNSCC and co-targeting SRC and IKK β could be an effective strategy to treat cisplatin-resistant HNSCC.

Introduction

Head-and-neck cancer originates in organs such as the larynx, pharynx, lips, mouth, nose, and salivary glands. Since most head-and-neck cancers begin in the squamous cells of these organs, these cancers are named head-and-neck squamous cell carcinoma (HNSCC). HNSCC is

currently the sixth cause of cancer-related deaths in the world and accounts for about 3 percent of all cancers in the United States^{1–3}. The main treatments for HNSCC include surgery, radiation therapy, chemotherapy, and immunotherapy, which are used singly or in combination for different stages of disease. Surgery or radiation therapies have proven very successful for treatment of early-stage HNSCC patients and, in combination with radiation therapy, achieve good survival rates in patients who develop loco-regional lymphoma metastasis. For all metastatic HNSCC and most recurrent HNSCC, however, chemotherapy is the only treatment option^{4–8}.

Correspondence: Hancai Dan (HDan@som.umaryland.edu)

¹Marlene and Stewart Greenebaum Comprehensive Cancer Center, University of Maryland School of Medicine, Baltimore, MD, USA

²Department of Pathology, University of Maryland School of Medicine, Baltimore, MD, USA

Edited by I. Amelio

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Cisplatin has been a major anti-cancer drug used in head-and-neck cancer therapy and is frequently combined with other chemotherapy drugs, such as Taxol and 5-Fluorouracil (5-FU)^{3,9,10}. Patients initially show a good response to cisplatin-combined chemotherapy, but nearly all patients eventually develop resistance to cisplatin and die within a year. In order to explore new therapies for cisplatin-resistant HNSCC, it is vital to define the mechanisms that confer cisplatin resistance and identify effective inhibitors to block crucial survival-signaling pathways that are elevated or activated upon cisplatin treatment^{11,12}.

Proto-oncogene tyrosine-protein kinase SRC plays an important role in HNSCC. It has been reported that, when activated, SRC promotes tumorigenesis through its downstream signaling pathways, including PI3 kinase/Akt/mTOR and MEK/ERK, to facilitate cancer growth, migration, invasion, and metastasis, as well as chemotherapy resistance^{13–15}. Therefore, SRC kinase has been an attractive target for cancer therapy, including those for HNSCC. However, the FDA approved SRC inhibitor Dasatinib shows very limited efficacy in HNSCC clinical trials, even though it completely inhibits SRC in the laboratory setting^{16–18}. This suggests that SRC inhibition can cause compensatory up-regulation and/or activation of other survival pathways, which means co-targeting SRC and other signaling pathways could improve Dasatinib efficacy.

Our long-term goal is to discover new therapeutics to treat cisplatin-resistant HNSCC through targeted therapies without the use of cisplatin. Patients who have progressed to cisplatin-resistant HNSCC may not continue to tolerate increasing doses of cisplatin. We previously reported that elevated IKK β /NF- κ B activity played an important role to control cell proliferation and cisplatin resistance¹⁹. Recently, we found that SRC signaling pathways were also up-regulated in cisplatin-resistant HNSCC, and SRC kinase regulated cisplatin-resistant HNSCC through regulation of transcription factor ETS-1²⁰. These results prompted us to investigate the molecular link between SRC and IKK β /NF- κ B in cisplatin-resistant HNSCC regulation.

In the current study, we found that treatment of cisplatin-resistant HNSCC with SRC inhibitor, Dasatinib, inhibited the SRC/ETS-1 signaling pathway, thereby leading to further elevation of the IKK β /NF- κ B pathway. Moreover, we found that IKK β interacted with ETS-1 to regulate its degradation in a manner independent of SRC and NF- κ B. Therefore, SRC and IKK β regulated ETS-1 in parallel. Subsequently, a combination of the SRC inhibitor, Dasatinib, and the IKK β inhibitor, CmpdA, led to complete inhibition of SRC, IKK β , NF- κ B, and ETS-1. Consistently, this combination synergistically suppressed cell proliferation and induced apoptosis.

Results

Inhibition of SRC and ETS-1 and induction of IKK β /NF- κ B by the SRC inhibitor Dasatinib in cisplatin-resistant HNSCC

We recently showed that Dasatinib treatment inhibited phosphorylation of SRC and decreased ETS-1 expression in cisplatin resistant HNSCC cells²⁰. Here, we wanted to determine the effect of Dasatinib treatment on IKK β /NF- κ B pathway activity. Consistent with our previous data, Dasatinib completely inhibited phosphorylation of SRC and led to a modest decrease of ETS-1 expression in cisplatin-resistant Cal27CP cells (Fig. 1a). Interestingly, phosphorylation of NF- κ B at Serine 536 and phosphorylation of IKK α / β at their activation loop increased, whereas the total levels of these proteins did not change with Dasatinib treatment (Fig. 1a). Similar results were found in cisplatin-resistant SCC25CP cell line originated from cisplatin-sensitive SCC25 cells and cisplatin-resistant FaDu-CP cell line originated from cisplatin-sensitive FaDu cells (Fig. 1b, c). In order to confirm that IKK/NF- κ B activity is elevated by Dasatinib treatment, we tested the gene expression of IL-6, a NF- κ B target gene^{21,22}. The results showed that Dasatinib significantly elevated IL-6 expression in Cal27CP cells (Fig. 1d). Our data indicate that Dasatinib inhibits SRC/ETS-1 expression but induces the IKK/NF- κ B pathway in cisplatin-resistant HNSCC.

IKK β inhibitor, CmpdA, treatment also inhibits ETS-1 expression

We treated Cal27CP, SCC25CP, and FaDu-CP cells with the IKK β inhibitor, CmpdA (Bay65-1942)²³, for 24 h and examined its effect on ETS-1 expression. CmpdA decreased phosphorylation of p65 in a dose-dependent manner, which indicated that IKK activity was inhibited. As predicted, CmpdA also decreased ETS-1 expression in a dose-dependent manner but had no effects on SRC phosphorylation (Fig. 2a–c). Our results confirm that ETS-1 is regulated by IKK through a mechanism dependent or independent of NF- κ B in cisplatin resistant HNSCC cells.

Depletion of IKK, but not NF- κ B, decreases ETS-1 expression in cisplatin-resistant HNSCC

In order to determine the molecular link between the IKK β /NF- κ B pathway and ETS-1, we used siRNA against IKK α , IKK β , or NF- κ B (p65) to decrease their expression, respectively, before examining their effects on ETS-1 expression in Cal27CP cells. Western blot results showed that these siRNA were effective at lowering the expression of their target proteins. As shown in Fig. 3a, knockdown of IKK α or IKK β , but not p65 (NF- κ B), led to decreased ETS-1 expression (Fig. 3a). Furthermore, we found that concurrent knockdown of IKK α and IKK β caused more significant decreases of ETS-1 compared to knockdown of




Fig. 1 Dasatinib inhibits SRC and ETS-1 but induces IKKβ/NF-κB in cisplatin resistant HNSCC. **a–c** Cell lysates were prepared from Cal27CP (**a**), SCC25C (**b**), and FaDu-CP (**c**) cells treated with Dasatinib for 8 h and phosphorylation and total levels of SRC, p65, IKK, and expression of ETS-1 and β-actin were detected by Western blot analysis. The density of P-p65, p65, and β-actin bands was quantitated and the level of phosphorylation of p65 was normalized by p65 and β-actin. **d** Cal27CP cells were treated with DMSO or 200 nM Dasatinib for 24 h, mRNA was extracted and the expression of IL-6 was detected by real time PCR.

either IKKα or IKKβ alone. These results suggested that both IKKα and IKKβ are involved in the regulation of ETS-1 expression (Fig. 3b) and IKKα and IKKβ regulation of ETS-1 is NF-κB independent. Moreover, we found that ETS-1 knockdown had no effect on NF-κB phosphorylation and IKK expression (Fig. 3c). Therefore, ETS-1 can act as a downstream target of IKK.



IKKβ interacts with ETS-1 and regulates its degradation

We examined whether IKK and ETS-1 could interact in Cal27CP cells. The cell lysates from Cal27CP cells were immune-precipitated with IgG control, IKKβ, or ETS-1, respectively. The immunoprecipitates were detected with antibodies against IKKβ or ETS-1, followed by Western blot analysis. The results showed that IKKβ was detected in ETS-1 antibody immunoprecipitates and ETS-1 was detected in IKKβ-antibody-immunoprecipitated lysates (Fig. 4a). Similar results were found in SCC25CP and FaDu-CP cells (Data not shown). These results suggested an interaction between IKKβ and ETS-1 in cisplatin-resistant HNSCC. We next determined whether IKKβ regulated ETS-1 through degradation control. Cal27CP and SCC25CP cells were treated with different doses of CmpdA for 24 hours prior to treatment with either DMSO control or the protease inhibitor MG-132 for two hours. Similar to the results shown in Fig. 2, CmpdA

caused dose-dependent decreases of ETS-1 expression in DMSO vehicle control-treated cells, but not in the cells treated with MG-132. As a control, the expression of p65 was not decreased by CmpdA treatment. These data indicate that IKKβ can help regulate ETS-1 degradation (Fig. 4d, e).

Dasatinib cooperates with the IKKβ inhibitor CmpdA to inhibit ETS-1 and NF-κB, as well as to induce caspase-3 cleavage

Both NF-κB and ETS-1 are involved in cell proliferation, survival, and resistance to chemo- and targeted therapies^{24–29}. We next determined whether simultaneous blockage of the SRC and IKKβ signaling pathways could lead to a significant increase in IKK/NF-κB and ETS-1 inhibition. Cal27CP cells were treated with CmpdA, Dasatinib, or a combination for 24 h. CmpdA inhibited phosphorylation of NF-κB and decreased ETS-1 expression (Fig. 5a, lane 1 versus 2). Dasatinib blocked SRC phosphorylation and decreased ETS-1 expression, while still inducing NF-κB phosphorylation (Fig. 5a, lane 1 versus lanes 2 and 4). The combination of Dasatinib and CmpdA more effectively inhibited SRC, NF-κB, and ETS-1, as well as induced significant caspase-3 cleavage (Fig. 5a, lanes 5 and 6) in comparison to either treatment alone. Similar results were found in SCC25CP cells (Fig. 5b).



These results suggest that Dasatinib cooperates with the IKKβ inhibitor to inhibit ETS-1 expression and NF-κB activity, as well as induce caspase-3 cleavage.

Dasatinib and CmpdA synergistically induce apoptosis in cisplatin-resistant HNSCC

The ability of Dasatinib and CmpdA in combination to increase caspase-3 cleavage prompted us to determine the effects of Dasatinib, CmpdA, or their combination, on apoptosis. Cal27CP cell were treated with either Dasatinib, CmpdA, or a combination for 48 h, and early and late-stage apoptosis was determined by Annexin V. Treatment of cells with 100 nM Dasatinib induced apoptosis by 11%, while treatment of cells with 5 μM CmpdA induced apoptosis by 17%; however, the combination induced apoptosis by 35% (Fig. 6a, b). Similar experiments were performed in SCC25CP cells, and the results showed that treatment with Dasatinib or CmpdA

alone induced apoptosis, whereas the combination treatment caused more (Fig. 6c, d). Our data indicate that Dasatinib and CmpdA synergistically induce apoptosis in cisplatin-resistant HNSCC.

Inhibition of IKKβ/NF-κB to improve the efficacy of Dasatinib to suppress cisplatin-resistant HNSCC

Next, we examined whether IKKβ inhibition enhanced the ability of Dasatinib to inhibit cell proliferation. Dasatinib inhibited Cal27CP cell proliferation in a dose-dependent manner. However, addition of 5 μM CmpdA to Dasatinib treatment led to increased inhibition of cell proliferation (Fig. 7a). The combination index values (CI) were analyzed according to the Chou–Talalay method³⁰ and the results showed that CI values from all of the combined inhibitor doses were less than 1 (Fig. 7a). Similar results from MTT assays were found in SCC25CP cells (Fig. 7b). We also performed colony formation assays




Fig. 4 IKKβ associates with ETS-1 and regulates its degradation. **a** The lysates from Cal27CP cells were immunoprecipitated with anti-IKKβ, anti-ETS-1, or IgG control, electrophoresed on an SDS gel, and detected with IKKβ and ETS-1 antibodies, respectively. *Note:* the red lines in panel A showed that blots were cut for detection of IKKβ (upper) and ETS-1 (bottom), respectively. **b, c** Cal27CP (**b**) and SCC25CP (**c**) cells treated with DMSO control or MG-132 for 2 h were incubated with media containing increasing concentrations of CmpdA for 24 h and lysed. The expression of ETS-1, p65, and β-actin was detected by Western blot analysis.




Fig. 5 Synergistic inhibition of IKKβ/NF-κB and ETS by combination of Dasatinib with IKKβ inhibitor, CmpdA. Cal27CP (**a**) and SCC25CP (**b**) cells were treated with vehicle control, CmpdA, Dasatinib, or a combination for 24 h, lysed, and phosphorylation and total levels of SRC and p65 and expression of ETS-1, cleaved-caspase-3, and β-actin were detected by Western blot analysis.


in Cal27CP and SCC25CP cells. Treatment of Cal27CP cells with Dasatinib or CmpdA inhibited colony formation, but the combination of Dasatinib and CmpdA significantly increased this inhibition (Fig. 7c, d). These data indicate that CmpdA improves the efficacy of Dasatinib to inhibit cell proliferation.

Discussion

We recently demonstrated that SRC/ETS-1 signaling was elevated in cisplatin-resistant HNSCC²⁰. Depletion of ETS-1 significantly impaired cell proliferation and survival, as

well as cisplatin resistance. Interestingly, inhibition of SRC by the SRC inhibitor Dasatinib only marginally diminished cell proliferation and survival. These results imply that it is important to identify the compensatory survival pathways that are up-regulated upon Dasatinib treatment.

In this study, we investigated the molecular link between SRC and the IKK/NF-κB pathway. We found that: (1) Inhibition of SRC by Dasatinib lowered ETS-1 expression but elevated the IKK/NF-κB pathway; (2) IKK interacted with ETS-1 and regulated its degradation independent of NF-κB and SRC; and (3) Combination of




Dasatinib and the IKKβ inhibitor CmpdA led to significant inhibition of ETS-1 expression, cell proliferation, and cell survival (Fig. 8). Our data explored the crucial role of IKK/NF-κB in conferring resistance to SRC inhibitors in cisplatin-resistant HNSCC.

Nuclear Factor κB (NF-κB) plays important roles in the regulation of cell proliferation and survival, as well as resistance to chemo- and targeted therapies^{31,32}. Many factors are involved in IKK/NF-κB activation^{33–35}. Our current study showed that inhibition of SRC kinase led to up-regulation of IKK/NF-κB, but we are not currently clear on the precise mechanism(s) by which Dasatinib activates IKKβ/NF-κB pathway. Our data are consistent with the results from the study by Wolf, et al., which showed that SRC inhibition activated IKK/NF-κB to increase IL-12 synthesis on TLR-mediated activation in dendritic cells (DCs)³⁶. It is important to further determine how Dasatinib can induce IKK/NF-κB and whether or not other SRC kinase family members such as Fgr, Hck, and Lyn are also involved in this process.

The transcription factor ETS-1 plays an important role in several cancers, including breast, lung, and head-and-neck^{25,27,37–39}. Previous studies have shown that SRC kinase inhibits ETS-1 degradation through phosphorylation and inhibition of the tumor suppressor gene protein FRWD2 (COP1)³⁸. In this study, we demonstrated that knockdown of IKKβ, but not NF-κB, decreased ETS-1 expression, but had no effect on SRC phosphorylation and expression. These data suggest that IKKβ regulates ETS-1 through mechanisms independent of SRC. It would be very interesting to define the more detailed mechanisms by which IKKβ regulates ETS-1 in cisplatin-resistant HNSCC.

The current study emphasized the functional interaction of SRC, ETS, and IKK/NF-κB to control proliferation and survival, as well as on the efficacy of combining Dasatinib with the IKKβ inhibitor, CmpdA, to inhibit cell proliferation through simultaneous inhibition of NF-κB in cisplatin-resistant HNSCC. It should be noted that Dasatinib also inhibits Bcl-Abl, which also contributes to



NF- κ B/IKK activation⁴⁰. It has been reported that blockade of Bcl-Abl-induced NF- κ B activation via IKK β inhibition is effective in suppression of chronic myelogenous leukemia. It might also be important to test if a combination of the Bcr-Abl inhibitor Imatinib with CmpdA synergistically inhibits cisplatin-resistant HNSCC.

Materials and methods

Cell culture

HNSCC cell lines, Cal 27, FaDu, and SCC25, were obtained from ATCC. Cisplatin-resistant Cal27 cells (Cal27CP), SCC25 cells (SCC25CP) and FaDu cells (FaDu-CP) were generated from parental Cal27 and SCC25 cells through long-term treatment with cisplatin (0.5–5 μ M) until the cells can grow normally in media with 5 μ M cisplatin. The cell lines were authenticated by short tandem repeat analysis (STR) and tested for mycoplasma contamination in the Translational Core Facility of the University of Maryland Marlene and Stewart Greenebaum Cancer Center. All cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2 mM glutamine, and 100 U/mL penicillin and streptomycin (Gibco).

Antibodies and inhibitors


The following antibodies were purchased from Cell Signaling Technology (CST): phospho-SRC-Y416 (CST-2101), SRC (CST-2123), ETS-1 (CST-14069), phospho-p65-S536 (CST-3033), p65 (CST-6956), phospho-IKK α (S176)/ β (S177) (CST-2697 and CST-2078), IKK α (CST-2682), IKK β (CST-8943), cleaved-caspase 3 (CST-9664), and β -Actin (CST-4967). Dasatinib was from Selleck Chemicals. IKK β inhibitor, CmpdA was a gift from Dr. Albert Baldwin (University of North Carolina at Chapel Hill, Chapel Hill, NC, USA).

Cell lysis and Western blot analysis

Cells were lysed and Western blot experiments were performed as described previously^{19,41}.

siRNA knockdown experiment

Nonspecific control siRNAs and siRNA SMARTpool IKK α , IKK β and NF- κ B (p65) were purchased from Dharmacon. Cells were transfected with non-target siRNA, and siRNA against IKK α , IKK β , or p65 using Lipofectamine Rnaimax Transfection Reagent (Thermo Scientific) according to the manufacturer's instructions.



Analyzing apoptosis by Annexin V/propidium iodide staining

Cells treated with inhibitor(s) for two days were trypsinized, washed with PBS and Annexin V binding buffer, and re-suspended in 1 mL Annexin V binding buffer. 2×10^5 cells were then stained with 0.5 μ L of Annexin V and 0.7 μ L of propidium iodide (PI) for 15 min at room temperature. Staining was then analyzed by flow cytometry on BD FACSCanto II™ Cell Analyzer (BD Biosciences). Results were analyzed by FCS Express 6. All experiments were performed in twice by triplicate and statistical analysis was performed (mean \pm SD).

Cell proliferation assays

Cell proliferation was assessed by MTS assay using the CellTiter 96 Aqueous ONE Solution kit (Promega) as described previously⁴¹. In brief, 5×10^4 cells/mL were seeded into 96-well plates for 24 h. The next day, media were replaced with fresh media that contained the indicated concentrations of Dasatinib, CmpdA, a combination of Dasatinib and CmpdA, or the vehicle control (DMSO). After an additional 72 h incubation, MTS reagent (20 μ L) was added to each well and cells with the reagent were incubated at 37°C for 2 h. Absorbance at 490 nm was measured using a microplate reader (Bio-Rad). Each experiment was performed in triplicate. In order to

determine synergy of drug combination, the combination index values were determined according to the Chou–Talalay method (26) using CalcuSyn software.

Colony formation assay

1000 cells were seeded in 12-well plates. The next day, cells were treated with vehicle control, Dasatinib, CmpdA or a combination for 48 hours, and then grown in normal media for 10 to 14 days. After gently washing once with 1 \times PBS, cells were fixed with methanol and stained with crystal violet.

Statistical analysis

All data are shown as mean \pm SD. Statistical analysis was performed using GraphPad Prism version 7.04 (GraphPad Software Inc.).

Acknowledgements

We thank Dr. Albert S. Baldwin for providing the IKKβ inhibitor, CmpdA. We also thank Drs. Antonino Passaniti and Amy Fulton for critical discussion. This study was supported, in part, by the grants from the NIH National Cancer Institute (NCI) to H.D. (R00CA149178 and R01CA212094), and the University of Maryland Marlene and Stewart Greenebaum Comprehensive Cancer Center.

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 2 September 2019 Revised: 17 January 2020 Accepted: 4 February 2020

Published online: 15 May 2020

References

- Vokes, E. E., Weichselbaum, R. R., Lippman, S. M. & Hong, W. K. Head and neck cancer. *N. Engl. J. Med.* **328**, 184–194 (1993).
- Seeburg, D. P., Baer, A. H. & Aygun, N. Imaging of patients with head and neck cancer: from staging to surveillance. *Oral Maxil. Surg. Clin. N. Am.*, <https://doi.org/10.1016/j.coms.2018.06.004> (2018).
- Pendleton, K. P. & Grandis, J. R. Cisplatin-based chemotherapy options for recurrent and/or metastatic squamous cell cancer of the head and neck. *Clin. Med. Insights Ther.*, <https://doi.org/10.4137/cmt.S10409> (2013).
- Du, Y., Peyser, N. D. & Grandis, J. R. Integration of molecular targeted therapy with radiation in head and neck cancer. *Pharmacol. Ther.* **142**, 88–98 (2014).
- Chai, R. L. & Grandis, J. R. Advances in molecular diagnostics and therapeutics in head and neck cancer. *Curr. Treat. Options Oncol.* **7**, 3–11 (2006).
- Fung, C. & Grandis, J. R. Emerging drugs to treat squamous cell carcinomas of the head and neck. *Expert Opin. Emerg. Drugs* **15**, 355–373 (2010).
- Santuray, R. T., Johnson, D. E. & Grandis, J. R. New therapies in head and neck cancer. *Trends Cancer* **4**, 385–396 (2018).
- Schick, U., Huguet, F., Pointreau, Y. & Pradier, O. [Radiotherapy for head and neck squamous cell carcinoma: state of the art and future directions]. *Cancer Radiother.* **21**, 498–504 (2017).
- Argiris, A. et al. Induction docetaxel, cisplatin, and cetuximab followed by concurrent radiotherapy, cisplatin, and cetuximab and maintenance cetuximab in patients with locally advanced head and neck cancer. *J. Clin. Oncol.* **28**, 5294–5300 (2010).
- Siegel, R., Naishadham, D. & Jemal, A. Cancer statistics, 2013. *CA Cancer J. Clin.* **63**, 11–30 (2013).

11. Guidi, A., Codeca, C. & Ferrari, D. Chemotherapy and immunotherapy for recurrent and metastatic head and neck cancer: a systematic review. *Med. Oncol.* **35**, 37 (2018).
12. Lo Nigro, C., Denaro, N., Merlotti, A. & Merlano, M. Head and neck cancer: improving outcomes with a multidisciplinary approach. *Cancer Manag. Res.* **9**, 363–371 (2017).
13. Morgan, S. & Grandis, J. R. ErbB receptors in the biology and pathology of the aerodigestive tract. *Exp. Cell Res.* **315**, 572–582 (2009).
14. Kundu, S. K. & Nestor, M. Targeted therapy in head and neck cancer. *Tumour Biol.* **33**, 707–721 (2012).
15. Montero, J. C., Seoane, S., Ocana, A. & Pandiella, A. Inhibition of SRC family kinases and receptor tyrosine kinases by dasatinib: possible combinations in solid tumors. *Clin. Cancer Res.* **17**, 5546–5552 (2011).
16. Bozec, A., Peyrade, F. & Milano, G. Molecular targeted therapies in the management of head and neck squamous cell carcinoma: recent developments and perspectives. *Anticancer Agents Med. Chem.* **13**, 389–402 (2013).
17. Zibelman, M. & Mehra, R. Overview of current treatment options and investigational targeted therapies for locally advanced squamous cell carcinoma of the head and neck. *Am. J. Clin. Oncol.* **39**, 396–406 (2016).
18. Gnoni, A., Marech, I., Silvestri, N., Vacca, A. & Lorusso, V. Dasatinib: an anti-tumour agent via Src inhibition. *Curr. Drug Targets* **12**, 563–578 (2011).
19. Li, Z. et al. IKK phosphorylation of NF-kappaB at serine 536 contributes to acquired cisplatin resistance in head and neck squamous cell cancer. *Am. J. Cancer Res.* **5**, 3098–3110 (2015).
20. Yang, Z. et al. Regulation of cisplatin-resistant head and neck squamous cell carcinoma by the SRC/ETS-1 signaling pathway. *BMC Cancer* **19**, 485 (2019).
21. Karin, M. NF-kappaB as a critical link between inflammation and cancer. *Cold Spring Harb. Perspect. Biol.* **1**, a000141 (2009).
22. He, G. & Karin, M. NF-kappaB and STAT3 - key players in liver inflammation and cancer. *Cell Res.* **21**, 159–168 (2011).
23. Ziegelbauer, K. et al. A selective novel low-molecular-weight inhibitor of IkkappaB kinase-beta (IKK-beta) prevents pulmonary inflammation and shows broad anti-inflammatory activity. *Br. J. Pharmacol.* **145**, 178–192 (2005).
24. Kato, T. et al. ETS1 promotes chemoresistance and invasion of paclitaxel-resistant, hormone-refractory PC3 prostate cancer cells by up-regulating MDR1 and MMP9 expression. *Biochem. Biophys. Res. Commun.* **417**, 966–971 (2012).
25. Dittmer, J. The role of the transcription factor Ets1 in carcinoma. *Semin. Cancer Biol.* **35**, 20–38 (2015).
26. Garrett-Sinha, L. A. Review of Ets1 structure, function, and roles in immunity. *Cell. Mol. Life Sci.* **70**, 3375–3390 (2013).
27. Shaikhibrahim, Z. & Wernert, N. ETS transcription factors and prostate cancer: the role of the family prototype ETS-1 (review). *Int. J. Oncol.* **40**, 1748–1754 (2012).
28. Wang, C. Y., Cusack, J. C. Jr., Liu, R. & Baldwin, A. S. Jr. Control of inducible chemoresistance: enhanced anti-tumor therapy through increased apoptosis by inhibition of NF-kappaB. *Nat. Med.* **5**, 412–417 (1999).
29. Wang, C. Y., Mayo, M. W. & Baldwin, A. S. Jr. TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF-kappaB. *Science* **274**, 784–787 (1996).
30. Chou, T. C. Drug combination studies and their synergy quantification using the Chou-Talalay method. *Cancer Res.* **70**, 440–446 (2010).
31. Orłowski, R. Z. & Baldwin, A. S. Jr. NF-kappaB as a therapeutic target in cancer. *Trends Mol. Med.* **8**, 385–389 (2002).
32. Basseres, D. S. & Baldwin, A. S. Nuclear factor-kappaB and inhibitor of kappaB kinase pathways in oncogenic initiation and progression. *Oncogene* **25**, 6817–6830 (2006).
33. Baldwin, A. S. Regulation of cell death and autophagy by IKK and NF-kappaB: critical mechanisms in immune function and cancer. *Immunol. Rev.* **246**, 327–345 (2012).
34. Karin, M. The IkkappaB kinase - a bridge between inflammation and cancer. *Cell Res.* **18**, 334–342 (2008).
35. Luo, J. L., Kamata, H. & Karin, M. IKK/NF-kappaB signaling: balancing life and death-a new approach to cancer therapy. *J. Clin. Invest* **115**, 2625–2632 (2005).
36. Wolf, M. et al. Src-kinase inhibitors sensitize human cells of myeloid origin to Toll-like-receptor-induced interleukin 12 synthesis. *Blood* **122**, 1203–1213 (2013).
37. Balko, J. M. et al. Activation of MAPK pathways due to DUSP4 loss promotes cancer stem cell-like phenotypes in basal-like breast cancer. *Cancer Res.* **73**, 6346–6358 (2013).
38. Lu, G. et al. Phosphorylation of ETS1 by Src family kinases prevents its recognition by the COP1 tumor suppressor. *Cancer Cell* **26**, 222–234 (2014).
39. Calli, A. O., Sari, A., Cakalagaoglu, F., Altinboga, A. A. & Oncel, S. ETS-1 proto-oncogene as a key newcomer molecule to predict invasiveness in laryngeal carcinoma. *Pathol. Res. Pract.* **207**, 628–633 (2011).
40. O'Hare, T. et al. Combined Abl inhibitor therapy for minimizing drug resistance in chronic myeloid leukemia: Src/Abl inhibitors are compatible with imatinib. *Clin. Cancer Res.* **11**, 6987–6993 (2005).
41. Zhang, Y. et al. Targeting IkkappaB kinase beta/NF-kappaB signaling in human prostate cancer by a novel IkkappaB kinase beta inhibitor CmpdA. *Mol. Cancer Therapeutics* **15**, 1504–1514 (2016).