

# Effects of Inappropriate Nurturing Experiences, Depressive Rumination, and Trait Anxiety on Depressive Symptoms

Yunosuke Usui<sup>1</sup>, Miki Ono<sup>1</sup>, Rintaro Nibuya<sup>1</sup>, Masayuki Kikkawa<sup>1</sup>, Shunichiro Ito<sup>1</sup>,  
Chihiro Morishita<sup>1</sup>, Mina Honyashiki<sup>1</sup>, Yu Tamada<sup>1,2</sup>, Takeshi Inoue<sup>1</sup>, Jiro Masuya<sup>1</sup>

<sup>1</sup>Department of Psychiatry, Tokyo Medical University, Shinjuku-ku, Tokyo, Japan; <sup>2</sup>Department of Psychiatry, Tokyo Medical University Hachioji Medical Center, Hachioji-shi, Tokyo, Japan

Correspondence: Jiro Masuya, Department of Psychiatry, Tokyo Medical University, 6-7-1 Nishi-shinjuku, Shinjuku-ku, Tokyo, 160-0023, Japan, Tel +81-(0)3-3342-6111, Email [j-masuya@tokyo-med.ac.jp](mailto:j-masuya@tokyo-med.ac.jp)

**Background:** Prior research has shown that inappropriate childhood nurturing experiences (low care and high overprotection), trait anxiety, and depressive rumination are risk factors for depression. However, no studies to date have analyzed the overall association between these factors and depressive symptoms. In the present study, we hypothesized that depressive rumination mediates the impacts of inappropriate childhood nurturing experiences on depressive symptoms, and that these mediating effects are moderated by trait anxiety, and tested these hypotheses in adult volunteers.

**Methods:** The subjects were adult volunteers who were investigated between April 2017 and April 2018. A self-report questionnaire on demographic data, childhood nurturing experiences, trait anxiety, depressive rumination, and depressive symptoms was distributed to conduct the survey, and written informed consent and valid responses were obtained from 585 subjects. Mediation and moderated-mediation analyses were performed by SPSS 28 and macro PROCESS 4.0 software. This study was approved by the Ethics Committee of Tokyo Medical University.

**Results:** Parental care showed a significant negative indirect effect on depressive symptoms via its effect on depressive rumination ( $p < 0.01$ ), whereas parental overprotection showed the opposite effect ( $p < 0.01$ ). Furthermore, the mediation effect of depressive rumination on depressive symptoms was increased by trait anxiety ( $p < 0.05$ ).

**Conclusion:** Our present study demonstrated that the main factor affecting depressive symptoms is inappropriate childhood nurturing experiences, which indirectly enhance depression by intensifying depressive rumination, and that depressive rumination and trait anxiety mutually reinforce each other to enhance depressive symptoms. These findings may be useful for the prevention of depressive symptoms. Large-scale prospective studies are needed to confirm the causal associations among these factors in the future.

**Keywords:** childhood parenting experiences, depressive rumination, trait anxiety, depressive symptoms, moderated-mediation analysis

## Introduction

Depression is a common psychiatric disorder that causes depressed mood, low motivation, anxiety, insomnia or hypersomnia, and a decrease or increase in appetite over a long period of time.<sup>1</sup> Although the causes of depression have not been fully clarified to date, environmental factors, such as inappropriate nurturing from parents (low care and high overprotection), abuse, and bullying victimization by peers during childhood and harassment in adulthood are reported risk factors for the onset of depression.<sup>2–8</sup> Regarding environmental factors, several studies have reported that childhood stress, caused by factors such as inappropriate parental nurturing and abuse experienced during childhood, influences depression in adulthood via their effects on children's personality traits.<sup>6,9–19</sup> Cross-sectional studies have suggested that inappropriate nurturing experienced in childhood has long-term effects on depression and depressive symptoms in adulthood, particularly through its effects on personality traits, such as trait anxiety, neuroticism, self-

esteem, interpersonal sensitivity, and affective temperaments.<sup>9,13,18,20–22</sup> The Parental Bonding Instrument (PBI), which is a widely used self-report questionnaire for the retrospective assessment of nurturing experiences, evaluates childhood nurturing experiences in terms of care and overprotection.<sup>23</sup> Previous studies using the PBI have reported that inappropriate parental nurturing during childhood influences depressive symptoms, and that its effects on depression vary depending on the type of nurturing experience.<sup>2,6,13,24</sup>

Depressive rumination is continuous negative thinking about one's own depressive symptoms, and the causes and consequences of depression.<sup>25</sup> Nolen-Hoeksema pointed out the importance of depressive rumination as a psychological pathogenic mechanism of depression and depressive symptoms, and rumination has recently received clinical attention and has been applied in the treatment of depression.<sup>25–30</sup> The influences of childhood experiences of abuse and peer victimization on depressive symptoms are mediated by their effects on depressive rumination.<sup>9,31,32</sup> Inappropriate childhood nurturing experiences are also known to affect depressive rumination in addition to depression.<sup>6,13,18,29</sup> A prospective study analyzed the effects of childhood experiences of inappropriate maternal nurturing on depression and depressive symptoms, and showed the mediation by depressive rumination.<sup>33</sup> However, a limitation of their study was that they did not analyze paternal nurturing. On the other hand, various factors may have a moderating effect on the mediating association of the inappropriate childhood nurturing experience-depressive rumination-depression triangle, but this has never been reported previously.<sup>34</sup>

Anxiety and depression are closely associated with each other.<sup>35</sup> In psychiatric disorders, anxiety and depression often coexist, but in most cases, anxiety precedes and depression follows.<sup>36,37</sup> In other words, anxiety is a risk factor for the development of depression.<sup>37</sup> High levels of trait anxiety, which is an anxiety-prone personality trait, make people more likely to develop depression and anxiety disorders.<sup>38</sup> The biological basis of trait anxiety is the well-studied neuroendocrine response of the hypothalamic-pituitary-adrenocortical system, which is associated with trait anxiety and an exaggerated response to stress load.<sup>38</sup> Psychologically, people who are more prone to trait anxiety may be more vulnerable to depression when exposed to a stressor, as the neurocognitive cascade associated with the body is stimulated, which increases susceptibility to stress, resulting in cognitive and behavioral changes.<sup>38,39</sup> Trait anxiety is also closely associated with depressive rumination, and trait anxiety and depressive rumination reportedly have a cascading mediating effect on the impacts of childhood abuse and bullying victimization on depressive symptoms.<sup>9,32</sup> Furthermore, trait anxiety is also associated with inappropriate childhood nurturing experience.<sup>40</sup> Because of the nature of trait anxiety as an individual stress response trait as described above, it is possible that trait anxiety may have a moderating effect on the mediating association of the inappropriate childhood nurturing experience-depressive rumination-depression triangle, but no studies have investigated the possibility of such a moderated-mediation effect.

As mentioned above, some of the associations among the 4 variables, ie, inappropriate childhood nurturing experience, trait anxiety, depressive rumination, and depression, have been reported in previous studies, but no studies to date have analyzed the overall associations among the 4 variables in terms of mediation and moderated-mediation effects. It is also necessary to analyze whether trait anxiety and depressive rumination differ in their impacts on depressive symptoms caused by the experience of inappropriate childhood nurturing, depending on the father and mother or care and overprotection, but this has not been fully clarified to date. For psychosocial analysis of these 4 complex variables, a combined model of mediation and moderation effects is useful, as described by Baron and Kenny.<sup>34</sup> In the present study, we conducted moderated-mediation analysis, hypothesizing that depressive rumination mediates the effects of inappropriate childhood nurturing experiences on depressive symptoms, and that trait anxiety moderates these mediating effects. This is because trait anxiety precedes depression and affects both depressive rumination and depression.<sup>9,32,36,37</sup>

## Subjects and Methods

### Subjects

A cross-sectional self-report questionnaire survey was conducted on adult volunteers between April 1, 2017 and April 30, 2018. The study subjects were recruited by convenience sampling. The study was part of a larger study.<sup>9,32</sup> Anonymous questionnaires were collected until September 30, manual data input from the paper questionnaires to a computer was completed on December 31, and databases were checked and cleaned up to April 30, 2018. A total of 597 adult volunteers participated in

this study, and 12 subjects were excluded owing to many missing data. A demographic questionnaire and 4 questionnaires on childhood nurturing, depressive rumination, trait anxiety, and depressive symptoms were administered. There were 585 subjects (249 men and 336 women; age:  $41.7 \pm 12.1$  years) whose written consent was obtained and for whom valid responses were obtained anonymously. The inclusion criterion was an age of 20 years or older, and the exclusion criteria were not having any serious mental or physical illnesses. The subjects were informed that their participation in the study was voluntary, that they would not be disadvantaged in any way by nonparticipation, that their personal information would be completely anonymized, that their data would be kept strictly confidential, and that their personal information would be handled with the utmost care. This study was conducted in compliance with the Declaration of Helsinki (amended in 2013) and with the approval of the Ethics Committee of Tokyo Medical University (study approval no.: SH3502).

## Questionnaires

### Parental Bonding Instrument (PBI)

Adults were asked to recall their experiences as children, and their childhood nurturing experiences were assessed by the PBI Japanese version of the self-administered questionnaire.<sup>23,41</sup> The following 2 factors were rated for each of the father and mother: overprotection (13 items with a total of 0 to 39 points) and care (12 items with a total of 0 to 36 points). A higher overprotection score means that the subjects perceive their parents as being controlling, interfering, and indifferent, whereas a higher care score means that they perceive their parents as giving emotional warmth, empathy, and closeness. In the present study, the father's and mother's care and overprotection scores were used in the analysis. Cronbach's  $\alpha$  coefficients calculated for the individual subscales of the PBI were found to be 0.866 for overprotection by the father, 0.920 for care by the father, 0.873 for overprotection by the mother, and 0.915 for care by the mother, and indicate excellent internal consistency.

### Ruminative Responses Scale (RRS)

The RRS is a self-report questionnaire that measures how frequently the subjects experience depressive rumination.<sup>42</sup> The questionnaire consists of 22 items, and each item is rated on a 4-point scale (1 to 4 points). Higher total scores indicate more frequent depressive rumination. The Japanese version of the RRS was translated by Hasegawa, and its reliability and validity have been verified.<sup>43</sup> The Cronbach's  $\alpha$  coefficient for the total score of this scale was 0.944, indicating excellent internal consistency.

### State-Trait Anxiety Inventory Form Y (STAI-Y)

The STAI-Y is a self-report questionnaire that evaluates trait and state anxiety. For the present analysis, trait anxiety was used, which is a relatively stable response tendency to anxiety-provoking experiences. The trait anxiety section consisted of 20 items. The respondents were asked to rate how they usually feel using a 4-point scale (1 to 4 points).<sup>44</sup> Higher total scores indicate higher trait anxiety. The Japanese version was translated by Hidano et al, and its validity and reliability have been verified.<sup>45</sup> The Cronbach's  $\alpha$  coefficient for the total score of the trait anxiety subscale was 0.925, indicating excellent internal consistency.


### Patient Health Questionnaire-9 (PHQ-9)

The PHQ-9 is a self-report depression rating scale.<sup>46</sup> Nine items of depressive symptoms experienced by the subject in the previous 2 weeks were rated using a 4-point scale (0 to 3 points). Total scores reflecting symptom severity were used for analysis in this study. The Japanese version of the PHQ-9 has been translated, and its validity and reliability have been verified.<sup>47</sup> The Cronbach's  $\alpha$  coefficient for the total score of this scale was 0.854, indicating good internal consistency.


## Statistical Analysis

In the present study, the effects of inappropriate childhood nurturing experiences (PBI) on depressive symptoms (PHQ-9) were hypothesized to be mediated by depressive rumination, and trait anxiety (STAI-Y) was hypothesized to moderate the effects of inappropriate childhood nurturing experiences and depressive rumination on depressive symptoms and the mediation effect of depressive rumination. The association and correlation of demographic information and questionnaire data, and the mediation and moderated-mediation analyses were conducted using SPSS 28 software (IBM, Armonk, NY, USA) and its macro PROCESS 4.0 (models 4 and 15). Model 4 was used for the mediation analysis, and Model 15 for the moderated-mediation analysis (Figure 1). In both models, age and sex were included in the analysis as covariates.

(A) Mediation model (Model 4)



(B) Moderated-mediation model (Model 15)



**Figure 1** Mediation model (A, model 4) and moderated-mediation model (B, model 15) of inappropriate childhood nurturing experiences (PBI), trait anxiety (STAI-Y), depressive rumination (RRS), and depressive symptoms (PHQ-9). The data of (A) are shown in Table 2. The data of (B) are shown in Table 3. PBI, Parental Bonding Instrument; PHQ-9, Patient Health Questionnaire-9; RRS, Ruminative Responses Scale; STAI-Y, State-Trait Anxiety Inventory form Y.

Standardized coefficients were used for Model 4, and unstandardized coefficients were used for Model 15. A *p*-value of less than 0.05 was considered to indicate a statistically significant difference between groups.

Regarding the handling of missing values, as subjects with many questions being left unanswered on their questionnaires were excluded, the subjects included in the analysis had only a few missing values and no missing values for the mediation and moderated-mediation analyses. Imputation methods were not used for the analysis.

This study was part of a larger study, which investigated stress, personality traits, affective symptoms, sleep, quality of life, well-being, resilience, etc., in adult volunteers by a questionnaire survey. In the original larger study, multivariable analysis of more than 30 independent variables was initially performed, and after consideration of missing data, 597 subjects were initially enrolled.<sup>48,49</sup> In our present study, subjects with a lot of missing data were excluded, and 585 of the subjects were analyzed. This is the reason for this sample size.

## Results

### Demographic Characteristics, and PHQ-9, PBI, Trait Anxiety of STAI-Y, and RRS Scores of the Subjects

Table 1 shows the results of the correlations and associations of demographic information and data from each questionnaire with depressive symptoms (PHQ-9 total scores) in the 585 subjects. Sex, marital status, and past history of psychiatric disease showed a significant association with PHQ-9 scores, but the other demographic characteristics

**Table 1** Demographic Characteristics, Scores of STAI-Y, PBI, RRS, and PHQ-9, and Their Correlation with or Effects on PHQ-9 Scores

Characteristic or Measure	Value (% or Mean ± SD)	Correlation with PHQ-9 score (r) or Effect on PHQ-9 Score (mean ± SD of PHQ-9 score, <i>t</i> -test)
Age (Years)	41.7 ± 12.1	<i>r</i> = -0.034, <i>p</i> = 0.406
Sex (Male: Female)	42.6: 57.4	Male 3.5 ± 4.1 vs female 4.5 ± 4.3, <i>p</i> = 0.006 ( <i>t</i> -test)
Education Years	14.6 ± 1.8	<i>r</i> = -0.068, <i>p</i> = 0.100
Marital Status (Married: Unmarried)	66.0: 34.0	Married 3.5 ± 3.9 vs unmarried 5.2 ± 4.7, <i>p</i> < 0.001 ( <i>t</i> -test)
Past History Of Psychiatric Disease (Yes: No)	11.6: 88.4	Yes 6.7 ± 5.4 vs no 3.7 ± 3.9, <i>p</i> < 0.001 ( <i>t</i> -test)
STAI-Y Score (Trait Anxiety)	42.9 ± 10.4	<i>r</i> = 0.639, <i>p</i> < 0.001

(Continued)

**Table 1** (Continued).

Characteristic or Measure	Value (% or Mean ± SD)	Correlation with PHQ-9 score (r) or Effect on PHQ-9 Score (mean ± SD of PHQ-9 score, t-test)
PBI Father Care Score	23.5 ± 8.2	$r = -0.178, p < 0.001$
PBI Father Overprotection Score	9.7 ± 6.9	$r = 0.209, p < 0.001$
PBI Mother Care Score	28.0 ± 7.0	$r = -0.273, p < 0.001$
PBI Mother Overprotection Score	9.7 ± 7.0	$r = 0.240, p < 0.001$
RRS Total Score	35.2 ± 11.4	$r = 0.500, p < 0.001$
PHQ-9 Score	4.1 ± 4.2	

**Notes:** Data are presented as the percentage or mean ± SD,  $r$  = Pearson's correlation coefficient.

**Abbreviations:** SD, standard deviation; STAI-Y, State-Trait Anxiety Inventory form Y; PBI, Parental Bonding Instrument; RRS, Ruminative Responses Scale; PHQ-9, Patient Health Questionnaire-9.

**Table 2** Mediation Effect of Depressive Rumination (RRS) on the Association Between Parenting (PBI) and Depressive Symptoms (PHQ-9) (Analysis by Model 4 of PROCESS)

PBI	PBI to RRS (a)	PBI to PHQ-9 (b)	RRS to PHQ-9 (c)	Mediation Effect of RRS (95% CI)	R <sup>2</sup>
Father Care	-0.206***	-0.083*	0.474***	-0.098** (-0.142, -0.056)	0.260
Mother Care	-0.275***	-0.147**	0.457***	-0.126** (-0.175, -0.082)	0.277
Father OP	0.218***	0.112**	0.466***	0.101** (0.060, 0.146)	0.264
Mother OP	0.298***	0.097*	0.467***	0.139** (0.094, 0.189)	0.266

**Notes:** (a), (b), and (c) are shown in the model in [Figure 1A](#). This analysis controlled for age and sex. Mediation effects indicate indirect effects of PBI on PHQ-9 via RRS. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ . R<sup>2</sup> indicates the coefficients of determination.


**Abbreviations:** CI, confidence interval; OP, overprotection; PBI, Parental Bonding Instrument; PHQ-9, Patient Health Questionnaire-9; RRS, Ruminative Responses Scale.

**Table 3** Moderated-Mediation Analysis: Mediating Effect of Depressive Rumination (RRS) on the Association Between Parenting (PBI) and Depressive Symptoms (PHQ-9), and Moderation by Trait Anxiety (STAI-Y) of the Effects of Parenting (PBI) and Depressive Rumination (RRS) on Depressive Symptoms (PHQ-9)

PBI	Effect of STAI-Y on PHQ-9	Moderation by STAI-Y on Effect of PBI on PHQ-9 (a)	Moderation by STAI-Y on Effect of RRS on PHQ-9 (b)	Index of Moderated Mediation (95% CI)	R <sup>2</sup>
Father Care	0.199***	0.000	0.005***	-0.001* (-0.003, -0.001)	0.465***
Mother Care	0.204***	-0.008***	0.004***	-0.002* (-0.003, -0.001)	0.504***
Father OP	0.195***	0.001	0.005***	0.002* (0.001, 0.003)	0.468***
Mother OP	0.207***	0.001	0.005***	0.003* (0.001, 0.004)	0.483***

**Notes:** [Figures 1B](#) and [2A-D](#) show this model. (a) and (b) are shown in [Figure 1B](#). This analysis controlled for age and sex. Coefficients were not standardized. R<sup>2</sup> indicates the coefficients of determination. \* $p < 0.05$ ; \*\*\* $p < 0.001$ .

**Abbreviations:** CI, confidence interval; OP, overprotection; PBI, Parental Bonding Instrument; PHQ-9, Patient Health Questionnaire-9; RRS, Ruminative Responses Scale.



**Figure 2** Effects of STAI-Y trait anxiety as a moderator on indirect effects of inappropriate nurturing experiences in childhood (PBI: **(A)** paternal care; **(B)**, maternal care; **(C)** paternal overprotection; **(D)** maternal overprotection;) on depressive symptoms (PHQ-9) via depressive rumination (RRS) as a mediator in the moderated-mediation model (model 15).

**Abbreviations:** CI, confidence interval; PBI, Parental Bonding Instrument; PHQ-9, Patient Health Questionnaire-9; RRS, Ruminative Responses Scale; STAI-Y, State-Trait Anxiety Inventory form Y.

were not correlated with PHQ-9 scores. STAI-Y score (trait anxiety), maternal and paternal PBI overprotection scores, and RRS total score were positively correlated with PHQ-9 scores, and maternal and paternal PBI care scores were negatively correlated with PHQ-9 scores.

## Direct Effects from Model 4 Mediation Analysis (Table 2)

Father's care and mother's care received in childhood showed a significantly negative effect on depressive rumination, whereas father's overprotection and mother's overprotection showed a significantly positive direct effect on depressive rumination. Father's care and mother's care in childhood showed a significantly negative direct effect on depressive symptoms in adulthood, whereas father's overprotection and mother's overprotection showed a significantly positive direct effect on depressive symptoms. Depressive rumination showed a significantly positive direct effect on depressive symptoms.

## Indirect Effects from Model 4 Mediation Analysis (Table 2)

Father's care and mother's care in childhood had significantly negative indirect effects on depressive symptoms via their effects on depressive rumination. Father's overprotection and mother's overprotection during childhood had significantly positive indirect effects on depressive symptoms via their effects on depressive rumination. That is, care and overprotection by parents during childhood were associated with a reduction and exacerbation of depressive symptoms,

respectively, in adulthood via their effects on depressive rumination. The overall effect of inappropriate parental nurturing on depressive symptoms was partially mediated by depressive rumination, with a mediating effect of 46% to 59%.

### Moderated-Mediation Analysis Using Model 15 (Table 3)

Trait anxiety moderated (increased) the effect of depressive rumination on depressive symptoms. Trait anxiety moderated (attenuated) only the effect of the mother's care on depressive symptoms among the inappropriate nurturing experiences, but did not significantly moderate the effects of the mother's overprotection and the father's care and overprotection on depressive symptoms. The moderated-mediation effect indices for the effects of parental overprotection and care on depressive symptoms through depressive rumination (RRS) in this model were all significant (Table 3, Figure 2A–D). That is, when trait anxiety was high, the mediation effect of depressive rumination on parental care and depressive symptoms was decreased, and the mediation effect of depressive rumination on parental overprotection and depressive symptoms was enhanced (Figure 2A–D).

## Discussion

The present study was conducted using moderated-mediation analysis to test the hypothesis that the effects of inappropriate childhood nurturing experiences on depressive symptoms were mediated by depressive rumination, and that the mediating effects of depressive rumination were moderated by trait anxiety. The results showed that depressive rumination partially mediates the effect of inappropriate childhood nurturing experiences on depressive symptoms in adulthood, and that high trait anxiety interacts with depressive rumination and maternal care to worsen depressive symptoms. Furthermore, trait anxiety was found to moderate the mediation effects of depressive rumination. To the best of our knowledge, this is the first report to date of such findings. In summary, regarding the association of the 3 factors of inappropriate nurturing experience, depressive rumination, and depression, depressive rumination demonstrated a mediating effect, whereas trait anxiety demonstrated a moderating effect.

As described in the Introduction section, depressive rumination is involved in the development of depression.<sup>25–30</sup> Consistent with these previous studies, depressive rumination moderately correlated with depressive symptoms in the present study. What is new in this study is the finding that the effects of the 4 different inappropriate nurturing factors, care and overprotection by the father or mother, on depressive symptoms were statistically significant, and depressive rumination mediated all 4 types of associations. A previous long-term prospective study reported that the effects of positive nurturing by mothers in childhood on depression in adolescent girls were mediated by depressive rumination.<sup>33</sup> However, this previous study by Gaté et al did not analyze the mediating effect of depressive rumination on paternal nurturing, and furthermore, the mediating effect of depressive rumination on maternal aggressive nurturing was not statistically significant.<sup>33</sup> Thus, in the present study we confirmed the mediating effect of depressive rumination found by Gaté et al, by extending it to paternal nurturing and negative nurturing.<sup>33</sup> Similar to childhood inappropriate nurturing experiences, the mediating effect of depressive rumination on the effects of childhood abuse and peer victimization experiences on depression has been reported previously, suggesting that it can be generalized that various childhood stress experiences, not limited to inappropriate nurturing experiences, indirectly enhance depressive symptoms in adults by exacerbating depressive rumination.<sup>9,32</sup>

In the present study, the effect of the interaction between depressive rumination and trait anxiety on depressive symptoms was significantly positive. This means that depressive rumination worsens depressive symptoms in individuals with high trait anxiety. Thus, when individuals have high levels of both trait anxiety and depressive rumination, interventions that decrease both are expected to synergistically improve depressive symptoms. Both trait anxiety and depressive rumination are risk factors for the onset or exacerbation of depressive symptoms.<sup>9,15,25–30,32,38,39</sup> A chain reaction mediating effect was reported in which trait anxiety enhances depressive rumination, which further enhances depressive symptoms.<sup>9,32</sup> In addition, as we showed in the present study, trait anxiety and depressive rumination interact to enhance depressive symptoms, suggesting that they have a mutually complex association for enhancing depression. In other words, because trait anxiety and depressive rumination form a vicious cycle or a negative spiral against depressive symptoms, interventions to decrease both are expected to be effective in clinical practice.

It is unclear why only the mother's care had a moderating effect together with trait anxiety on depressive symptoms, whereas all 4 nurturing factors, paternal or maternal care and overprotection, had significant effects on depressive

symptoms in adulthood. In our recent study, among parental care and overprotection, maternal care in particular significantly affected hopeful thinking in adults, and reduced depressive symptoms through its effect on hopeful thinking.<sup>50</sup> Positive parental nurturing is considered to be a resilience factor for children,<sup>51</sup> and hopeful thinking is also considered to be a resilience-enhancing factor.<sup>52,53</sup> Similarly, some reports indicated the importance of maternal factors, particularly in influencing children's personality development, resilience, and interpersonal sensitivity during adolescence.<sup>17,54,55</sup> Thus, among the various parental nurturing factors, maternal care strongly influences some psychological mechanisms, which may explain why trait anxiety only interacted with maternal care in the present study.

Previous neuroimaging studies have shown that depressive rumination is associated with activation and connectivity within the amygdala, medial prefrontal cortex, and posterior cingulate cortex.<sup>29</sup> These brain regions include the default mode network, which is typically activated during resting states, when an individual is focused internally rather than on external tasks or environmental stimuli.<sup>29</sup> Individuals who have trait rumination show increased activity within the default mode network. On the other hand, trait anxiety is also closely associated with these brain regions.<sup>38,56</sup> The above neural mechanisms may account for the relevance of depressive rumination and trait anxiety to depression, because depression was associated with these regions, including the default mode network.<sup>57–59</sup>

The results of the present study indicate that depressive symptoms are exacerbated in the context of childhood experiences of inappropriate nurturing, which enhance depressive symptoms by intensifying depressive rumination, and that depressive rumination and trait anxiety mutually reinforce each other and enhance depressive symptoms. These results provide clinically useful suggestions. Preventing low care and overprotection in childhood is expected to prevent depressive symptoms, but it is not easy for parents or third parties to intervene preventively in inappropriate parental nurturing. However, even if a person has experienced low care and overprotection, if depressive rumination as a mediating factor and trait anxiety as a moderating factor are recognized through careful clinical assessment as in this study, interventions for these factors can reduce depressive symptoms and prevent the onset or exacerbation of depression. Interventions for depressive rumination are considered clinically important in terms of reducing depressive symptoms, and cognitive-behavioral therapies focused on depressive rumination have recently been developed.<sup>29</sup> Mindfulness-based cognitive therapy was shown to be effective in lowering depressive rumination.<sup>30</sup> In addition, selective serotonin reuptake inhibitors, the transcendental meditation technique, and exercise therapy have been reported to attenuate trait anxiety.<sup>60–62</sup> Future development of evaluations that encompass a variety of factors, similar to the Habit development, Executive control, Abstract processing, Goal discrepancies, Negative bias (H-EX-A-GO-N) model is needed.<sup>29</sup> The results obtained in the present study indicate that elucidating the mechanisms of the association among experiences of inappropriate nurturing in childhood, trait anxiety, and depressive rumination may contribute to the development of interventions for depression. Furthermore, future studies are necessary to investigate the efficacy of interventions for these factors in clinical settings.

## Limitations

Because this study relied on the memories of the subjects and used a self-report questionnaire, memory bias may have confounded the results. However, the validity of the retrospective evaluation by the PBI used in the present study has been confirmed in previous studies.<sup>41</sup> Adverse childhood experiences (ACEs) have been reported to affect mental and physical health in adulthood.<sup>63</sup> Numerous retrospective studies on ACEs have been conducted around the world, and a prospective study using the Dunedin cohort confirmed the validity of the retrospective assessment of ACEs.<sup>64</sup> Therefore, we did not consider memory bias to be a serious problem in the present study. Additionally, as the study was conducted on general adult volunteers, the present findings may not apply to patients with depression. There are cultural differences in parenting styles,<sup>65</sup> and these differences might influence the effects of parenting on depressive symptoms. As this study was conducted in Japan, cultural differences may influence the generalizability of our findings to other populations. Therefore, replicating the study in different cultural and geographic contexts would enhance the generalizability of the findings. As the design of the present study is cross-sectional, a prospective study, which can be retested for example in 1 year, is necessary for concluding a causal association. In the present study, we hypothesized that trait anxiety, which is a type of personality trait that causes anxiety, influences rumination. However, a prospective verification study is needed in the future, as higher rumination may result in higher trait anxiety. In addition to the inappropriate nurturing experiences analyzed in the present study, other environmental factors, including abuse and



bullying victimization in childhood and harassment experienced in adulthood may influence trait anxiety and depressive rumination. Therefore, a more comprehensive analysis including these multiple factors is needed in the future.

## Conclusions

The present study is the first report to our knowledge that depressive rumination mediates the effects of inappropriate nurturing experiences in childhood on depressive symptoms, and that these mediating effects are moderated by trait anxiety. The interrelationships among inappropriate childhood nurturing experiences, trait anxiety, depressive rumination, and depressive symptoms were demonstrated in a moderated-mediation model. In the future, based on the results of this study, a large-scale prospective study of community residents is needed to confirm a causal association.

## Acknowledgments

We thank Dr. Nobutada Takahashi of Fuji Psychosomatic Rehabilitation Institute Hospital, Dr. Hiroshi Matsuda of Kashiwazaki Kosei Hospital, Dr. Yasuhiko Takita (deceased) of Maruyamasou Hospital, and Dr. Yoshihide Takaesu of Izumi Hospital for their collection of subject data. We thank Dr. Helena Popiel of the Center for International Education and Research, Tokyo Medical University, for editorial review of the manuscript.

## Author Contributions

All authors made a significant contribution to the work reported, whether in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

This work was partly supported by a Grant-in-Aid for Scientific Research (no. 21K07510, to TI) from the Japanese Ministry of Education, Culture, Sports, Science and Technology and by AMED under Grant Number JP23rea522113.

## Disclosure

Chihiro Morishita has received personal fees from Otsuka Pharmaceutical, Sumitomo Pharma, and Meiji Seika Pharma. Yu Tamada has received personal fees from Otsuka Pharmaceutical, Sumitomo Pharma, Eisai, MSD, and Meiji Seika Pharma. Takeshi Inoue has received personal fees from Mochida Pharmaceutical, Takeda Pharmaceutical, Janssen Pharmaceutical, Novartis Pharma, MSD, Yoshitomiyakuhin, Nipro, Kyowa Pharmaceutical Industry, Viatrix, Lundbeck, Boehringer Ingelheim, Ono Pharmaceutical, and Meiji Seika Pharma; grants from Daiichi Sankyo, and Tsumura; and grants and personal fees from Shionogi, Otsuka Pharmaceutical, Sumitomo Pharma, Mitsubishi Tanabe Pharma, and Eisai; and is a member of the advisory boards of Luye, Shionogi, GlaxoSmithKline, Viatrix, and Otsuka Pharmaceutical. Jiro Masuya has received personal fees from Otsuka Pharmaceutical, Eli Lilly, Astellas, and Meiji Yasuda Mental Health Foundation, as well as grants from Pfizer. The other authors do not have any actual or potential conflicts of interest to declare for this work.

## References

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-5*. Washington, D.C: American Psychiatric Publication Inc.; 2013.
2. Alloy LB, Abramson LY, Smith JM, Gibb BE, Neeren AM. Role of parenting and maltreatment histories in unipolar and bipolar mood disorders: mediation by cognitive vulnerability to depression. *Clin Child Fam Psych*. 2006;9(1):23–64. doi:10.1007/s10567-006-0002-4
3. Kendler KS, Kuhn J, Prescott CA. The interrelationship of neuroticism, sex, and stressful life events in the prediction of episodes of major depression. *Am J Psychiatry*. 2004;161(4):631–636. doi:10.1176/appi.ajp.161.4.631
4. Kessler RC, Magee WJ. Childhood adversities and adult depression: basic patterns of association in a US national survey. *Psychol Med*. 1993;23(3):679–690. doi:10.1017/S0033291700025460
5. Lereya ST, Copeland WE, Costello EJ, Wolke D. Adult mental health consequences of peer bullying and maltreatment in childhood: two cohorts in two countries. *Lancet Psychiatry*. 2015;2(6):524–531. doi:10.1016/S2215-0366(15)00165-0
6. Parker G, Hadzi-Pavlovic D, Greenwald S, Weissman M. Low parental care as a risk factor to lifetime depression in a community sample. *J Affect Disord*. 1995;33(3):173–180. doi:10.1016/0165-0327(94)00086-O

7. Tachi S, Asamizu M, Uchida Y, et al. Victimization in childhood affects depression in adulthood via neuroticism: a path analysis study. *Neuropsychiatr Dis Treat*. 2019;15:2835–2841. doi:10.2147/NDT.S220349
8. Lange S, Burr H, Rose U, Conway PM. Workplace bullying and depressive symptoms among employees in Germany: prospective associations regarding severity and the role of the perpetrator. *Int Arch Occup Environ Health*. 2020;93(4):433–443. doi:10.1007/s00420-019-01492-7
9. Deguchi A, Masuya J, Naruse M, et al. Rumination mediates the effects of childhood maltreatment and trait anxiety on depression in non-clinical adult volunteers. *Neuropsychiatr Dis Treat*. 2021;17:3439–3445. doi:10.2147/NDT.S332603
10. Higashiyama M, Hayashida T, Sakuta K, et al. Complex effects of childhood abuse, affective temperament, and subjective social status on depressive symptoms of adult volunteers from the community. *Neuropsychiatr Dis Treat*. 2019;15:2477–2485. doi:10.2147/NDT.S209100
11. Nakai Y, Inoue T, Toda H, et al. The influence of childhood abuse, adult stressful life events and temperaments on depressive symptoms in the nonclinical general adult population. *J Affect Disord*. 2014;158:101–107. doi:10.1016/j.jad.2014.02.004
12. Ono K, Takaesu Y, Nakai Y, et al. Associations among depressive symptoms, childhood abuse, neuroticism, and adult stressful life events in the general adult population. *Neuropsychiatr Dis Treat*. 2017;13:477–482. doi:10.2147/NDT.S128557
13. Ono Y, Takaesu Y, Nakai Y, et al. The influence of parental care and overprotection, neuroticism and adult stressful life events on depressive symptoms in the general adult population. *J Affect Disord*. 2017;217:66–72.
14. Toda H, Inoue T, Tsunoda T, et al. The structural equation analysis of childhood abuse, adult stressful life events, and temperaments in major depressive disorders and their influence on refractoriness. *Neuropsychiatr Dis Treat*. 2015;11:2079–2090. doi:10.2147/NDT.S82236
15. Uchida Y, Takahashi T, Katayama S, et al. Influence of trait anxiety, child maltreatment, and adulthood life events on depressive symptoms. *Neuropsychiatr Dis Treat*. 2018;14:3279–3287. doi:10.2147/NDT.S182783
16. Hayashi Y, Okamoto Y, Takagaki K, et al. Direct and indirect influences of childhood abuse on depression symptoms in patients with major depressive disorder. *BMC Psychiatry*. 2015;15(1):244. doi:10.1186/s12888-015-0636-1
17. Sakado K, Kuwabara H, Sato T, Uehara T, Sakado M, Someya T. The relationship between personality, dysfunctional parenting in childhood, and lifetime depression in a sample of employed Japanese adults. *J Affective Disorders*. 2000;60(1):47–51. doi:10.1016/S0165-0327(99)00150-0
18. Ito S, Morishita C, Masuya J, et al. Moderating and mediating effects of resilience together with neuroticism on depressive symptoms in adult volunteers. *Neuropsychiatr Dis Treat*. 2022;18:1751–1761. doi:10.2147/NDT.S370201
19. Boyce P, Hickie I, Gordon P. Parents partners or personality? Risk factors for post-natal depression. *J Affective Disorders*. 1991;21(4):245–255. doi:10.1016/0165-0327(91)90004-C
20. Hayashida T, Higashiyama M, Sakuta K, et al. Subjective social status via mediation of childhood parenting is associated with adulthood depression in non-clinical adult volunteers. *Psychiatry Res*. 2019;274:352–357. doi:10.1016/j.psychres.2019.02.061
21. Otsuka A, Takaesu Y, Sato M, et al. Interpersonal sensitivity mediates the effects of child abuse and affective temperaments on depressive symptoms in the general adult population. *Neuropsychiatr Dis Treat*. 2017;13:2559–2568. doi:10.2147/NDT.S144788
22. Toyoshima K, Inoue T, Masuya J, Fujimura Y, Higashi S, Kusumi I. Associations among childhood parenting, affective temperaments, depressive symptoms, and cognitive complaints in adult community volunteers. *J Affect Disord*. 2020;276:361–368. doi:10.1016/j.jad.2020.07.107
23. Parker G, Tupling H, Brown LB. A parental bonding instrument. *Br J Med Psychol*. 1979;52(1):1–10. doi:10.1111/j.2044-8341.1979.tb02487.x
24. Parker G. Parental ‘affectionless control’ as an antecedent to adult depression. A risk factor delineated. *Arch Gen Psychiatry*. 1983;40(9):956–960. doi:10.1001/archpsyc.1983.01790080038005
25. Nolen-Hoeksema S. The response styles theory. In: Papageorgiou C, Wells A, editors. *Depressive Rumination: Nature, Theory, and Treatment*. West Sussex: John Wiley & Sons; 2004:107–123.
26. Nolen-Hoeksema S. The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *J Abnorm Psychol*. 2000;109(3):504–511. doi:10.1037/0021-843X.109.3.504
27. Nolen-Hoeksema S, Morrow J, Fredrickson BL. Response styles and the duration of episodes of depressed mood. *J Abnormal Psychol*. 1993;102(1):20–28. doi:10.1037/0021-843X.102.1.20
28. Visted E, Vollestad J, Nielsen MB, Schanche E. Emotion regulation in current and remitted depression: a systematic review and meta-analysis. *Front Psychol*. 2018;9:756. doi:10.3389/fpsyg.2018.00756
29. Watkins ER, Roberts H. Reflecting on rumination: consequences, causes, mechanisms and treatment of rumination. *Behav Res Ther*. 2020;127:103573. doi:10.1016/j.brat.2020.103573
30. Perestelo-Perez L, Barraca J, Peñate W, Rivero-Santana A, Alvarez-Perez Y. Mindfulness-based interventions for the treatment of depressive rumination: systematic review and meta-analysis. *Int J Clin Health Psychol*. 2017;17(3):282–295. doi:10.1016/j.ijchp.2017.07.004
31. Raes F, Hermans D. On the mediating role of subtypes of rumination in the relationship between childhood emotional abuse and depressed mood: brooding versus reflection. *Depress Anxiety*. 2008;25(12):1067–1070. doi:10.1002/da.20447
32. Masuya J, Morishita C, Higashiyama M, et al. Depressive rumination and trait anxiety mediate the effects of childhood victimization on adulthood depressive symptoms in adult volunteers. *PLoS One*. 2023;18(5):e0286126. doi:10.1371/journal.pone.0286126
33. Gaté MA, Watkins ER, Simmons JG, et al. Maternal Parenting Behaviors and Adolescent Depression: the Mediating Role of Rumination. *J Clin Child Adolesc Psychol*. 2013;42(3):348–357. doi:10.1080/15374416.2012.755927
34. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. 1986;51(6):1173–1182. doi:10.1037/0022-3514.51.6.1173
35. Clark LA, Watson D. Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications. *J Abnorm Psychol*. 1991;100(3):316–336. doi:10.1037/0021-843X.100.3.316
36. Lamers F, van Oppen P, Comijs HC, et al. Comorbidity patterns of anxiety and depressive disorders in a large cohort study: the Netherlands Study of Depression and Anxiety (NESDA). *J Clin Psychiatry*. 2011;72(03):341–348. doi:10.4088/JCP.10m06176blu
37. Wittchen HU, Kessler RC, Pfister H, Lieb M, Lieb R. Why do people with anxiety disorders become depressed? A prospective-longitudinal community study. *Acta Psychiatr Scand Suppl*. 2000;102(s406):14–23. doi:10.1111/j.0065-1591.2000.acp29-03.x
38. Weger M, Sandi C. High anxiety trait: a vulnerable phenotype for stress-induced depression. *Neurosci Biobehav Rev*. 2018;87:27–37. doi:10.1016/j.neubiorev.2018.01.012
39. Sandi C, Richter-Levin G. From high anxiety trait to depression: a neurocognitive hypothesis. *Trends Neurosci*. 2009;32(6):312–320. doi:10.1016/j.tins.2009.02.004

40. Nakajima A, Uchida Y, Katayama S, Inoue T. Influence of parenting quality, trait anxiety, and adulthood life events on subjective well-being in adult volunteers in Japan. *J Tokyo Med Univ.* 2022;80:29–38.
41. Kitamura T, Suzuki T. A validation study of the Parental Bonding Instrument in a Japanese population. *Jpn J Psychiatry Neurol.* 1993;47:29–36. doi:10.1111/j.1440-1819.1993.tb02026.x
42. Treynor W, Gonzalez R, Nolen-Hoeksema S. Rumination reconsidered: a psychometric analysis. *Cognit Ther Res.* 2003;27(3):247–259. doi:10.1023/A:1023910315561
43. Hasegawa A. Translation and initial validation of the Japanese version of the ruminative responses scale. *Psychol Rep.* 2013;112(3):716–726. doi:10.2466/02.08.PR0.112.3.716-726
44. Spielberger CD. *Manual for the State-Trait Anxiety Inventory STAI (Form Y)*. Palo Alto: Consulting Psychologists Press; 1983.
45. Hidano N, Fukuhara M, Iwakaki M, Soga S, Spielberger CD. *State-Trait Anxiety Inventory-Form JYZ*. Tokyo: Jitsumu Kyoiku Shuppan; 2000.
46. Spitzer RL, Kroenke K, Williams JB, the Patient Health Questionnaire Primary Care Study Group. Validation and Utility of a Self-report Version of PRIME-MD The PHQ Primary Care Study. *JAMA.* 1999;282(18):1737–1744. doi:10.1001/jama.282.18.1737
47. Muramatsu K, Miyaoka H, Kamijima K, et al. The patient health questionnaire, Japanese version: validity according to the mini-international neuropsychiatric interview-plus. *Psychol Rep.* 2007;101(3 Pt 1):952–960. doi:10.2466/pr0.101.3.952-960
48. Peduzzi P, Concato J, Feinstein AR, Holford TR. Importance of events per independent variable in proportional hazards regression analysis. II. Accuracy and precision of regression estimates. *J Clin Epidemiol.* 1995;48(12):1503–1510. doi:10.1016/0895-4356(95)00048-8
49. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol.* 1996;49(12):1373–1379. doi:10.1016/S0895-4356(96)00236-3
50. Mutoh K, Ono M, Masuya J, et al. Hope mediates the effects of maternal care in childhood, neuroticism, educational level, and subjective social status on depression in adult volunteers. *J Tokyo Med Univ.* 2023;81(1):39–48.
51. Fritz J, de Graaff AM, Caisley H, van Harmelen A-L, Wilkinson PO. A systematic review of amenable resilience factors that moderate and/or mediate the relationship between childhood adversity and mental health in young people. *Frontiers in Psychiatry.* 2018;9:230. doi:10.3389/fpsy.2018.00230
52. Ritschel LA, Sheppard CS. Hope and depression. In: Gallagher MW, Lopez SJ, editors. *Handbook of Hope New York*. Oxford University Press. 2018:209–219.
53. Ong AD, Standiford T, Deshpande S. Hope and Stress Resilience. In: Gallagher MW, Lopez SJ, editors. *The Oxford Handbook of Hope New York*. Oxford University Press. 2018:255–284.
54. Fujioka H, Murayama M. The relationship between university student's resilience and parenting attitude of parents. *Kobe Gakuin Univ J Psychol.* 2019;2(1):31–36.
55. Kobayashi M. The influence of parents' nurturing attitudes in middle school on adolescent friendship: self-concept as a mediating variable. *Toyama hattatu hukushi gakunenpou.* 2011;2:21–28.
56. Bishop S, Forster S. Trait anxiety, neuroticism, and the brain basis of vulnerability to affective disorder. In: Armony J, Vuilleumier P, editors. *The Cambridge Handbook of Human Affective Neuroscience*. Cambridge: Cambridge University Press; 2013:553–574.
57. Zhou HX, Chen X, Shen YQ, et al. Rumination and the default mode network: meta-analysis of brain imaging studies and implications for depression. *Neuroimage.* 2020;206:116287. doi:10.1016/j.neuroimage.2019.116287
58. Imperatori C, Farina B, Adenzato M, et al. Default mode network alterations in individuals with high-trait-anxiety: an EEG functional connectivity study. *J Affect Disord.* 2019;246:611–618. doi:10.1016/j.jad.2018.12.071
59. Li C, Hu Q, Zhang D, Hoffstaedter F, Bauer A, Elmenhorst D. Neural correlates of affective control regions induced by common therapeutic strategies in major depressive disorders: an activation likelihood estimation meta-analysis study. *Neurosci Biobehav Rev.* 2022;137:104643. doi:10.1016/j.neubiorev.2022.104643
60. 2018 Physical Activity Guidelines Advisory Committee. 2018 physical activity guidelines advisory committee scientific report. 2018.
61. Nagata T, van Vliet I, Yamada H, Kataoka K, Iketani T, Kiriike N. An open trial of paroxetine for the “offensive subtype” of taijin kyofusho and social anxiety disorder. *Depress Anxiety.* 2006;23(3):168–174. doi:10.1002/da.20153
62. Orme-Johnson DW, Barnes VA. Effects of the transcendental meditation technique on trait anxiety: a meta-analysis of randomized controlled trials. *J Altern Complement Med.* 2013;20(5):330–341. doi:10.1089/acm.2013.0204
63. Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med.* 1998;14(4):245–258. doi:10.1016/S0749-3797(98)00017-8
64. Reuben A, Moffitt TE, Caspi A, et al. Lest we forget: comparing retrospective and prospective assessments of adverse childhood experiences in the prediction of adult health. *J Child Psychol Psychiatr.* 2016;57(10):1103–1112. doi:10.1111/jcpp.12621
65. Neoh MJY, Carollo A, Bonassi A, Mulatti C, Lee A, Esposito G. A cross-cultural study of the effect of parental bonding on the perception and response to criticism in Singapore, Italy and USA. *PLoS One.* 2021;16:e0257888.

## Neuropsychiatric Disease and Treatment

Dovepress

### Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>