


Continuation of common bile duct clearance with gallbladder stenting after duodenal perforation with subsequent treatment for tension pneumoperitoneum and pneumothorax


OPEN
ACCESS




▶ **Video 1** Endoscopic retrograde cholangiopancreatography with gallbladder stenting completed after treatment of iatrogenic duodenal perforation, and subsequent management of tension pneumoperitoneum.



▶ **Fig. 1** Guidewire insertion into the duodenal lumen to prevent accidental luminal closure during clipping of perforation.



▶ **Fig. 2** First over-the-scope clip deployment by mucosal suction over the lacerating tissue at the caudal area of the defect.




▶ **Fig. 3** Contrast enterography revealed no contrast leakage into the peritoneum.


An 88-year-old woman underwent endoscopic retrograde cholangiopancreatography (ERCP) because of septic cholangitis with acute calculous cholecystitis. Unfortunately, incidental duodenal perforation (Stapfer classification type 1) [1] occurred during duodenoscope intubation by our trainee endoscopist. Gastroscopy with a transparent cap revealed a linear 4-cm defect with active oozing at the duodenal apex (▶ **Video 1**). Perforation closure was attempted with preceding guidewire insertion into the downstream duodenal lumen (▶ **Fig. 1**) to prevent accidental luminal closure [2]. The first traumatic type over-the-scope clip (OTSC, 12/6t; Ovesco) was deployed by suction on the lacerating tissue at the caudal side of the defect (▶ **Fig. 2**). However, the defect did not close completely; therefore, a second OTS clip was deployed using twin graspers to appose the edges of the defect. Contrast enterography revealed no intraperitoneal leakage (▶ **Fig. 3**). Immediately after closure, ERCP with stones removal and transpapil-

lary gallbladder stenting with a double-pigtail plastic stent to prevent recurrent cholecystitis was successfully performed. At almost 40 minutes before completion of the procedure, the patient developed marked abdominal distension, hypotension, and desaturation. Tension pneumoperitoneum with right pneumothorax was confirmed by fluoroscopy (▶ **Fig. 4**). Emergency needle decompression was performed using an 18 G needle to release the tension pneumoperitoneum, and the patient was then intubated. Her abdomen gradually softened with an improvement in oxygen saturation. At 4 hours later, plain radiography showed no free air (▶ **Fig. 5**). The patient was extubated and resumed oral intake the following day. Duodenal wall perforation can occur during ERCP, especially when performed by a less experienced endoscopist. ERCP can be completed safely if the patient is stable after early detection of the perforation with immediate endoscopic closure. The OTS clip is preferred for a defect

larger than 1 cm, and more than one clip may be required if the defect is larger than 3 cm [3]. Tension pneumoperitoneum after endoscopy-related perforation is life-threatening, and early detec-



► **Fig. 4** Fluoroscopy showed right pneumothorax.



► **Fig. 5** Plain radiography 4 hours later showed no free air.

tion with emergency needle decompression is the key to saving the patient's life [4].

Endoscopy_UCTN_Code_CPL_1AK_2AC

Competing Interest

The authors declare that they have no conflict of interest.

The authors

Natee Faknak^{1,2,3}, Santi Kulpatcharapong^{1,2}, Salin Samutrangsi^{1,2}, Parit Mekaroonkamol^{1,2}, Wiriyaoporn Ridtitid^{1,2}, Rungsun Rerknimitr^{1,2}

- 1 Division of Gastroenterology, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand
- 2 Excellence Center for Gastrointestinal Endoscopy, King Chulalongkorn Memorial Hospital, Bangkok, Thailand
- 3 Division of Gastroenterology, Department of Medicine, Sawanpracharak Hospital, Nakhonsawan, Thailand

Corresponding author

Rungsun Rerknimitr, MD

Division of Gastroenterology, Department of Internal Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok 10310, Thailand

Fax: +66-2-2527839

ERCP@live.com

References

- [1] Stapfer M, Selby RR, Stain SC et al. Management of duodenal perforation after endoscopic retrograde cholangiopancreatography and sphincterotomy. *Ann Surg* 2000; 232: 191–198
- [2] Martínez-Alcalá García Á, Martínez-Alcalá F, Mönkemüller K. Use of a biliary guidewire to assist placement of two over-the-scope clips for deep ulcer bleeding while preventing duodenal closure. *Endoscopy* 2022; 54: E1–E9
- [3] Piyachaturawat P, Mekaroonkamol P, Rerknimitr R. Use of the over the scope clip to close perforations and fistulas. *Gastrointest Endosc Clin N Am* 2020; 30: 25–39

- [4] Chiapponi C, Stocker U, Körner M et al. Emergency percutaneous needle decompression for tension pneumoperitoneum. *BMC Gastroenterol* 2011; 11: 48

Bibliography

Endoscopy 2023; 55: E125–E126

DOI 10.1055/a-1949-0494

ISSN 0013-726X

published online 17.10.2022

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>