


ORIGINAL RESEARCH

Decreased EDHF-mediated relaxation is a major mechanism in endothelial dysfunction in resistance arteries in aged mice on prolonged high-fat sucrose diet

Shannon M. Dunn^{1,3} & Kumuda C. Das² 

1 Department of Pharmacology & Neuroscience, Texas Tech University Health Sciences Center, Lubbock, Texas

2 The Department of Translational & Vascular Biology, University of Texas Health Sciences Center at Tyler, Tyler, Texas

Keywords

Aging, endothelial dysfunction, endothelium-dependent hyperpolarizing factor, high-fat sucrose diet, obesity, peroxynitrite, reactive oxygen species, vascular relaxation.

Correspondence

Kumuda C. Das, The Department of Translational & Vascular Biology, The University of Texas Health Sciences Center at Tyler, Tyler, TX 75708. Tel: 903-877-8122; Fax: 903-877-7558; E-mail: kumuda.das2@uthct.edu

Current address

Shannon Dunn, 685 W. Baltimore Street, MSTF Room, 357-B Baltimore, Maryland

Funding Information

The work presented here is supported by National Heart, Lung and Blood Institutes grants, HL 1R01HL107885, 3R01HL107885-05S1 and 1R01HL109397 to KCD.

Received: 8 August 2017; Revised: 20 October 2017; Accepted: 23 October 2017

doi: 10.14814/phy2.13502

Physiol Rep, 5 (23), 2017, e13502.
<https://doi.org/10.14814/phy2.13502>

Abstract

High-fat sucrose (HFS) diet in aged individuals causes severe weight gain (obesity) with much higher risk of cardiovascular diseases such as hypertension or atherosclerosis. Endothelial dysfunction is a major contributor for these vascular disorders. We hypothesize that prolonged ingestion of HFS diet by aged mice would accentuate endothelial dysfunction in the small resistance arteries. Male C57BL/6J mice at 12 weeks of age were divided into four groups and fed either normal chow (NC) or high-fat sucrose diet (HFS). Young group received NC for 4 months, and high-fat diet (HFD) for 3 months and 1 month HFS + 10% Sucrose (HFS diet). Aged mice received NC for 12 months. Aged HFS group received HFD for 4 months + 1 month HFD + 10% sucrose + 8 months HFD. Total body weight, plasma blood glucose levels, and glucose tolerance were determined in all groups. Isolated mesenteric arteries were assessed for arterial remodeling, myogenic tone, and vasomotor responses using pressure and wire myography. Both young and aged HFS mice showed impaired glucose tolerance (Y-NC, 137 ± 8.5 vs. Y-NC HFS, 228 ± 11.71 ; A-NC, 148 ± 6.42 vs. A-HFS, 225 ± 10.99), as well as hypercholesterolemia (Y-NC 99.50 ± 6.35 vs. Y-HFS 220.40 ± 16.34 mg/dL; A-NC $108.6 \pm$ vs. A-HFS 279 ± 21.64) and significant weight gain (Y-NC 32.13 ± 0.8 g vs. Y-HFS 47.87 ± 2.18 g; A-NC 33.72 vs. A-HFS 56.28 ± 3.47 g) compared to both groups of mice on NC. The mesenteric artery from mice with prolonged HFS diet resulted in outward hypertrophic remodeling, increased stiffness, reduced myogenic tone, impaired vasodilation, increased contractility and blunted nitric oxide (NO) and EDH-mediated relaxations. Ebselen, a peroxynitrite scavenger rescued the endothelium derived relaxing factor (EDHF)-mediated relaxations. Our findings suggest that prolonged diet-induced obesity of aged mice can worsen small resistance artery endothelial dysfunction due to decrease in NO and EDHF-mediated relaxation, but, EDHF-mediated relaxation is a major contributor to overall endothelial dysfunction.

Introduction

Obesity is linked to the onset of several cardiovascular diseases and metabolic complications such as hypertension, atherosclerosis, and Type-2 diabetes (Li et al. 2017). In addition, obesity is also linked to the aging process, as

a major proportion of aged individuals are obese (Bailey-Downs et al. 2013). Consequently, the risk of cardiovascular and metabolic disorders is considerably higher in these individuals. Obesity-mediated dysfunction of the vascular endothelium has been implicated as an underlying mechanism for all of these cardiovascular and

metabolic disorders (Bailey-Downs et al. 2013). Endothelial cells line the lumen of vascular tissue and secrete several active compounds that is essential to promote vascular flow and maintain healthy arterial structure and function. However, normal function of endothelium is impaired in obese individuals (Sloboda et al. 2012). In addition, hyperglycemia induces endothelial dysfunction, which is a major vascular complication in diabetes (Sharma et al. 2017). This dysfunction of endothelium also occurs in aged individuals (Seals et al. 2011). Endothelium-dependent vasodilation has been shown to be impaired after 40 years of age and progressively decline with increasing age (Seals et al. 2011).

Endothelial dysfunction is primarily characterized by an impaired endothelium-dependent dilation due to a reduction in NO bioavailability (Hamilton et al. 2002; Crabtree et al. 2013). In obesity NO production decreases due to increased oxidative stress and inflammation (Bailey-Downs et al. 2013). In hyperglycemia EC apoptosis occurs, which has been shown to cause decreased NO production (Lan et al. 2015). Furthermore, during aging the eNOS becomes dysfunctional and produces superoxide anion ($O_2^{\bullet-}$) instead of NO causing decreased NO availability (Hilgers et al. 2017). Endothelial dysfunction in small resistance arteries is an early marker of developing vascular disorders such as atherosclerosis (Hilgers and Das 2015). Small vessels contribute substantially to the vascular resistance, and are key determinant of impaired microcirculatory function (Bender et al. 2015). Impaired vasodilation has been shown to occur in peripheral vascular disease, diabetes, hypercholesterolemia, hypertension, chronic renal failure and atherosclerosis, aortic aneurism and other vascular complications (Abularrage et al. 2005). Endothelial dysfunction can also occur in a similar fashion in multiple vascular beds throughout the body (Higashi et al. 2012). However, the structural and functional differences in vascular beds critically impact aged-related flow-mediated dilation resulting in on set of cardiovascular diseases (CVD). For example, resistance artery relaxation is largely mediated by endothelium derived hyperpolarizing factor (EDHF), however, NO is the principal relaxing factor in large conduit arteries such as aortae (Matoba et al. 2000).

Impaired flow-mediated dilation in young adults has been observed in obesity due to a “western-type” high-fat/high-sucrose diet (Keogh et al. 2005). Studies using isolated arteries from animals on a HFD have shown endothelial dysfunction in both large conduit (Woodman et al. 2005) and small resistance arteries (Erdei et al. 2006; Donato et al. 2012). However, structural and functional alterations of resistance arteries in response to prolonged HFS in aging remains understudied. Both, obesity and aging are accompanied by arterial stiffening of large

conduit arteries involving vascular remodeling processes including vascular cell proliferation, migration, hypertrophy, and changes in the composition of elastin/collagen in the extracellular matrix (Chatterjee et al. 2014). However, microvascular remodeling associated with obesity and insulin resistance is characterized by medial hypertrophy and increased extracellular matrix deposition, which may be vascular-bed specific (Briones et al. 2014). Previous studies have shown that a high-fat, low-nutrient diet increases the likelihood and advancement of obesity and diabetes (Willett et al. 1990; Giovannucci and Willett 1994; Petrofsky et al. 2013). Obesity and aging are risk factors for developing Type II diabetes (T2D), with obesity being one of the strongest risk markers (Cassano et al. 1992; Chan et al. 1994; Carey et al. 1997). There are virtually no studies addressing the effects of obesity in aged resistance artery endothelial dysfunction. Furthermore, the specific mechanism of endothelial dysfunction in these arteries has not been addressed.

Endothelium derived relaxing factor has been shown to be important in modulating vasomotor tone in microvessels (Shimokawa et al. 1996; Urakami-Harasawa et al. 1997). Although EDHF is very distinct from NO, studies suggest that both factors are functionally similar in maintaining vascular relaxations (Nagao et al. 1992). For example, both NO and EDHF are synthesized by the endothelial cells (Takaki et al. 2008). In conditions where NO-mediated relaxation is decreased (e.g., hypertension, hyperlipidemia) EDHF compensates for NO to cause endothelium-dependent relaxations (Takaki et al. 2008). For example, a dysfunctional nitric oxide synthase (eNOS) in aged animals produces $O_2^{\bullet-}$ instead of NO due either to oxidation of Tetrahydrobiopterin (BH_4) or depletion of L-arginine (Chen et al. 2013). The $O_2^{\bullet-}$ produced by eNOS is readily converted to H_2O_2 , and H_2O_2 function as an EDHF in this setting (Matoba et al. 2000). Although endothelial dysfunction has been reported in aged animals and also due to HFD fed aged animals, the exact mechanism of endothelial dysfunction in these animals remains understudied. For example, although decreased NO production has been attributed to endothelial dysfunction due to HFD in aged mice, the role of EDHF in HFD fed mice and HFD fed aged mice remains unclear. In addition, the role of HFS diet or increased blood glucose in EDHF-mediated endothelial dysfunction remains poorly understood.

In this study we hypothesized that HFS diet in aged mice would promote endothelial dysfunction in the resistance arteries due to impairment of EDHF-dependent vascular relaxation. Our study design incorporates both, an aging model and a diet-induced obesity model. We utilized a mice model with near lifelong HFS diet consumption to evaluate age-related vascular dysfunction.

We studied the effects of HFD on bodyweight, glucose metabolism, cholesterol levels, epididymal white adipose tissue (EWAT) deposition, microvascular reactivity and structure in young and aged mice. We also evaluated whether HFS diet in aged mice resistance arteries would show endothelial dysfunction due to impairment of EDHF. We show here that HFS diet caused severe obesity, and impaired glucose tolerance in mice that resembles a T2D phenotype. Furthermore, HFS exacerbated age-related endothelial dysfunction in the small resistance arteries that is dependent on EDHF-mediated relaxation since loss of EDHF in the resistance arteries caused endothelial dysfunction due to HFS in aged mice.

Materials and Methods

Animals and treatments

Male C57BL/6J mice at 3 months were fed a normal chow (NC) diet (12% kcal from fat, Harlan Laboratories, Inc.; Cat# 2020X) or HFD (42% kcal from fat, Harlan Laboratories, Inc., Envigo, Cat #88137). Sucrose (10% diluted in tap water, Sigma Aldrich) was given in addition to the HFD for a period of 1 month in the HFS group. All mice enter into the study at 3 months age. Mice were divided into four groups; group 1, young mice with normal chow (NC) diet for 4 months; group 2, young mice with high-fat sucrose (HFS) diet (3 months with HFD plus 1 month

in 10% sucrose), group 3, aged (12 months) mice with NC, and group 4, aged mice with HFS diet (12 months). Mice in the young group were fed NC for 4 months or HFD for 3 months followed by 1 month of HFD plus 10% sucrose. Mice in the NC aging group were placed on a NC chow until an age of 15 months. Mice in the aged HFS group were placed on HFD for 3 months and HFD+10% sucrose for 1 month. These mice were further placed on HFD for an additional 8 months (Fig. 1A). Bodyweight was monitored weekly in the two young mice groups. All procedures were approved by the IACUC at the Texas Tech University Health Sciences Center and were consistent with the *Guide for the Care and Use of Laboratory Animals* published by the National Institute of Health. All animals were maintained on a standard 12-h light/12-h dark cycle, in a temperature-controlled barrier facility.

Fasting glucose and insulin measurement

Glucose and serum insulin levels were measured after 4–6 h of fasting from mice receiving 4 months of normal chow (NC young) or 12 months of NC (NC aged); 4 months of HFS (HFS young) or 12 months of HFS (HFS aged). The tail was snipped and fasting blood glucose levels were measured using the OneTouch Ultra glucose monitor (LifeScan Inc.). Blood was drawn from the orbital sinus, spun down at 3000g for 15 min and serum




Figure 1. Effect of HFS on body weight of young and aged mice: Study timeline of feeding (A), weekly time log of body weight increases in young mice (B), final body weight of young and aged mice on NC or HFS (C), weight of epididymal fat pads for young and aged mice on NC or HFS (D), derived from mice receiving 6 months of normal chow (NC young; white bars), 16 months of NC (NC aged; dark gray bars), 3 months of a high-fat diet followed by 4 weeks of 10% sucrose in drinking water (HFS young; black bars), and 12 months of HFS (HFS aged; black bar). Values are shown as means \pm SEM ($n = 6$); * $P < 0.05$ versus young NC; ** $P < 0.05$ versus aged NC.

was separated and insulin was measured using Rat/Mouse Insulin ELISA as per instructions (EMD Millipore, MA). The means were compared using one-way ANOVA. Values are shown as standard error means \pm (SEM)

Intraperitoneal glucose and insulin tolerance tests (GTT/ITT)

Mice were anesthetized by brief inhalation of 2% isoflurane in pure oxygen, followed by an intraperitoneal injection of glucose (2.0 g/kg) or insulin (Humulin R; 0.75 U/kg), and blood was drawn from the tail at 0 (time point taken before injection), 15, 30, 60, and 120 min after injection with the OneTouch Ultra glucose monitor (LifeScan Inc.).

Cholesterol measurements

At the end of the study, mice were euthanized and blood was collected via retro-orbital sinus puncture. Cholesterol and triglycerides were measured from whole blood samples with a CardioCheck Plus Analyzer (PTS Diagnostics, IN). Epididymal white adipose tissue (EWAT) was removed after each mouse was sacrificed by cutting the tissue distal to the epididymal blood vessel. The adipose tissues were then weighed.

Wire-myography

Mice were killed by an overdose of CO₂ inhalation and mesentery were removed and placed in cold Krebs's ringer buffer (KRB) with the following composition (in mmol/L): 118.5 NaCl, 4.7 KCl, 2.5 CaCl₂, 1.2 MgSO₄, 1.2 KH₂PO₄, 25.0 NaHCO₃, and 5.5 D-glucose. From the mesentery, three segments (2 mm) of second order mesenteric artery (MA₂) from each mouse were carefully dissected. Segments were mounted in a wire myograph (model 620M; Danish Myotechnology, Aarhus, Denmark) for the recording of isometric force development. Segments were passively stretched according to a procedure first described by Halpern and Mulvany (Halpern and Mulvany 1977). In brief, segments were distended stepwise, in 50 μ m increments to their optimal lumen diameter for active tension development. Segments were stretched to a passive wall tension of 90% of the internal circumference of that was achieved when they were exposed to a passive tension yielding a transmural pressure of 100 mmHg, which is referred to as its "optimal diameter." At this passive wall tension, arteries were contracted with high K⁺ KRB (60 mmol/L KCl in KRB solution; replacing equimolar NaCl with KCl), thus generating a stable contraction that reached a plateau after 10–15 min. This active wall tension was set to a

100% contraction level. Contractile responses in MA₂ were assessed by analyzing cumulative concentration-response curves (CRCs) with the α_1 -adrenergic agonist Phenylephrine (PHE; 0.01–30 μ mol/L). Relaxing responses were determined in PHE (3–10 μ mol/L) contracted segments by performing CRCs to Acetylcholine (ACh; 0.01–10 μ mol/L). In a subset of MA₂, NO-dependent relaxations were analyzed in conditions where NO release and the endothelium-dependent hyperpolarization (EDH) relaxing response was pharmacologically blocked by a cocktail of the cyclooxygenase blocker indomethacin (INDO; 10 μ mol/L) and the endothelial K_{Ca} channel blockers TRAM-34 and UCL1684 (both 1 μ mol/L). EDH relaxing responses were assessed in the presence of the non-selective NO synthase inhibitor L-NAME and INDO. To study the role of endogenous peroxynitrite (OONO⁻) on EDH responses, MA₂ were incubated with ebselen, a peroxynitrite scavenger (Masumoto and Sies 1996). To assess the effect of HFS diet on endothelium-independent relaxations we first incubated MA₂ with L-NAME and INDO for 30 min to block endothelium-dependent relaxations. Next, we used NO donor SNP (0.0001–1 μ mol/L) in these arteries to determine ACh-mediated relaxing responses in smooth muscle cells that might have been modulated due to HFS diet.

Pressure myography

A segment (4 mm) of the MA₂ was cannulated between two small glass micropipettes (O.D. 120–150 μ m) in a pressure-myograph chamber (model 110P, DMT-USA, Inc., Aarhus, Denmark). The ends of the segment were tied with two small nylon knots (17 μ m thin). The chamber was filled with warm Krebs's ringer buffer (KRB) and continuously gassed with carbogen. Intraluminal pressure was set to 60 mmHg and the artery was allowed to adjust in KRB containing 2.5 mmol/L CaCl₂ for 30 min. The internal lumen diameter and wall thickness were continuously monitored using the MYOVUE II software (DMT-USA, Inc.). Viability was confirmed by adding the α_1 -adrenergic agonist (PHE; 0.1–1 μ mol/L) to the myograph chamber. After a stable diameter was reached, cumulative concentrations of ACh (0.001–100 μ mol/L) were added to the chamber. The chamber was washed three times with KRB and myogenic tone was assessed by monitoring internal diameter during stepwise (10 mmHg) increases in intraluminal pressures from 10 to 140 mmHg. At each pressure step, the artery was allowed to adjust its internal diameter for at least 2 min or until a stable diameter was obtained. Elastic characteristics of the vessel wall were determined under passive (calcium-free) conditions in KRB without CaCl₂ and containing 2.5 mmol/L ethylene glycol tetracetic acid (EGTA) and

the NO donor SNP (10 $\mu\text{mol/L}$) to allow complete relaxation of the smooth muscle. According to Laplace's law, wall tension (T in mN/mm) depends upon transmural pressure (P_t in mN/mm^2 , where 100 mmHg equals 13.33 mN/mm^2) and the radius of the artery (r in mm): $T = P_t \cdot r$. Circumferential wall stress (σ in mN/mm^2) is related to wall thickness (wt in mm) and wall tension (T): $\sigma = T/wt$. Circumferential wall strain was calculated as $(D_i - D_{10})/D_{10}$ where D_i is the lumen diameter at any intraluminal pressure and D_{10} the lumen diameter at 10 mmHg pressure. The incremental elastic modulus (E_{inc}) was calculated according to Bergel (Bergel 1961). $E_{\text{inc}} = 1.5 \cdot r_o^2 \cdot r_i \cdot \Delta P \cdot (r_o^2 - r_i^2) \cdot \Delta r_i^2$, where r_o = outer radius, r_i = inner radius, and $\Delta P = 10$ mmHg or 1.333 mN/mm^2 . Cross-sectional compliance (CC in $\mu\text{m}^2/\text{kPa}$) was calculated as the change in inner lumen area (ΔA_i) induced by a pressure change (ΔP_i).

Statistical analysis

Data were analyzed using one-way analysis of variance (ANOVA) for comparison of multiple means followed by Tukey's post hoc test. Two-way ANOVA, was used for concentration-response curves followed by Bonferroni post hoc test (Prism-Graphpad v6.0). Student's t -test was performed for comparison of two means using Prism v 6.00 (GraphPad Prism software, La Jolla, CA). Data were represented as standard error of mean (SEM). The minimum number of animals used for experiments performed during this study was 6.

Results

Effect of HFS diet on weight gain, glucose tolerance, cholesterol, and triglyceride levels

The duration of feeding to four mice groups are presented in Figure 1A. Young mice on a NC diet for 4 months had significantly lower bodyweight (BW) than mice fed a HFS diet (Fig. 1B). There was no significant difference in the initial BW of both mice groups, but BW gradually increased and reached statistical significance ($P < 0.01$) after 6 weeks on a HFS diet (Fig. 1B). The BW measured after the end of the experiment was significantly higher ($P < 0.01$) in young or aged mice on HFS compared to young or aged mice on NC (Fig. 1C). Mice fed with HFS diet showed significant increase in EWAT (Fig. 1D) and also increased fat accumulation in the visceral cavity (not shown). Since obesity is directly linked to glucose homeostasis, we determined glucose tolerance levels between young HFS mice and age-matched control NC fed mice, and also between aged

NC mice and aged HFS mice with a glucose tolerance test (GTT). As shown in Figure 2A, HFS fed mice had impaired glucose tolerance compared with their age-matched controls on NC. Additionally, aged HFS mice showed a higher glucose retention level compared to aged NC mice (Fig. 2A). However, there was no significant difference in glucose levels between young NC or aged NC mice, indicating no effect of age on glucose tolerance. We further determined insulin sensitivity in HFS or NC mice with an insulin tolerance test (ITT). When young or aged mice were given 0.75 U/kg of insulin after a 6-h fasting period, the blood glucose of mice that were on a HFS diet (both young and aged) showed higher blood glucose retention at 60 and 120 min after the insulin injection compared with mice that were fed a NC diet (Fig. 2B), suggesting that young or aged HFS mice have developed significant insulin resistance. Supporting this finding we found that young or aged HFS mice had increased fasting glucose (Fig. 2C) and serum insulin levels (Fig. 2D) compared with age-matched control young or aged NC mice. We also determined whether HFS diet would impact the cholesterol and triglyceride levels, as their increase is implicated in cardiovascular diseases. As shown in Figure 2E, young or aged HFS mice had an increase in cholesterol level compared with young or aged mice on NC respectively. However, there was no significant difference in the total triglycerides levels (Fig. 2F).

Effect of HFS diet on small mesenteric artery remodeling and vasodilation

We used pressure myography to determine the effect of HFS on resistance artery wall stress and wall thickness since lumen diameter and arterial wall remodeling would impact vascular flow dynamics. We determined the lumen diameter and wall thickness of MA_2 that were cannulated and pressurized in the pressure-myograph chamber. As shown in Figure 3A, the lumen diameters were significantly larger in MA_2 from aged mice on a HFS diet compared to MA_2 from young mice on HFS. Aging alone did not alter lumen diameter in mice on a normal chow diet (Fig. 3A). Similarly, wall thickness was larger in MA_2 from aged HFS mice compared to their younger counterparts (Fig. 3B). Consequently, wall cross-sectional area (CSA) of the MA_2 was significantly increased in aged HFS group compared with younger HFS group (Fig. 3C), which indicated outward hypertrophic remodeling in response to a 1-year HFS diet. Aging alone did not cause any significant changes in wall CSA in MA_2 from NC mice (Fig. 3C). We calculated circumferential wall strain and circumferential wall stress values and plotted them against each other. Figure 3D showed that the slope of




Figure 2. Effect of HFS on glucose intolerance, insulin resistance and cholesterol levels in young and aged mice: Glucose tolerance test at 4 months (young) and at 15 months (aged) (A), Insulin tolerance test at same age mentioned for 'A' (B), fasting blood glucose level of young and aged mice, (C), serum insulin levels (D), total cholesterol in whole blood (E), total triglycerides in whole blood (F), derived from mice receiving 6 months of normal chow (NC young; white bars), 16 months of NC (NC aged; dark gray bars), 3 months of HFD followed by 4 weeks of 10% sucrose in drinking water (HFS young; black bars), and 12 months of HFS (HFS aged; black bars). Values are shown as means \pm SEM ($n = 6$); * $P < 0.05$ versus young NC; ** $P < 0.05$ versus aged NC. $n = 6$; * $P < 0.05$ Young NC versus young HFS, ** Aged NC versus aged HFS.

the strain–stress curves increased in MA_2 from aged mice on a HFS diet compared to young mice on a HFS diet, indicating decreased arterial elasticity in the former. There was no significant difference between the stress strain curves from young or aged mice on a NC diet (Fig. 3D). At higher circumferential wall stress levels incremental elastic modulus (E_{inc}) values were higher for MA_2 from aged mice on a HFS diet compared to the other groups, again indicating an increased arterial stiffness during a near-lifelong HFS diet (Fig. 3E). Due to the observed outward remodeling in MA from aged mice on a HFS diet, the cross-sectional compliance was higher at low pressures, but dropped faster after 50 mmHg due to increased stiffness compared to the other groups (Fig. 3F). We also determined endothelium-dependent ACh-induced vasodilation in these arteries using wire myography. As shown in Figure 4A, ACh-induced vasodilation was significantly impaired in MA_2 from young HFS compared with young NC mice. However, impairment in ACh-induced vasodilation in aged NC mice was similar to young HFS mice (Fig. 4A). In addition, aged HFS mice showed significantly decreased vasodilation compared to aged NC mice (Fig. 4A). These studies indicate that increased arterial stiffness and decreased relaxing responses in these arteries

could have caused impaired vasodilation. Next, we determined the myogenic tone (pressure-induced constriction) in these arteries using pressure myography to evaluate the effect of HFS on myogenic tone which might contribute to impaired vasorelaxation in these arteries. As shown in Figure 4B, myogenic tone was decreased in MA_2 from aged HFS compared with young HFS, but did not reach statistical significance. This indicates a tendency toward impairment of constrictor tone (Fig. 4B). Taken together, decreased elasticity and increased stiffness of MA_2 was observed in mice with HFS, indicating that structural elements of resistance arteries might undergo HFS-mediated remodeling.

Effect of HFS on small mesenteric artery endothelial dysfunction in young and aged mice

Since increased arterial stiffening and decreased arterial elasticity of MA_2 was observed in response to HFS diet, we determined the impact of these arterial structural changes on endothelial dysfunction in HFS diet. Vascular contractile and relaxing responses were measured using wire myography. Vascular contractile responses to




Figure 3. Age and HFS –mediated structural wall remodeling of small mesenteric arteries: Pressure–lumen diameter relationships (A), pressure – wall thickness relationships (B), pressure–wall cross-sectional area (CSA) relationships (C), circumferential wall strain – circumferential wall stress relationships (D), incremental elastic modulus – circumferential wall stress relationships (E), pressure – cross-sectional compliance (F) for small mesenteric arteries derived from mice receiving 4 months of normal chow (NC young; open circles), 12 months of NC (NC aged; open squares), HFS young, closed circles, and 12 months of HFS (HFS aged; closed squares). Values are shown as means \pm SEM ($n = 6$); * $P < 0.05$ versus young NC; # $P < 0.05$ versus young HFS.




Figure 4. Effect of age and HFS on vasodilatation and myogenic tone in small mesenteric arteries: Endothelium-dependent ACh-mediated vasodilatation (A), and myogenic constriction (B) for small mesenteric arteries derived from mice receiving 4 months of normal chow (NC young; open circles), 12 months of NC (NC aged; open squares), HFS young (closed circles), and HFS aged (closed squares). Values are shown as means \pm SEM ($n = 6$); * $P < 0.05$ versus young NC; # $P < 0.05$ versus young HFS.

60 mmol/L KCl were decreased in MA₂ from young HFS mice compared with young NC (Fig. 5A). However, contractile tension in response to this depolarizing solution was significantly increased in MA₂ from aged HFS compared with young HFS (Fig. 5A). Contractile responses to the α_1 -adrenergic agonist PHE were significantly increased in MA from aged mice on a HFS diet compared to the other groups (Fig. 5B). Similar to observations in the pressure-myograph, relaxations to ACh were significantly blunted in MA₂ from young HFS compared with age-matched NC, whereas a combination of aging and HFS diet had a more severe inhibitory effect on endothelium-dependent ACh-mediated relaxations (Fig. 6A). Sensitivity (pEC_{50}) and maximal relaxing responses (E_{max}) to ACh are summarized in Table 1. Interestingly, endothelium-dependent relaxing responses from MA₂ of aged NC mice did not differ from young HFS mice (Fig. 6A), suggesting that a HFS diet can accelerate endothelial dysfunction in young mice similar to aged mice in NC. The endothelial dysfunction caused by a four-month HFS diet was due to increased release of vaso-contractile prostaglandins because the nonselective cyclooxygenase blocker




Figure 5. Effect of age and HFS on contractile response of small mesenteric arteries: Active tension (mN/mm) in response to 60 mmol/L KCl in KRB (A), cumulative concentration-response curves (CRC) to the α_1 -adrenergic receptor agonist phenylephrine (B) for small mesenteric arteries derived from mice receiving 4 months of normal chow (NC young; open circles), 12 months of NC (NC aged; open squares), HFS young (closed circles), and HFS aged (closed squares). Values are shown as means \pm SEM ($n = 6-9$); * $P < 0.05$ versus young NC; # $P < 0.05$ versus young HFS.

indomethacin restored ACh-induced relaxing responses in MA₂ from young HFS (Fig. 6B and Table 1). Surprisingly, prolonged HFS diet for 12 months caused an irreversible endothelial dysfunction that could not be restored by indomethacin (Fig. 6B and Table 1). Microvessels such as MA₂ are known to show EDHF-mediated vasorelaxation, and to a lesser degree, the NO-dependent vasodilation (Takaki et al. 2008). We sought to determine the relative contribution of NO or EDHF on endothelial dysfunction in HFS fed mice. We first used a cocktail of Indomethacin, TRAM34 and UCL1684 to inhibit prostaglandins and endothelial calcium activated K⁺ channels, respectively to block EDHF-mediated relaxation. As shown in Figure 6D, endothelium-dependent (NO-dependent) ACh-mediated relaxations were impaired in aged HFS mice, but not in young HFS mice. Next, we blocked NO-mediated relaxing responses using eNOS blocker, L-NAME and non-selective cyclooxygenase inhibitor indomethacin, and determined ACh-mediated relaxations to determine the contribution of EDHF in small vessel relaxation in HFS. As shown in Figure 6D, an acute impairment of EDHF-dependent relaxation was observed in aged HFS fed mice, but not in aged NC or young NC mice. Young HFS mice showed a mild decrease in relaxation compared to young NC mice, but this was not statistically significant. These data show that although age and HFS-induced endothelial dysfunction was characteristic of both, an impaired NO release (Fig. 6C) and a reduced endothelium-dependent hyperpolarization (Fig. 6D), EDHF response was a major relaxing factor in HFS fed aged mice.




Figure 6. Effect of age and HFS on endothelium-dependent relaxation of small mesenteric arteries: Endothelium-dependent ACh-mediated relaxations in the absence of any inhibitors (A), endothelium-dependent ACh-mediated relaxations in the presence of the non-selective cyclooxygenase blocker indomethacin (B), NO-dependent ACh-mediated relaxations in the presence of indomethacin, TRAM-34 and UCL 1684 (C), endothelium-dependent hyperpolarizing (EDH) relaxing responses in the presence of indomethacin and the non-selective NOS blocker L-NAME (D) for small mesenteric arteries derived from mice receiving 3 months of normal chow (NC young; open circles), NC aged (open squares), HFS young (closed circles), and HFS aged (closed squares). Values are shown as means \pm SEM ($n = 6-9$); * $P < 0.05$ versus young NC; # $P < 0.05$ versus young HFS.

Effect HFS diet on EDHF-mediated endothelial dysfunction in young and aged mice

In a recent publication we have shown increased production of O₂^{•-} in the thoracic artery of aged mice (Hilgers et al. 2017). Dysfunctional eNOS was the major contributor for enhanced O₂^{•-} generation. Others and we have also shown increased ROS generation in response to hyperglycemia and mice fed with HFD (Patel et al. 2013; Heinonen et al. 2014). We reasoned that O₂^{•-} and NO produced in the aged HFS fed mice would react to produce OONO⁻ that might decrease the levels of H₂O₂ causing an impairment of EDHF mediated vasorelaxation, as

Table 1. Effect of pharmacological inhibitors on sensitivity (pEC_{50}) and maximal relaxing responses (E_{max}) to ACh in mesenteric resistance arteries from young and aged mice on a normal chow (NC) and a high fat sucrose (HFS) diet.

Treatment	Parameter	NC young	NC aged	HFS young	HFS aged
No inhibitors	pEC_{50} (-log M)	6.60 ± 0.05	6.05 ± 0.09*	6.04 ± 0.07*	4.66 ± 0.14*. [#]
	E_{max} (%)	91 ± 2	79 ± 8	75 ± 4*	37 ± 9*. [#]
I	pEC_{50} (-log M)	6.19 ± 0.06**	5.90 ± 0.13	6.13 ± 0.08	4.59 ± 0.21*. [#]
	E_{max} (%)	86 ± 4	74 ± 12	78 ± 5	32 ± 10*. [#]
T+U+I	pEC_{50} (-log M)	5.79 ± 0.05**	5.60 ± 0.07**	5.54 ± 0.07**	4.67 ± 0.23*. [#]
	E_{max} (%)	67 ± 3**	66 ± 6**	60 ± 4**	37 ± 9*. [#]
L-N+I	pEC_{50} (-log M)	5.49 ± 0.11**	5.36 ± 0.18**	4.96 ± 0.09**	4.27 ± 0.90*. [#]
	E_{max} (%)	55 ± 6**	54 ± 14**	42 ± 7**	13 ± 11*. [#]
L-N+I+ebselen	pEC_{50} (-log M)	5.74 ± 0.13**	5.97 ± 0.19	6.13 ± 0.06	5.61 ± 0.15**
	E_{max} (%)	72 ± 7**	67 ± 16	73 ± 4	63 ± 9*

The sensitivity (pEC_{50}) of ACh is calculated with GraphPad Prism6 software as described in the methods and denotes the negative logarithmic concentration of ACh that induces 50% relaxation compared to the maximal relaxation (E_{max}). Abbreviations used: I (indomethacin), L-N (L-NAME), T (TRAM-34), and U (UCL1684). See text for inhibitor concentrations used. Values are shown as means ± SEM. * $P < 0.05$ versus NC Young, [#] $P < 0.05$ versus HFS Young, ** $P < 0.05$ versus no inhibitors.

H₂O₂ has been shown as an EDHF in mesenteric vascular bed (Matoba et al. 2000). Therefore, if OONO⁻ was a major inhibitor of EDH response, then we could rescue this response using ebselen, a potent peroxynitrite scavenger (Masumoto and Sies 1996). As shown in Figure 7A, ebselen (0.2 μmol/L) did not have a significant effect in improving EDH responses in MA₂ from young or aged mice on NC diet. However, ebselen markedly ameliorated EDH responses in MA₂ from both young and aged mice on a HFS diet (Fig. 7B). In fact, EDH responses were normalized for all 4 mice groups (Table 1), suggesting that ONOO⁻ production might have contributed to endothelial dysfunction in HFS fed mice. We also determined the endothelium-independent relaxations to NO donor SNP to demonstrate the effect of HFS-mediated

endothelial dysfunction specifically in the endothelium, but not in the smooth muscle cells. As shown in Figure 7C, relaxing responses to the NO donor SNP were unchanged in either group, demonstrating normal smooth muscle NO sensitivity as SNP produces NO via soluble guanylyl cyclase (sGC) in the smooth muscle cells.

Discussion

In this study we have utilized an obesity model (young NC vs. young HFS), an aging model (young NC vs. aged NC), and an aging plus obesity model (young HFS vs. aged HFS). The goal was to determine the effect of obesity on young, as well as aged using the same group of




Figure 7. Effect of ebselen on EDH relaxing responses in small mesenteric arteries from young and aged mice in HFS: (A) Endothelium-dependent hyperpolarizing (EDH) relaxing responses in the absence (L-N+I) or presence of ebselen for small mesenteric arteries derived from young and aged mice receiving normal chow. (B) Endothelium-dependent hyperpolarizing (EDH) relaxing responses in the absence (L-N+I) or presence of ebselen for small mesenteric arteries derived from young and aged mice receiving a high-fat and high-sucrose diet. Values are shown as means ± SEM ($n = 6$); * $P < 0.05$ aged L-N+I versus aged + ebselen; ** $P < 0.05$ young L-N+I versus young + ebselen; (C) Endothelium-independent sodium nitroprusside (SNP; 0.1 nmol/L–1 μmol/L)-mediated relaxations in the presence of L-NAME and indomethacin for small mesenteric arteries derived from mice receiving 3 months of normal chow (NC young; open circles), NC aged (open squares), HFS young (closed circles), and HFS aged (closed squares). Values are shown as means ± SEM ($n = 6–9$).

mice as they age. We have shown that metabolic disorder (such as T2D) induced by HFS-mediated obesity adversely affects vascular functions, which is accentuated as the animal ages. We have shown that obesity is associated with: (1) impaired glucose and insulin tolerance with high blood glucose, high cholesterol levels. (2) metabolic impairment resulting in vascular endothelial dysfunction in resistance arteries, and (3) decreased myogenic tone in resistance arteries. We also showed that aging is associated with impaired ACh-mediated vasodilation. However, there were no significant changes in weight gain, glucose or insulin levels, and glucose tolerance in young and aged groups. Furthermore, there was no alteration in myogenic tone or EDHF response between these groups. However, the aged obese mice showed significant: (1) increase in lumen diameter, arterial wall thickness and increased CSA; (2) decrease in arterial elasticity and increased arterial stiffness; (3) blunted vasodilation, decreased myogenic tone; (4) increase in contractile responses; (5) irreversible endothelial dysfunction that could not be restored by indomethacin; (6) impairment of EDHF-dependent vasorelaxation that could be restored by ebselen.

Our study addressed both functional (contractile and relaxing responses) and structural (arterial remodeling and elasticity) characteristics of small mesenteric arteries derived from young or aged mice on a normal or high-fat diet. Consumption of HFS diet resulted in significant weight gain, hypercholesterolemia, glucose intolerance, and insulin resistance. Indeed, C57BL/6J mice have been shown to be very sensitive to the development of obesity and T2D when placed on high-fat diet (Rebuffe-Scrive et al. 1993; Surwit et al. 1995; Collins et al. 2004). HFS diet in C57BL/6J mice resembles diet-induced obesity and T2D as observed in humans. T2D has also an increasing correlation with age that was recapitulated in our aged mice with HFS diet. We studied C57BL/6J mice that received a normal diet and assessed microvascular reactivity and structure in young (7 months) and aged (15 months) mice. The purpose of these two groups was to dissociate the aging effect from the diet-induced obesity effects.

Short-term effects of HFS diet on microvascular structure and function

Pressure myography was applied to study pressure-induced myogenic constriction and arterial structural parameters, such as inner diameter, wall thickness, media CSA, strain–stress relationships, incremental elastic modulus, and cross-sectional compliance. None of the above-mentioned parameters were significantly altered in small mesenteric arteries from mice after a four-month HFS diet compared with mice on a regular diet. We observed impaired endothelium-dependent ACh-induced relaxing

responses in small mesenteric arteries that were fed a HFS diet for four months. This endothelial dysfunction is in agreement with other experimental obesity rodent models studying this vascular bed (O'Brien et al. 1998; Naderali et al. 2001; Subramanian and MacLeod 2003; Oltman et al. 2006; Young et al. 2008). Our results are in contrast to the study by Ellis and colleagues, who did not observe endothelial dysfunction in small mesenteric arteries in mice on a western diet for 20 weeks (Ellis et al. 2008). We assessed NO-mediated relaxing responses in the presence of the cyclooxygenase blocker indomethacin (to rule out vasoactive prostaglandins) and the endothelial calcium-activated K^+ channel blockers TRAM 34 and UCL 1684 to inhibit EDH and subsequent smooth muscle relaxation. We did not observe a statistically significant reduction in NO-mediated relaxing responses after four months of HFS diet. Besides NO, other important mechanisms contribute to relaxation in small resistance arteries, such as the EDH response. In the mesenteric vascular bed, the Ca^{2+} -activated small ($K_{Ca2.3}$)- and intermediate ($K_{Ca3.1}$)-conductance K^+ channels play a crucial role in the EDH response (Burnham et al. 2002; Hilgers and Webb 2007). Studies have shown that this EDH response mediated by $K_{Ca3.1}$ and $K_{Ca2.3}$ channels remains unaltered or even increased in obesity (Chadha et al. 2010; Climent et al. 2014). When small mesenteric arteries were treated with the NO synthase blocker L-NAME and indomethacin, the residual EDH response tended to be lower after four months of HFS diet, but this did not reach statistical significance compared with mice given a normal diet. Endothelium-dependent ACh-induced relaxation was restored in the presence of indomethacin alone, suggesting an enhanced contribution of vasocontractile prostaglandins released in response to a HFS diet. Indeed, diminished release of arachidonic acid-induced prostacyclin (PGI_2) and increased thromboxane (TXA_2)-mediated contraction has been observed in obesity animal models (Hodnett et al. 2009; Gamez-Mendez et al. 2015). Endothelium-independent relaxations to the NO donor SNP were similar for all groups suggesting no alteration in smooth muscle function. Thus, HFS impairs vasorelaxation in an endothelial-specific manner.

Effects of aging on microvascular structure and reactivity

We observed endothelial dysfunction in small mesenteric arteries from aged mice on a normal diet compared with young mice on a normal diet, as characterized by a reduced sensitivity to ACh, but this impaired NO or EDH-mediated relaxing responses was not statistically significant. Indomethacin potentiated the ACh-induced response in small mesenteric arteries from aged mice on a

NC diet, suggesting generation of vasocontractile prostanoids with aging, as shown by others in rat mesenteric arteries (Matz et al. 2000; Stewart et al. 2000).

It is known that ROS and cytokines, factors that are increased with aging, can induce the expression of inducible prostaglandin H synthase, which leads to vasoconstriction with aging rats (Stewart et al. 2000). However, we did not observe hyper reactivity to phenylephrine in aged mice, suggesting that the functional differences between young and aged vessels were the result of altered endothelial metabolism rather than changes in smooth muscle responsiveness, which was confirmed by the similar sensitivity to the NO donor SNP and comparable structural characteristics between small mesenteric arteries from young and aged mice on a normal diet. Endothelial dysfunction has been described with aging in both animals and humans in response to agonists or hyperemia. A reduced ACh-mediated response was observed in both small and large arteries with aging (Hongo et al. 1988; Atkinson et al. 1994; Kung and Luscher 1995; Taddei et al. 1995). In our study using resistance arteries, which are also considered as microvessels, endothelium-dependent ACh-induced relaxing responses were similar between aged mice on a regular diet and young mice on the HFS diet, suggesting that HFS induces endothelial dysfunction in microvessels in young mice. Consistent with this observation using large conduit arteries such as aortae, Bailey-Downs et al. (2013) have reported endothelial dysfunction in young mice on HFD, which was similar to aged mice on a standard diet.

Long-term deleterious effects of HFS diet

In response to long-term HFS diet, small mesenteric arteries showed an outward hypertrophic remodeling characteristic of both an increased wall thickness and wall cross-sectional area when compared with age-matched mice fed a regular diet. In addition, a decrease in arterial elasticity was observed as evidenced by significant leftward shift of the stress-strain curve and increased incremental elastic modulus in aged mice fed on HFS diet compared with aged mice on a normal diet. Furthermore, small mesenteric arteries from aged mice on a HFS diet showed a tendency for decreased myogenic tone compared to small mesenteric arteries from age-matched NC mice. Previous studies have reported that loss of myogenic tone is associated with altered EDHF response (McSherry et al. 2006), and therefore EDHF could be a major contributor of vasorelaxation in aged HFS mice. Similar myogenic response has also been observed by Ogalla, et al. in their studies with resistance arteries in ApoE^{-/-} mice fed with HFD (Ogalla et al. 2015)

Active tension in response to a depolarizing potassium chloride solution was increased in small mesenteric arteries from aged mice on HFD compared to their counterparts from aged-matched mice on a NC. It is likely that the increase in lumen diameter and wall mass in the former generated a greater contractile force in response to the depolarization. We observed a severely blunted endothelium-dependent ACh-induced relaxing response in small mesenteric arteries from mice fed a HFS diet for 12 months. This was observed in both, pressurized arteries and stretched arteries. This endothelial dysfunction was characteristic of both an impaired NO- and EDH-mediated response. The reduced NO bioavailability is characteristic of obesity (Naderali et al. 2001; Erdos et al. 2002; Frisbee et al. 2002). Increased production of vascular O₂^{•-} has been shown to lead to reduced NO bioavailability, resulting in reduced agonist-induced relaxing responses of mesenteric and skeletal muscle microvessels of obese mice and rats (Bohlen and Lash 1993; Frisbee and Stepp 2001; Bagi et al. 2003). This increase in O₂^{•-} production might be responsible for the observed increase in sensitivity to phenylephrine in small mesenteric arteries from aged mice on a HFS.

Our study also showed a severely impaired EDH response in small mesenteric arteries from aged mice on a HFS diet compared with age-matched mice on a regular diet. Other studies have observed unaltered or even a compensatory increase in the EDH response during obesity (Burnham et al. 2002; Ellis et al. 2008). These conflicting results may be vascular-bed specific and could be explained by the fact that our experimental obesity model is accompanied by hyperglycemia, hypercholesterolemia, and insulin resistance; all of which may have additive deleterious effects on the endothelium. In addition, our temporal study shows the progressive nature of the severity of endothelial dysfunction, hypercontractility, and degree of outward hypertrophic remodeling in small mesenteric arteries. The peroxynitrite scavenger, ebselen, restored the EDH-relaxing response, confirming enhanced ROS production during obesity by other studies (Erdei et al. 2006; Oltman et al. 2006). We propose to conduct future experiments to precisely understand the mechanism by which endothelial dysfunction occurs in a metabolic disease. We have already shown that vascular redox state is critical to endothelial function, and a shift of vascular redox state from reducing state to oxidizing state occurs in aging (Hilgers et al. 2017). This alteration causes age-related hypertension. We hypothesize that metabolic disease such as T2D causes altered vascular redox conditions, which may produce a dysfunctional eNOS and also an impairment of EDHF (Hilgers and Das 2015)

One of the limitations in this study was a lack of ROS detection in the arteries of NC or HFS mice. However, we have recently shown that aged MA₂ generates significant amounts of superoxide anion via a dysfunctional eNOS using electron paramagnetic resonance spectrometry (Hilgers et al. 2017). In addition to eNOS, activated NADPH oxidases (Nox) or mitochondria could also generate HFS-mediated O₂^{•-} (Garcia-Ruiz et al. 2016; Zeng et al. 2017). The other limitation is the specificity of ebselen for scavenging OONO⁻ as ebselen can scavenge other radicals.

Conclusion

In summary, our study presents evidence that prolonged feeding of HFS diet promoted a T2D phenotype in young, as well as aged mice with glucose intolerance and higher blood glucose levels. These changes are associated with blunted endothelial dysfunction in young HFS mice. Additionally, HFS diet with aging exacerbated endothelial dysfunction. Furthermore, we showed that decreased EDHF in aged HFS fed mice is a major contributor to endothelial dysfunction as antioxidant ebselen restored the endothelial function in these mice. We conclude that prolonged HFS diet in aged mice causes endothelial dysfunction in small resistance arteries due to loss of EDHF, complicating glucose tolerance and glucose utilization that accelerates a T2D phenotype.

Acknowledgments

The authors gratefully acknowledge the excellent technical assistance by Dr. R.H Hilgers. The authors gratefully acknowledge the technical assistance of Dr. V.K Sridharan during the course of this study.

References

- Abularrage, C. J., A. N. Sidawy, G. Aidinian, N. Singh, J. M. Weiswasser, and S. Arora. 2005. Evaluation of the microcirculation in vascular disease. *J. Vasc. Surg.* 42:574–581.
- Atkinson, J., R. Tatchum-Talom, and C. Capdeville-Atkinson. 1994. Reduction of endothelial function with age in the mesenteric arterial bed of the normotensive rat. *Br. J. Pharmacol.* 111:1184–1188.
- Bagi, Z., A. Koller, and G. Kaley. 2003. Superoxide-NO interaction decreases flow- and agonist-induced dilations of coronary arterioles in Type 2 diabetes mellitus. *Am. J. Physiol. Heart Circ. Physiol.* 285:H1404–H1410.
- Bailey-Downs, L. C., Z. Tucsek, P. Toth, D. Sosnowska, T. Gautam, W. E. Sonntag, et al. 2013. Aging exacerbates obesity-induced oxidative stress and inflammation in perivascular adipose tissue in mice: a paracrine mechanism contributing to vascular redox dysregulation and inflammation. *J. Gerontol. A Biol. Sci. Med. Sci.* 68:780–792.
- Bender, S. B., J. A. Castorena-Gonzalez, M. Garro, C. C. Reyes-Aldasoro, J. R. Sowers, V. G. DeMarco, et al. 2015. Regional variation in arterial stiffening and dysfunction in Western diet-induced obesity. *Am. J. Physiol. Heart Circ. Physiol.* 309:H574–H582.
- Bergel, D. H. 1961. The static elastic properties of the arterial wall. *J. Physiol.* 156:445–457.
- Bohlen, H. G., and J. M. Lash. 1993. Topical hyperglycemia rapidly suppresses EDRF-mediated vasodilation of normal rat arterioles. *Am. J. Physiol.* 265(1 Pt 2):H219–H225.
- Briones, A. M., R. Aras-Lopez, M. J. Alonso, and M. Salaices. 2014. Small artery remodeling in obesity and insulin resistance. *Curr. Vasc. Pharmacol.* 12:427–437.
- Burnham, M. P., R. Bychkov, M. Feletou, G. R. Richards, P. M. Vanhoutte, A. H. Weston, et al. 2002. Characterization of an apamin-sensitive small-conductance Ca(2+)-activated K(+) channel in porcine coronary artery endothelium: relevance to EDHF. *Br. J. Pharmacol.* 135:1133–1143.
- Carey, V. J., E. E. Walters, G. A. Colditz, C. G. Solomon, W. C. Willett, B. A. Rosner, et al. 1997. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. *Am. J. Epidemiol.* 145:614–619.
- Cassano, P. A., B. Rosner, P. S. Vokonas, and S. T. Weiss. 1992. Obesity and body fat distribution in relation to the incidence of non-insulin-dependent diabetes mellitus. A prospective cohort study of men in the normative aging study. *Am. J. Epidemiol.* 136:1474–1486.
- Chadha, P. S., R. E. Haddock, L. Howitt, M. J. Morris, T. V. Murphy, T. H. Grayson, et al. 2010. Obesity up-regulates intermediate conductance calcium-activated potassium channels and myoendothelial gap junctions to maintain endothelial vasodilator function. *J. Pharmacol. Exp. Ther.* 335:284–293.
- Chan, J. M., E. B. Rimm, G. A. Colditz, M. J. Stampfer, and W. C. Willett. 1994. Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes Care* 17:961–969.
- Chatterjee, S., D. Bedja, S. Mishra, C. Amuzie, A. Avolio, D. A. Kass, et al. 2014. Inhibition of glycosphingolipid synthesis ameliorates atherosclerosis and arterial stiffness in apolipoprotein E^{-/-} mice and rabbits fed a high-fat and -cholesterol diet. *Circulation* 129:2403–2413.
- Chen, C. A., F. De Pascali, A. Basye, C. Hemann, and J. L. Zweier. 2013. Redox modulation of endothelial nitric oxide synthase by glutaredoxin-1 through reversible oxidative post-translational modification. *Biochemistry* 52:6712–6723.

- Climont, B., U. Simonsen, and L. Rivera. 2014. Effects of obesity on vascular potassium channels. *Curr. Vasc. Pharmacol.* 12:438–452.
- Collins, S., T. L. Martin, R. S. Surwit, and J. Robidoux. 2004. Genetic vulnerability to diet-induced obesity in the C57BL/6J mouse: physiological and molecular characteristics. *Physiol. Behav.* 81:243–248.
- Crabtree, M. J., R. Brixey, H. Batchelor, A. B. Hale, and K. M. Channon. 2013. Integrated redox sensor and effector functions for tetrahydrobiopterin- and glutathionylation-dependent endothelial nitric-oxide synthase uncoupling. *J. Biol. Chem.* 288:561–569.
- Donato, A. J., G. D. Henson, R. G. Morgan, R. A. Enz, A. E. Walker, and L. A. Lesniewski. 2012. TNF-alpha impairs endothelial function in adipose tissue resistance arteries of mice with diet-induced obesity. *Am. J. Physiol. Heart Circ. Physiol.* 303:H672–H679.
- Ellis, A., Z. J. Cheng, Y. Li, Y. F. Jiang, J. Yang, M. Pannirselvam, et al. 2008. Effects of a Western diet versus high glucose on endothelium-dependent relaxation in murine micro- and macro-vasculature. *Eur. J. Pharmacol.* 601:111–117.
- Erdei, N., A. Toth, E. T. Pasztor, Z. Papp, I. Edes, A. Koller, et al. 2006. High-fat diet-induced reduction in nitric oxide-dependent arteriolar dilation in rats: role of xanthine oxidase-derived superoxide anion. *Am. J. Physiol. Heart Circ. Physiol.* 291:H2107–H2115.
- Erdos, B., A. W. Miller, and D. W. Busija. 2002. Impaired endothelium-mediated relaxation in isolated cerebral arteries from insulin-resistant rats. *Am. J. Physiol. Heart Circ. Physiol.* 282:H2060–H2065.
- Frisbee, J. C., and D. W. Stepp. 2001. Impaired NO-dependent dilation of skeletal muscle arterioles in hypertensive diabetic obese Zucker rats. *Am. J. Physiol. Heart Circ. Physiol.* 281:H1304–H1311.
- Frisbee, J. C., K. G. Maier, and D. W. Stepp. 2002. Oxidant stress-induced increase in myogenic activation of skeletal muscle resistance arteries in obese Zucker rats. *Am. J. Physiol. Heart Circ. Physiol.* 283:H2160–H2168.
- Gamez-Mendez, A. M., H. Vargas-Robles, A. Rios, and B. Escalante. 2015. Oxidative stress-dependent coronary endothelial dysfunction in obese mice. *PLoS ONE* 10:e0138609.
- Garcia-Ruiz, I., P. Solis-Munoz, D. Fernandez-Moreira, M. Grau, T. Munoz-Yague, and J. A. Solis-Herruzo. 2016. NADPH oxidase is implicated in the pathogenesis of oxidative phosphorylation dysfunction in mice fed a high-fat diet. *Sci. Rep.* 6:23664.
- Giovannucci, E., and W. C. Willett. 1994. Dietary factors and risk of colon cancer. *Ann. Med.* 26:443–452.
- Halpern, W., and M. J. Mulvany. 1977. Tension responses to small length changes of vascular smooth muscle cells [proceedings]. *J. Physiol.* 265:21P–23P.
- Hamilton, C. A., M. J. Brosnan, S. Al-Benna, G. Berg, and A. F. Dominiczak. 2002. NAD(P)H oxidase inhibition improves endothelial function in rat and human blood vessels. *Hypertension* 40:755–762.
- Heinonen, I., P. Rinne, S. T. Ruohonen, S. Ruohonen, M. Ahotupa, and E. Savontaus. 2014. The effects of equal caloric high fat and western diet on metabolic syndrome, oxidative stress and vascular endothelial function in mice. *Acta Physiol. (Oxf)* 211:515–527.
- Higashi, Y., Y. Kihara, and K. Noma. 2012. Endothelial dysfunction and hypertension in aging. *Hypertens. Res.* 35:1039–1047.
- Hilgers, R. H., and K. C. Das. 2015. Role of in vivo vascular redox in resistance arteries. *Hypertension* 65:130–139.
- Hilgers, R. H., and R. C. Webb. 2007. Reduced expression of SKCa and IKCa channel proteins in rat small mesenteric arteries during angiotensin II-induced hypertension. *Am. J. Physiol. Heart Circ. Physiol.* 292:H2275–H2284.
- Hilgers, R. H., V. Kundumani-Sridharan, J. Subramani, L. C. Chen, L. G. Cuello, N. J. Rusch, et al. 2017. Thioredoxin reverses age-related hypertension by chronically improving vascular redox and restoring eNOS function. *Sci. Transl. Med.* 9:eaaf6094.
- Hodnett, B. L., J. A. Dearman, C. B. Carter, and R. L. Hester. 2009. Attenuated PGI2 synthesis in obese Zucker rats. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 296:R715–R721.
- Hongo, K., T. Nakagomi, N. F. Kassell, T. Sasaki, M. Lehman, D. G. Vollmer, et al. 1988. Effects of aging and hypertension on endothelium-dependent vascular relaxation in rat carotid artery. *Stroke* 19:892–897.
- Keogh, J. B., J. A. Grieger, M. Noakes, and P. M. Clifton. 2005. Flow-mediated dilatation is impaired by a high-saturated fat diet but not by a high-carbohydrate diet. *Arterioscler. Thromb. Vasc. Biol.* 25:1274–1279.
- Kung, C. F., and T. F. Luscher. 1995. Different mechanisms of endothelial dysfunction with aging and hypertension in rat aorta. *Hypertension* 25:194–200.
- Lan, K. C., C. Y. Chiu, C. W. Kao, K. H. Huang, C. C. Wang, K. T. Huang, et al. 2015. Advanced glycation end-products induce apoptosis in pancreatic islet endothelial cells via NF-kappaB-activated cyclooxygenase-2/prostaglandin E2 up-regulation. *PLoS ONE* 10:e0124418.
- Li, M., M. Qian, and J. Xu. 2017. Vascular endothelial regulation of obesity-associated insulin resistance. *Front. Cardiovasc. Med.* 4:51.
- Masumoto, H., and H. Sies. 1996. The reaction of ebselen with peroxynitrite. *Chem. Res. Toxicol.* 9:262–267.
- Matoba, T., H. Shimokawa, M. Nakashima, Y. Hirakawa, Y. Mukai, K. Hirano, et al. 2000. Hydrogen peroxide is an endothelium-derived hyperpolarizing factor in mice. *J. Clin. Invest.* 106:1521–1530.
- Matz, R. L., C. Schott, J. C. Stoclet, and R. Andriantsitohaina. 2000. Age-related endothelial dysfunction with respect to nitric oxide, endothelium-derived hyperpolarizing factor and cyclooxygenase products. *Physiol. Res.* 49:11–18.

- McSherry, I. N., S. L. Sandow, W. B. Campbell, J. R. Falck, M. A. Hill, and K. A. Dora. 2006. A role for heterocellular coupling and EETs in dilation of rat cremaster arteries. *Microcirculation* 13:119–130.
- Naderali, E. K., L. C. Pickavance, J. P. Wilding, and G. Williams. 2001. Diet-induced endothelial dysfunction in the rat is independent of the degree of increase in total body weight. *Clin. Sci. (Lond.)* 100:635–641.
- Nagao, T., S. Illiano, and P. M. Vanhoutte. 1992. Calmodulin antagonists inhibit endothelium-dependent hyperpolarization in the canine coronary artery. *Br. J. Pharmacol.* 107:382–386.
- O'Brien, S. F., J. D. McKendrick, M. W. Radomski, S. T. Davidge, and J. C. Russell. 1998. Vascular wall reactivity in conductance and resistance arteries: differential effects of insulin resistance. *Can. J. Physiol. Pharmacol.* 76:72–76.
- Ogalla, E., C. Claro, M. Alvarez de Sotomayor, M. D. Herrera, and R. Rodriguez-Rodriguez. 2015. Structural, mechanical and myogenic properties of small mesenteric arteries from ApoE KO mice: characterization and effects of virgin olive oil diets. *Atherosclerosis* 238:55–63.
- Oltman, C. L., L. L. Richou, E. P. Davidson, L. J. Coppey, D. D. Lund, and M. A. Yorek. 2006. Progression of coronary and mesenteric vascular dysfunction in Zucker obese and Zucker diabetic fatty rats. *Am. J. Physiol. Heart Circ. Physiol.* 291:H1780–H1787.
- Patel, H., J. Chen, K. C. Das, and M. Kavdia. 2013. Hyperglycemia induces differential change in oxidative stress at gene expression and functional levels in HUVEC and HMVEC. *Cardiovasc. Diabetol.* 12:142.
- Petrofsky, J. S., F. Alshammari, G. S. Bains, I. A. Khowailed, H. Lee, Y. N. Kuderu, et al. 2013. What is more damaging to vascular endothelial function: Diabetes, age, high BMI, or all of the above? *Med. Sci. Monit.* 19:257–263.
- Rebuffe-Scrive, M., R. Surwit, M. Feinglos, C. Kuhn, and J. Rodin. 1993. Regional fat distribution and metabolism in a new mouse model (C57BL/6J) of non-insulin-dependent diabetes mellitus. *Metabolism* 42:1405–1409.
- Seals, D. R., K. L. Jablonski, and A. J. Donato. 2011. Aging and vascular endothelial function in humans. *Clin. Sci. (Lond.)* 120:357–375.
- Sharma, A., L. Rizky, N. Stefanovic, M. Tate, R. H. Ritchie, K. W. Ward, et al. 2017. The nuclear factor (erythroid-derived 2)-like 2 (Nrf2) activator dh404 protects against diabetes-induced endothelial dysfunction. *Cardiovasc. Diabetol.* 16:33.
- Shimokawa, H., H. Yasutake, K. Fujii, M. K. Owada, R. Nakaike, Y. Fukumoto, et al. 1996. The importance of the hyperpolarizing mechanism increases as the vessel size decreases in endothelium-dependent relaxations in rat mesenteric circulation. *J. Cardiovasc. Pharmacol.* 28:703–711.
- Sloboda, N., B. Feve, S. N. Thornton, R. Nzietchueng, V. Regnault, G. Simon, et al. 2012. Fatty acids impair endothelium-dependent vasorelaxation: a link between obesity and arterial stiffness in very old Zucker rats. *J. Gerontol. A Biol. Sci. Med. Sci.* 67:927–938.
- Stewart, K. G., Y. Zhang, and S. T. Davidge. 2000. Aging increases PGHS-2-dependent vasoconstriction in rat mesenteric arteries. *Hypertension* 35:1242–1247.
- Subramanian, R., and K. M. MacLeod. 2003. Age-dependent changes in blood pressure and arterial reactivity in obese Zucker rats. *Eur. J. Pharmacol.* 477:143–152.
- Surwit, R. S., M. N. Feinglos, J. Rodin, A. Sutherland, A. E. Petro, E. C. Opara, et al. 1995. Differential effects of fat and sucrose on the development of obesity and diabetes in C57BL/6J and A/J mice. *Metabolism* 44:645–651.
- Taddei, S., A. Virdis, P. Mattei, L. Ghiadoni, A. Gennari, C. B. Fasolo, et al. 1995. Aging and endothelial function in normotensive subjects and patients with essential hypertension. *Circulation* 91:1981–1987.
- Takaki, A., K. Morikawa, M. Tsutsui, Y. Murayama, E. Tekes, H. Yamagishi, et al. 2008. Crucial role of nitric oxide synthases system in endothelium-dependent hyperpolarization in mice. *J. Exp. Med.* 205:2053–2063.
- Urakami-Harasawa, L., H. Shimokawa, M. Nakashima, K. Egashira, and A. Takeshita. 1997. Importance of endothelium-derived hyperpolarizing factor in human arteries. *J. Clin. Invest.* 100:2793–2799.
- Willett, W. C., M. J. Stampfer, G. A. Colditz, B. A. Rosner, and F. E. Speizer. 1990. Relation of meat, fat, and fiber intake to the risk of colon cancer in a prospective study among women. *N. Engl. J. Med.* 323:1664–1672.
- Woodman, C. R., M. A. Thompson, J. R. Turk, and M. H. Laughlin. 2005. Endurance exercise training improves endothelium-dependent relaxation in brachial arteries from hypercholesterolemic male pigs. *J. Appl. Physiol.* (1985) 99:1412–1421.
- Young, E. J., M. A. Hill, W. B. Wiehler, C. R. Triggie, and J. J. Reid. 2008. Reduced EDHF responses and connexin activity in mesenteric arteries from the insulin-resistant obese Zucker rat. *Diabetologia* 51:872–881.
- Zeng, J., S. Deng, Y. Wang, P. Li, L. Tang, and Y. Pang. 2017. Specific inhibition of Acyl-CoA oxidase-1 by an acetylenic acid improves hepatic lipid and reactive oxygen species (ROS) metabolism in rats fed a high fat diet. *J. Biol. Chem.* 292:3800–3809.