

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Rational hand hygiene during the coronavirus 2019 (COVID-19) pandemic



To the Editor: The enhanced preventive measures during the SARS-CoV-2 coronavirus disease 2019 (COVID-19) pandemic include proper hand hygiene. Health care workers (HCWs) may perform frequent handwashing with water and soap, leading to the potential complication of skin damage. In Lan et al's survey¹ of 526 front-line COVID-19 HCWs, 74.5% reported damage to hand skin from enhanced infection prevention measures. HCWs who washed their hands more than 10 times per day reported more damage to hand skin (odds ratio, 2.17). This skin damage creates a route of entry for COVID-19, and basic skincare measures should be taken after handwashing to address this.

Notably, the cell receptor for SARS-CoV-2 entry, angiotensin-converting enzyme 2 (ACE2), is abundantly present in blood vessels/capillaries of the skin, the basal layer of the epidermis, and hair follicles. ACE2 is also present in eccrine glands.²

Yan et al.³ discuss several measures that can be done to avoid hand eczema, maceration, and erosion due to PPE and handwashing during the pandemic. Qualified sanitizers with ethanol as the main component are prioritized for hand decontamination, and the application of hand cream/moisturizers on intact skin after hand washing is instructed.³ During handwashing, overly-hot water can lead to contact dermatitis and should therefore be avoided.

Hypoallergenic moisturizers, gloves, and alcoholic hand sanitizers are vital to prevent hand eczema. Educational intervention about handeczema risk factors—such as frequent hand washing, surgical scrubbing, and prolonged glove use—was shown to promote hand-skin health among HCWs.⁴

Due to the risk of hand-skin damage, rational hand-hygiene measures respectful of the skin along with proper use of protective gloves and moisturizers must be instructed to HCWs during the fight against COVID-19.

- Gregory Cavanagh, BS, and Carlos Gustavo Wambier, MD, PhD
- From the Department of Dermatology, The Warren Alpert Medical School of Brown University, Providence, Rhode Island.
- Funding sources: None.
- Conflicts of interest: None disclosed.
- IRB approval status: Not applicable.
- Reprints not available from the authors.
- Correspondence to: Carlos Gustavo Wambier, MD, PhD, 65 Village Square Dr, Ste 201, South Kingstown, RI 02879
- E-mail: carlos_wambier@brown.edu

REFERENCES

- 1. Lan J, Song Z, Miao X, et al. Skin damage among healthcare workers managing coronavirus disease-2019. J Am Acad Dermatol. 2020;82(5):1215-1216.
- Hamming I, Timens W, Bulthuis MLC, Lely AT, Navis GJ, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. J Pathol. 2004;203: 631-637.
- 3. Yan Y, Chen H, Chen L, et al. Consensus of Chinese experts on protection of skin and mucous membrane barrier for health-care workers fighting against coronavirus disease 2019. [Epub ahead of print]. *Dermatol Ther.* 2020:e13310. https://doi.org/10.1111/dth.13310. Accessed March 13, 2020.
- Gasparini G, Carmisciano L, Giberti I, Murgioni F, Parodi A, Gallo R. "HEALTHY HANDS". A pilot study for the prevention of chronic hand eczema in healthcare workers of an Italian University Hospital. [Epub ahead of print]. *G Ital Dermatol Venereol*. 2019. https://doi.org/10.23736/S0392-0488.19.06220-5. Accessed June 12, 2019.

https://doi.org/10.1016/j.jaad.2020.03.090