

Neonatal innate immunity and Toll-like receptor

Hye Sun Yoon, M.D.

Department of Pediatrics, Eulji Hospital, Eulji University School of Medicine, Seoul, Korea

Received: 15 November 2010,
Accepted: 30 November 2010
Corresponding author: Hye Sun Yoon, M.D.
Department of Pediatrics, Eulji Hospital, 280-1 Hageye-1 dong, Nowon-gu, Seoul 139-711, Korea
Tel: +82.2-970-8228, Fax: +82.2-976-5441
Email : yhs3211@eulji.ac.kr

Copyright © 2010 by The Korean Pediatric Society

The innate immune response is the first line of defense against microbial infections. Innate immunity is made up of the surface barrier, cellular immunity and humoral immunity. In newborn, immunologic function and demands are different to adults. Neonatal innate immunity specifically suppresses Th1-type immune responses, and not Th2-type immune responses, which are enhanced. And the impaired response of macrophages is associated with the defective innate immunity in newborn period. Toll-like receptors (TLRs) play a key roles in the detection of invading pathogens and in the induction of innate immune responses. In newborn, the expression of TLRs is age dependent, so preterm has low expression of TLRs. Also, there are defects in signaling pathways downstream of TLRs. As a consequence, the defects of TLRs activity cause the susceptibility to infection in the neonatal period.

Key words: Innate immunity, Toll-like receptor, Newborn

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Neonatal immunity is immature. Both defective adaptive immunity and innate immunity contribute to the impaired neonatal host defense. Adaptive immune response is impaired at birth due minimal preexisting antigen exposure and due to the impaired functions of effector B and T cells¹. And adaptive immunity is slowly developing system after exposure of microorganisms. In contrast, innate immune response does not require immunologic experience to function well. And innate immunity is the critical first-line barrier of host defense, bridging the interval between exposure to the pathogen and the specific response of the adaptive immune system². Therefore, a neonate is largely dependent on passively acquired antibodies and defective innate immune responses in the defense against microorganisms³. This review describes the current understanding of neonatal innate immunity and its important receptors, Toll-like receptors (TLRs).

Neonatal innate immunity

Innate immunity is a natural host defense mechanism that operates effectively without exposure to a microorganism or its antigen. This system is made up of the surface barrier (such as skin and mucous membrane), many cells (such as neutrophil, monocyte, macrophage, dendritic cell, and natural killer cell) and humoral factors (such as complement)⁴. There are limitations of exposure to antigens in utero and defective adaptive immunity in the newborn period, newborn must rely on their innate immune system for protection⁵.

1. Surface barrier

For the immune system of surface barrier, the mucous membrane and skin have tissue macrophages and dendritic cells act as the first-line defense mechanism against the entry of microorganism. When microorganism enters to the body, these cells under the skin first recognize pathogen and are activated for secretion of

inflammatory cytokines and recruitment of many immune cells. After birth, the neonatal skin and gut are rapidly colonized with microbial floras. The newborn skin has protective mechanism including vernix caseosa and protective epithelial cells. The vernix caseosa consists of lysozyme, α -defensins, ubiquitin, psoriasin and antimicrobial free fatty acids. It can act with immune cells to kill the microorganisms⁶.

The gastrointestinal tract is an important immune organ. It could secrete peptides and proteins and other host defenses. Innate immunity plays a central role in intestinal immune defense against invading pathogens. Also, it serves as a bridge to the activation of the adaptive immune system. Some kinds of intestinal mucosal surface receptors serve as the pathogen recognition receptors in the innate immune defense system. Secreted bactericidal peptides or defensins produced by the intestinal epithelia represent another crucial element of innate mucosal immune defense. Mutations in pattern recognition receptors and dysfunction of secretory bactericidal peptides may impair host immune defenses leading to an invasion of pathogens resulting in chronic inflammation of the gut⁷. Neonatal intestinal immunity can be significantly modified by breastfeeding. Breast milk contains diverse immunological factors, including innate immune molecules such as the antimicrobial proteins and peptides, lactoferrin and lysozyme⁸.

2. Cellular immunity

Mononuclear inflammatory cells, particularly mast cells and tissue macrophages, play a major role in the cellular immunity of host defense against any microbes that break the physical barriers such as intact skin. Mast cells act by releasing tumor necrosis factor- α (TNF- α). TNF- α recruits other innate host defense cells such as polymorphonuclear leukocytes (PMNs), monocytes and dendritic cells. Also, it can instruct dendritic cells to process and present microbial antigenic material to T lymphocytes. This is an initial step in the generation of a specific immune response⁴.

Circulating monocytes differentiate into mononuclear macrophages which can release many proinflammatory cytokines. In the newborn period, the function and the number of monocyte and mononuclear macrophage are normal. In vitro studies of cord blood from healthy term baby have effective phagocytosis activity similar to adult. But the functional maturation of TLR expression on the immune cell is a main problem for the defective immune system.

Also, selective impairment of specific immune system existed. For example, interferon- γ (IFN- γ) production is decreased in the newborn period⁹. Overall, neonatal cellular immune system expresses selective impairments in Th1-type immune response production and Th2-type immune response enhancement. Th1

and Th2-type immune response derived from proliferating helper T cells. Proliferating helper T cells differentiate into two major subtypes of cells known as Th1 and Th2 cells (also known as type 1 and type 2 helper T cells, respectively). Th1 cells produce IFN- γ , interleukin (IL)-2, and tumor necrosis factor- β (TNF- β), which activate macrophages and are responsible for cell-mediated immunity and phagocyte-dependent protective responses. By contrast, Th2 cells produce IL-4, IL-5, IL-10, and IL-13, which are responsible for strong antibody production, eosinophil activation, and inhibition of several macrophage functions, thus providing phagocyte-independent protective responses. Th1-type immune system relate to the alloimmune reaction between mother and fetus, and major cause of fetal loss. So, during pregnancy, placenta-derived Th2-type immune cytokines antagonize Th1-type immune responses, there are neonatal Th2-type immune polarization¹⁰. Therefore, during the newborn period, Th1-type immune cytokines such as TNF, IFN- α , IFN- γ , IL-12, IL-1b are decreased. In contrast, Th2-type immune cytokines such as IL-6, IL-8, IL-10, and IL-23 increased during newborn period.

Also, macrophages from newborn are hyporesponsive to activation by IFN- γ . The impaired production of Th1-type immune cytokines and the impaired response of macrophages are associated to the defective innate immunity in newborn period.

Toll like receptor

In mammals, a key factor of the innate immune system is a family of TLRs. Toll receptors were originally identified in *Drosophila*¹¹ and not only function in determination of dorsal-ventral development in the fly embryo but also for antimicrobial function. In adult *Drosophila*, the Toll receptor recognize microorganism and activate a signaling pathway for induction of antifungal and antibacterial peptides about infection. In 1997, Janeway et al. identified TLR4, a human homologue of Toll receptor¹². They found out that the activation of TLR4 leads to activation of various inflammatory cytokines such as nuclear factor- κ B (NF- κ B), IL-1, IL-6, and IL-8. He first suggested the existence of a set of factors which he named pattern recognition receptor (PRR), were capable of detecting pathogen-associated molecular patterns (PAMPs). TLRs act as a main PRR to the various PAMPs of microorganisms¹³. If a pathogen enters to the body, TLR recognizes microorganism PAMPs, and activates the signal pathway by utilization of an adaptor protein and results in activation of the transcription factor NF- κ B and expression of inflammatory cytokine genes^{14,15}. As a result, various cytokines, chemokines and colony stimulating factors produced and these factors have a crucial role in the induction of immune response.

Another key role is a link between detection and recognition of microbial pathogens and the activation of the adaptive immune system¹⁶. TLRs are one kind of transmembrane receptors composed of extracellular leucine-rich repeat (LRR) motifs and a cytoplasmic Toll/interleukin-1 receptor (TIR) homology domain. They are expressed on the surface of monocytes, macrophages, dendritic cells and epithelial cells or in the cytoplasm of cells from different tissues. In humans, 10 TLRs have emerged to mediate the activation of innate effector cells. Some of the TLRs reside at the cell surface, and some of them are located in the endoplasmic reticulum. TLR1, 2, 4, 5 and 6 are present on the cell membrane which recognize mainly microbial membrane components such as lipids, lipoproteins and proteins, whereas TLR3, 7, 8 and 9 appear to be confined to intracellular compartments where they recognize microbial nucleic acid. Each TLR has different ligand recognition and has a distinct function in terms of PAMP recognition and immune response (Table 1).

TLR4 responds to bacterial lipopolysaccharide (LPS), an outer membrane component of Gram-negative bacteria. In addition to LPS, TLR4 is involved in recognition of respiratory syncytial virus fusion proteins¹⁷. TLR2 form a heterodimer with TLR1 or TLR6 recognize various microbial components, such as peptidoglycans, lipoprotein, and lipoteichoic acid from Gram-positive bacteria. TLR5 is involved in the recognition of flagellin protein component of bacterial flagella. TLR5 is expressed and functional to the basolateral site of the epithelial cell in intestine. TLR3, 7, 9 recognize the genomic RNA of virus including respiratory syncytial virus. TLR3 triggers mainly antiviral immune responses¹⁷.

In newborn, basal expression of TLR, the accessory protein, and

the adaptor protein are equivalent to adult level, but activation by the TLR appears to be reduced¹⁸. This reduction is age dependent pattern. One study with monocytes from very low birth weight infant (VLBWI) found that decreased surface expression of TLR4, lower mRNA expression of TLR4 and reduced cytokine production after LPS stimulation in gestational age less than 30 weeks¹⁹. Thus, monocytes in premature infants can be considered phenotypically and functionally immature. The other characteristic is impaired TLR4-mediated signaling in LPS-exposed newborn monocytes. Recent study provided evidence that TLR4-mediated, NF- κ B dependent transcriptional activation of neonatal monocytes macrophages was depressed²⁰. As a consequence, the defects of TLR activity cause the susceptibility to infection in the neonatal period.

TLR associated neonatal researches

So far, there are little human researches about association with neonatal diseases and TLRs. Most researches are animal based. Among them, the relation with TLRs and neonatal sepsis is most advanced part²¹⁻²³. TLR2 and TLR4 expression is enhanced in preterm delivery with chorioamnionitis, and recent studies showed an association between preterm birth and TLR polymorphisms²⁴. Recent study showed polymorphisms in TLR2, TLR5, IL10, and PLA2G2A genes were associated with preterm sepsis²⁵.

The direct role of TLRs in neonatal brain injury is suspected^{26, 27}. The strong innate immune response in the central nervous system (CNS) is critical for pathogen elimination and vital to host survival. However, it is evident that chronic or acute dysregulated inflammation in the CNS can cause tissue damage and neurodegeneration. Animal studies described a relation between the TLR4 agonist LPS and brain injury in newborn animals²⁸. In necrotizing enterocolitis (NEC), increased proinflammatory cytokines are found in intestinal samples, suggesting that these mediators play a role in NEC development. One study in a rat NEC model showed overexpression of TLR2, as well as NF- κ B in intestinal epithelial cells that was correlated with the severity of mucosal damage^{29,30}.

Also, the intestinal expression of TLRs and cytokines precedes histological injury in the experimental NEC³¹. NEC is also associated with increased expression of TLR4 in the intestinal mucosa and physiological stressors such as exposure to LPS and hypoxia. These findings demonstrate a critical role for TLR4 in the development of NEC through effects on enterocyte injury and repair³².

Table 1. Ligands for Various Toll Like Receptors in Human

Toll like receptor (TLR)	Ligands
TLR1 (with TLR2) TLR2	Triacyllipopeptides (bacteria) Peptidoglycan (bacteria) Lipoteichoic acid Diacyllipopeptides (mycoplasma) Zymosan (fungi) Atypical lipopolysaccharide (bacteria) Lipoprotein HSP70
TLR3	Double-stranded RNA (virus)
TLR4	Lipopolysaccharide (bacteria) HSP60 (chlamydia) F protein (respiratory syncytial virus)
TLR5	Flagellin (bacteria)
TLR6	Diacyllipopeptides (mycoplasma)
TLR7	Single-stranded RNA (virus)
TLR8	Single-stranded RNA (virus)
TLR9	CpGDNA (bacteria, virus)
TLR10	Unknown

Conclusion

The innate immune system in neonate is immature which results in increased susceptibility to various infections. On the other hand, newborn innate immunity has a characteristic feature that the balance of Th1 and Th2 type immunity and impairment of TLRs signaling downstream pathway. However, the functional state of the various components of innate immunity in newborn largely unknown and only recently a number of studies have assessed this feature of the innate immune system. Thus, more researches need to include the discovery of neonatal immune mechanism so that prevention and treatment of neonatal infections can more safely be targeted.

References

- 1) Adkins B, Leclerc C, Marshall-Clarke S. Neonatal adaptive immunity comes of age. *Nat Rev Immunol* 2004;4:553-64.
- 2) Marodi L. Neonatal innate immunity to infectious agents. *Infect Immun* 2006;74:1999-2006.
- 3) Belderbos M, Levy O, Bont L. Neonatal innate immunity in allergy development. *Curr Opin Pediatr* 2009;21:762-9.
- 4) Clapp DW. Developmental regulation of the immune system. *Semin Perinatol* 2006;30:69-72.
- 5) Kenzel S, Henneke P. The innate immune system and its relevance to neonatal sepsis. *Curr Opin Infect Dis* 2006;19:264-70.
- 6) Tollin M, Bergsson G, Kai-Larsen Y, Lengqvist J, Sjövall J, Griffiths W, et al. Vernix caseosa as a multi-component defence system based on polypeptides, lipids and their interactions. *Cell Mol Life Sci* 2005;62:2390-9.
- 7) Levy O. Innate immunity of the newborn: basic mechanisms and clinical correlates. *Nat Rev Immunol* 2007;7:379-90.
- 8) Maródi L, Káposzta R, Campbell DE, Polin RA, Csongor J, Johnston RB Jr. Candidacidal mechanisms in the human neonate. Impaired INF-gamma activation of macrophages in newborn infants. *J Immunol* 1994; 153:5643-9.
- 9) Marodi L. Innate cellular immune responses in newborns. *Clin Immunol* 2006;118:137-44.
- 10) Belderbos ME, van Bleek GM, Levy O, Blanken MO, Houben ML, Schuijff L, et al. Skewed pattern of toll-like receptor 4-mediated cytokine production in human neonatal blood: low LPS-induced IL-12p70 and high IL-10 persist throughout the first month of life. *Clin Immunol* 2009;133:228-37.
- 11) Anderson KV, Jurgens G, Nusslein-Volhard C. Establishment of dorsal-ventral polarity in the *Drosophila* embryo: genetic studies on the role of the Toll gene product. *Cell* 1985;42:779-89.
- 12) Medzhitov R, Preston-Hurlburt P, Janeway CA Jr. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 1997;388:394-7.
- 13) Uematsu S, Akira S. Toll-like receptors and innate immunity. *J Mol Med* 2006;84:712-25.
- 14) Flier A, Krediet TG. Innate immunity: Toll-like receptors and some more. *Neonatology* 2007;92:145-57.
- 15) Lakhani SA, Bogue CW. Toll-like receptor signaling in sepsis. *Curr Opin Pediatr* 2003;15:278-82.
- 16) Lee JY, Zhao L, Hwang DH. Modulation of pattern recognition receptor-mediated inflammation and risk of chronic diseases by dietary fatty acids. *Nutr Rev* 2010;68:38-61.
- 17) Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 2010;11:373-82.
- 18) Kollmann TR, Crabtree J, Rein-Weston A, Blimkie D, Thommai F, Wang XY, et al. Neonatal innate TLR-mediated responses are distinct from those of adults. *J Immunol* 2009;183:7150-60.
- 19) Föster-Waldl E, Sadeghi K, Tamandl D, Gerhold B, Hallwirth U, Rohrmeister K, et al. Monocyte Toll-like receptor 4 expression and LPS-induced cytokine production increase during gestational aging. *Pediatr Res* 2005;58:121-4.
- 20) Hasegawa K, Ichiyama T, Isumi H, Nakata M, Sase M, Furukawa S. NF-kappa B activation in peripheral blood mononuclear cells in neonatal asphyxia. *Clin Exp Immunol* 2003;132:261-4.
- 21) Viemann D, Dubbel G, Schlieffenbaum S, Harms E, Sorg C, Roth J. Expression of toll-like receptors in neonatal sepsis. *Pediatr Res* 2005;58:654-9.
- 22) Levy E, Xanthou G, Petrakou E, Zacharioudaki V, Tsatsanis C, Fotopoulos S, et al. Distinct roles of TLR4 and CD14 in LPS-induced inflammatory responses of neonates. *Pediatr Res* 2009;66:179-84.
- 23) Harju K, Ojaniemi M, Rounioja S, Glumoff V, Paananen R, Vuolteenaho R, et al. Expression of Toll-like receptor 4 and endotoxin responsiveness in mice during perinatal period. *Pediatr Res* 2005;57:644-8.
- 24) Krediet TG, Wiertsema SP, Vossers MJ. Toll-like receptor 2 polymorphism is associated with preterm birth. *Pediatr Res* 2007;62:474-6.
- 25) Abu-Maziad A, Schaa K, Bell EF, Dagle JM, Cooper M, Marazita ML, et al. Role of polymorphic variants as genetic modulators of infection in neonatal sepsis. *Pediatr Res* 2010;68:323-9.
- 26) Mallard C, Wang X, Hagberg H. The role of Toll-like receptors in perinatal brain injury. *Clin Perinatol* 2009;36:763-72.
- 27) Hagberg H, Mallard C, Jacobsson B. Role of cytokines in preterm labour and brain injury. *BJOG* 2005;112 Suppl 1:16-8.
- 28) Hagberg H, Peebles D, Mallard C. Models of white matter injury; comparison of infectious, hypoxic-ischemic, and excitotoxic insults. *Ment Retard Dev Disabil Res Rev* 2002;8:30-8.
- 29) Le Mandat Schultz A, Bonnard A, Barreau F, Aigrain Y, Pierre-Louis C, Berrebi D, et al. Expression of TLR-2, TLR-4, NOD2 and pNFkB in a neonatal rat model of necrotizing enterocolitis. *PLoS One* 2007;2:e1102.
- 30) Martin CR, Walker WA. Intestinal immune defences and the inflammatory response in necrotizing enterocolitis. *Semin Fetal Neonatal Med* 2006;11:369-77.
- 31) Liu Y, Zhu L, Fatheree NY, Liu X, Pacheco SE, Tatevian N, et al. Changes in intestinal Toll-like receptors and cytokines precede histological injury in a rat model of necrotizing enterocolitis. *Am J Physiol Gastrointest Liver Physiol* 2009;297:G442-50.
- 32) Leapheart CL, Cavallo J, Gripar SC, Cetin S, Li J, Branca MF, et al. A critical role for TLR4 in the pathogenesis of necrotizing enterocolitis by modulating intestinal injury and repair. *J Immunol* 2007;179:4808-20.