

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Case Report

Computed tomography features of supracardiac total anomalous pulmonary venous connection in an infant

Tariq Alam MMBS, FCPS^a, Hidayatullah Hamidi MD^{b,*}, Mer Mahmood Shah Hoshang MD^b

^a Department of Radiology, Northwest General Hospital Peshawar, KPK, Pakistan

^b Department of Radiology, French Medical University for Children (FMIC), Kabul, Fifth district, 1006 Kabul, Afghanistan

ARTICLE INFO

Article history:

Received 20 February 2016

Received in revised form

8 March 2016

Accepted 17 April 2016

Available online 14 May 2016

Keywords:

anomalous pulmonary venous connection
congenital heart disease
cyanosis

ABSTRACT

Total anomalous pulmonary venous connection (TAPVC) is a rare congenital anomaly of the pulmonary veins drainage. In this entity, the pulmonary veins, instead of draining to left atrium, connect abnormally to the systemic venous circulation. A right-to-left shunt is obligatory for survival. Based on its type and degree of pulmonary venous obstruction, TAPVC may result in pulmonary hypertension and congestive heart failure. In severe cases, urgent diagnosis and surgical correction is essential to reduce morbidity and mortality. Echocardiography as the first and safest imaging modality for cardiovascular abnormalities may fail in complete depiction of some complex feature of TAPVC. Computed tomography angiography is then a noninvasive and sensitive choice for mapping the pulmonary veins without the need for invasive cardiac catheterization. Contrast-enhanced MR angiography can be a radiation-free alternative. Authors present a computed tomography-detected supracardiac TAPVC with small patent ductus arteriosus in a 2 months cyanotic infant.

© 2016 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Case presentation

A 2-month-old female infant was sent to radiology department to undergo chest computed tomography (CT) angiography for evaluation of pulmonary veins (PVs). The child developed cyanosis in her first week of life. Unfortunately, neither prenatal ultrasound nor postnatal echocardiography reports were available.

Chest CT angiography was performed with 128-slice Siemens scanner after administration of 8 mL of nonionic water-soluble contrast material.

The CT findings were as follows:

All 4 PVs were joining together behind the small left atrium (LA; Fig. 1) and their confluence converging to a large vertical vein (VV) that was ascending lateral to the pulmonary trunk. LA was smaller in size and no PV entrance was noted into it (Fig. 2). The VV was draining into the dilated left innominate vein and subsequently to enlarged superior vena cava (SVC) and right atrium (RA; Fig. 3). The RA and right ventricle (RV) were also dilated (Fig. 3).

A small (approximately 6.5 mm) atrial septal defect (ASD) was detected (Fig. 2). The small ASD was functioning as right-

The work was primarily carried out in French Medical University for Children (FMIC); Kabul, Afghanistan.

Competing Interests: The authors have declared that no competing interests exist.

* Corresponding author.

E-mail address: Hidayatullah.hamidi@gmail.com (H. Hamidi).
<http://dx.doi.org/10.1016/j.radcr.2016.04.005>

1930-0433/© 2016 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – Axial (right) and coronal (left) postcontrast computed tomography sections: the right pulmonary veins (RPVs) and left pulmonary veins (LPVs) join all together posterior to the left atrium (LA) and left ventricular outlet (LVO). The confluence of PVs (red star) converges into a large left vertical vein (LVV) that is ascending lateral to the pulmonary trunk (PT). The superior vena cava (SVC), right atrium (RA), right ventricle (RV), and even left internal jugular vein (IJV) are dilated. The pulmonary trunk (PT) is prominent compared to the ascending aorta (AA) at the same level.

to-left shunt and made this supracardiac total anomalous pulmonary venous connection (TAPVC) compatible with life.

No evidence of patent ductus arteriosus (PDA) or ventricular septal defect (VSD) was observed.

Discussion

Background

TAPVC is a rare congenital anomaly of the PVs drainage that connects abnormally to the systemic venous circulation instead of LA. [1] The incidence of this rare entity is

approximately 1 of 17,000 live births [2] or 1%-5% of all cardiovascular congenital anomalies [3].

TAPVC is caused by nonfusion of pulmonary venous confluence with LA, thus, the oxygenated pulmonary blood is redirected to right heart and pumped once again to the lungs [4]. A right-to-left shunt is obligatory for survival that typically occurs at atrial level through either an ASD, patent foramen ovale [3] complete absence of the atrial septum [4] or less commonly a PDA [3]. Some of the mixed blood of the RA crosses through this shunt to the left heart to supply the systemic circulation [4].

Cases of tetralogy of Fallot and double-outlet RV are reported in association with TAPVC [5]. Higher frequency of TAPVC is observed in patients with heterotaxy syndromes especially asplenia [6].




Fig. 2 – Reformatted axial image at atrial level: the right atrium (RA) is dilated, whereas the left atrium (LA) has smaller size. The small atrial septal defect (ASD) is well depicted. The congested lung vessels can also be seen.

Clinical perspective

TAPVC is a cause of neonatal cyanosis, and if right-to-left shunt is not present, it may rapidly result in death [3]; hence, being a surgical emergency in neonates [2].

In severe cases, urgent diagnosis and surgical correction is essential to reduce morbidity and mortality [2]. However, if there is no obstruction and a sufficient shunt is present, patients may not present to clinic urgently [7].

TAPVC can occasionally be misdiagnosed as persistent pulmonary hypertension of the newborn and even echocardiographic findings may overlap [7].

Classification

Based on the location of pulmonary venous drainage, TAPVC is classified into 4 major types [1,3,4]. In all 4 types, complete drainage of pulmonary venous blood is directed to the right heart [4].




Fig. 3 – VRT (right) and MPR (left) reformatted coronal images through the abnormal pulmonary venous pathway: the left vertical vein (LVV), the left brachiocephalic (innominate) vein (BCV), and the superior vena cava (SVC) as well as the right atrium (RA), and right ventricle (RV) are well depicted. This whole pathway is dilated and even early reflux of contrast to the dilated inferior vena cava (IVC) is also seen. The cardiac apex is elevated representing dilatation of the right heart.

Type I: supracardiac (approximately 50%): As in our case, PVs coalesce in the midline, forming a transverse confluence just behind and slightly above the small LA. The confluence drains into the remnant of left cardinal vein (VV), then into the left innominate vein, subsequently crossing the midline toward SVC and finally flows in to the RA [4].

Rarely the confluence may drain directly to right SVC, left SVC, or azygous system [8].

Type II: cardiac (approximately 25%): the pulmonary venous confluence drains either to the coronary sinus or directly into the RA [3,4].

Type III: infracardiac (approximately 25%): the pulmonary venous confluence may drain into IVC, hepatic vein, azygous system, or portal venous system [3,4].

Type IV: mixed type (approximately 9%) [3]: usually no midline pulmonary confluence is present; thus, the right and left PVs may have different drainages. Almost any combination of drainage may occur into SVC, innominate veins, coronary sinus, RA, azygous vein [4], or infradiaphragmatic veins [3].

Pulmonary venous obstruction may occur in any type, but it is most commonly seen in infracardiac type which may be present in up to 78% of cases [3,8].

Imaging perspective

The diagnostic imaging findings are those described previously corresponding to each type.

Echocardiography as the first and safest imaging modality for cardiovascular abnormalities may fail in complete depiction of some complex feature of TAPVC [3]. CT angiography is then a noninvasive and sensitive choice for mapping the PVs without need for invasive cardiac catheterization [2,9].

Contrast-enhanced magnetic resonance (MR) angiography [3] is the radiation-free alternative for CT angiography but needs general anesthesia in younger children.

In a case report, Masaki Ogawa et al suggested that electrocardiography-gated CT and phase-contrast cine MR imaging (PC-MRI) are more useful than non-electrocardiography-gated CT for differentiating the VV in TAPVC from persistent left SVC in adult patients [1].

Outcomes

Based on its type and degree of pulmonary venous obstruction, TAPVC may result in pulmonary hypertension and congestive heart failure [1]. Approximately 50% of patients develop symptoms in neonatal period and the majority may expire during infancy without surgical repair [10].

Unobstructed type I TAPVC has very similar pathophysiology to a large ASD, so lesions not detected in infancy may be discovered later [4]. Several cases of adult untreated TAPVC have been reported, even few cases were diagnosed after 60 years of age [1,11,12].

Generally, patients with type I and II TAPVC can live for older age than those with type III and IV lesions [13]. The most important factors affecting the survival period are large ASDs and normal to near normal pulmonary artery pressure [1].

As precise diagnosis is essential and surgery is the mainstay of treatment, improvements in diagnostic imaging resulted in accurate depiction of this complex entity and advanced surgical techniques decreased the postoperative mortality rates. [3]

REFERENCES

- [1] Ogawa M, Nakagawa M, Hara M, Ito M, Goto T, Ohte N, et al. Total anomalous pulmonary venous connection in a 64-year-old man: a case report. *Ann Thorac Cardiovasc Surg* 2013;19(1):46–8.
- [2] Lakshminrusimha S, Wynn RJ, Youssfi M, Pabalan MJ, Bommaraju M, Kirmani K, et al. Use of CT angiography in the diagnosis of total anomalous venous return. *J Perinatol* 2009;29(6):458–61.

- [3] Dillman JR, Yarram SG, Hernandez RJ. Imaging of pulmonary venous developmental anomalies. *AJR Am J Roentgenol* 2009;192(5):1272–85.
- [4] Hines MH, Hammon JW. Anatomy of total anomalous pulmonary venous connection. *Oper Tech Thorac Cardiovasc Surg* 2001;6(1):2–7.
- [5] DeLeon SY, Gidding SS, Ilbawi MN, Idriss FS, Muster AJ, Cole RB, et al. Surgical management of infants with complex cardiac anomalies associated with reduced pulmonary blood flow and total anomalous pulmonary venous drainage. *Ann Thorac Surg* 1987;43(2):207–11.
- [6] Karamlou T, Gurofsky R, Al Sukhni E, Coles JG, Williams WG, Caldarone CA, et al. Factors associated with mortality and reoperation in 377 children with total anomalous pulmonary venous connection. *Circulation* 2007;115(12):1591–8.
- [7] Allan LD, Sharland GK. The echocardiographic diagnosis of totally anomalous pulmonary venous connection in the fetus. *Heart* 2001;85(4):433–7.
- [8] White CS, Baffa JM, Haney PJ, Pace ME, Campbell AB. MR imaging of congenital anomalies of the thoracic veins. *Radiographics* 1997;17(3):595–608.
- [9] Hu ZH, Wang SY, Yang YY. Diagnosis of total anomalous pulmonary venous connection using multi-slice spiral CT. *Nan Fang Yi Ke Da Xue Xue Bao* 2009;29(9):1922–4.
- [10] Nurkalem Z, Gorgulu S, Eren M, Bilal MS. Total anomalous pulmonary venous return in the fourth decade. *Int J Cardiol* 2006;113(1):124–6.
- [11] Meng F, Sun JP, Chen M, Lee AP, Yu CM. Supracardiac total anomalous pulmonary venous connection. *Int J Cardiol* 2014;174(1):141–2.
- [12] McMullan MH, Fyke FE. Total anomalous pulmonary venous connection: surgical correction in a 66-year-old man. *Ann Thorac Surg* 1992;53(3):520–1.
- [13] Feng Q, Wu S, Yu G. Surgical treatment of a 56-year-old woman with an intracardiac type of total anomalous pulmonary venous connection. *Thorac Cardiovasc Surg* 2010;58(3):175–6.