

POSTER PRESENTATION

Open Access


# Reproduction of EEG power spectrum over frontal region during the propofol-induced general anesthesia

Meysam Hashemi<sup>1\*</sup>, Axel Hutt<sup>1</sup>, Jamie Sleigh<sup>2</sup>, Peter beim Graben<sup>3</sup>

From The Twenty Third Annual Computational Neuroscience Meeting: CNS\*2014  
Québec City, Canada. 26-31 July 2014

The present work aims to reproduce certain changes observed experimentally in the EEG power spectrum over the frontal head region during general anesthesia induced by propofol. These observations include increased delta (0-4 Hz) and alpha (8-12 Hz) activities [1]. We extend a previous cortical model [2] and study a neuronal population model of a single thalamocortical module consisting of three different populations of neurons, namely cortical excitatory neurons, thalamocortical relay neurons and inhibitory thalamic reticular neurons (Fig. 1). Each module obeys a neural mass model. The cortical inhibitory population is neglected in our model to reveal the effect of propofol action in the thalamus. Our model describes well the characteristic spectral changes observed experimentally within the delta- and alpha- frequency bands in frontal and occipital electrodes with increasing concentration of propofol.

This shows that neglecting inhibitory action in the cortex but considering thalamic GABAergic action suffices to reproduce the data. From a modeling point of view, our reduced mathematical model is low dimensional and remains analytically treatable while still being adequate to reproduce observed changes in EEG rhythms. Moreover, it is shown that the propofol concentration acts as a control parameter of the system and that propofol-induced changes in the stationary states of the model system lead to changes in the corresponding nonlinear gain function that result in EEG power modulation: increases of power over the frontal region can be caused by an increase in the gain function of thalamocortical network. The results suggest that intra-thalamic inhibition from reticular neurons to relay cells plays an important role in the generation of



the characteristic EEG patterns seen during general anesthesia.

**Authors' details**

<sup>1</sup>INRIA CR Nancy - Grand Est, Villers-les-Nancy, France. <sup>2</sup>Department of Anaesthetics, Waikato Hospital, Hamilton, New Zealand. <sup>3</sup>Department of German Language and Linguistic, Humboldt-Universität zu Berlin, Germany.

Published: 21 July 2014

**References**

1. Cimener A, et al: Tracking brain states under general anesthesia by using global coherence analysis. *PNAS* 2011, 108:8832-8837.
2. Hutt A: The anaesthetic propofol shifts the frequency of maximum spectral power in EEG during general anaesthesia: analytical insights from a linear model. *Front. Comput. Neurosci* 2013, 7:2.

\* Correspondence: meysam.hashemi@inria.fr

<sup>1</sup>INRIA CR Nancy - Grand Est, Villers-les-Nancy, France

Full list of author information is available at the end of the article

doi:10.1186/1471-2202-15-S1-P211

**Cite this article as:** Hashemi et al.: Reproduction of EEG power spectrum over frontal region during the propofol-induced general anesthesia. *BMC Neuroscience* 2014 **15**(Suppl 1):P211.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

