

LETTER

Climate warming can reduce biocontrol efficacy and promote plant invasion due to both genetic and transient metabolomic changes

Yan Sun^{1,2}  | Tobias Züst³ | Daniele Silvestro^{2,4,5} | Matthias Erb⁶  |
Oliver Bossdorf⁷  | Pierre Mateo⁶ | Christelle Robert⁶ | Heinz Müller-Schärer² 

¹College of Resources and Environment, Huazhong Agricultural University, Wuhan, China

²Department of Biology/Ecology & Evolution, University of Fribourg, Fribourg, Switzerland

³Institute of Systematic and Evolutionary Botany, University of Zürich, Zürich, Switzerland

⁴Swiss Institute of Bioinformatics, Fribourg, Switzerland

⁵Department of Biological and Environmental Sciences and Global Gothenburg Biodiversity Centre, University of Gothenburg, Gothenburg, Sweden

⁶Institute of Plant Sciences, University of Bern, Bern, Switzerland

⁷Plant Evolutionary Ecology, Institute of Evolution & Ecology, University of Tübingen, Tübingen, Germany

Correspondence

Yan Sun, College of Resources and Environment, Huazhong Agricultural University, 430070 Wuhan, China.
Email: yan.sun@mail.hzau.edu.cn

Funding information

Swiss National Science R'EQUIP, Grant/Award Number: 157884; Novartis Foundation, Grant/Award Number: #17B083; German Research Foundation, Grant/Award Number: INST 37/935-1 FUGG; Swiss National Science Foundation, Grant/Award Number: 31003A_166448, P300PA_161014 and PCEFP3_187012; Scientific Research Foundation for Returned Scholars, Huazhong Agricultural University, Grant/Award Number: 11042110026

Editor: Montserrat Vila

Abstract

Climate change may affect plant–herbivore interactions and their associated ecosystem functions. In an experimental evolution approach, we subjected replicated populations of the invasive *Ambrosia artemisiifolia* to a combination of simulated warming and herbivory by a potential biocontrol beetle. We tracked genomic and metabolomic changes across generations in field populations and assessed plant offspring phenotypes in a common environment. Using an integrated Bayesian model, we show that increased offspring biomass in response to warming arose through changes in the genetic composition of populations. In contrast, increased resistance to herbivory arose through a shift in plant metabolomic profiles without genetic changes, most likely by transgenerational induction of defences. Importantly, while increased resistance was costly at ambient temperatures, warming removed this constraint and favoured both vigorous and better defended plants under biocontrol. Climate warming may thus decrease biocontrol efficiency and promote *Ambrosia* invasion, with potentially serious economic and health consequences.

KEY WORDS

Ambrosia artemisiifolia, biological control, biological invasions, climate change, common ragweed, experimental evolution, genomics, hierarchical Bayesian model, metabolomics, *Ophraella communa*

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Ecology Letters* published by John Wiley & Sons Ltd.

INTRODUCTION

Plants have evolved a wide array of resistance traits that reduce the attack rates or development of antagonists (Kessler & Baldwin, 2001). Changes in abiotic factors such as climate can impact these plant–herbivore interactions, by selecting for particular plant phenotypes that differ in suitability for herbivores (Descombes et al., 2020). Understanding of the indirect abiotic effects on plant–herbivore interactions is essential and timely in the context of climate change (Burkpile & Parker, 2017), as shifting abiotic selection may alter crop–pest dynamics in agricultural systems, or impact the efficacy of biocontrol agents against invasive alien plants (Müller-Schärer et al., 2020). In a recent review, Sun, Ding, et al. (2020) found evidence for both positive and negative direct and plant-mediated effects of climate change on the performance of biocontrol agents on invasive plants. However, the mechanisms by which changing abiotic environments interact with plant traits to affect herbivores remain poorly understood, posing significant challenges for predicting biocontrol efficacy under future climate scenarios.

Experimental warming frequently enhances plant biomass and productivity (Dawes et al., 2015) as a result of increased soil organic matter mineralisation and nutrient availability at elevated temperatures (Rustad et al., 2001; Shaver et al., 2000), or because smaller maladapted genotypes are outcompeted (Sun, Bossdorf, et al., 2020). Warming can also modify leaf physiological traits (Hudson et al., 2011) or modulate the nutritional content and the concentrations of resistance compounds in plants (Descombes et al., 2020). These changes can then in turn affect herbivory rates (Lemoine et al., 2013; Pellissier et al., 2018) or herbivore feeding preferences (Descombes et al., 2020).

The production and maintenance of plant defensive traits is generally expected to be costly (Züst & Agrawal, 2017). One of the key hypotheses for the evolution of plant resistance is the resource availability hypothesis

(RAH, Coley et al., 1985), which posits that different resource environments select for different plant strategies along the growth–resistance spectrum. The RAH has received broad empirical support to explain interspecific patterns of plant resistance (Endara & Coley, 2011), but it is still unclear how resources influence the evolution of resistance within species. A recent review by Hahn and Maron (2016) on intraspecific variation in plant resistance found little support for the RAH.

Here, we studied the effects of warming on common ragweed (*Ambrosia artemisiifolia* L., Asteraceae, *Ambrosia* in the following), one of the most noxious plant invaders in Europe, and of its main antagonist, the accidentally introduced *Ophraella communa* LeSage (Coleoptera: Chrysomelidae, *Ophraella* in the following) (Figure 1). Where *Ambrosia* and *Ophraella* co-occur, beetles significantly impact *Ambrosia* and have been shown to reduce pollen production by more than 80% since 2013 (Bonini et al., 2015), with predicted potential reductions in health costs in Europe by more than 1 billion Euro per year (Schaffner et al., 2020). Unfortunately, species distribution models predict a faster northward spread of *Ambrosia* than for *Ophraella* under climatic change both in the introduced European and East Asian ranges, resulting in less overlap of the two species distributions and likely reduced biocontrol efficacy in the future (Sun et al., 2017, 2018). However, evolutionary responses have so far largely been ignored in species distribution models (Lavergne et al., 2010; Sun, Ding, et al., 2020), and both *Ambrosia* and *Ophraella* may evolve novel traits in response to future climatic conditions and ongoing antagonistic selection.

To incorporate rapid evolutionary responses into our predictions of future biocontrol efficacy, we performed an experimental evolution field study in which replicated populations of *Ambrosia* with identical initial genetic composition were subjected to two generations of simulated climate warming, herbivory by *Ophraella* or both treatments together, and were compared to beetle-free control populations at ambient climate. We




FIGURE 1 The plant invader and its potential biocontrol agent. (a) *Ambrosia artemisiifolia* causes high economic costs by producing highly allergenic pollen and as a crop weed; (b) all three larval stages and adults of the leaf beetle *Ophraella communa* feed on *Ambrosia*; (c) our field experimental evolution study in Northern Italy (top, 20 cages in the field; bottom-left, field warming treatment; bottom-right, field beetle treatment; for more details see Figure S1); (d) Example of a host preference test

monitored the outcomes of selection in these populations using -omics tools and with bioassays under controlled conditions. This allowed us to ask (1) whether warming and herbivory applied for two generations altered growth- or resistance-related traits in experimental *Ambrosia* populations, (2) whether changes in traits were associated with changes in metabolite production, and/or (3) in genetic composition of the studied populations, and finally through an integrative hierarchical model (4) to what extent genomics and metabolomics mediate and explain the effect of field treatments on the offspring plants' growth and resistance, and their growth–resistance correlation. We discuss our findings in the light of the potential use of *Ophraella* to control invasive *Ambrosia* populations and explore whether climate change is likely to increase or decrease the efficacy of biocontrol against *Ambrosia*.

MATERIALS AND METHODS

The North American native annual *Ambrosia artemisiifolia* has become a problematic alien invasive plant on many continents, including Asia, Oceania and Europe (Chapman et al., 2014; Essl et al., 2015). It causes great damage to human society due to its highly allergenic pollen and because it is an important and hard-to-control crop weed (Hamaoui-Laguel et al., 2015; Müller-Schärer et al., 2018) (Figure 1a). Previous studies showed high genetic variation within introduced *Ambrosia* populations in Europe (Genton et al., 2005; van Boheemen et al., 2017) due to multiple introductions and pre-admixture (Gaudeul et al., 2011), resulting in increased standing genetic variation in quantitative traits for selection to act upon, when compared to native populations (van Boheemen, Atwater, et al., 2019; van Boheemen et al., 2017). *Ambrosia* produces a range of chemical compounds that contribute to herbivore resistance (van Boheemen, Bou-Assi, et al., 2019), including sesquiterpene lactones (Tagliatalata-Scafati et al., 2012).

An important herbivore antagonist of *Ambrosia* is *Ophraella communa*, an oligophagous leaf beetle native to North America that prefers *Ambrosia* as its host plant (Augustinus, Gentili, et al., 2020; Futuyma et al., 1995) (Figure 1b). Following its accidental introductions, it is currently the most successful biocontrol agent against *Ambrosia* in China (Ma et al., 2008; Zhou et al., 2011) and also causes considerable damage to *Ambrosia* in Japan (Fukano & Doi, 2013). Müller-Schärer et al. (2014) reported a first record of *Ophraella* in Europe (Southern Switzerland and Northern Italy) in 2013 following an accidental introduction. Under favourable conditions, *Ophraella* can complete 4–7 generations per year (Augustinus, Sun, et al., 2020; Zhou et al., 2014), which allows it to build up high local densities during the second half of the *Ambrosia* growing season, leading

to complete defoliation and death of *Ambrosia* (Müller-Schärer et al., 2018; Zhou et al., 2014).

Field experimental evolution study

In April of 2016, we set up five blocks in a fenced park in Magnago, Northern Italy, with four cages ($2 \times 2 \times 2$ m) in each block (Figure 1c, Figure S1). Each experimental population was founded by planting 120 seedlings (F_1) evenly spaced in a randomised grid directly into the ground in the central 1×1.2 m of a cage. We recorded their positions so that the few dead seedlings could be replaced accordingly until May 2016, i.e. before the treatments were applied. In each cage, we transferred two seedlings grown from each of 60 maternal families that had previously been sampled from 19 invasive natural field *Ambrosia* populations (F_0) in 2013–2015 (2–4 maternal families per population; Figure S1). We expected no *Ambrosia* in the soil seed bank as *Ambrosia* had been observed only very rarely before at the experimental site. The geographic region selected for the experiment, as one of the most *Ambrosia*-infested areas of Europe and with the *Ophraella* beetle widely distributed since 2013, provides a suitable environment for our field study. We employed a factorial design, with two factors each with two levels, to evaluate the effects of climate warming and *Ophraella* herbivory on population demography and evolution. In each block, one of the experimental populations was subjected to simulated climate warming, one to *Ophraella* herbivory, one to both treatments and the remaining one served as a control (Figure 1c). Herbivory by *Ophraella* was applied by releasing 30 adults in mid-June 2016 and 2017 to each field cage. By adding or removing beetles, we maintained c. 50% visual leaf damage from August (after c. 2 beetle generations) to September (after c. 3 beetle generations). Damage levels are variable in the field, but often reach levels of 50% leaf damage by mid-season (Augustinus, Lommen, et al., 2020). The *Ophraella* beetles used in all our experiments were from a mixture of five populations collected on repeated occasions within 30 km of the field experimental site. To simulate climate warming, we constructed open-top Plexiglas chambers (1.5×1.8 m) that were placed over the plants. These increased daily mean temperature by 2.2°C , matching the predicted future change in climate obtained from www.worldclim.org for the current distribution of *Ambrosia* (Sun et al., 2017). The cages were hand weeded to maintain monocultures of *Ambrosia*, an ecological situation that is not uncommon when *Ambrosia* invades disturbed sites (Essl et al., 2015; Savić et al., 2021; Zhao et al., 2021). *Ambrosia* plants were allowed to shed seeds and regenerate naturally within each field cage. The same treatments were applied again to naturally established F_2 offspring populations in 2017 (Appendix A provides further details about the field cages and experimental treatments).

Common garden experiments for offspring phenotyping

To assess phenotypic divergence among experimental field populations after two generations of field treatments, we collected a pooled seed sample from field cage in September–October 2017. In each cage, we collected 5–10 mature F_3 seeds from each branch of each F_2 plant growing in the central 1×1.2 m area. The branch-based harvesting and subsequent mixing ensured that the proportional contribution of each genotype to follow-up experiments reflected its relative fitness on the field plots. We used the collected F_3 seeds to assess the preference and performance of *Ophraella* and the performance of *Ambrosia* (described below), as well as seed morphology and germination (see [Appendix B](#) for details) for each experimental field population.

Ophraella preference and performance

In May 2018, 50 randomly selected cold stratified F_3 seeds from each of the 20 seed pools collected in the experimental populations in 2017 were germinated and grown in a common environment at the University of Fribourg (as described in [Appendix B](#)). In the quarantine lab, we placed four similar sized (10–12 leaves) plants, one from each of the four treatments within the same field block (Figure [S1](#)) into one insect nylon netting assay cage ($24.5 \times 24.5 \times 63$ cm; BugDorm-4M2260) (Figure [1d](#)). We assembled 10 groups of 4 plants from each of the 5 field blocks, resulting in a total of 50 assay cages and placed 1 *Ophraella* pair in the centre of each assay cage. We daily recorded the position of new egg batches on the host plants and visually assessed the percentage of feeding damage of the plants every week. The experiment ended after 30 days, or after the 15th egg batch was laid, to prevent avoidance effects by egg batch accumulation on plants.

For the performance experiment, we used 10 F_3 plants from the same seed germination batch as above. Hatched larvae (L_1 instar) were isolated and transferred with a paint brush onto cut leaves (10–15 days old) from a specific *Ambrosia* population placed in moistened floral foam and into a Petri dish on a clean filter paper (\varnothing 90 mm). We used a single leaf for L_1 and L_2 instar larvae and replaced it with a new leaf from the same *Ambrosia* individual as soon as larvae reached L_3 instar to avoid food limitation. We conducted at least 20 replicate bioassays per population (each individual F_3 plant assayed using two larvae), for a total of 474 tests. We recorded survival and developmental time of each stage until emergence to adult, leaf area consumption, sex and weight and water content of emerging adults (see [Appendix B](#) for further details).

Ambrosia performance

In 2018, we grew 15–25 *Ambrosia* individuals per field cage from seeds collected in 2017, resulting in a total of 432 pots that were placed in a greenhouse at University of Fribourg (16/8h day/night cycle at 28/18°C). Regional health regulations prohibited us to let plants flower in the greenhouse. Therefore, we use biomass at harvest as a proxy for plant fitness, as *Ambrosia* biomass was found to be highly correlated with seed and pollen production in a field study across 39 sites in Europe (Lommen et al., 2017). To further characterise plant differences in functional traits, we calculated the relative water content (RWC) and specific leaf area (SLA) using two leaves of each plant (see [Appendix B](#) for further details).

Statistical analyses

We analysed the *Ophraella* preference and performance data and *Ambrosia* growth data with (generalised) linear mixed models, using the functions *glmer/lmer* function in the R package *lme4*, which uses maximum likelihood to estimate model parameters (Bates et al., 2014). The models included the warming treatment and beetle treatment in the field experiment, and their interactions, as fixed factors, with initial plant height, or days to egg batch laying or adult sex (for performance experiment only) as a covariate, and field experiment cages and cohort (for preference experiment only) as random effects (see [Appendix B](#) for details). For plotting figures, the fitted means and standard errors of fixed effects parameters were extracted using the *fixef* and *devfun2* functions with *stderr* from the R package *lme4*.

Metabolomics analyses

To assess the chemical diversity within and among generations and experimental treatments, we performed untargeted metabolomics analyses of plants growing in the field experiment in Italy. From each of the 20 caged experimental field populations, and for each generation separately, we collected paired opposite leaves from all 120 F_1 plants in June 2016, and then again from 120 randomly selected F_2 plants in each plot in June 2017 (one individual per cell in a 1×1.2 m area with 10×10 cm grid cells in the centre of each plot). We collected undamaged young leaves (8- to 10-leaf stage) in both years before *Ophraella* was released into the cages, thus our analyses reflect the constitutive levels of plant chemicals. Of each leaf pair, one was used for metabolomic analyses and the other one for a genome scan (see below). As our analyses were focused on population differences in chemical compounds, we pooled all 120 leaves from an

individual population and generation into a single composite sample.

To characterise the overall chemical diversity of each composite sample (= population \times generation), we calculated (1) the chemical richness as the number of individual mass features obtained from positive mode of MS detection, (2) the total chemical abundance as the summed abundances of individual mass features and (3) the inverse Simpson diversity based on the relative abundances of individual mass features (see [Appendix C](#) for details on sample collection, subsequent processing using an Acquity ultra-performance liquid chromatography (UPLC) system coupled to a G2-XS QTOF mass spectrometer and on data analyses).

Genomic analyses

To assess molecular diversity and differentiation among experimental generations and treatments, we collected 40 pooled samples (20 pools of F_1 populations and 20 pools of F_2 populations) analogous to the metabolomics pooled samples. Each pool contained equal amounts of tissue from individual plant (~1 mg from a cut leaf from a total of 120 individuals per experimental field population). Samples were processed for DNA sequencing on an Illumina HiSeq3000 platform at the Max Planck Institute for Developmental Biology in Tübingen (see [Appendix D](#) for details on DNA extraction, library preparation and processing of the raw data). After the pipeline processing of the sequencing data from the 40 population pools, we retained 3'317'743'837 paired-end reads (97.9%) with a median Phred score of 37. Duplicate filtering resulted in removal of 8.5%–11.2% of the mapped reads (Bieker et al., 2022), and only 2%–2.6% were considered missing data. The remaining reads corresponded to an average 30x coverage per experimental field population pool.

Statistical analyses

To understand how the field selection treatments affected the genetic structure of populations across generations, we ran redundancy analysis (RDA) to the raw chemical abundance profiles and the untransformed allele frequencies, and correspondence analysis (CA) on presence/absence data of mass features in each composite sample using *vegan* package (Oksanen et al., 2020). We used permutational multivariate analysis of variance (PERMANOVA) to test for significant differences between generations and among treatments using the function *adonis* with 9999 permutations and including cage in the strata option in the package *vegan* in R (R Core Team, 2019).

Integrating field treatment, -omics and phenotypic response data

We used an integrative hierarchical Bayesian model to infer response to field beetle and climate warming treatments from the measurements obtained through phenotyping, genotyping and metabolomics. Specifically, we modelled the genotypic and metabolomic data as linear responses to different treatments and phenotypic measurements (growth and resistance) as linear responses to the genotypic and metabolomic data (fixed effects). In our model, we jointly estimate all the fixed effects as well as cage-specific and block-specific random effects. The results are posterior estimates of (1) the effect of the treatments on genotypic and metabolomic values based on the replicated measurements derived from the five blocks in our experimental design, and (2) the effect of genotypic and metabolomic values on the phenotypes based on multiple measurements across all 20 cages. Our model thus controls for confounding effects and repeated measures. In our hierarchical model, we use hyper-priors to relax the subjectivity of choices in the prior distribution and reduce the risks of overparameterisation (Gelman et al., 2013). We sampled all parameters, including fixed and random effects and the covariance between growth and resistance, from their joint posterior distribution using a Metropolis–Hastings Markov Chain Monte Carlo (MCMC) estimator. We then summarised the posterior samples by computing the mean and 95% credible intervals of all parameters. The hierarchical Bayesian model developed here was implemented in Python (www.python.org) (see [Appendix E](#) for further details).

RESULTS

Ophraella preference and performance

In the *Ophraella* preference study ([Figure 1d](#)), we found *Ophraella* egg batch numbers ($\chi^2 = 16.74$, $p < 0.001$), total numbers of eggs ($\chi^2 = 14.06$, $p < 0.001$) and numbers of feeding larvae at the end of the experiment ($\chi^2 = 7.26$, $p = 0.007$) to differ significantly between the offspring of herbivore-damaged field plants and of herbivore-free plants ([Figure 2](#), [Table S1](#)). *Ambrosia* offspring (F_3) of plants from the beetle and warming+beetle treatments in the field experiment had fewer egg batches, a lower total number of eggs and fewer feeding *Ophraella* larvae in the preference experiments than *Ambrosia* offspring originating from the control or warming field treatments (Tukey $p \leq 0.009$; [Figure 2](#), [Table S1](#)). Other preference measurements such as the numbers of eggs per batch, the hatching rates of *Ophraella* beetles or the final damage of *Ambrosia* offspring were not affected by herbivory in the experimental field population ($\chi^2 \leq 1.8$, $p \geq 0.18$;




FIGURE 2 Preference and performance of *Ophraella* beetles on (upper four panels) and plant phenotypic performance traits of (lower four panels) *Ambrosia* offspring (F_2) from the four field selection treatments when compared in a common environment. Blue and red points represent ambient and warming, and with and without black circles indicating with and without beetle field selection treatments; the small transparent points give the mean of each population. Treatment means \pm SE, indicated by larger points, and population means are extracted from (generalised) linear mixed models (see M&M for details). The error bars are \pm SE

for details see Table S1). We also found no effects of the field warming treatment, or its interaction with the beetle treatment, on any of the preference measurements ($\chi^2 \leq 2.15$, $p \geq 0.14$; Table S1).

In the *Ophraella* performance experiment, the treatments in the experimental field populations significantly influenced the mortality ($\chi^2 = 42.55$, $p < 0.001$) and water content of adult beetles ($\chi^2 = 4.12$, $p = 0.04$; Table S1) feeding on the offspring plants, with more than double the mortality in beetles feeding on offspring of plants from cages exposed to herbivory compared to cages not exposed to herbivory. For all other *Ophraella* performance traits, there were no significant differences between plants from different beetle or warming treatments, and no interaction between the two treatments ($\chi^2 \leq 3.33$, $p \geq 0.07$; Table S1).

Ambrosia performance

The germination rate, dry weight, seed coat ratio and seedling length of F_3 seeds did not differ among any of the field selection treatments ($\chi^2 \leq 3.88$, $p \geq 0.27$; Figure S2).

Comparison of *Ambrosia* F_3 offspring in the standardised greenhouse environment showed that both warming and beetle herbivory in previous generations affected offspring phenology and reproductive output ($\chi^2 \geq 3.67$, $p \leq 0.05$; Table S1), with later flowering and more inflorescences in offspring from warming and/or beetle treatments (Figure 2). Moreover, offspring plants that experienced warming generally had a larger biomass and lower soil moisture content ($\chi^2 \geq 14.56$, $p < 0.001$; Figure 2). There were also significant interactions between warming and beetle treatments for plant biomass and relative water content ($\chi^2 \geq 3.81$, $p \leq 0.05$; Table S1). Under ambient temperature conditions, previous exposure to *Ophraella* beetles decreased the average biomass of *Ambrosia* plants, whereas under warming conditions, it increased it. For all other phenotypic traits, there were no significant differences among the offspring from different experimental treatments ($\chi^2 \leq 2.08$, $p \geq 0.15$; see details in Table S1).

Metabolomics

We detected a total of 6043 metabolomic features across all plants using mass spectrometry with positive ionisation. For both richness and diversity of the metabolomic features, we found no significant differences among treatments and generations ($p \geq 0.21$). However, metabolomic diversity was associated with several plant resistance traits. We found negative correlations between the inverse Simpson's chemical diversity of populations and the number of *Ophraella* egg batches and larvae observed on offspring from these populations (Table 1).

There were also negative correlations between the chemical richness of plant populations and total numbers of eggs and larvae, as well as the per cent damage observed in the *Ophraella* preference experiment (Table 1).

A RDA based on all identified metabolomic features showed that there was no clustering by treatments in the parental generation, but that the two generations separated strongly along the first RDA axis ($F = 12.06$, $p < 0.001$), and that there was a strong separation of beetle and non-beetle *Ambrosia* offspring along the second axes ($F = 2.43$, $p = 0.01$), resulting in a significant generation \times beetle interaction ($F = 1.87$, $p = 0.04$; Figure 3a). In contrast, there was no significant RDA differentiation of *Ambrosia* offspring by the temperature treatments ($F = 0.83$, $p = 0.58$), i.e. the metabolomic profiles of warming and non-warming plants were generally very similar in both generations ($F = 0.73$, $p = 0.73$), and with or without beetles ($F = 0.60$, $p = 0.91$; Figure 3a, Table S2). Moreover, we found negative correlations between the first ordination axis of RDA (RDA1) and total *Ophraella* mortality as well as the numbers of *Ambrosia* inflorescences, while RDA1 was strongly positively correlated with the number of *Ophraella* egg batches and larvae, and CA1 with the number of egg batches (Table 1).

When comparing the metabolite profiles of F_2 plants from the beetle versus the non-beetle treatments, we found 695 up-regulated and 175 down-regulated features in positive mode (Figure S3). An automated database comparison of these differential features putatively identified four sesquiterpene lactones. Four of these responded positively to the beetle treatment, and one negatively (Table S3). Since for each of these significant features there were several additional adducts, isotopes and fragments that were recorded as separate features, the four compounds were in fact responsible for approximately 30% of all 870 significant mass features. We manually identified an additional 12 sesquiterpene lactones (Table S3), which were not significantly different between beetle and control treatments, but together with the significant 4, they constituted all major peaks in the chromatograms of *Ambrosia* (Figure S4). Despite the lack of individual significant effects for several of the identified sesquiterpene lactones, all compounds showed remarkably synchronised and similar responses to the two beetle treatments, with approximately half of the compounds increasing and the other half decreasing their relative expression in response to these treatments, resulting in very different sesquiterpene lactone profiles for the experimental treatments (Figure S5).

Genomics

The RDA visualising genetic dissimilarities among pooled population samples showed a strong genetic separation between the parental and offspring generation, with little treatment differences in the parental generation, but




FIGURE 3 Redundancy analysis (RDA) of the results of untargeted metabolomics analysis (a: showing RDA of metabolomics dissimilarity, using the chemical abundance of individual mass features) and of the genomic analyses (b: showing RDA of genetic dissimilarity, using allele frequencies of all polymorphic SNP markers) using leaves of F_1 and F_2 *Ambrosia* plants in the field experiment. Each symbol represents one experimental population (cage), and the ellipses are 95% SE around the group means. For details on the statistical tests, see Table S2

a clear separation of F_2 plants from warming (warming and warming+beetle treatment) versus ambient (control and beetle treatments) conditions (Figure 3b). In contrast, there were no systematic genetic differences related to the beetle treatments (Figure 3b). These results were verified by PERMANOVA (Table S4), with significant effects of generation and a significant interaction between generation and warming ($p \leq 0.04$; Table S4), but no significant main effect of the beetle treatment or beetles \times generation interaction ($p \geq 0.3$, Table S4).

Integrating components of the field selection treatments, multi-omics and phenotypic responses

Our hierarchical Bayesian model (Figure 4a) showed high linear correlations between the observed and posterior predicted plant genomic, metabolomic and phenotypic means ($\text{RMSE} \geq 0.19$, $p < 0.001$; Figure S6), indicating that the model adequately captures their variation. Our analyses confirmed and strengthened the above results, revealing a strong effect of the field beetle treatment on metabolomic changes, and of the field warming treatment and its interaction with the beetle treatment on plant genetic changes (Figure 4b and c). We also found strong effects of both the genetic and metabolomic composition on *Ambrosia* offspring growth and a strong effect of the metabolomics on the *Ambrosia* offspring resistance (Figure 4d and e). Most interestingly, our model showed a significant difference in the *Ambrosia* offspring growth–resistance relationship between two field

warming treatments (Welch's t -test: $p < 0.001$; Figure 5), i.e. a negative correlation resulting from the field ambient treatment, but not so when selected through the field warming treatment, which showed both vigorous and well-defended plant genotypes. Importantly, our hierarchical modelling approach clearly provided evidence that the field treatment effects, i.e. climate warming and biocontrol herbivory, are differently mediated by the *Ambrosia* phenotypes, either mainly through genomic or transient metabolomic changes respectively.

DISCUSSION

Invasive plants may evolve rapidly to a changing biotic and abiotic environment. By hierarchically analysing all data of the genetic and metabolomic structures across generations from the field experiment, and of the phenotypic data from offspring plants in the common garden in a unified statistical framework (Figure 4a), we could use the covariance between estimated mean growth and resistance to show that warming of the climate selects for plants that show both faster growth and higher resistance.

More specifically, we found that exposure to beetles selected for increased herbivore resistance, independent of the warming treatment. The increased resistance was correlated with phenotypic shifts in secondary metabolites and sesquiterpene lactone profiles, but not with changes in the overall genetic composition of plant populations. Even though we cannot rule out minor changes in genotype or allele frequencies, our results indicate




FIGURE 4 (a) Outcomes of a hierarchical Bayesian model applied to our experiment, integrating the field selection treatments, multi-omics and phenotypic responses under controlled environmental conditions, with the effect size of each parameter (see M&M) and their posterior probability $P(E > 0)$. Lower p -values indicate stronger negative effects and higher p -values stronger positive effects; solid lines represent strong and dotted lines weak effects. (b) The posterior probability distribution for the relative importance of field beetle and warming treatments on genotypic changes and (c) on metabolomic changes; and (d) the relative importance of genomes and (e) metabolomes on growth and resistance. (b)-(e) refer to the lines in (a)




FIGURE 5 (a) Relationships between growth and resistance of *Ambrosia artemisiifolia* offspring from different field selection treatments in our common environment experiment. The small transparent points give the mean of each population, and the larger points the treatment means (corresponding to Figure 2). Error bars are \pm SE. The arrows indicate the direction of changes in response to herbivory, with a shift towards increased resistance at the cost of reduced growth under ambient conditions, but a shift towards increased resistance and increased growth under warmer conditions. (b) Significant difference in the relationship between the estimated mean growth and resistance in the response to climate treatments (ambient vs. warming) based on the hierarchical Bayesian model ($t = -27.09, p < 0.001$)

that the observed increased resistance primarily resulted from transgenerational induction. The increased resistance was associated with lower biomass of common garden plants, indicating costs of resistance, for plants exposed to beetles under field ambient temperatures, but not under field elevated temperatures (Figure 5a). Warming generally selected for more vigorous plants, and unlike the beetle treatment it was also associated with strong changes in the genetic composition of plant populations. Surprisingly, while offspring plants from the field warming+beetle treatment were as resistant to *Ophraella* herbivory as plants from beetle selection alone, they did not experience the same biomass reduction as the latter (Figure 5a). Note, however, that this pattern was largely driven by a few outlier populations. Nonetheless, our integrated modelling approach confirmed and strengthened the surprising result of a differential effect of the two field temperature treatments on the growth–resistance relationship (Figure 5b). Such a difference suggests that the warming-induced selection may purge *Ambrosia* genotypes with growth–resistance trade-offs and result in populations that both grow and defend well, and continued selection may strengthen these effects.

Genetic and metabolomic differentiation between generations and selection treatments

The starting populations of our experiment harboured substantial standing genetic variation. In a companion study, we showed the maternal families used here differed significantly in all plant phenotypic traits measured (Sun, Bossdorf, et al., 2020), indicating that there was plenty of variation for selection to act upon (Figure S7). Interestingly, our pool-seq results indicate strong genetic changes under abiotic (warming) selection, but not under biotic (herbivore) selection. The results for abiotic selection are in line with previous studies demonstrating trait differentiation of *Ambrosia* populations in response to climate change (van Boheemen, Atwater, et al., 2019; Sun, Bossdorf, et al., 2020). The lack of genetic response to biotic selection on the other hand is surprising, as similar herbivore pressure has resulted in rapid genetic divergence over three to five generations in other experimental evolution studies (Agrawal et al., 2012; Züst et al., 2012). Our metabolomic analysis was targeted towards the detection of small organic molecules of intermediate polarity ('secondary' or specialised plant metabolites), which we found to mainly consist of sesquiterpene lactones in *Ambrosia*.

Sesquiterpene lactones are biologically active compounds that are often involved in plant resistance. Rather than altering the absolute amounts of sesquiterpene lactones, biotic selection induced a change in the sesquiterpene lactone profiles of experimental field populations, while abiotic (warming) selection had no effect

on the expression of these compounds. Even though we do not yet understand sesquiterpene lactone synthesis in *Ambrosia*, our correlative results suggest that not all compounds are equal in this plant, but rather that only a subset of compounds provide effective resistance against *Ophraella*. This subset appears to be costly to produce, and to trade-off with a set of less effective, but also less costly sesquiterpene lactone compounds. Surprisingly, field warming selection did not favour genotypes producing these low-cost, low-efficiency compounds, contrary to what the RAH would have predicted. Instead, expected higher resource availability under warming appeared to ameliorate costs of production for efficient compounds, providing further evidence of the limited use of the RAH for intraspecific predictions.

Transgenerationally induced resistance?

In response to biotic selection, we observed a strong shift in chemical profiles, but no parallel shift in genetic differentiation. A possible explanation is that the chemical changes were induced in the parents and carried over to the next generation, i.e. they represent a case of transgenerational plasticity. Many previous studies on various plant species have demonstrated such transgenerational effects for plant resistance to herbivores or pathogens (Agrawal, 2002; Colicchio, 2017), but also for responses to abiotic stress (Bonduriansky et al., 2012). Transgenerational effects can be mediated by epigenetic processes (Rasmann et al., 2012), maternal nutrient provisioning and seed modifications (Galloway, 2001). In our study, we found no significant reduction in seed size, but substantial changes in the metabolomic composition, especially of several sesquiterpene lactones, in *Ambrosia* offspring from the two beetle selection treatments as compared to the non-beetle treatments (Figures S2 and S5). The dissimilarity in secondary compounds between offspring from beetle and non-beetle treatments might have been the cause of the 30% fewer eggs laid and 50% increased mortality of *Ophraella* on plants from beetle treatments. Although the precise mechanisms underlying transgenerational plant resistance are not well understood, previous studies suggested that small RNA, heritable DNA methylation changes and/or histone modifications could be involved (Gutzat & Scheid, 2012; Rasmann et al., 2012).

CONCLUDING REMARKS

Our experimental evolution study predicts a decreased biocontrol efficiency and an increased *Ambrosia* invasion ability under climate warming, with the plant invader thus gaining an advantage under climate change. Understanding the evolutionary responses of plants to biotic interactions such as herbivore pressure under

warming conditions is becoming increasingly important and urgent in the light of the rapidly changing climate (Sun, Ding, et al., 2020). Invasive alien plants with their biocontrol organisms offer excellent opportunities for exploring growth–resistance strategies of plants, and their rapid evolution and adaptation, under novel abiotic (climate) and biotic (escape of specialist herbivores) conditions. Rapid evolutionary changes in invasive alien plants post-introduction have been well documented, but there are yet few experimental tests of evolutionary responses to biocontrol organisms when invasive species become re-associated with their key natural enemies (Müller-Schärer et al., 2020). The results of our experimental evolution study provide a step towards a better understanding of the potential of a biocontrol agent to influence plant evolution, and how this interaction may be modulated by indirect effects of climate warming on plant traits that feed back on plant suitability for the herbivores.

The beetle *Ophraella* can strongly reduce the reproduction and population growth rate of *Ambrosia* in Europe (Augustinus, Lommen, et al., 2020; Augustinus, Sun, et al., 2020; Schaffner et al., 2020). However, based on our findings, we predict a reduced biocontrol efficacy under climate warming, highlighting the importance of accounting for the interplay between ecological and evolutionary processes in predicting biotic interactions in changing environments. We acknowledge that post-release adaptation of the biocontrol agent may further affect growth–resistance relations in the target plant and thereby biocontrol efficacy, but plants too may evolve further in the longer term in response to the herbivores (Sun, Ding, et al., 2020; Wright & Bennett, 2018). It would now be important to establish whether our results are valid also for other plant species, such as rare and endangered species, but especially for other invasive alien plants that harbour large standing genetic variation for selection to act upon (Dlugosch et al., 2015).

ACKNOWLEDGEMENTS

YS was supported by an Advanced Postdoc. Mobility fellowship from the Swiss National Science Foundation (SNSF project P300PA_161014), with additional support from the Novartis Foundation (#17B083 to HMS and YS). YS acknowledges funding by Scientific Research Foundation for Returned Scholars, Huazhong Agricultural University (11042110026 to YS). HMS acknowledges funding through the Swiss National Science Foundation (SNSF project 31003A_166448). ME acknowledges Swiss National Science R'EQUIP grant 157884 that provided funding for the mass spectrometer. YS and OB acknowledge support by Maximilian Hanussek from the High Performance and Cloud Computing Group at the Zentrum für Datenverarbeitung of the University of Tübingen, the state of Baden-Württemberg through bwHPC and the German Research Foundation (DFG) through grant no INST 37/935-1 FUGG. DS received funding from the Swiss National Science Foundation (PCEFP3_187012) and from

the Swedish Research Council (VR: 2019-04739). We are grateful to the Comune di Magnago for providing the experimental field site and to the many people who helped with the field and common garden studies, especially to Carine Beuchat and Noemi Verdicchia, and the local gardener Giuseppe Giambrone. The DNA sequencing was supported by Detlef Weigel, Gautam Shirsekar, Julia Hildebrandt and Ilja Bezrukov (all MPI for Developmental Biology Tübingen), and the pool-seq analyses by Martin Kapun and Daniel Wegmann. We also thank Kay Hodgins and Michael Martin for providing the draft genome of *Ambrosia artemisiifolia*. Open Access Funding provided by Université de Fribourg. [Correction added on 23 May 2022, after first online publication: CSAL funding statement has been added.]

AUTHOR CONTRIBUTION

YS, HMS and OB designed the study, YS and HMS conducted the field experimental evolutionary study, YS conducted the common garden experiments, YS performed all genomic and phenotypic statistical analyses, PM and CR measured metabolomic components, TZ and ME performed metabolomic analyses, DS developed the hierarchical Bayesian model, YS wrote the first draft of the manuscript, YS, HMS, TZ, OB, DS and ME contributed substantially to the interpretation and the final version.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/ele.14000>.

OPEN RESEARCH BADGES



This article has earned Open Data and Open Materials Design badges. Data and materials design and analysis plan are available at: <http://doi.org/10.6084/m9.figshare.14199593>.

DATA AVAILABILITY STATEMENT

Data and hierarchical Bayesian model associated with the manuscript are available as supplementary materials and are deposited in Figshare digital repository (<http://doi.org/10.6084/m9.figshare.14199593>).

ORCID

Yan Sun <https://orcid.org/0000-0002-6439-266X>
 Matthias Erb <https://orcid.org/0000-0002-4446-9834>
 Oliver Bossdorf <https://orcid.org/0000-0001-7504-6511>
 Heinz Müller-Schärer <https://orcid.org/0000-0003-0936-1470>

REFERENCES

Agrawal, A.A. (2002) Herbivory and maternal effects: mechanisms and consequences of transgenerational induced plant resistance. *Ecology*, 83, 3408–3415.

- Agrawal, A.A., Hastings, A.P., Johnson, M.T.J., Maron, J.L. & Salminen, J.P. (2012) Insect herbivores drive real-time ecological and evolutionary change in plant populations. *Science*, 338, 113–116.
- Augustinus, B.A., Gentili, R., Horvath, D., Naderi, R., Sun, Y., Tournet, A.-M. et al. (2020) Assessing the risks of non-target feeding by the accidentally introduced ragweed leaf beetle, *Ophraella communa*, to native European plant species. *Biological Control*, 150, 104356.
- Augustinus, B.A., Lommen, S.T.E., Fogliatto, S., Vidotto, F., Smith, T., Horvath, D. et al. (2020) In-season leaf damage by a biocontrol agent explains reproductive output of an invasive plant species. *NeoBiota*, 55, 117–146.
- Augustinus, B., Sun, Y., Beuchat, C., Schaffner, U. & Müller-Schärer, H. (2020) Predicting impact of a biocontrol agent: Integrating distribution modelling with climate-dependent vital rates. *Ecological Applications*, 30, e02003.
- Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H. et al. (2014) *Package 'lme4'*. Vienna: R foundation for statistical computing, p. 12.
- Bieker, V.C., Battlay, P., Petersen, B., Sun, X., Wilson, J., Brealey, J.C., et al. (2022). Uncovering the hologenomic basis of an extraordinary plant invasion. *bioRxiv*, 2022.2002.2003.478494.
- Bonduriansky, R., Crean, P., Day, T. (2012) The implications of non-genetic inheritance for evolution in changing environments. *Evolutionary Applications*, 5, 192–201.
- Bonini, M., Šikoparija, B., Prentović, M., Cislighi, G., Colombo, P., Testoni, C. et al. (2015) A follow-up study examining airborne *Ambrosia* pollen in the Milan area in 2014 in relation to the accidental introduction of the ragweed leaf beetle *Ophraella communa*. *Aerobiologia*, 32, 371–374.
- Burkepile, D.E. & Parker, J.D. (2017) Recent advances in plant-herbivore interactions. *F1000Research*, 6, 119.
- Chapman, D.S., Haynes, T., Beal, S., Essl, F. & Bullock, J.M. (2014) Phenology predicts the native and invasive range limits of common ragweed. *Global Change Biology*, 20, 192–202.
- Coley, P.D., Bryant, J.P. & Chapin Iii, F.S. (1985) Resource availability and plant antiherbivore defense. *Science*, 230(4728), 895–899.
- Colicchio, J. (2017) Transgenerational effects alter plant defence and resistance in nature. *Journal of Evolutionary Biology*, 30, 664–680.
- Dawes, M.A., Philipson, C.D., Fonti, P., Bebi, P., Hättenschwiler, S., Hagedorn, F. et al. (2015) Soil warming and CO₂ enrichment induce biomass shifts in alpine tree line vegetation. *Global Change Biology*, 21, 2005–2021.
- Descombes, P., Kergunteuil, A., Glauser, G., Rasmann, S. & Pellissier, L. (2020) Plant physical and chemical traits associated with herbivory in situ and under a warming treatment. *Journal of Ecology*, 108, 733–749.
- Dlugosch, K.M., Anderson, S.R., Braasch, J., Cang, F.A. & Gillette, H.D. (2015) The devil is in the details: genetic variation in introduced populations and its contributions to invasion. *Molecular Ecology*, 24, 2095–2111.
- Endara, M.J. & Coley, P.D. (2011) The resource availability hypothesis revisited: a meta-analysis. *Functional Ecology*, 25, 389–398.
- Essl, F., Biró, K., Brandes, D., Broennimann, O., Bullock, J.M., Chapman, D.S. et al. (2015) Biological Flora of the British Isles: *Ambrosia artemisiifolia*. *Journal of Ecology*, 103, 1069–1098.
- Fukano, Y. & Doi, H. (2013) Population abundance and host use pattern of *Ophraella communa* (Coleoptera: Chrysomelidae) in its native and introduced range. *Biocontrol Science and Technology*, 23, 595–601.
- Futuyma, D.J., Keese, M.C. & Funk, D.J. (1995) Genetic constraints on macroevolution: the evolution of host affiliation in the leaf beetle genus *Ophraella*. *Evolution*, 49, 797–809.
- Galloway, L.F. (2001) The effect of maternal and paternal environments on seed characters in the herbaceous plant *Campanula americana* (Campanulaceae). *American Journal of Botany*, 88, 832–840.
- Gaudeul, M., Giraud, T., Kiss, L. & Shykoff, J.A. (2011) Nuclear and chloroplast microsatellites show multiple introductions in the worldwide invasion history of common ragweed, *Ambrosia artemisiifolia*. *PLoS One*, 6, e17658.
- Gelman, A., Carlin, J., Stern, H., Dunson, D., Vehtari, A. & Rubin, B. (2013) *Bayesian data analysis*, 3rd edn. London: Chapman & Hall/CRC Texts in Statistical Science.
- Genton, B.J., Shykoff, J.A. & Giraud, T. (2005) High genetic diversity in French invasive populations of common ragweed, *Ambrosia artemisiifolia*, as a result of multiple sources of introduction. *Molecular Ecology*, 14, 4275–4285.
- Gutzat, R. & Scheid, O.M. (2012) Epigenetic responses to stress: triple defense? *Current Opinion in Plant Biology*, 15, 568–573.
- Hahn, P.G. & Maron, J.L. (2016) A framework for predicting intraspecific variation in plant defense. *Trends in Ecology & Evolution*, 31, 646–656.
- Hamaoui-Laguel, L., Vautard, R., Liu, L.I., Solmon, F., Viovy, N., Khvorostyanov, D. et al. (2015) Effects of climate change and seed dispersal on airborne ragweed pollen loads in Europe. *Nature Climate Change*, 5, 766–771.
- Hudson, J., Henry, G. & Cornwell, W. (2011) Taller and larger: shifts in Arctic tundra leaf traits after 16 years of experimental warming. *Global Change Biology*, 17, 1013–1021.
- Kessler, A. & Baldwin, I.T. (2001) Defensive function of herbivore-induced plant volatile emissions in nature. *Science*, 291, 2141–2144.
- Lavergne, S., Mouquet, N., Thuiller, W. & Ronce, O. (2010) Biodiversity and climate change: integrating evolutionary and ecological responses of species and communities. *Annual Review of Ecology, Evolution, and Systematics*, 41, 321–350.
- Lemoine, N.P., Drews, W.A., Burkepile, D.E. & Parker, J.D. (2013) Increased temperature alters feeding behavior of a generalist herbivore. *Oikos*, 122, 1669–1678.
- Lommen, S.T.E., Hallmann, C.A., Jongejans, E., Chauvel, B., Leitsch-Vitalos, M., Aleksanyan, A. et al. (2017) Explaining variability in the production of seed and allergenic pollen by invasive *Ambrosia artemisiifolia* across Europe. *Biological Invasions*, 20, 1475–1491.
- Ma, J., Guo, J., Wan, F., Hu, X., Wan, F., Li, B. et al. (2008) Biological control of *Ambrosia artemisiifolia* and *A. trifida*. In: Wan, F., Li, B. & Guo, J. (Eds.) *Biological invasions: biological control theory and practice*. China: Science Press Beijing, pp. 157–185.
- Müller-Schärer, H., Bouchemousse, S., Litto, M., McEvoy, P.B., Roderick, G.K. & Sun, Y. (2020) How to better predict long-term benefits and risks in weed biocontrol: an evolutionary perspective. *Current Opinion in Insect Science*, 38, 84–91.
- Müller-Schärer, H., Lommen, S.T.E., Rossinelli, M., Bonini, M., Boriani, M., Bosio, G. et al. (2014) *Ophraella communa*, the ragweed leaf beetle, has successfully landed in Europe: fortunate coincidence or threat? *Weed Research*, 54, 109–119.
- Müller-Schärer, H., Sun, Y., Chauvel, B., Karrer, G., Kazinczi, G., Kudsk, P. et al. (2018) Cross-fertilizing weed science and plant invasion science to improve efficient management: a European challenge. *Basic and Applied Ecology*, 33, 1–13.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGinn, D. et al. (2020) *Vegan: Community ecology package*. R package version, 2. 5-7. <https://CRAN.R-project.org/package=vegan>
- Pellissier, L., Descombes, P., Hagen, O., Chalmandrier, L., Glauser, G., Kergunteuil, A. et al. (2018) Growth-competition-herbivore resistance trade-offs and the responses of alpine plant communities to climate change. *Functional Ecology*, 32, 1693–1703.
- R Core Team. (2019) *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rasmann, S., De Vos, M., Casteel, C.L., Tian, D., Halitschke, R., Sun, J.Y. et al. (2012) Herbivory in the previous generation primes plants for enhanced insect resistance. *Plant Physiology*, 158, 854–863.
- Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A. et al. (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia*, 126, 543–562.

- Savić, A., Oveisi, M., Božić, D., Pavlović, D., Saulić, M., Schärer, H.M. et al. (2021) Competition between *Ambrosia artemisiifolia* and *Ambrosia trifida*: is there a threat of a stronger competitor? *Weed Research*, 61(4):298–306.
- Schaffner, U., Steinbach, S., Sun, Y., Skjøth, C.A., de Weger, L.A., Lommen, S.T. et al. (2020) Biological weed control to relieve millions from *Ambrosia* allergies in Europe. *Nature Communications*, 11, 1745.
- Shaver, G.R., Canadell, J., Chapin, F.S., Gurevitch, J., Harte, J., Henry, G. et al. (2000) Global Warming and Terrestrial Ecosystems: A Conceptual Framework for Analysis: Ecosystem responses to global warming will be complex and varied. Ecosystem warming experiments hold great potential for providing insights on ways terrestrial ecosystems will respond to upcoming decades of climate change. Documentation of initial conditions provides the context for understanding and predicting ecosystem responses. *BioScience*, 50, 871–882.
- Sun, Y., Bossdorf, O., Grados, R.D., Liao, Z. & Müller-Schärer, H. (2020) Rapid genomic and phenotypic change in response to climate warming in a widespread plant invader. *Global Change Biology*, 26, 6511–6522.
- Sun, Y., Brönnimann, O., Roderick, G.K., Poltavsky, A., Lommen, S.T. & Müller-Schärer, H. (2017) Climatic suitability ranking of biological control candidates: a biogeographic approach for ragweed management in Europe. *Ecosphere*, 8, e01731.
- Sun, Y., Ding, J., Siemann, E. & Keller, S.R. (2020) Biological control of invasive weeds under climate change: progress, challenges and management implications. *Current Opinion in Insect Science*, 38, 72–78.
- Sun, Y., Zhou, Z., Wang, R. & Müller-Schärer, H. (2018) Biological control opportunities of ragweed are predicted to decrease with climate change in East Asia. *Biodiversity Science*, 25, 1285–1284.
- Tagliatalata-Scafati, O., Pollastro, F., Minassi, A., Chianese, G., De Petrocellis, L., Di Marzo, V. et al. (2012) Sesquiterpenoids from common ragweed (*Ambrosia artemisiifolia* L.), an invasive biological pollutant. *European Journal of Organic Chemistry*, 2012, 5162–5170.
- van Boheemen, L.A., Atwater, D.Z. & Hodgins, K.A. (2019) Rapid and repeated local adaptation to climate in an invasive plant. *New Phytologist*, 222, 614–627.
- van Boheemen, L.A., Bou-Assi, S., Uesugi, A. & Hodgins, K.A. (2019) Rapid growth and defence evolution following multiple introductions. *Ecology and Evolution*, 9, 7942–7956.
- van Boheemen, L.A., Lombaert, E., Nurkowski, K.A., Gauffre, B., Rieseberg, L.H. & Hodgins, K.A. (2017) Multiple introductions, admixture and bridgehead invasion characterize the introduction history of *Ambrosia artemisiifolia* in Europe and Australia. *Molecular Ecology*, 26, 5421–5434.
- Wright, M.G. & Bennett, G.M. (2018) Evolution of biological control agents following introduction to new environments. *BioControl*, 63, 105–116.
- Zhao, W., Liu, T., Liu, Y., Wang, H., Wang, R., Ma, Q. et al. (2021) The significance of biomass allocation to population growth of the invasive species *Ambrosia artemisiifolia* and *Ambrosia trifida* with different densities. *BMC Ecology and Evolution*, 21, 1–13.
- Zhou, Z.S., Chen, H.S., Zheng, X.W., Guo, J.Y., Guo, W., Li, M. et al. (2014) Control of the invasive weed *Ambrosia artemisiifolia* with *Ophraella communa* and *Epiblema strenuana*. *Biocontrol Science and Technology*, 24, 950–964.
- Zhou, Z.S., Guo, J.Y., Zheng, X.W., Luo, M., Chen, H.-S. & Wan, F.H. (2011) Reevaluation of biosecurity of *Ophraella communa* against sunflower (*Helianthus annuus*). *Biocontrol Science and Technology*, 21, 1147–1160.
- Züst, T. & Agrawal, A.A. (2017) Trade-offs between plant growth and defense against insect herbivory: an emerging mechanistic synthesis. *Annual Review of Plant Biology*, 68, 513–534.
- Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D.J. & Turnbull, L.A. (2012) Natural enemies drive geographic variation in plant defenses. *Science*, 338, 116–119.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Sun, Y., Züst, T., Silvestro, D., Erb, M., Bossdorf, O., Mateo, P., et al. (2022) Climate warming can reduce biocontrol efficacy and promote plant invasion due to both genetic and transient metabolomic changes. *Ecology Letters*, 25, 1387–1400. Available from: <https://doi.org/10.1111/ele.14000>