

# Properties of the Low Threshold Ca Current in Single Frog Atrial Cardiomyocytes

## *A Comparison with the High Threshold Ca Current*

JULIO L. ALVAREZ and GUY VASSORT

From the Laboratorio de Electrofisiologia, Instituto de Cardiología y Cirugía Cardiovascular, La Habana, Cuba 10600; and Unité de Recherches de Physiologie Cellulaire Cardiaque, Institut National de la Santé et de la Recherche Médicale U-241, Université Paris-Sud, Orsay, France

**ABSTRACT** The properties of the low threshold Ca current ( $I_{CaT}$ ) in bullfrog (*Rana catesbeiana*) isolated atrial cardiomyocytes were studied using the whole-cell recording patch-clamp technique and compared with those of the high threshold Ca current ( $I_{CaL}$ ). In 91% of atrial cells we observed both  $I_{CaT}$  and  $I_{CaL}$  when collagenase and trypsin were used to dissociate the cells. But when pronase was used, only 30% of the cells exhibited  $I_{CaT}$ .  $I_{CaT}$  was never found in ventricular cells.  $I_{CaT}$  could be investigated more easily when  $I_{CaL}$  was inhibited by Cd ions (50  $\mu$ M). Its kinetics were unchanged by substituting Ba for Ca, or in the presence of high concentrations of Ba. Both  $I_{CaT}$  and  $I_{CaL}$  exhibited reduced inactivation after high depolarizing prepulses.  $I_{CaT}$  was found to be sensitive to dihydropyridines: 1  $\mu$ M nifedipine decreased this current while 1  $\mu$ M BAY K 8644 increased it; this occurred without significant variations in the steady-state inactivation curve.  $I_{CaT}$  was more sensitive than  $I_{CaL}$  to  $\alpha_1$ -adrenergic and  $P_2$ -purinergic stimulations, while  $I_{CaL}$  was more sensitive to  $\beta$ -adrenergic stimulation. Isoproterenol was still able to increase  $I_{CaT}$  in the presence of high intracellular cAMP. Both currents were increased by 1  $\mu$ M ouabain (although  $I_{CaL}$  only transiently) and decreased by 10  $\mu$ M ouabain. It is concluded that the two types of Ca channels can be observed in bullfrog atrial cells and that they are specifically altered by pharmacological agents and neuromediators. This may have implications for cardiac behavior.

### INTRODUCTION

Calcium entry through voltage-gated Ca channels is essential for several cellular functions including excitability, contraction, secretion, transmitter release, metabolic processes, and regulation of Ca channel kinetics themselves. In cardiac cells, two types of Ca channels have been identified: the predominant type activates around

Address reprint requests to Dr. G. Vassort, Physiologie Cellulaire Cardiaque, INSERM U-241, Université Paris-Sud, Bât. 443, F-91405 Orsay Cedex, France.

-30 mV, inactivates slowly when Ba is the charge carrier, and has a large unitary conductance (Bean, 1985*b*; Nilius, Hess, Lansman, and Tsien, 1985; Hagiwara, Irisawa, and Kameyama, 1988). The current through these channels is commonly termed L-type Ca current ( $I_{CaL}$ ) and corresponds to the slow inward current described many years ago by Rougier, Vassort, Garnier, Gargouil, and Coraboeuf (1969).  $I_{CaL}$  is more depressed by Cd than by Ni (Nilius et al., 1985; Mitra and Morad, 1986; Bonvallet, 1987; Bonvallet and Rougier, 1989; Hirano, Fozzard, and January, 1989*a*; but see Hagiwara et al., 1988). Its activity is modulated by  $\alpha_1$ -adrenergic (Alvarez, Mongo, and Vassort, 1987),  $\beta$ -adrenergic (Brum, Osterreider, and Trautwein, 1984), and  $P_2$ -purinergic (Alvarez, Mongo, Scamps, and Vassort, 1990; Scamps, Legssyer, Mayoux, and Vassort, 1990) stimulation.  $I_{CaL}$  is sensitive to dihydropyridines (Bean, 1985*b*; Nilius et al., 1985; Mitra and Morad, 1986; Hagiwara et al., 1988; Bonvallet and Rougier, 1989; Hirano et al., 1989*a*).

The other cardiac Ca channel ( $I_{CaT}$ ) has a low activation threshold, inactivates quickly even when Ba is the charge carrier, and has a smaller unitary conductance (Bean, 1985*b*; Nilius et al., 1985; Hagiwara et al., 1988; Droogmans and Nilius, 1989). In contrast to  $I_{CaL}$ , this current is more depressed by Ni than by Cd (Nilius et al., 1985; Mitra and Morad, 1986; Bonvallet, 1987; Bonvallet and Rougier, 1989; Hirano et al., 1989*a*) and is reported to be insensitive to dihydropyridines (Bean, 1985*b*; Nilius et al., 1985; Hagiwara et al., 1988; Bonvallet and Rougier, 1989; Hirano et al., 1989*a*; Bois and Lenfant, 1991; Charnet, Richard, Gurney, Ouadid, Tiaho, and Nargeot, 1991; but see Tseng and Boyden, 1989, and Akaike, Kostyuk, and Osipchuk, 1989).  $I_{CaT}$  is also reported to be insensitive to neurotransmitters (Bean, 1985*b*; Hagiwara et al., 1988; Tytgat, Nilius, Vereecke, and Carmeliet, 1988; Bonvallet and Rougier, 1989; Bois and Lenfant, 1991; Charnet et al., 1991). However, Mitra and Morad (1986) showed an increase in  $I_{CaT}$  by isoproterenol in guinea pig ventricular myocytes. Tseng and Boyden (1988) reported that  $\beta$ -stimulation increased  $I_{CaT}$  in canine ventricular myocytes but not in Purkinje cells, whereas  $\alpha_1$ -adrenergic stimulation increased  $I_{CaT}$  in both cell types. In a subsequent paper, Tseng and Boyden (1989) confirmed the increase in  $I_{CaT}$  by  $\alpha_1$ - but not by  $\beta$ -adrenergic stimulation in canine Purkinje cells. Furthermore, Tseng and Boyden (1991) suggested that activation of protein kinase C decreases  $I_{CaT}$  in cardiac Purkinje cells. In addition, Bonvallet and Rougier (1989) presented evidence that angiotensin II increases and atrial natriuretic factor decreases  $I_{CaT}$ . Thus, there seems to exist an agreement in the literature regarding the kinetics, selectivity, modulation, and pharmacology of  $I_{CaL}$ , but incomplete or contradictory results are reported for  $I_{CaT}$ . Even its presence in frog atrium remains a matter of debate (compare Bonvallet and Rougier, 1989, with Campbell, Giles, Hume, and Shibata, 1988*a*, and Campbell, Giles, and Shibata, 1988*b*).

Because each type of calcium channel might have different functional implications, an investigation of their pharmacological profiles would contribute to the understanding of their specialized physiological role. In this study, using the whole-cell patch-clamp technique, we investigated the characteristics and pharmacological properties of  $I_{CaT}$  in bullfrog atrial cells and compared them with those of  $I_{CaL}$  of atrial and ventricular cells.

## MATERIALS AND METHODS

Single atrial and ventricular cells were dissociated from bullfrog hearts (*Rana catesbeiana*) by a method similar to that described by Arrio-Dupont and De Nay (1985). In brief, hearts were mounted in a Langendorff column and washed at 30°C with 100 ml of a standard Ringer solution (see below) containing only 50  $\mu$ M  $CaCl_2$  (low-Ca Ringer). Thereafter, 20 ml of low-Ca Ringer supplemented with 5 mM creatine, 1  $\mu$ l/ml of nonessential amino acids and vitamins medium (MEM; Boehringer Mannheim Corp., Indianapolis, IN), 1 mg/ml bovine serum albumin (Sigma Chemical Co., St. Louis, MO), 1.5 mg/ml collagenase (Boehringer Mannheim Corp.), and 0.4 mg/ml trypsin (Merck-Clévenot S.A., Nogent-sur-Marne, France), was recirculated for 45 min. Atria and ventricle were then separated, cut into small pieces, and agitated in the same Ringer without enzymes for 20 min. Supernatants were filtered through a nylon gauze (250  $\mu$ m) and centrifugated at 900 revolutions/min for 5 min. Pellets of cells were then "diluted" in fresh Ringer, supplemented with MEM containing 0.9 mM  $CaCl_2$ , and stored in a refrigerator. Cells could be used for up to 36 h after isolation.

In some experiments, pronase E (1.5–1.75 mg/ml; Merck-Clévenot S.A.) was used instead of collagenase and trypsin. In these cases, the enzymatic dissociation period could be shortened to 20–25 min with a similar cell yield. However, as the low threshold Ca current ( $I_{CaT}$ ) was often missing in pronase dissociation, this method was finally discarded. It is to be noted that in agreement with Dowell and Tarr (1988), single myocytes obtained from pronase dissociations were superior in overall viability and could be used for up to 72 h after isolation.

For electrophysiological recording, cells were left to lie in Petri dishes (type 1008; Falcon, Becton Dickinson and Co., Plymouth, UK) on the stage of an inverted microscope and perfused by gravity (0.65 ml/min) with a standard Ringer solution containing 20 mM CsCl, no KCl, and 1  $\mu$ M tetrodotoxin (TTX; Sigma, L'Isle d'Abeau, Chesnes, France) to ensure complete blockade of K and Na currents. Cells were positioned at the extremity of one of six microcapillaries (Tygon microbore tubing, i.d. = 250  $\mu$ m, o.d. = 750  $\mu$ m; Norton Performance Plastics, Wayne, NJ) independently perfused by gravity (16  $\mu$ l/min) allowing rapid changes ( $\leq 1$  s) of the extracellular medium. Patch electrodes (1–3 M $\Omega$ ) were made from glass capillaries (Microcaps; Drummond Scientific Co., Broomall, PA). A patch-clamp system (RK-300; Biologic, Claix, France) was used which permitted liquid junction potential compensation. After a gigaseal was obtained, the membrane patch was broken and whole-cell membrane capacitance and series resistance were measured and compensated. The intracellular solution in the patch electrode contained (mM): 120 CsCl, 5 K<sub>2</sub>-EGTA, 4 MgCl<sub>2</sub>, 5 Na<sub>2</sub>-creatine phosphate, 3 Na<sub>2</sub>-ATP, 0.4 Na<sub>2</sub>-GTP, and 10 HEPES (pH 7.1, adjusted with KOH or CsOH). Under these experimental conditions, Ca currents can be measured as the difference between peak inward current and the current level at the end of the voltage-clamp pulse (Alvarez et al., 1990).

Voltage-clamp protocols were delivered by a SEN-7103 electronic stimulator (Nihon-Kohden, Tokyo, Japan) coupled to a S-8300 step-pulse generator (Nihon-Kohden). Membrane potential was routinely depolarized for 200 ms every 4 s from a holding potential (HP) of  $-100$  to  $-50$  mV for the study of the high threshold Ca current ( $I_{CaT}$ ) or to 0 mV for the study of the low threshold Ca-current ( $I_{CaL}$ ). In most ventricular cells studied, the HP was  $-80$  mV. Availability curves for  $I_{CaT}$  and  $I_{CaL}$  were obtained from double-pulse protocols: every 4 s a 200-ms prepulse to different potentials was followed by a 200-ms pulse to  $-50$  mV for  $I_{CaT}$  or to 0 mV for  $I_{CaL}$ . Prepulses and test pulses were separated by a 3-ms return to the holding potential. The amplitude of  $I_{Ca}$  elicited by the test pulse is plotted as a percentage of the  $I_{Ca}$  evoked in the absence of a prepulse. From prepulse depolarizations current-to-voltage relationships were obtained. Reactivation (recovery from inactivation) was analyzed by depolarizing the cell every 8 s from the HP to  $-50$  mV for  $I_{CaT}$  or to 0 mV for  $I_{CaL}$ , returning to the HP for 13 intervals

from 3 ms to 3 s. Current recordings were stored on FM tape (DFR 3415; Sony, Clichy, France) and simultaneously displayed on a storage oscilloscope (VC-10; Nihon-Kohden).

### *Solutions and Drugs*

The composition of the standard Ringer solution was (mM): 88.4 NaCl, 2.5 KCl, 23.8 NaHCO<sub>3</sub>, 0.5 NaH<sub>2</sub>PO<sub>4</sub>, 1.8 MgCl<sub>2</sub>, 1.8 CaCl<sub>2</sub>, 5.0 glucose, and 5.0 Na<sub>2</sub>-pyruvate; pH was adjusted to 7.4. Inorganic or organic compounds were added to this Ringer as indicated and all solutions were gassed with 95% O<sub>2</sub>/5% CO<sub>2</sub>. In a few experiments NaCl, NaHCO<sub>3</sub>, and NaH<sub>2</sub>PO<sub>4</sub> were substituted by 73.3 mM BaCl<sub>2</sub> and 10 mM HEPES; the pipette solution then contained 5 mM BaATP and 400 μM LiGTP to substitute all Na salts. All experiments were carried out at room temperature, 20–23°C.

The following drugs were used: isoproterenol, phenylephrine, and propranolol (Sigma Chemical Co.), nifedipine and BAY K 8644 both dissolved in ethanol (Bayer, Wuppertal, Germany), ATP and 3', 5' cAMP (Merck-Clévenot), prazosin (Pfizer, Orsay, France), and ouabain (BDH Chemicals Ltd., Dorset, UK).

Results were analyzed by a paired *t* test and are expressed as means and standard error of the mean.

## RESULTS

### *Two Components of Calcium Current, I<sub>CaT</sub> and I<sub>CaL</sub>*

In an atrial cell held at –100 mV, clamp steps above –70 mV activated a time-dependent inward current which became larger with increasing depolarizations. When the HP was set at –50 mV, clamp steps up to –35 mV did not activate any current. Depolarizations to more positive potentials elicited another inward current (Fig. 1A). Fig. 1B shows the voltage dependence of the averaged peak inward currents relative to cell membrane capacity obtained in most of the atrial cells. At –100 mV HP, an activation threshold of –75 mV and two maxima were clearly distinguished in the current–voltage (*I*-*V*) relationship. As TTX was present at a concentration of 1 μM (or 3 μM in some experiments), a concentration that induced a complete blockade of the Na current (see Hume and Giles, 1983), it follows that the maximum current at –30 mV in the *I*-*V* curve was probably not due to the inflow of Na through fast Na channels. Moreover, reducing Na to 30 mM (substituted by choline chloride) did not significantly affect the kinetics or the amplitude of the current elicited by membrane depolarizations up to –30 mV. However, the current evoked at 0 mV was increased by 15 ± 3% (*n* = 7) in Na-poor solutions. Finally, 3 mM MnCl<sub>2</sub> depressed the low and high threshold currents by ~40 and 60%, respectively (data not shown). These observations reinforce the view that the low and high threshold currents correspond to the T- and L-type Ca<sup>2+</sup> currents, respectively.

In a series of experiments conducted on 34 atrial cells in which both currents were recorded over the full range of voltages, five cells exhibited *I*<sub>CaT</sub> whose peak amplitude was larger than *I*<sub>CaL</sub> (Fig. 1C). Notice that in these five cells, *I*<sub>CaL</sub> was much weaker and *I*<sub>CaT</sub> slightly larger than in the other cells. Their shape under the microscope and their averaged membrane capacitance (68.6 ± 6.8 pF) were not different from other cells (see Table I). Finally, one cell exhibited only *I*<sub>CaT</sub>; its peak amplitude was 1.4 pA/pF at –30 mV depolarization.

The characteristics of both *I*<sub>CaT</sub> and *I*<sub>CaL</sub> from a total of 156 cells are given in Table

I, which also compares  $I_{CaL}$  elicited in either atrial or ventricular cells. Membrane capacitance was the same in both cell types, indicating that we used atrial and ventricular cells of similar size. Peak  $I_{CaL}$  elicited by a depolarization to 0 mV had the same amplitude and kinetics in both cell types and was independent of the presence or absence of  $I_{CaT}$ . The maximal amplitude of  $I_{CaT}$  estimated at -30 mV was close to half-peak  $I_{CaL}$ .  $I_{CaT}$  kinetics were markedly accelerated by depolarizing to -30 mV instead of -50 mV, and were then similar to those of  $I_{CaL}$  at 0 mV.




FIGURE 1. Inward currents activated by depolarizing steps from in frog atrial cell. (A) Current recordings at HPs of -100 mV (*top*) and -50 mV (*bottom*) elicited in a cell by depolarizing pulses as indicated. Horizontal lines indicate zero current level. (B and C) Calcium current density-voltage relationships averaged on two pools of atrial cells which exhibit either  $I_{CaT} < I_{CaL}$  (B;  $n = 29$ ) or  $I_{CaT} > I_{CaL}$  (C;  $n = 5$ ). Results are expressed as means  $\pm$  SE.

These two types of Ca currents,  $I_{CaT}$  and  $I_{CaL}$ , were found in 107 of 117 atrial cells when collagenase and trypsin were used for dissociating the cells. However, in pronase dissociations only 30% of atrial cells (4 of 12) presented both currents. Nevertheless, the characteristics of  $I_{CaT}$ , when present, were not affected by pronase.

In 35 ventricular cells examined, no inward currents were activated between -70 and -30 mV, indicating that  $I_{CaT}$  was absent, in agreement with the contention that ventricular cells exhibited only the high threshold current (see also Argibay, Fischmeister, and Hartzell, 1988).

TABLE I  
 Characteristics of the T- and L-Type Ca Currents

	$I_{Ca}$	$C_m$	Density	Time to peak	Time to half-inactivation
$I_{CaT}$	$pA$	$pF$	$pA/pF$	$ms$	$ms$
-50 mV ( $n = 78$ )	$40.1 \pm 2.3$	$79.8 \pm 2.1$	$0.50 \pm 0.03$	$13.3 \pm 0.4$	$20.1 \pm 0.6$
-30 mV ( $n = 29$ )	$67.8 \pm 4.9$	$70.1 \pm 2.6$	$0.97 \pm 0.07$	$4.9 \pm 0.3$	$14.6 \pm 0.5$
$I_{CaL}$					
Atrial ( $n = 69$ )	$180.0 \pm 15.7$	$81.1 \pm 2.1$	$2.22 \pm 0.17$	$5.4 \pm 0.1$	$16.9 \pm 0.3$
Ventricular ( $n = 35$ )	$188.0 \pm 17.4$	$80.2 \pm 3.3$	$2.34 \pm 0.24$	$5.8 \pm 0.1$	$16.3 \pm 0.5$

Notice that  $I_{CaL}$  peak and density were  $215.8 \pm 44.5$  pA and  $2.25 \pm 0.47$  pA/pF, respectively, in 14 atrial cells that did not show  $I_{CaT}$ . Amplitude, time to peak, and time to half-inactivation of  $I_{CaT}$  and  $I_{CaL}$  were estimated on traces elicited at -50 or -30 mV and at 0 mV, respectively, from a HP of -100 mV for atrial cells and -80 mV for ventricular cells.

When atrial cells were held at -50 mV, the threshold potential was more positive (-30 mV) and only one maximum was seen between 0 and +10 mV. The difference current obtained by subtracting peak current recorded for the same depolarizations applied from either -100 or -50 mV HP displayed a maximum at -30 mV (see Fig. 2). However, in ~70% of the cells, both atrial and ventricular, the amplitude of the inward current at potentials positive to -20 mV was significantly reduced when the HP was -50 instead of -100 mV. However, our standard double-pulse protocol did not demonstrate that this resulted from the voltage-dependent inactivation of  $I_{CaL}$  in these cases since identical steady-state inactivation curves were obtained at the two HPs. Such a reduction could be attributed to the slow inactivation of  $I_{CaL}$  as recently described by Schouten and Morad (1989). This phenomenon prevented us from




FIGURE 2. Effects of divalent cations on  $I_{CaT}$  and  $I_{CaL}$  in atrial cells. (A) The upper traces show the effects on  $I_{CaT}$  elicited by a 200-ms depolarizing step from -100 to -50 mV of Cd (50  $\mu$ M) and Ni (50  $\mu$ M) ions. The lower traces show the effects of the same two cations on  $I_{CaL}$  elicited by a depolarizing step from -100 to 0 mV. Horizontal lines at the left indicate zero current. Two different cells. (B) Current density-voltage relationships established in the presence of 50  $\mu$ M Cd (●,  $n = 10$ , solid line) or in control solution for the current obtained by subtracting peak currents elicited by the same depolarization range from two HPs (-100 and -50 mV) averaged in four cells that did not exhibit slow inactivation (▲, dotted line).

estimating  $I_{CaT}$  at membrane potentials positive to  $-30$  mV and from using such a high HP during pharmacological studies of  $I_{CaL}$ .

In the following studies, for routine monitoring of these currents, we measured  $I_{CaT}$  and  $I_{CaL}$  elicited by depolarizing pulses at  $-50$  and  $0$  mV, respectively, from a HP of  $-100$  mV. A depolarization to  $-50$  mV allowed us to study changes in  $I_{CaT}$  without contamination of  $I_{CaL}$ , even under the influence of compounds that shifted the voltage dependence of  $I_{CaL}$  to more negative potentials (see below). Estimation of  $I_{CaL}$  at  $0$  mV in cells that possessed  $I_{CaT}$  was difficult. We chose to ignore the contribution of  $I_{CaT}$  to total inward current at  $0$  mV, since on the cells that did not show slow inactivation, the estimated  $I_{CaT}$  amplitude obtained from the difference current (see Fig. 2) was on average only 9% of total inward current ( $n = 13$ ). Inward current at  $0$  mV in atrial cells will then be referred to as  $I_{CaL}$ .

#### *Effects of Divalent Cations*

Raising the extracellular Ca concentration to  $3.6$  mM significantly enhanced both currents:  $I_{CaT}$  and  $I_{CaL}$  were increased by  $52 \pm 5$  and  $31 \pm 5\%$ , respectively ( $P < 0.05$ ;  $n = 6$ ).

In another series of experiments we studied the effects of the inorganic blockers Cd and Ni, as well as the use of Ba as charge carrier on  $I_{CaT}$  and  $I_{CaL}$ . Typical examples of the inhibitory effects of Cd and Ni ions are shown in Fig. 2A. In the presence of  $50$   $\mu$ M Cd ions, averaged  $I_{CaT}$  amplitude elicited by a  $-50$ -mV depolarization was  $47.7 \pm 2.6\%$  ( $n = 23$ ) of control. It was also noticeable that the  $I_{CaT}$  kinetics were slowed down. Time to peak current was increased from  $12.7 \pm 0.2$  to  $15.0 \pm 0.5$  ms and time of  $I_{CaT}$  half-inactivation ( $19.9 \pm 0.9$  ms at  $-50$  mV) was significantly prolonged by  $28.3 \pm 3.0\%$  ( $n = 23$ ). The average  $I_{CaT}$  density-voltage relationship as shown in Fig. 2B demonstrated that the reduction in peak current induced by Cd was seen at each potential; this occurred with a minor shift toward the left when compared with the  $I_{CaT}$  density-voltage relationship obtained after estimating  $I_{CaT}$  with the difference method in four control cells. Such an effect suggests that the Cd block is weakly voltage dependent.

Ni ions ( $50$   $\mu$ M) abolished  $I_{CaT}$  and slightly decreased  $I_{CaL}$  ( $9.2 \pm 0.4\%$ ;  $n = 8$ ). In the presence of  $50$   $\mu$ M Ni the whole  $I_{CaL}$ - $V$  relationship was shifted by  $\sim 10$  mV toward hyperpolarizing potentials with a weak reduction in the maximal inward current; the threshold for the inward current was now  $-35$  mV (not shown). A similar shift occurred on the availability curve. These effects were observed in the five cells investigated and were already reported by Mitra and Morad (1986).

The substitution of  $Ba^{2+}$  for  $Ca^{2+}$  significantly increased  $I_{CaL}$  by  $64 \pm 8\%$  ( $n = 6$ ) and markedly delayed its inactivation as already reported (Argibay et al., 1988); this left  $I_{CaT}$  amplitude and time course unaffected except in a few cells in which  $I_{CaT}$  amplitude was slightly reduced (not shown). Five experiments were performed in the total absence of Na ions, which were replaced by  $Ba^{2+}$  as is often done during single-channel patch-clamp experiments. Fig. 3 shows the currents elicited by depolarizing to  $-20$  and  $+20$  mV and the  $I$ - $V$  relationships established under these conditions ( $73.3$  mM Ba), as well as after adding  $50$   $\mu$ M Cd ions. Except that the voltage dependence of both currents was shifted by  $\sim 30$  mV in the depolarizing direction as expected from the screening effect of large divalent concentration,

roughly similar current kinetics were observed as with 1.8 mM Ba as the charge carrier (Fig. 3A). Cd ions suppressed the large Ba current carried by L-type channels, but in the lower range of depolarizations left a significant current with typical T-type kinetics (Fig. 3B). Furthermore, Ni ions (50  $\mu$ M) abolished the Ba current carried through T-type channels ( $n = 3$ ; see Fig. 8B).

*Inactivation and Recovery from Inactivation of  $I_{CaT}$  and  $I_{CaL}$*

The amplitude of  $I_{CaT}$  elicited by depolarizations to  $-50$  mV was dependent upon the short-term previous history of the membrane potential. Prepulses of different magnitudes applied with different delays reduced  $I_{CaT}$  to different extents during the test pulse. A 200-ms prepulse to  $-50$  mV fully suppressed the current during the test pulse applied after a 3-ms interval; however, after prepulse depolarization to  $+50$




FIGURE 3. Inward currents in the presence of high Ba concentration and effects of Cd ions. (A) Representative recordings of currents elicited at  $-20$  and  $+20$  mV from a  $-100$  mV HP in the presence of 73.3 mM Ba or after adding 50  $\mu$ M Cd ions ( $\blacktriangle$ );  $C_m = 70$  pF. (B)  $I$ - $V$  relationships established on five cells in high Ba solution (73.3 mM,  $\bullet$ ) or after adding 50  $\mu$ M Cd ( $\blacktriangle$ ). Peak Ba current amplitude was estimated by the algebraic difference between the peak inward current in Ba and the current recorded in the presence of Cd ions at the end of the 200-ms pulse of the same voltage.

mV,  $I_{CaT}$  partially recovered to reach  $\sim 40\%$  of its full value (Fig. 4A). Its kinetics were not markedly changed.  $I_{CaT}$  recorded during the test pulse was markedly reduced when increasing the interval to 50 ms (Fig. 4B). Finally,  $I_{CaT}$  was not significantly altered by increasing the prepulse from 200 to 500 ms (Fig. 4C).

Fig. 5A shows averaged inactivation curves for both Ca currents. On individual cells, the inactivation curve could be described in part by the theoretical Boltzmann distribution function, assuming that the decrease in availability from 1 to 0 when increasing prepulse amplitude is only sensitive to membrane depolarization. The mean values for the half-maximal potentials,  $V_{1/2}$ , and the slope factors,  $k$ , were  $-66.6 \pm 0.5$  mV and  $6.4 \pm 0.2$  for  $I_{CaT}$  ( $n = 29$ ) and  $-29.9 \pm 0.9$  mV and  $7.1 \pm 0.2$  for  $I_{CaL}$  ( $n = 36$ ), respectively. Inactivation curves for  $I_{CaT}$  were thus displaced  $\sim 35$






FIGURE 4. Influence of the conditioning pulses on the current recorded during a test pulse to  $-50$  mV. (A) Currents ( $\bullet$ ,  $\blacktriangle$ ,  $\blacklozenge$ ) recorded after prepulses of different amplitude: 0,  $-50$ , and  $+50$  mV, respectively. (B) Currents ( $\blacktriangle$ ,  $\bullet$ ) recorded after the same prepulse was applied 3 or 50 ms before the test pulse. (C) Currents ( $\bullet$ ,  $\blacktriangle$ ) recorded after a prepulse of 200 or 500 ms duration. HP was  $-100$  mV.

mV to more negative potentials and displayed a steeper voltage dependence when compared with  $I_{CaL}$  inactivation curves.

Both currents showed less inactivation with positive prepulse potentials. This is a well-known phenomenon for  $I_{CaL}$ , which was related to Ca-dependent Ca current inactivation (Mentrard, Vassort, and Fischmeister, 1984; Lee, Marban, and Tsien, 1985). A similar relationship with membrane potential was found for  $I_{CaT}$  inactivation. Increasing TTX concentrations up to  $3 \mu\text{M}$  did not abolish  $I_{CaT}$  recovery after high prepulse potentials. This indicates that it was not an artefact due to  $I_{Na}$ . Nor was




FIGURE 5. Inactivation curves for  $I_{CaT}$  and  $I_{CaL}$ . (A) Voltage dependence of steady-state inactivation (availability) of  $I_{CaT}$  ( $\blacktriangle$ ) and  $I_{CaL}$  ( $\bullet$ ), obtained with the protocol described in Materials and Methods. Solid lines represent Boltzmann's fits of averaged values with  $V_f = -66.2$  and  $-25.4$  mV and  $k = 5.2$  and  $5.8$  for  $I_{CaT}$  and  $I_{CaL}$ , respectively. (B) Voltage dependence of steady-state inactivation (availability) of  $I_{CaT}$  established with a 3-ms interval between the test pulse to  $-50$  mV and the prepulse (200 ms) in control ( $\bullet$ ),  $50 \mu\text{M Cd}$  ( $\blacktriangle$ ), and when Ba ( $1.8 \text{ mM}$ ) was the charge carrier ( $\blacksquare$ ); HP was  $-100$  mV ( $n = 4$ ). Curves fitted by the Boltzmann equation up to  $-50$  mV.

it the result of a Ca-mediated inactivation of  $I_{CaT}$ , since similar availability curves for  $I_{CaT}$  were obtained in the presence of Cd or when Ba replaced Ca. Only a minor leftward shift ( $-4.6 \pm 0.6$  mV;  $n = 5$ ) was determined on the availability curves of  $I_{CaT}$  when comparing  $I_{CaT}$  recorded in cells before and in the presence of  $50 \mu\text{M}$  Cd (Fig. 5 B). Equimolar Ba substitution did not significantly affect the  $I_{CaT}/V$  relationship (not shown) or the availability curve (Fig. 5 B). Availability curves of  $I_{CaT}$  with similar characteristics including recovery from inactivation for high prepulses were




FIGURE 6. Reactivation curves for  $I_{CaT}$ . (A) Reactivation (recovery from inactivation) curves for  $I_{CaT}$  ( $\blacktriangle$ ) and  $I_{CaL}$  ( $\bullet$ ) obtained from an HP of  $-100$  mV. Stimulation protocols were as described in Materials and Methods. (B) Reactivation (recovery from inactivation) curves established in control ( $\bullet$ ) or in the presence of  $50 \mu\text{M}$  Cd and  $1.8$  mM Ba ( $\blacktriangle$ ) with a  $200$ -ms conditioning prepulse to  $+70$  mV followed at increasing intervals by a  $-50$ -mV test pulse; HP was  $-100$  mV as indicated in the inset. Stimulation frequency was  $1/8$  s ( $n = 6$ ). (C) Reactivation (recovery from inactivation) curves established in the presence of  $50 \mu\text{M}$  Cd and  $1.8$  mM Ba on three other cells held successively at  $-100$  ( $\bullet$ ),  $-70$  ( $\blacksquare$ ), and  $-120$  mV ( $\blacktriangle$ ), otherwise using the same protocol as in B.

obtained in the high Ba (0 Na) solution and after the further addition of  $50 \mu\text{M}$  Cd (not shown). The whole curves were, however, shifted to the right by  $\sim 30$  mV; half-inactivation was  $-37.1 \pm 1.5$  or  $-40.6 \pm 1.4$  mV ( $n = 5$ ) in  $73.3$  or in  $73.3$  mM Ba and  $50 \mu\text{M}$  Cd, respectively. T-type currents recorded in Ba solutions at  $-20$  mV showed similar time courses whether or not the test pulse was preceded by a prepulse as previously observed under control conditions.

$I_{CaT}$  and  $I_{CaL}$  showed differences in their recovery characteristics (Fig. 6 A). Average

time of half-recovery from inactivation was  $71.3 \pm 0.9$  ms for  $I_{CaL}$  ( $n = 15$ ) and  $171.0 \pm 1.8$  ms for  $I_{CaT}$  ( $n = 10$ ). In agreement with the observations of Argibay et al. (1988), cells with  $I_{CaL}$  density  $< 2.0$  pA/pF exhibited an overshoot of up to 120% on the  $I_{CaL}$  reactivation curves. Such a phenomenon was never seen with  $I_{CaT}$ ; rather, the slow recovery of  $I_{CaT}$  inactivation seemed to imply two kinetic processes as already characterized by Bossu and Feltz (1986).

In each ionic experimental condition, the  $I_{CaT}$  availability curve bent up. However, this apparent reduction of current inactivation with depolarization was obtained with a 3-ms interval between the high depolarizing prepulse and the test pulse and at  $-100$  mV HP. On establishing a reactivation curve with a test pulse at  $-50$  mV and a prepulse at  $+70$  mV, the prolongation of the interval above 3 ms initially decreased current amplitude during the test pulse so that it was nearly abolished within 25 ms (Fig. 6 B; see also Fig. 4 B). Further increases in the interval between the conditioning pulse at high voltage and the test pulse allowed for full recovery from inactivation. Similar results were obtained in the presence of Cd ions and when Ba ions were the charge carrier as shown in Fig. 6 B. Recoveries from inactivation of both  $I_{CaT}$  and  $I_{CaL}$  were already shown to be voltage dependent (Argibay et al., 1988; Tseng, 1988; Tseng and Boyden, 1989; Hirano, Fozzard, and January, 1989b). Furthermore, for  $I_{CaT}$ , the time course of recovery did not depend on the ion carrier. In the presence of  $50 \mu\text{M}$  Cd and  $1.8$  mM Ba, and after a 200-ms prepulse to  $+70$  mV, some recovery from inactivation of  $I_{CaT}$  was seen only after at least 300 ms at  $-70$  mV HP, while recovery was already significant after only 10 ms when HP was  $-120$  mV (Fig. 6 C). Average times for half-recovery from inactivation were  $65 \pm 17$ ,  $90 \pm 18$ , and  $1,150 \pm 237$  ms at  $-120$ ,  $-100$ , and  $-70$  mV HP, respectively ( $n = 3$ ). These values were independent of the charge carrier and of the presence or absence of Cd.

Fig. 6 shows that T-type current did not recover from inactivation in a simple manner regardless of whether Ca or Ba was the main charge carrier. The amplitude of the current during the test pulse depended on both the prepulse amplitude (see Fig. 5; cf. Fig. 6, A and B) and the interpulse duration (Fig. 6 B). Fig. 6 C further suggests that after 3 ms at a  $-100$  mV interpulse HP, a significant part of the current had already deactivated when compared with the results at  $-70$  mV HP, a situation in which the test pulse current was still 85% of peak current after 3 ms. Indeed, the decrease in the remaining  $I_{CaT}$  (deactivation) after a high depolarizing prepulse was strongly voltage dependent.  $I_{CaT}$  deactivated exponentially with a time constant of 29 ms at  $-70$  mV. Deactivation of  $I_{CaT}$  was completed within 25 ms at  $-100$  mV and within a few milliseconds at  $-120$  mV. This was also true in the high-Ba solution; taking into account a 30-mV shift, deactivation was also completed in  $< 10$  ms at  $-100$  mV. The deactivation rate was not dependent on the prepulse amplitude. These results are reminiscent of our previous results obtained with the double sucrose gap method on the slow inward Ca current (see Fig. 14 in Mentrard et al., 1984). They led us to reinvestigate the time dependence of inactivation at several voltages for both  $I_{CaT}$  and  $I_{CaL}$ .

The effects of increasing the prepulse duration on both  $I_{CaL}$  and  $I_{CaT}$  are illustrated in Fig. 7, at different voltages and when the current carrier was either Ca or Ba, using a 3-ms interpulse. Inactivation of  $I_{CaL}$  increased with prepulse duration. The effects were more marked at 0 mV prepulse potential and in the presence of Ca ions than at

higher prepotentials and in the presence of Ba ions (Fig. 7 *A*). When the prepulse potential was +40 mV, intermediate values were obtained in either Ca or Ba solutions, except that the initial reduction of  $I_{CaL}$  occurred more rapidly than with the two other prepulse voltages. These observations are in agreement with the fact that some part of the Ca current inactivation is mediated by Ca ions entering through the channel. The time dependence of  $I_{CaT}$  inactivation exhibited much more complex characteristics. A prepulse potential at -50 mV inactivated  $I_{CaT}$  within 200 ms. With larger prepulse depolarizations, inactivation was never completed (Fig. 7 *B*). A prepulse at +40 mV reduced  $I_{CaT}$  to 45% of its control amplitude within 20 ms; longer prepulses did not significantly change the inactivation level. A prepulse to +70 mV also reduced  $I_{CaT}$  by about half, but with a slower kinetic; however,




FIGURE 7. Effects of increasing prepulse duration on  $I_{CaT}$  and  $I_{CaL}$ . (*A*) Inactivation of  $I_{CaL}$  recorded during a depolarizing pulse at 0 mV by prepulse depolarizations to 0 mV ( $\circ$ ,  $\bullet$ ) or to +70 mV ( $\Delta$ ,  $\blacktriangle$ ) whose durations increased up to 500 ms in control conditions (open symbols) or in 1.8 mM Ba (closed symbols) ( $n = 3-5$ ). (*B*) Inactivation of  $I_{CaT}$  recorded during a depolarizing pulse to -50 mV by prepulse depolarizations to -50 ( $\bullet$ ), +40 ( $\blacktriangledown$ ), and +70 mV ( $\blacktriangle$ ), whose durations increased up to 500 ms in control conditions ( $n = 3-5$ ). Curves obtained in 1.8 mM Ba were identical to the ones obtained in control conditions at each of three potentials and are not shown for clarity.

prolonging it > 50 ms induced less inactivation. These effects were similar whether Ca or Ba was the charge carrier and in the presence or absence of Cd ions.

#### *Effects of Dihydropyridines*

BAY K 8644 at 1  $\mu$ M increased both  $I_{CaT}$  and  $I_{CaL}$  under our experimental conditions (Fig. 8). However, there were significant differences in its action on the two current types. The effects of BAY K 8644 on  $I_{CaT}$  seemed to occur at lower concentrations and were less than on  $I_{CaL}$ ; at 10 nM the increases in  $I_{CaT}$  and  $I_{CaL}$  were 1.5- and 1.4-fold the control value, respectively, instead of 1.8 and 2.7 at 1  $\mu$ M (Fig. 8 *B*). The relative increase in  $I_{CaT}$  induced by 1  $\mu$ M BAY K 8644 was the same whether HP was -100 or -80 mV; but  $I_{CaL}$ , which was increased twofold or more when HP was -100




FIGURE 8. Effects of  $1 \mu\text{M}$  BAY K 8644 on  $I_{CaT}$  and  $I_{CaL}$ . (A) The tracings (a) show recordings of  $I_{CaT}$  (top, elicited at  $-50 \text{ mV}$ ) and  $I_{CaL}$  (bottom, elicited at  $0 \text{ mV}$ ), evoked from a HP of  $-100 \text{ mV}$  in control conditions ( $\blacktriangledown$ ) and after perfusion with BAY K 8644 at  $1 \mu\text{M}$  ( $\blacklozenge$ ). The tracings (b) show the effects of  $50 \mu\text{M}$  Cd on  $I_{CaT}$  elicited at  $-50 \text{ mV}$  (upper part) and the effects of a further addition of BAY K 8644 (lower part) ( $\blacktriangledown$ , control;  $\blacksquare$ , Cd;  $\bullet$ , BAY K). (B) Current density-voltage relationships in the control solution ( $\blacktriangledown$ ) and during the action of  $1 \mu\text{M}$  BAY K 8644 ( $\blacklozenge$ ). Inward currents were increased over the whole range of potentials by BAY K 8644 when HP was  $-100 \text{ mV}$  ( $n = 6$ ). The addition of  $50 \mu\text{M}$  Ni inhibited current below  $-40 \text{ mV}$  and reduced the BAY K 8644 effect on  $I_{CaL}$  ( $\bullet$ ,  $n = 3$ ). (C) Inactivation curves for  $I_{CaT}$  ( $\blacksquare$ ,  $\bullet$ ) and  $I_{CaL}$  ( $\blacktriangledown$ ,  $\blacklozenge$ ) in control conditions ( $\blacktriangledown$ ,  $\blacksquare$ ) and under the action of BAY K 8644 ( $\bullet$ ,  $\blacklozenge$ ) ( $n = 6$ ). The normalization factors used to scale the curves after the effects of BAY K 8644 were 0.6 and 0.5 for  $I_{CaT}$  and  $I_{CaL}$ , respectively, as a consequence of the dihydropyridine-induced increased in currents elicited at  $-50$  and  $0 \text{ mV}$ . (D) Current density-voltage relationships for  $I_{CaT}$  in atrial cells in the presence of  $50 \mu\text{M}$  Cd ( $\blacksquare$ ) and after the addition of  $1 \mu\text{M}$  BAY K 8644 ( $\bullet$ ) ( $n = 4$ ).

$\text{mV}$ , was inhibited when HP was  $-50 \text{ mV}$  (not shown). Such an effect has already been reported by others (Sanguinetti, Krafte, and Kass, 1986; Kass, 1987). Note that holding the cell at either  $-80$  or  $-50 \text{ mV}$  should similarly affect the voltage-dependent kinetics of  $I_{CaT}$  or  $I_{CaL}$ , respectively, according to the voltage range of their availability curves. Fig. 8 B also shows that in the presence of BAY K 8644, peak  $I_{CaT}$  was not shifted, while peak  $I_{CaL}$  occurred at  $\sim 10 \text{ mV}$  lower depolarizations. This  $10\text{-mV}$  leftward shift was also seen when HP was set at  $-50 \text{ mV}$  (see also Sanguinetti et al., 1986). However, such a shift of  $I_{CaL}$  voltage dependence could not affect

estimation of  $I_{CaT}$  elicited at  $-50$  mV since there was no current measurable below  $-40$  mV in the presence of  $50 \mu\text{M}$  Ni (Fig. 8 *B*). Note that Ni induced a further leftward shift and slightly reduced the BAY K 8644-enhanced  $I_{CaL}$ . Also, there was no current elicited by depolarizations below  $-40$  mV in four other cells when Ba ( $1.8$  mM) was the charge carrier in the presence of  $50 \mu\text{M}$  Ni. BAY K 8644 significantly shortened the time of  $I_{CaT}$  half-inactivation by  $24.7 \pm 3.8\%$  (from  $21.2 \pm 2.1$  to  $16.3 \pm 2.3$  ms;  $P < 0.05$ ,  $n = 9$ ) at  $10$  nM and by  $23.6 \pm 4.1\%$  (to  $17.4 \pm 2.3$  ms;  $P < 0.05$ ,  $n = 9$ ) at  $1 \mu\text{M}$ . This was also true for  $I_{CaL}$  but only at  $1 \mu\text{M}$ ; time of  $I_{CaL}$  half-inactivation was decreased by  $29.1 \pm 8.2\%$  (from  $17.3 \pm 1.0$  to  $12.0 \pm 1.0$  ms,  $P < 0.05$ ,  $n = 6$ ). The steady-state inactivation curve for  $I_{CaT}$  was not shifted (except by  $\sim 5$  mV in one cell) under the action of  $1 \mu\text{M}$  BAY K 8644, while  $I_{CaL}$  inactivation was significantly displaced by  $10.9 \pm 0.9$  mV ( $P < 0.01$ ,  $n = 6$ ) toward more negative potentials (Fig. 8 *C*).

That BAY K 8644 increased  $I_{CaT}$  at every potential independent of its effects on  $I_{CaL}$  was reinforced by the following two observations. First, the fast inactivating T-type current was similarly increased by BAY K 8644 whether Ba or Ca ions were the charge carrier, without affecting current kinetics. Second, when  $I_{CaT}$  was recorded after inhibition of  $I_{CaL}$  by  $50 \mu\text{M}$  Cd ions, the addition of the dihydropyridine agonist ( $1 \mu\text{M}$ ) induced an increase in current amplitude with roughly the same magnification at each depolarizing test pulse (Fig. 8 *D*). As in control conditions, this increase was also unaffected by holding the cells at  $-80$  mV. Typical current traces show that BAY K 8644 very markedly shortened time to peak and accelerated inactivation in the presence of Cd ions (Fig. 8 *A, b*). Note that the stimulatory effect of BAY K 8644 on peak  $I_{CaT}$  recorded in the presence of Cd ions was also much larger.

BAY K 8644 delayed the recoveries of inactivation of both  $I_{CaT}$  and  $I_{CaL}$ , the effect being more marked on  $I_{CaL}$  (Fig. 9, *A* and *B*). In addition, BAY K 8644 suppressed the overshoot in the recovery of inactivation curve of  $I_{CaL}$  when it occurred under control conditions. Time of  $I_{CaT}$  half-recovery of inactivation was increased by  $54.5 \pm 8.6$  ms ( $P < 0.01$ ,  $n = 3$ ) and that of  $I_{CaL}$  by  $117.5 \pm 12.5$  ms ( $P < 0.01$ ,  $n = 4$ ) with  $1 \mu\text{M}$  BAY K 8644. In Fig. 9 *C* it is shown that the negative staircase of  $I_{CaT}$ , after an abrupt increase in stimulation rate, was only slightly enhanced in the presence of  $1 \mu\text{M}$  BAY K 8644, also suggesting a weak use-dependent action of the drug on this current. On the other hand,  $I_{CaL}$  was markedly depressed in a use-dependent fashion by BAY K 8644 as reported in Fig. 9 *C* on a ventricular cell which showed a slight positive staircase of  $I_{CaL}$  under control conditions.

Nifedipine was used at  $10$  nM and  $1 \mu\text{M}$ . Either increases or decreases in  $I_{CaT}$  and  $I_{CaL}$  were observed with nifedipine at  $10$  nM, but at  $1 \mu\text{M}$  nifedipine decreased  $I_{CaT}$  and  $I_{CaL}$  to  $82.5 \pm 2.6$  and  $51.4 \pm 4.1\%$  ( $P \leq 0.01$ ,  $n = 7$ ) of their respective controls. Inactivation of both currents was also altered. Times of half-inactivation were decreased by  $15.0 \pm 3.6$  and  $22.7 \pm 7.0\%$  for  $I_{CaT}$  and  $I_{CaL}$ , respectively. The availability curve of  $I_{CaT}$  was not affected by nifedipine ( $1 \mu\text{M}$ ), while a marked shift to more negative potentials ( $10.3 \pm 0.7$  mV;  $P < 0.01$ ,  $n = 7$ ) was found for  $I_{CaL}$ . The nifedipine-induced inhibition of  $I_{CaT}$  had similar relative magnitude when HP was less negative ( $-90$  or  $-80$  mV). All of these effects, including small increases in  $I_{CaT}$  amplitude induced by  $10$  nM nifedipine, were also observed when applying the dihydropyridine in the presence of Cd ions.

*Effects of Neuromediators*

The effects of  $\beta$ - and  $\alpha_1$ -adrenergic and  $P_2$ -purinergic stimulations were investigated only on cells obtained from collagenase–trypsin dissociations; they are summarized in Table II. Under our experimental conditions, both  $I_{CaT}$  and  $I_{CaL}$  were sensitive to the three neuromediators. The agonistic effects of  $\alpha_1$ -adrenergic and  $P_2$ -purinergic stimulations were even larger on  $I_{CaT}$  than on  $I_{CaL}$ ; however, the increases were limited when compared with the large increase eventually elicited on  $I_{CaL}$  by  $\beta$ -adrenergic stimulation (Table II, Fig. 10A, and see Fig. 11). In the presence of 1  $\mu$ M propranolol, the  $\alpha_1$ -adrenergic agonist phenylephrine (10  $\mu$ M) was effective in increasing  $I_{CaT}$  by 48%, and although the effect was variable from cell to cell




FIGURE 9. Effects of 1  $\mu$ M BAY K 8644 on  $I_{CaT}$  and  $I_{CaL}$  characteristics. The effects of BAY K 8644 are shown (dotted lines) on reactivation curves for  $I_{CaT}$  (A:  $\blacktriangle$ ,  $n = 3$ ) and  $I_{CaL}$  (B:  $\bullet$ ,  $n = 4$ ). (C) Use-dependent effects of BAY K 8644 on  $I_{CaT}$  and  $I_{CaL}$ . In both control conditions and after the addition of BAY K 8644, stimulation was stopped and resumed after 1 min at a cycle length of 400 ms. Note that BAY K 8644 (1  $\mu$ M, closed symbols) changed the staircase pattern of  $I_{CaL}$  (O,  $\bullet$ ) from slightly positive to negative. The drug only slightly enhanced the negative staircase of  $I_{CaT}$  ( $\Delta$ ,  $\blacktriangle$ ) after the increase in rate. HP =  $-100$  mV and  $I_{CaT}$  and  $I_{CaL}$  were elicited at  $-50$  and  $0$  mV, respectively, on a given cell.

(22–100% increase), it was seen in each of the seven cells studied.  $I_{CaL}$ , on the other hand, was increased at most by 20%, and in half of the cells this was only a transient effect. These effects were blocked by 0.1  $\mu$ M prazosin in the three cells investigated. At 1  $\mu$ M, ATP exerted a strong agonistic effect on  $I_{CaT}$ . This effect was seen in 12 of 17 cells. We already reported that ATP was not always active on  $I_{CaL}$  (Alvarez et al., 1990). As on  $I_{CaL}$ , the agonistic potency of ATP on  $I_{CaT}$  is limited to a very narrow range of concentrations; higher concentrations were less efficient or even reduced the current (Table II). No significant changes were detected in the kinetics of  $I_{CaT}$  or  $I_{CaL}$  under both  $\alpha_1$ - and  $\beta$ -adrenergic or  $P_2$ -purinergic stimulations. The effects of the three neuromediators were rapidly reversible. Further results, obtained in the

TABLE 11  
Effects of  $\alpha_1$ - and  $\beta$ -adrenergic and Purinergic Stimulations on  $I_{CaT}$  and  $I_{CaL}$   
in Frog Cardiac Cells

	Isoproterenol		Phenylephrine	ATP		
	0.1 $\mu$ M	1 $\mu$ M	10 $\mu$ M	0.1 $\mu$ M	1 $\mu$ M	10 $\mu$ M
$I_{CaT}$	133.0 $\pm$ 13.3 <sup>‡</sup> (5/8)	195.0 $\pm$ 27.5 <sup>‡</sup> (4/8)	148.2 $\pm$ 10.4* (7/7)	100 (0/4)	172.5 $\pm$ 26.3 <sup>‡</sup> (4/7)	121.5 $\pm$ 5.3 <sup>‡</sup> (3/4)
$I_{CaT}$ (Cd)		218.0 $\pm$ 18.1* (11/12)	217.5 $\pm$ 31.1* (4/4)		222.8 $\pm$ 27.5* (8/10)	
$I_{CaL}$	463.6 $\pm$ 27.1* (14/14)	562.8 $\pm$ 23.2* (14/14)	120.1 $\pm$ 5.5 <sup>‡</sup> (9/9)	123.0 $\pm$ 6.9 <sup>‡</sup> (5/6)	140.0 $\pm$ 4.6 <sup>‡</sup> (5/6)	112.8 $\pm$ 3.7 (4/5)

Values are percent of control peak current amplitudes taking into account only cells that responded to the agonists (indicated in parenthesis). The experiments with phenylephrine were performed in the presence of 1  $\mu$ M propranolol; the increase in  $I_{CaL}$  was transient in five of the nine cells and was estimated at peak. Currents were measured when depolarizing from a HP of -100 mV to -50 mV ( $I_{CaT}$ ) or to 0 mV ( $I_{CaL}$ ). In ventricular cells (eight with isoproterenol, two with phenylephrine, and two with ATP) HP was -80 mV.  $I_{CaT}$  (Cd) refers to Ca current recorded in the presence of 50  $\mu$ M Cd in another series of experiments. Means  $\pm$  SE. \* $P \leq 0.05$ . <sup>‡</sup> $P \leq 0.05$ .




FIGURE 10. Modulation of  $I_{CaT}$  and  $I_{CaL}$  by neurotransmitters. (A) Effects of 1  $\mu$ M isoproterenol and 10  $\mu$ M phenylephrine in the presence of 1  $\mu$ M propranolol and 1  $\mu$ M ATP on  $I_{CaT}$  and  $I_{CaL}$  elicited by depolarizations to -50 or 0 mV from HP = -100 mV in control conditions (left) or on  $I_{CaT}$  in the presence of 50  $\mu$ M Cd. Note that phenylephrine effect on  $I_{CaL}$  was maximal after 1 min (as shown here);  $I_{CaL}$  then recovered its control value within 2 min. Horizontal lines denote zero current level and filled circles indicate the presence of the neuromediators. (B) Current-voltage relationship established in control (○) and in the presence of 1  $\mu$ M ATP (●) on an atrial cell that exhibited an unusually small  $I_{CaL}$ .



presence of Cd ions and reported in Fig. 10A and Table II, also show that the amplitude of  $I_{CaT}$  could be modulated by neuromediators. Under this experimental condition essentially similar relative increases in  $I_{CaT}$  were observed with each of the three agonists. Fig. 10B reports the effects of ATP obtained on one of the five atrial cells that exhibited an unusually small  $I_{CaL}$ . At each membrane depolarization both  $I_{CaT}$  and  $I_{CaL}$  were markedly increased by ATP.

The effects of  $\beta$ -adrenergic stimulation on  $I_{CaT}$  and  $I_{CaL}$  were further characterized in another experimental series by applying alternative depolarizations to  $-50$  and  $0$  mV every 4 s from  $-100$  mV HP. The results of such an experiment are shown on Fig. 11A, which reports the relative increases in both currents after adding  $1 \mu\text{M}$




FIGURE 11. Effects of isoproterenol on  $I_{CaT}$  and  $I_{CaL}$ . (A) Relative increase in both  $I_{CaT}$  (●) and  $I_{CaL}$  (○) induced by the addition of  $1 \mu\text{M}$  isoproterenol. The further addition of  $1 \mu\text{M}$  propranolol antagonized these effects.  $I_{CaT}$  and  $I_{CaL}$  were 20 and 64 pA in control conditions at  $-50$  and  $0$  mV, respectively.  $C_m = 58$  pF. (B) Breaking the patch with a pipette containing  $50 \mu\text{M}$  cAMP hardly increased  $I_{CaT}$  (●) as compared with the markedly enhanced  $I_{CaL}$  (○). The later addition of  $1 \mu\text{M}$  isoproterenol did not affect  $I_{CaL}$  but increased  $I_{CaT}$ . The whole experiment was performed in the presence of  $100 \mu\text{M}$  IBMX and  $0.1 \mu\text{M}$  prazosin.  $I_{CaT}$  and  $I_{CaL}$  were 24 and 170 pA at  $-50$  and  $0$  mV, respectively, just after breaking the patch ( $C_m = 100$  pF).

isoproterenol. The isoproterenol-induced increase in  $I_{CaT}$  was much weaker and reached its maximum much faster than the  $I_{CaL}$  increase. Increases in both Ca currents were antagonized by  $1 \mu\text{M}$  propranolol, with a much faster recovery of  $I_{CaT}$ . Prazosin was without effect. However, the  $\beta$ -adrenergic-induced increase in  $I_{CaT}$  was not related to a cAMP increase; evidence is provided in Fig. 11B. When breaking the patch after a gigaseal was made with a pipette containing  $50 \mu\text{M}$  cAMP,  $I_{CaL}$  increased markedly; there was also a small sustained increase in  $I_{CaT}$  which could be

TABLE III  
Effects of Ouabain on  $I_{CaT}$  and  $I_{CaL}$  in Frog Cardiac Cells

	0.1 $\mu$ M	1 $\mu$ M (1 min)	1 $\mu$ M (5 min)	10 $\mu$ M
$I_{CaT}$ ( $n = 6$ )	94.0 $\pm$ 8.3	127.3 $\pm$ 10.8	139.6 $\pm$ 14.9 <sup>†</sup>	82.5 $\pm$ 4.9 <sup>†</sup>
$I_{CaL}$ ( $n = 8$ )	102.0 $\pm$ 2.0	137.6 $\pm$ 1.9*	88.8 $\pm$ 9.1	51.6 $\pm$ 2.1*

Values are percent of controls before the application of ouabain and are estimated from  $I_{CaT}$  and  $I_{CaL}$  elicited at  $-50$  and  $0$  mV, respectively, from a HP of  $-100$  mV but for three ventricular cells in which HP was  $-80$  mV. Means  $\pm$  SE. \* $P \leq 0.01$ . <sup>†</sup> $P \leq 0.05$ .

related to the usual "run up" of current during the stabilizing period. After  $I_{CaL}$  had reached its steady state, the further application of  $1 \mu$ M isoproterenol (in the presence of  $100 \mu$ M IBMX and  $0.1 \mu$ M prazosin) had no effect on  $I_{CaL}$  but rapidly and significantly increased  $I_{CaT}$ . On wash-out of isoproterenol,  $I_{CaT}$  recovery was also very rapid. In this experimental series the isoproterenol-induced increases in  $I_{CaT}$  were  $63.8 \pm 10.0$  and  $62.8 \pm 16.0\%$  of  $I_{CaT}$  control, and their times to half-maximal amplitude were  $13.5 \pm 4.3$  and  $23.2 \pm 5.3$  s in control ( $n = 8$ ) and cAMP-containing ( $n = 5$ ) internal solutions, respectively. For comparison, on the same cells the increases in  $I_{CaL}$  were  $405.6 \pm 95.1$  and  $687.2 \pm 120.6\%$  of  $I_{CaL}$  control and times to half-maximal amplitude were  $78.5 \pm 11.1$  and  $66.0 \pm 9.3$  s on applying isoproterenol or after perfusing with cAMP, respectively.

#### Effects of Ouabain

It is well known that cardiac glycosides can, under some circumstances, increase (Marban and Tsien, 1982), decrease (Deslauriers, Ruiz-Ceretti, Schanne, and Payet, 1982), or exert a dual effect (Fischmeister, Brocas-Randolph, Lechene, Argibay, and Vassort, 1986) on  $I_{CaL}$ . The effects of ouabain on the two Ca currents in frog atrial cells are reported in Table III. At  $1 \mu$ M, ouabain induced a sustained increase in  $I_{CaT}$  and exerted a biphasic effect on  $I_{CaL}$  as previously reported in frog ventricular cells (Fischmeister et al., 1986). Times of half-inactivation of both currents were significantly decreased with  $1$  and  $10 \mu$ M ouabain, but without a clear relation to concentration. Time of  $I_{CaT}$  half-inactivation was reduced by  $29.9 \pm 9.5\%$ , while that of  $I_{CaL}$  was reduced by  $19.0 \pm 5.3\%$  ( $n = 6$ ). In three of six cells the increase in  $I_{CaT}$  was preceded by oscillatory variations in amplitude (Fig. 12).




FIGURE 12. Effects of  $1 \mu$ M ouabain on  $I_{CaT}$ . Note that after the beginning of ouabain perfusion,  $I_{CaT}$  showed transient variations in amplitude which stabilized within 1 min to a steady-state value greater than control. The inset shows current recordings. HP =  $-100$  mV;  $I_{CaT}$  was evoked at  $-50$  mV for 200 ms every 4 s.

## DISCUSSION

The present results confirm that in bullfrog atrial, but not ventricular cells, two types of Ca currents,  $I_{CaT}$  and  $I_{CaL}$ , coexist, which can be recorded at physiological extracellular Ca concentrations. Following previous studies, several criteria were used to define the T-type Ca current. It activates and inactivates at relatively negative voltages, it is unchanged when substituting Ba for Ca; and it is partially inhibited by 50  $\mu$ M Cd and blocked by 50  $\mu$ M Ni but is insensitive to TTX. The possibility that the inward current recorded during depolarizations between  $-65$  and  $-30$  mV was a residual Na current was excluded because it appears unchanged when reducing Na to 30 mM and still exists when Ba is the only charge carrier, although within a different potential range as a result of surface charge screening. The most salient features of this work are the complex voltage dependence of  $I_{CaT}$  kinetics and the sensitivity of  $I_{CaT}$  to several pharmacological agents as well as to neuromediators.

Previous results obtained in cardiac cells isolated from the same (Bean, 1985a; Alvarez and Vassort, 1991) and another (Bonvallet, 1987; Bonvallet and Rougier, 1989) species of the same genus already suggested the existence of two Ca currents. On the other hand, Campbell et al. (1988a, b) found no evidence of the existence of  $I_{CaT}$  in the bullfrog atrial cells. As pointed out by Bonvallet and Rougier (1989), this could be due to species specificity or to the use of different recording techniques (dialyzed vs. nondialyzed cells). To these proposals we can add another one: the lack of recording  $I_{CaT}$  in bullfrog atrial cells might be related to the enzymatic dissociation procedure. Indeed, when we used pronase E, only 30% of atrial cells showed  $I_{CaT}$ , compared with 91% when collagenase and trypsin were used. A possible relationship between  $I_{CaT}$  and Ca paradox phenomenon might, a priori, be ruled out since the dissociation period in the low-Ca medium was 20–25 min when pronase was used, and 45 min with collagenase-trypsin. It has long been known that proteases affect ionic channels (see review by Narahashi, 1974). Thus, it has been shown that neuraminidase selectively enhances the transient Ca current in guinea pig myocytes (Yee, Weiss, and Langer, 1989). In squid giant axon, pronase specifically destroys the Na channel inactivation mechanism (Rojas and Armstrong, 1971). A pronase-derived alkaline proteinase b modifies the voltage dependence of the K channel from sarcoplasmic reticulum (Miller and Rosenberg, 1979). It has been recently shown in GH<sub>3</sub> cells that a protease-sensitive site within the Ca channel proteins must be phosphorylated to open the channel (Armstrong and Eckert, 1987). Finally, proteolytic enzymes such as trypsin and pronase markedly reduced the binding sites to indolizine sulfone, a new class of Ca antagonist (Chatelain, Beaufort, Meymans, and Clinet, 1991). It is then possible that pronase, in our case, or the rather long periods of enzymatic dissociation (up to 2.5 h) needed by Campbell et al. (1988a, b; see also Hume and Giles, 1981) might induce cellular changes leading to the disappearance of T-type channels. As for pronase, we cannot discard the possibility that collagenase and trypsin exert deleterious actions on Ca channels; however, 91% of atrial cells showed  $I_{CaT}$  when these enzymes were used. The small percent of atrial cells lacking  $I_{CaT}$  in these dissociations could also be related to the fact that these cells probably have their origins near the atrio-ventricular ring and are more "ventricular-like,"

having only  $I_{CaL}$  (Bonvallet and Rougier, 1989). As shown by Argibay et al. (1988) and the present results, frog ventricular cells do not exhibit  $I_{CaT}$ .

It is not easy to isolate  $I_{CaT}$  from  $I_{CaL}$  at every membrane potential. As already reported by others (e.g., Bean, 1989) Cd and Ni ions are far from being specific inhibitors of  $I_{CaL}$  and  $I_{CaT}$ , respectively; moreover, Cd block was shown to be weaker at more negative test potentials (Lansman, Hess, and Tsien, 1986). The difference  $I$ - $V$  relationship method (at two HPs) is also not fully reliable since in many cells a HP of  $-50$  mV (or even  $-60$  mV) inactivates a substantial fraction of the inward current flowing at potentials positive to  $-20$  mV (see also Schouten and Morad, 1989). Nevertheless, a reasonable picture can be formed on the basis of combining the results obtained with Cd, Ni, and the difference current method. In bullfrog atrial cells at physiological Ca concentrations,  $I_{CaT}$  and  $I_{CaL}$  activate at around  $-65$  and  $-30$  mV, respectively. Maximal  $I_{CaT}$  occurs around  $-35$  mV and maximal  $I_{CaL}$  at 0 to  $+10$  mV. The reversal potential for  $I_{CaL}$  is estimated to be near to  $+50$  mV and that for  $I_{CaT}$  between  $+20$  and  $+30$  mV. These values are very similar to those reported by others in frog atrial and ventricular cells (Argibay et al., 1988; Campbell et al., 1988b; Bonvallet and Rougier, 1989; Alvarez and Vassort, 1991).  $I_{CaL}$  inactivates between  $-40$  and 0 mV, with a potential for half-inactivation around  $-30$  mV. At positive prepulse potentials, a facilitation of  $I_{CaL}$  occurs which was attributed to Ca-dependent Ca inactivation (Mentrard et al., 1984; Lee et al., 1985; Argibay et al., 1988).  $I_{CaT}$ , on the other hand, inactivates in the potential range from  $-80$  to  $-50$  mV, with a potential for half-inactivation around  $-65$  mV; it also has a steeper voltage dependence than  $I_{CaL}$ . These observations are in full agreement with those of Droogmans and Nilius (1989) using single channel analysis. Ba substitution of Ca ions did not affect the inactivation time course of  $I_{CaT}$  and did not increase its amplitude. The voltage dependence of  $I_{CaT}$  availability remained unchanged in the presence of Ba. Reactivation kinetics (recovery from inactivation) of  $I_{CaT}$  was much slower and never showed an overshoot, as was commonly seen for  $I_{CaL}$ . These results indicate that inactivation of  $I_{CaT}$  is only voltage dependent as previously reported by others (Bean, 1985b; Bonvallet and Rougier, 1989; Hirano et al., 1989b).

Altogether the above reported characteristics of  $I_{CaT}$  confirm and extend previous results. T-type current amplitude and density in our cells were comparable to those reported by others in frog atrial cells (e.g., Bonvallet, 1987; Bonvallet and Rougier, 1989). Although variable, the ratio of  $I_{CaT}$  at  $-50$  mV to  $I_{CaL}$  at 0 mV was 1:5 as an average, a value that was similar to that reported by Tseng and Boyden (1989) for canine ventricular myocytes. Peak  $I_{CaT}$  at  $-30$  mV could be as large as half-peak  $I_{CaL}$ . On the other hand, a few cells exhibited very large  $I_{CaT}$  relative to  $I_{CaL}$ . They could be compared to embryonic cells (Kawano and de Haan, 1989) or to the sinus venosus cells (Bois and Lenfant, 1991), although they were not spontaneously beating. In most aspects our results obtained on  $I_{CaT}$  are similar to those reported in rabbit sinoatrial node cells by Hagiwara et al. (1988), in canine Purkinje cardiac cells by Hirano et al. (1989a), and in frog cardiac sinus venosus cells by Bois and Lenfant (1991). As indicated above, they documented that  $I_{CaT}$  kinetics were insensitive to Ca ions but depended on voltage (Bean, 1985b; Hirano et al., 1989b). However, our results show a specific characteristic of  $I_{CaT}$ : namely, that high prepolarizing steps did not fully inactivate  $I_{CaT}$  so that the availability curves bent up and the recoveries

from inactivation were biphasic. This remaining activation could be attributed to an additional, long-lasting open state induced by high depolarizations as early described for the Na current in nerve cells (Bezanilla and Armstrong, 1977) and recently reported for the cardiac L-type Ca channel (Pietrobon and Hess, 1990). Under their experimental conditions, the L-type current carried by Ba ions decayed exponentially with a time constant of  $\sim 100$  ms. In our experiments (Fig. 6 C), after high depolarizations peak  $I_{CaT}$  decayed with increasing test pulse intervals; the time constant of decays was markedly voltage dependent, varying from a few milliseconds at  $-120$  mV to 29 ms at  $-70$  mV; it was not significantly changed in the presence of Ba at low or high concentrations, or in the presence of Cd ions. The activation that remains after a high depolarization might have a second origin. In a recent publication (Slesinger and Lansman, 1991), neuronal L-type Ca channels were shown to reopen upon return to resting potential, which suggests that some channels recover from inactivation by passing through an open state. Extending this complementary hypothesis, it can be proposed that high depolarizations allow both cardiac T- and L-type Ca channels to reopen from inactivation. Reopenings would not occur below  $-30$  and  $+10$  mV for T- and L-type channels, respectively; however, above these values reopenings would increase with increasing membrane depolarizations. Such reopenings also depend on the duration of the prepulse as particularly illustrated by the reduction in  $I_{CaT}$  inactivation which occurred during predepolarization to  $+70$  mV lasting  $> 50$  ms (Fig. 7). In several aspects, including independence of ion carrier species and insensitivity to dihydropyridine Ca channel agonists, these results are comparable to the recently described double-pulse  $Ca^{2+}$  current facilitation in rat sympathetic neurons (Ikeda, 1991). This phenomenon thus represents a new characteristic of Ca currents.

A second specific observation relative to current kinetics and which also requires single-channel analysis is that Cd ions increased  $I_{CaT}$  time to peak and slowed its inactivation time course. The best available interpretation would be that Cd ions increased the first latency of openings of the T-type channel which has been proposed to control inactivation time course of the macroscopic T-type current (Droogmans and Nilius, 1989). Such a hypothesis would also account for the larger increase in peak current by BAY K 8644 in the presence of Cd, assuming that dihydropyridine antagonized the effects of Cd as suggested by the recovery of  $I_{CaT}$  inactivation kinetics under these conditions.

Another salient feature of this work is the finding that  $I_{CaT}$  was found to be more sensitive to dihydropyridines and neurotransmitters than previously reported (see, for example, Bean, 1985b; Nilius et al., 1985; Hagiwara et al., 1988; Bonvallet and Rougier, 1989; Hirano et al., 1989a; Tytgat et al., 1988; but see Tseng and Boyden, 1988, 1989, 1991), although a marked inhibition to 20% of  $I_{CaT}$  control by  $1 \mu M$  nifedipine was recently communicated (Romanin, Seydl, Glossmann, and Schindler, 1992). A decrease in both currents (18% for  $I_{CaT}$  and 49% for  $I_{CaL}$ ) is recorded with  $1 \mu M$  nifedipine under our experimental conditions. However, as previously reported for the effects of nisoldipine on canine Purkinje cells (Tseng and Boyden, 1989), the steady-state inactivation of  $I_{CaT}$  was barely affected by nifedipine, while that of  $I_{CaL}$  was displaced by 10 mV toward more negative potentials. This suggests different mechanisms of action of this dihydropyridine on the two Ca currents. The same is

true with the dihydropyridine agonist BAY K 8644, which induced a 10-mV shift of  $I_{CaL}$  kinetics and increased or decreased  $I_{CaL}$  according to the holding membrane potential (Sanguinetti et al., 1986; Kass, 1987). Such an effect was not observed on  $I_{CaT}$ . The absence of shifts on the  $I-V$  and availability curves and the fact that the increase in  $I_{CaT}$  was unchanged by varying HP between  $-100$  and  $-80$  mV suggest that the action of BAY K 8644 on  $I_{CaT}$  is not voltage dependent. Moreover, the fact that the  $I_{CaT}$ -availability curve was not shifted after addition of BAY K 8644, as with the inhibitory dihydropyridines, argues against the view that under such a condition the increase in  $I_{CaT}$  is artefactually related to  $I_{CaL}$  increase, since in such a case the  $I_{CaT}$ -availability curve would be shifted close to the  $I_{CaL}$ -availability curve. Our results reinforce the observations of Tseng and Boyden (1989) mentioned above, and of Mitra and Morad (1986), who noted that at high concentrations Ca agonists and antagonists also interact with the low threshold Ca current. Furthermore, Akaike et al. (1989) have shown that the low threshold Ca current in hypothalamic neurons is extremely sensitive to dihydropyridines and other organic blockers. They found, however, only a blocking effect of BAY K 8644 (at high concentrations,  $10 \mu\text{M}$ ) as already reported for  $I_{CaL}$  in cardiac cells (Bechem and Schramm, 1987). Finally, in a recent report (Wu, Wang, Karpinski, and Pang, 1992) it was shown that in neuroblastoma cells the effects of BAY K 8644 depend on the solvent; a 50% decrease of the T-type Ca current was observed in DMSO but not in ethanol with no shift of the availability curves in either case.

More surprising results were the specific responses of  $I_{CaT}$  and  $I_{CaL}$  to the neuromediators. As previously reported in frog cells, extracellular ATP increased  $I_{CaL}$  (Alvarez et al., 1990). The present results show that ATP also enhanced  $I_{CaT}$ . The effects were marked but not observed in every cell; they were obtained in a very narrow range of concentration.  $I_{CaT}$  was more sensitive to  $\alpha_1$ -adrenergic stimulation, while  $I_{CaL}$  was more sensitive to  $\beta$ -adrenergic stimulation. Such a small increase in  $I_{CaL}$  after  $\alpha_1$ -adrenergic stimulation has been previously reported in frog ventricular cells (Alvarez et al., 1987). As well, an increase in  $I_{CaT}$  by  $\alpha_1$  stimulation has been found in canine ventricular and Purkinje cells (Tseng and Boyden, 1988, 1989) and in smooth muscle cells of rat portal vein (Pacaud, Loirand, Mironneau, and Mironneau, 1987). Regarding the increase of  $I_{CaT}$  by  $\beta$ -adrenergic stimulation, our results are in agreement with those of Tseng and Boyden (1988) on canine ventricular myocytes and of Mitra and Morad (1986) in guinea pig ventricular cells, but differ from other reports on various cell types (Bean, 1985b; Tytgat et al., 1988; Bonvallet and Rougier, 1989). It should be noted that under our experimental conditions not only was  $I_{CaT}$  less sensitive to  $\beta$  stimulation than  $I_{CaL}$ , but the effects of  $\beta$  stimulation on  $I_{CaT}$  as well as those of the other neuromediators were not seen in every cell studied. These variations may arise from species and/or tissue specificity as well as from alterations in the state of the membrane following dissociation procedure. As we discussed above, T channels in frog atrial cells can be affected by the enzymatic dissociation to a larger extent than  $I_{CaL}$ . Whatever the case, with each of the three agonists the observed increases in  $I_{CaT}$  should not be artefacts related to increases in  $I_{CaL}$ : first, a depolarizing pulse to  $-50$  mV was used to activate this current, well below  $I_{CaL}$  threshold, and second, similar relative increases were also observed after inhibition of  $I_{CaL}$  by Cd ions. The exact mechanisms of action of the

neuromediators on  $I_{CaT}$  require further investigation. Evidence exists in canine Purkinje cells that  $I_{CaT}$  can be transiently increased by raising intracellular Ca concentration (Tseng and Boyden, 1991). Such a mechanism might account for the weak increase in  $I_{CaT}$  seen in some experiments after adding intracellular cAMP (Fig. 11 B). However, it could hardly be involved in most aspects of this study on the positive effects of neuromediators as well as on the partial inactivation after high depolarizing prepulses, since very similar effects were observed when  $I_{CaL}$ , the major origin of intracellular Ca load, was inhibited by Cd (see Figs. 6 B, 8 D, and 10 A) or when  $I_{CaL}$  had been markedly increased by cAMP (Fig. 11 B). According to the rather fast changes in peak current, intracellular messengers might not be involved, but direct coupling by G proteins should be considered (Brown and Birnbaumer, 1988) as well as membrane perturbation similar to that obtained with alcohols or anesthetics (Mongo and Vassort, 1990).

Our experiments with ouabain confirm those of Fischmeister et al. (1986) on frog (*R. esculenta*) single ventricular cells and of Legrand, Deroubaix, Coulombe, and Coraboeuf (1990) on guinea pig ventricular myocytes. They indicate transient effects of cardiac glycosides on  $I_{CaL}$  while the increase in  $I_{CaT}$  was sustained. As postulated by Fischmeister et al. (1986), the inhibition of  $I_{CaL}$  by cardiac glycosides could be caused by an increase in free intracellular Ca (consecutive to Na-K pump inhibition) leading to a decrease in availability of  $I_{CaL}$ . The difference in behavior of  $I_{CaT}$  under the action of ouabain may then be related to the fact that  $I_{CaT}$  does not exhibit a Ca-dependent inactivation mechanism.

#### *Physiological Consequences*

The physiological role of  $I_{CaT}$  in heart cells remains uncertain. However, it might have major electrophysiological implications. From a mathematical simulation model, Nilius (1986) suggested that  $I_{CaT}$  plays an essential role in the pacemaker potential.  $I_{CaT}$  activation is considered to be responsible for the last part of the diastolic depolarization (Hagiwara et al., 1988). It has also been suggested that  $I_{CaT}$  participates in certain arrhythmias related to reentry due to slowly propagating action potential (Carmeliet, 1988) or to delayed after-depolarizations (Hirano et al., 1989a).

It is well known that ouabain and other cardiac glycosides may induce early and delayed after-depolarizations (Cranefield, 1975). Although the cellular basis for their initiation is not yet well established (for review, see January and Fozzard, 1988), we may suggest that ouabain-induced changes in  $I_{CaL}$  and  $I_{CaT}$  could, in part, contribute to the development of early and delayed after-depolarizations, respectively. It has been similarly shown that  $I_{CaL}$  plays a role in the early after-depolarizations induced by BAY K 8644 (January and Riddle, 1989).

$\alpha_1$ -Adrenergic stimulation whose responsivity is increased during myocardial ischemia and reperfusion contributes to the development of malignant ventricular arrhythmias. Receptor blockade has been shown to be effective in restoring sinus rhythm after such arrhythmias have been experimentally induced (Sheridan, Penkoske, Sobel, and Corr, 1980; Davey, 1986). Such arrhythmias may be consequent to an inward current activated by increased cytoplasmic calcium; the latter could be triggered following the activation of the phosphoinositol turnover by  $\alpha_1$ -adrenergic

agonists. Arrhythmias will be facilitated by the fact that these agonists enhanced the T-type calcium current as described above. Furthermore, since P<sub>2</sub>-purinergic agonists also increase phosphoinositol turnover (Legssyer, Poggioli, Renard, and Vassort, 1988) as well as facilitate  $I_{CaT}$ , it should be anticipated that P<sub>2</sub>-purinergic stimulation could also lead to arrhythmia.

In conclusion, our study reports the existence of T-type Ca current on frog atrial cells. This current differs from the L-type Ca current by the voltage dependence of its kinetics, by the fact that Ba does not alter its kinetics, and by its lower sensitivity to Cd. Both currents are sensitive to pharmacological agents and neuromediators, but each with their own characteristics. In a few cells, T-type Ca current was larger than the L-type, but generally its maximal amplitude was about half; nevertheless, this suggests that it might have a major influence on electrical cell activity. Some atrial cells did not exhibit T-type Ca current under whole-cell patch-clamp conditions; in most cases this could be attributed to the enzymatic treatment during the dissociation procedure.

Many thanks are due to Dr. Lidia Rodriguez-Nande for her skillful assistance, to Dr. Lourdes Rubio for participating in some experiments, to Drs. Rodolphe Fischmeister, Pierre-François Mery, and Frédérique Scamps for their helpful suggestions, to Dr. Ian Findlay for advice on the manuscript, and to Denise Reuter for secretarial help.

This work was supported by an exchange program between the University of La Habana and the University of Orsay.

*Original version received 21 October 1991 and accepted version received 5 June 1992.*

#### REFERENCES

- Akaike, N., P. G. Kostyuk, and Y. V. Osipchuk. 1989. Dihydropyridine-sensitive low-threshold calcium channels in isolated rat hypothalamic neurones. *Journal of Physiology*. 412:333–349.
- Alvarez, J. L., K. G. Mongo, F. Scamps, and G. Vassort. 1990. Effects of purinergic stimulation on the Ca current in single frog cardiac cells. *Pflügers Archiv*. 416:189–195.
- Alvarez, J. L., K. G. Mongo, and G. Vassort. 1987. Effects of  $\alpha_1$ -adrenergic stimulation on the Ca current in single ventricular frog cells. *Journal of Physiology*. 390:66P. (Abstr.)
- Alvarez, J. L., and G. Vassort. 1991. Dual action of prajmalium on the Ca currents in frog isolated cardiomyocytes. *Journal of Molecular and Cellular and Cardiology*. 23:627–638.
- Argibay, J. A., R. Fischmeister, and H. C. Hartzell. 1988. Inactivation, reactivation and pacing dependence of calcium current in frog cardiocytes: correlation with current density. *Journal of Physiology*. 401:202–226.
- Armstrong, D., and R. Eckert. 1987. Voltage-activated calcium channels that must be phosphorylated to respond to membrane depolarization. *Proceedings of the National Academy of Sciences, USA*. 84:2518–2522.
- Arrio-DuPont, M., and D. De Nay. 1985. High yield preparation of calcium tolerant myocytes from frog ventricles: some properties of the isolated cells. *Biology of the Cell*. 54:163–170.
- Bean, B. P. 1985a. Two kinds of calcium channels in atrial cells from dog and frog hearts. *Biophysical Journal*. 47:498a. (Abstr.)
- Bean, B. P. 1985b. Two kinds of calcium channels in canine atrial cells. Differences in kinetics, selectivity and pharmacology. *Journal of General Physiology*. 86:1–30.
- Bean, B. P. 1989. Classes of calcium channels in vertebrate cells. *Annual Review of Physiology*. 51:367–384.



- Bechem, M., and M. Schramm. 1987. Calcium agonists. *Journal of Molecular and Cellular Cardiology*. 19:63–76.
- Bezanilla, F., and C. M. Armstrong. 1977. Inactivation of the sodium channel. *Journal of General Physiology*. 70:549–566.
- Bois, P., and J. Lenfant. 1991. Evidence for two types of calcium currents in frog cardiac sinus venosus cells. *Pflügers Archiv*. 417:591–596.
- Bonvallet, R. 1987. A low threshold calcium current recorded at physiological Ca concentrations in single frog atrial cells. *Pflügers Archiv*. 408:540–542.
- Bonvallet, R., and O. Rougier. 1989. Existence of two calcium currents recorded at normal calcium concentrations in single frog atrial cells. *Cell Calcium*. 10:499–509.
- Bossu, J.-L., and A. Feltz. 1986. Inactivation of the low-threshold transient calcium current in rat sensory neurones: evidence for a dual process. *Journal of Physiology*. 376:341–357.
- Brown, A. M., and L. Birnbaumer. 1988. Direct G protein gating of ion channels. *American Journal of Physiology*. 254:H401–H410.
- Brum, G., W. Osterreider, and W. Trautwein. 1984.  $\beta$ -adrenergic increase in the calcium conductance of cardiac myocytes studied with the patch-clamp. *Pflügers Archiv*. 406:340–342.
- Campbell, D. L., W. R. Giles, J. R. Hume, and E. F. Shibata. 1988a. Inactivation of calcium current in bull-frog atrial myocytes. *Journal of Physiology*. 403:287–315.
- Campbell, D. L., W. R. Giles, and E. F. Shibata. 1988b. Ion transfer characteristics of the calcium current in bull-frog atrial myocytes. *Journal of Physiology*. 403:239–266.
- Carmeliet, E. 1988. Role of calcium channels in cardiac arrhythmias. In *The Calcium Channel: Structure, Functions and Implications*. M. Morad, W. Nayler, S. Kazda, and M. Schramm, editors. Springer Verlag, Berlin. 310–316.
- Charnet, P., S. Richard, A. M. Gurney, H. Ouadid, F. Tiaho, and J. Nargeot. 1991. Modulation of Ca currents in isolated frog atrial cells studied with photosensitive probes. Regulation by cAMP and  $Ca^{2+}$ : a common pathway? *Journal of Molecular and Cellular Cardiology*. 23:343–356.
- Chatelain, P., P. Beaufort, L. Meymans, and M. Clinet. 1991. Characterization of the slow calcium channel binding sites for [ $^3H$ ] SR 33557 in rat heart sarcolemmal membranes. *Molecular Pharmacology*. 39:64–71.
- Cranefield, P. F. 1975. *The Conduction of the Cardiac Impulse: The Slow Response and Cardiac Arrhythmias*. Futura Publishing Co., Mount Kisco, NY. 1–404.
- Davey, M. J. 1986. Alpha-adrenoceptors: an overview. *Journal of Molecular and Cellular Cardiology*. 18:1–15.
- Deslauriers, Y., E. Ruiz-Ceretti, O. F. Schanne, and M. D. Payet. 1982. The toxic effects of ouabain: a voltage clamp study. *Canadian Journal of Physiology and Pharmacology*. 60:1153–1159.
- Dowell, R. T., and M. Tarr. 1988. Contractile and biochemical properties of rat cardiac myocytes isolated using nonspecific protease. In *Biology of Isolated Adult Cardiac Myocytes*. W. A. Clark, K. S. Decker, and T. K. Borg, editors. Elsevier Science Publishing Co., Inc., New York. 390–393.
- Droogmans, G., and B. Nilius. 1989. Kinetic properties of the cardiac T-type calcium channel in the guinea-pig. *Journal of Physiology*. 419:627–650.
- Fischmeister, R., M. Brocas-Randolph, P. Lechene, J. A. Argibay, and G. Vassort. 1986. A dual effect of cardiac glycosides on Ca current in single cells of frog heart. *Pflügers Archiv*. 406:340–342.
- Hagiwara, N., H. Irisawa, and M. Kameyama. 1988. Contribution of two types of calcium currents to the pacemaker potentials of rabbit sino-atrial node cell. *Journal of Physiology*. 395:233–253.
- Hirano, Y., H. A. Fozzard, and C. T. January. 1989a. Characteristics of L- and T-type  $Ca^{2+}$  currents in canine cardiac Purkinje cells. *American Journal of Physiology*. 256:H1478–H1492.
- Hirano, Y., H. A. Fozzard, and C. T. January. 1989b. Inactivation properties of T-type calcium current in canine cardiac Purkinje cells. *Biophysical Journal*. 56:1007–1016.

- Hume, J. R., and W. Giles. 1981. Active and passive electrical properties of single bullfrog atrial cells. *Journal of General Physiology*. 78:18–43.
- Hume, J. R., and W. Giles. 1983. Ionic currents in single isolated bullfrog atrial cells. *Journal of General Physiology*. 81:153–194.
- Ikeda, S. R. 1991. Double-pulse calcium channel current facilitation in adult rat sympathetic neurones. *Journal of Physiology*. 439:181–214.
- January, C. T., and H. A. Fozzard. 1988. Delayed afterdepolarizations in heart muscle: mechanisms and relevance. *Pharmacological Reviews*. 40:219–227.
- January, C. T., and J. M. Riddle. 1989. Early after depolarizations: mechanism of induction and block. A role for L-type  $\text{Ca}^{2+}$  current. *Circulation Research*. 64:977–990.
- Kass, R. S. 1987. Voltage-dependent modulation of cardiac calcium channel current by optical isomers of BAY K 8644: implications for channel gating. *Circulation Research*. 6:11–15.
- Kawano, S., and R. L. De Haan. 1989. Low threshold current is the major calcium-current in chick ventricle cells. *American Journal of Physiology*. 256:H1505–H1508.
- Lansman, J. B., P. Hess, and R. W. Tsien. 1986. Blockade of current through single calcium channels by  $\text{Cd}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{Ca}^{2+}$ . Voltage and concentration dependence of calcium entry into the pore. *Journal of General Physiology*. 88:321–347.
- Lee, K. S., E. Marban, and R. W. Tsien. 1985. Inactivation of calcium channels in mammalian heart cells: joint dependence on membrane potential and intracellular calcium. *Journal of Physiology*. 364:395–411.
- Legrand, B., E. Deroubaix, A. Coulombe, and E. Coraboeuf. 1990. Stimulatory effect of ouabain on T- and L-type calcium currents in guinea pig cardiac myocytes. *American Journal of Physiology*. 258:H1620–H1623.
- Legssyer, A., J. Poggioli, D. Renard, and G. Vassort. 1988. ATP and other adenine compounds increase mechanical activity and inositol trisphosphate production in rat heart. *Journal of Physiology*. 401:185–199.
- Marban, E., and R. W. Tsien. 1982. Enhancement of calcium current during digitalis inotropy in mammalian heart: positive feed-back regulation by intracellular calcium? *Journal of Physiology*. 329:589–614.
- Mentrard, D., G. Vassort, and R. Fischmeister. 1984. Calcium-mediated inactivation of the Ca conductance in cesium-loaded frog heart cells. *Journal of General Physiology*. 83:105–131.
- Miller, C., and R. L. Rosenberg. 1979. Modification of a voltage-gated  $\text{K}^{+}$  channel from sarcoplasmic reticulum by a pronase-derived specific endopeptidase. *Journal of General Physiology*. 74:457–478.
- Mitra, R., and M. Morad. 1986. Two types of calcium channels in guinea pig ventricular myocytes. *Proceeding of the National Academy of Sciences, USA*. 83:5340–5344.
- Mongo, K. G., and G. Vassort. 1990. Inhibition by alcohols, halothane and chloroform of the Ca current in single frog ventricular cells. *Journal of Molecular and Cellular Cardiology*. 22:939–953.
- Narahashi, T. 1974. Chemicals as tools in the study of excitable membranes. *Physiological Reviews*. 54:813–889.
- Nilius, B. 1986. Possible functional significance of a novel type of cardiac Ca channel. *Biomedica et Biochemica Acta*. 45:K37–45.
- Nilius, B., P. Hess, J. B. Lansman, and R. W. Tsien. 1985. A novel type of cardiac calcium channel in ventricular cells. *Nature*. 316:443–446.
- Pacaud, P., G. Loirand, C. Mironneau, and J. Mironneau. 1987. Opposing effects of noradrenaline on the two classes of voltage-dependent calcium channels of single vascular smooth muscle cells in short term primary culture. *Pflügers Archiv*. 410:557–559.
- Pietrobon, D., and P. Hess. 1990. Novel mechanism of voltage-dependent gating in L-type calcium channels. *Nature*. 346:651–655.

- Rojas, E., and C. Armstrong. 1971. Sodium conductance activation without inactivation in pronase-perfused axons. *Nature New Biology*. 229:177–178.
- Romanin, C., K. Seydl, H. Glossmann, and H. Schindler. 1992. The dihydropyridine nifedipine inhibits T-type Ca<sup>2+</sup> currents in atrial myocytes. *Pflügers Archiv*. 420:410–412.
- Rougier, O., G. Vassort, D. Garnier, Y. M. Gargouil, and E. Coraboeuf. 1969. Existence and role of a slow inward current during the frog atrial action potential. *Pflügers Archiv*. 308:91–110.
- Sanguinetti, M. C., D. S. Krafte, and R. S. Kass. 1986. Voltage-dependent modulation of Ca channel current in heart cells. *Journal of General Physiology*. 88:369–392.
- Scamps, F., A. Legssyer, E. Mayoux, and G. Vassort. 1990. The mechanism of positive inotropy induced by adenosine triphosphate in rat heart. *Circulation Research*. 67:1007–1016.
- Schouten, V. J. A., and M. Morad. 1989. Regulation of Ca current in frog ventricular myocytes by the holding potential, c-AMP and frequency. *Pflügers Archiv*. 415:1–11.
- Sheridan, D. J., P. A. Penkoske, B. E. Sobel, and P. B. Corr. 1980. Alpha-adrenergic contributions to dysrhythmia during myocardial ischemia and reperfusion in cats. *Journal of Clinical Investigation*. 65:161–171.
- Slesinger, P. A., and J. B. Lansman. 1991. Reopening of Ca<sup>2+</sup> channels in mouse cerebellar neurons at resting membrane potentials during recovery from inactivation. *Neuron*. 7:755–762.
- Tseng, G. N. 1988. Calcium current restitution in mammalian ventricular myocytes is modulated by intracellular calcium. *Circulation Research*. 63:468–482.
- Tseng, G. N., and P. A. Boyden. 1988. Heterogeneous properties of Ca currents in isolated canine Purkinje and ventricular cells. Symposium on Molecular and Cellular Mechanisms of Antiarrhythmic Agents. Nashville, TN. T21-T22. (Abstr.)
- Tseng, G. N., and P. A. Boyden. 1989. Multiple types of Ca<sup>2+</sup> currents in single canine Purkinje cells. *Circulation Research*. 65:1735–1750.
- Tseng, G. N., and P. A. Boyden. 1991. Different effects of intracellular Ca and protein kinase C on cardiac T and L Ca currents. *American Journal of Physiology*, 261:H364–H379.
- Tytgat, J., B. Nilius, J. Vereecke, and B. Carmeliet. 1988. The T-type Ca channel in guinea pig ventricular myocytes is insensitive to isoproterenol. *Pflügers Archiv*. 411:704–706.
- Wu, L., R. Wang, E. Karpinski, and P. K. T. Pang. 1992. Bay K-8644 in different solvents acts as a transient calcium channel antagonist and a long-lasting calcium channel agonist. *The Journal of Pharmacology and Experimental Therapeutics*. 260:966–973.
- Yee, H. F., Jr., J. N. Weiss, and G. A. Langer. 1989. Neuraminidase selectively enhances transient Ca<sup>2+</sup> current in cardiac myocytes. *American Journal of Physiology*. 256:C1267–C1272.