

ERRATUM**Erratum to: The *in vitro* toxicity of venoms from South Asian Hump-nosed pit vipers (Viperidae: *Hypnale*)**

Kalana Maduwage^α, Wayne C Hodgson^β, Nicki Konstantakopoulos^β, Margaret A O'Leary^λ, Indika Gawarammana^φ, Geoffrey K Isbister^{β,λ,ψ,*}

^αDepartment of Biochemistry, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka, ^βMonash Venom Group, Department of Pharmacology, Monash University, Victoria, 3800 Australia, ^λDepartment of Clinical Toxicology and Pharmacology, Calvary Mater Newcastle, Newcastle, New South Wales, Australia, ^φDepartment of Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka, ^ψDiscipline of Clinical Pharmacology, University of Newcastle, New South Wales, Australia

*Correspondence to: Geoffrey Isbister, Email: Geoff.isbister@gmail.com, Tel: +612 4921 1211, Fax: +612 4921 1870

Erratum published online: 18 June 2011

© Copyright The Author(s) Published by Library Publishing Media. This is an open access article, published under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.5>). This license permits non-commercial use, distribution and reproduction of the article, provided the original work is appropriately acknowledged with correct citation details.

Erratum

Some labels were omitted in the published Figure 1, which have been included in this erratum.




Figure 1. Reverse phase high-performance liquid chromatogram (RP-HPLC) chromatogram of (A) *H. hypnale*, (B) *H. nepa* and (C) *H. zara* using a Phenomenex Jupiter C18 column (5 μ 300A 250 \times 4.6mm) with mobile phase 15% (v/v) acetonitrile containing 0.1% (v/v) trifluoroacetic acid, increasing to 70% acetonitrile by t=100min, at a flow rate of 0.5ml/min.