

INVITED SPEAKER PRESENTATION

Open Access

S76. Adoptive cell therapy of melanoma with autologous tumour infiltrating lymphocytes

I Svane*, R Andersen, M Donia

From 1st Immunotherapy of Cancer Conference (ITOC1)
Munich, Germany. 12-14 March 2014

Background

Adoptive T-cell therapy (ACT) with tumor infiltrating lymphocytes (TILs) is a personalized treatment for cancer defined as the infusion of T cells isolated from the patient's own tumor tissue after ex vivo activation and several rounds of expansion. This treatment has achieved impressive clinical results in several single institution phase I/II clinical trials performed outside Europe, and holds the promise to enter the mainstream of standard melanoma care in the near future. However, although transient, the toxicities associated with high-dose IL-2 classically administered together with TILs are severe and recent data have questioned its use.

Material and methods

In an ongoing phase I/II study, we have enrolled patients with progressive metastatic melanoma. TILs infusion was preceded by standard lymphodepleting chemotherapy but followed by low-dose subcutaneous IL-2 for 14 days or by an intravenous intermediate dose IL-2 decrescendo regimen.

Results

The lower doses of IL-2 considerably decreased the toxicity of the treatment while PET-CT imaging showed a preserved objective response rate of 48% including long-lasting complete responders. The absolute number of tumour specific T-cells infused was significantly associated to clinical response, with induction of peripheral tumour reactive T cells. To characterize the fate of TILs after infusion, we performed a longitudinal analysis of bulk tumor-reactive T cells from in vitro cultured TILs and found durable persistent T-cells after up to 1.5 years after infusion. Extensive analysis of disease relapse in two patients revealed multiple potential mechanisms of

tumor recurrence, linked either to tumor-changes or immune response decline.

Conclusion

Despite its clinical efficacy, with impressive response rates and a significant fraction of long surviving complete responders, the implementation of TIL based ACT into current practice has been severely hampered by the technical complexity of cell production, the toxicity profile demanding treatment at specialize cancer centers, and lack of investment from the pharmaceutical industry. Next step should be a pivotal phase III trial in melanoma which is required for regulatory approval. Further improvement of the therapy could also be pursued through combination treatment.

Published: 12 March 2014

doi:10.1186/2051-1426-2-S2-114

Cite this article as: Svane et al.: S76. Adoptive cell therapy of melanoma with autologous tumour infiltrating lymphocytes. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 2):114.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Copenhagen University Hospital, Department of Oncology Center for Cancer Immune Therapy, Herlev, Denmark



© 2014 Svane et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.