

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## (3*aRS*,7*aSR*)-7*a*-Methoxy-2-oxo-2,3,3*a*,4,5,6,7,7*a*-octahydro-1-benzofuran-4,4-dicarbonitrile

María González,<sup>a\*</sup> Andrea Martínez,<sup>a</sup> Marcos L. Rivadulla<sup>a</sup> and Berta Covelo<sup>b</sup>

<sup>a</sup>Dpto. Química Orgánica, Faculdade de Química, Universidade de Vigo, E-36310 Vigo, Spain, and <sup>b</sup>Unidad de Difracción de Raios X de Monocristal, Servicio Determinación Estructural, Proteómica e Xenómica, CACTI-Universidade de Vigo, E-36310 Vigo, Spain

Correspondence e-mail: magonpe@uvigo.es


Received 6 November 2012; accepted 26 November 2012

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.048;  $wR$  factor = 0.143; data-to-parameter ratio = 13.0.

The racemic title compound,  $\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_3$ , contains a [4.3.0]bicyclic unit in which the shared C—C bond adopts a *cis* configuration. The five- and six-membered rings are in twisted envelope (with the bridgehead C atom bearing the methoxy substituent as the flap) and distorted chair conformations, respectively. In the crystal, the molecules are linked *via* weak C—H $\cdots$ O interactions, forming ladder-like chains along [010].

### Related literature

For related syntheses of natural products, see: Jones & Goodbrand (1977). For details of a synthesis using different starting materials, see: Alonso *et al.* (2005); Pérez *et al.* (2004, 2005). For a related structure, see: Grudniewska *et al.* (2011). For puckering parameters, see, Cremer & Pople (1975).



### Experimental

#### Crystal data

$\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_3$   
 $M_r = 220.23$   
 Monoclinic,  $P2_1/n$

$a = 11.816$  (3) Å  
 $b = 7.228$  (2) Å  
 $c = 13.017$  (4) Å

$\beta = 104.250$  (5)°  
 $V = 1077.6$  (5) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.10$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.49 \times 0.24 \times 0.19$  mm

#### Data collection

Bruker SMART 1000 CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.697$ ,  $T_{\max} = 0.745$

5473 measured reflections  
 1893 independent reflections  
 1244 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.143$   
 $S = 1.03$   
 1893 reflections

146 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.45$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.19$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

D—H $\cdots$ A	D—H	H $\cdots$ A	D $\cdots$ A	D—H $\cdots$ A
C3—H3A $\cdots$ O3 <sup>i</sup>	0.97	2.59	3.290 (3)	129
C8—H8C $\cdots$ O2 <sup>ii</sup>	0.96	2.46	3.219 (4)	136

Symmetry codes: (i)  $-x + 1, -y + 2, -z + 1$ ; (ii)  $x, y + 1, z$ .

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2011) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

This work was supported financially by the Spanish Ministry of Foreign Affairs and Cooperation (PCIA/030052/10) and the Xunta de Galicia (INCITE845B-2010/020, INCI-TE08PXIB314255PR).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2158).

### References

- Alonso, D., Pérez, M., Gómez, G., Covelo, B. & Fall, Y. (2005). *Tetrahedron*, **61**, 2021–2026.
- Bruker (1998). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Grudniewska, A., Dancewicz, K., Bialońska, A., Ciunik, Z., Gabrys, B. & Wawrzęczyk, C. (2011). *Advances*, **1**, 498–510.
- Jones, J. B. & Goodbrand, H. B. (1977). *Can. J. Chem.* **55**, 2685–2691.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Pérez, M., Canoa, P., Gómez, G., Teijeira, M. & Fall, Y. (2005). *Synthesis*, pp. 411–414.
- Pérez, M., Canoa, P., Gómez, G., Terán, C. & Fall, Y. (2004). *Tetrahedron Lett.* **45**, 5207–5209.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supplementary materials

*Acta Cryst.* (2012). E68, o3496 [doi:10.1107/S1600536812048519]

**(3a*RS*,7a*SR*)-7a-Methoxy-2-oxo-2,3,3a,4,5,6,7,7a-octahydro-1-benzofuran-4,4-dicarbonitrile**

**María González, Andrea Martínez, Marcos L. Rivadulla and Berta Covelo**

**Comment**

The racemic title compound is a [4.3.0] bicyclic  $\gamma$ -lactone obtained through an intramolecular Michael addition. The construction of carbocyclic systems is of paramount importance in organic synthesis since they are intermediate compounds for the preparation of interesting natural products (Jones *et al.*, 1977) as carbocyclic nucleosides and vitamin D analogues. We have described a new methodology for the synthesis of oxacyclic compounds using either methoxyallene (Alonso *et al.*, 2005) or furan (Pérez *et al.*, 2005) as a starting material. In the title compound (Fig. 1), the C—C share bond of the bicyclic moiety adopts a *cis* configuration. The 5-membered ring adopts a twisted envelope conformation with puckering parameters  $Q = 0.358$  (2) Å and  $\varphi = 126.7$  (4)° (Cremer & Pople, 1975) and the 6-membered ring adopts a distorted chair conformation with puckering parameters  $Q = 0.531$  (3) Å,  $\theta = 19.3$  (3)° and  $\varphi = 123.3$  (9)° (Cremer & Pople (1975). All bond lengths and bond angles are normal comparable to those observed in similar crystal structures (Grudniewska *et al.*, 2011). In the crystal structure, the molecules are self-assembled *via* weak C—H $\cdots$ O intermolecular interactions (Table 1) to form a ladderlike chain structure along [010] (Fig. 2).

**Experimental**


To a solution of 2-(3-(2-methoxy-5-oxo-2,5-dihydrofuran-2-yl)propyl)malononitrile (0.38 mmol) in DMF (3 ml) was added DBU (0.19 mmol, 0.5 eq) and the mixture was stirred at room temperature. At the end of the reaction (TLC), EtOAc (20 ml) was added and the organic layers washed with water (3 x 20 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to afford a residue, which was chromatographed on silica gel giving the title compound. It was then recrystallized using EtOAc/Hexane.

**Refinement**


All H atoms were positioned geometrically and refined using a riding model with C—H = 0.96–0.98 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ .

**Computing details**

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINTE* (Bruker, 1998); data reduction: *SAINTE* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2011) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound. Non-H atoms are present as displacement ellipsoids at the 30% probability level.

**Figure 2**

The crystal structure of the title compound. Weak C—H...O intermolecular interactions link the molecules into chains along [001]. H atoms not involved in hydrogen bonds (dashed lines) have been omitted for clarity.

(3aRS,7aSR)-7a-Methoxy-2-oxo-2,3,3a,4,5,6,7,7a-octahydro- 1-benzofuran-4,4-dicarbonitrile

Crystal data

$C_{11}H_{12}N_2O_3$	$F(000) = 464$
$M_r = 220.23$	$D_x = 1.357 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2_1n$	Cell parameters from 1285 reflections
$a = 11.816 (3) \text{ \AA}$	$\theta = 2.7\text{--}22.8^\circ$
$b = 7.228 (2) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 13.017 (4) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 104.250 (5)^\circ$	Prism, colourless
$V = 1077.6 (5) \text{ \AA}^3$	$0.49 \times 0.24 \times 0.19 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART 1000 CCD diffractometer	5473 measured reflections
Radiation source: fine-focus sealed tube	1893 independent reflections
Graphite monochromator	1244 reflections with $I > 2\sigma(I)$
$\varphi$ and $\omega$ scans	$R_{\text{int}} = 0.030$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 25.0^\circ$ , $\theta_{\text{min}} = 2.1^\circ$
$T_{\text{min}} = 0.697$ , $T_{\text{max}} = 0.745$	$h = -13 \rightarrow 14$
	$k = -8 \rightarrow 8$
	$l = -14 \rightarrow 15$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.143$	$w = 1/[\sigma^2(F_o^2) + (0.0722P)^2 + 0.3369P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
1893 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
146 parameters	$\Delta\rho_{\text{max}} = 0.45 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

**Experimental.** [mp: 175–177 °C; IR (neat): 2248, 1800, 1744  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$ : 3.38 (3H, s, OMe); 3.21 (1H, dd,  $J = 7.60, 17.78$ ); 2.86 (1H, dd,  $J = 3.53, 7.60$ ); 2.70 (1H, dd,  $J = 3.53, 17.78$ ); 2.39–2.33 (2H, m); 2.16–1.80 (4H, m);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$ : 171.69 (C=O); 114.67 (CN); 112.89 (CN); 105.51 (C); 50.05 (OMe); 45.74 (CH); 35.79 (CH<sub>2</sub>); 33.59 (CH<sub>2</sub>); 31.89 (CH<sub>2</sub>); 28.56 (CH<sub>2</sub>); 17.68 (CH<sub>2</sub>); HRMS: calcd for  $C_{11}H_{13}N_2O_3 [M+1H]^+$  221.0926, found 221.0583].

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.67913 (13)	0.7856 (2)	0.36232 (13)	0.0423 (5)

O2	0.67361 (18)	0.5064 (3)	0.43402 (17)	0.0660 (6)
O3	0.63413 (13)	1.0294 (2)	0.45900 (12)	0.0401 (4)
N41	0.2043 (2)	0.7955 (4)	0.1943 (2)	0.0671 (7)
N42	0.5146 (2)	0.5424 (4)	0.1555 (2)	0.0700 (8)
C2	0.6249 (2)	0.6485 (4)	0.4044 (2)	0.0454 (6)
C3	0.5029 (2)	0.7069 (3)	0.40461 (19)	0.0415 (6)
H3A	0.4961	0.7318	0.4761	0.050*
H3B	0.4467	0.6125	0.3731	0.050*
C3A	0.48499 (18)	0.8826 (3)	0.33778 (17)	0.0344 (6)
H3	0.4371	0.9708	0.3658	0.041*
C4	0.42864 (19)	0.8445 (3)	0.21834 (18)	0.0394 (6)
C5	0.4493 (2)	1.0080 (4)	0.14904 (19)	0.0464 (7)
H5A	0.4119	0.9828	0.0753	0.056*
H5B	0.4150	1.1195	0.1698	0.056*
C6	0.5794 (2)	1.0369 (4)	0.1619 (2)	0.0508 (7)
H6A	0.5916	1.1360	0.1155	0.061*
H6B	0.6137	0.9250	0.1415	0.061*
C7	0.6392 (2)	1.0854 (4)	0.27642 (19)	0.0453 (6)
H7A	0.7230	1.0856	0.2845	0.054*
H7B	0.6164	1.2097	0.2911	0.054*
C7A	0.61086 (19)	0.9552 (3)	0.35683 (18)	0.0352 (6)
C8	0.7522 (2)	1.0884 (4)	0.5017 (2)	0.0520 (7)
H8A	0.8048	0.9939	0.4902	0.078*
H8B	0.7645	1.1105	0.5764	0.078*
H8C	0.7662	1.2004	0.4672	0.078*
C42	0.4769 (2)	0.6739 (4)	0.1824 (2)	0.0483 (7)
C41	0.3020 (2)	0.8146 (4)	0.2047 (2)	0.0460 (6)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0349 (9)	0.0400 (10)	0.0545 (10)	0.0055 (7)	0.0155 (8)	-0.0029 (8)
O2	0.0677 (13)	0.0411 (12)	0.0881 (16)	0.0172 (10)	0.0173 (11)	0.0123 (10)
O3	0.0359 (9)	0.0462 (10)	0.0375 (9)	-0.0036 (7)	0.0078 (7)	-0.0089 (7)
N41	0.0405 (14)	0.086 (2)	0.0718 (17)	-0.0063 (13)	0.0091 (12)	-0.0035 (14)
N42	0.0580 (16)	0.0740 (19)	0.0745 (18)	0.0034 (14)	0.0096 (13)	-0.0302 (15)
C2	0.0500 (15)	0.0367 (15)	0.0489 (15)	0.0027 (12)	0.0113 (12)	-0.0013 (12)
C3	0.0409 (14)	0.0411 (14)	0.0440 (14)	-0.0036 (11)	0.0133 (11)	-0.0006 (12)
C3A	0.0316 (12)	0.0379 (13)	0.0357 (13)	0.0007 (10)	0.0122 (10)	-0.0021 (10)
C4	0.0320 (12)	0.0472 (15)	0.0388 (13)	0.0008 (10)	0.0080 (10)	-0.0043 (11)
C5	0.0462 (15)	0.0574 (17)	0.0358 (14)	0.0016 (12)	0.0102 (11)	0.0048 (11)
C6	0.0494 (16)	0.0640 (19)	0.0424 (15)	-0.0038 (13)	0.0177 (12)	0.0042 (13)
C7	0.0438 (14)	0.0470 (15)	0.0468 (15)	-0.0060 (12)	0.0146 (11)	0.0027 (12)
C7A	0.0330 (12)	0.0357 (13)	0.0384 (13)	0.0012 (10)	0.0113 (10)	-0.0041 (10)
C8	0.0449 (15)	0.0523 (16)	0.0527 (16)	-0.0055 (13)	0.0003 (12)	-0.0030 (13)
C42	0.0392 (14)	0.0572 (18)	0.0473 (15)	-0.0049 (13)	0.0081 (12)	-0.0164 (14)
C41	0.0390 (15)	0.0535 (17)	0.0446 (15)	-0.0012 (12)	0.0086 (11)	-0.0042 (12)

Geometric parameters (Å, °)

O1—C2	1.366 (3)	C4—C42	1.482 (4)
O1—C7A	1.460 (3)	C4—C5	1.542 (3)
O2—C2	1.194 (3)	C5—C6	1.519 (4)
O3—C7A	1.397 (3)	C5—H5A	0.9700
O3—C8	1.433 (3)	C5—H5B	0.9700
N41—C41	1.137 (3)	C6—C7	1.525 (4)
N42—C42	1.141 (3)	C6—H6A	0.9700
C2—C3	1.503 (3)	C6—H6B	0.9700
C3—C3A	1.524 (3)	C7—C7A	1.505 (3)
C3—H3A	0.9700	C7—H7A	0.9700
C3—H3B	0.9700	C7—H7B	0.9700
C3A—C7A	1.539 (3)	C8—H8A	0.9600
C3A—C4	1.558 (3)	C8—H8B	0.9600
C3A—H3	0.9800	C8—H8C	0.9600
C4—C41	1.479 (3)		
C2—O1—C7A	108.73 (17)	H5A—C5—H5B	108.2
C7A—O3—C8	115.36 (17)	C5—C6—C7	110.6 (2)
O2—C2—O1	121.2 (2)	C5—C6—H6A	109.5
O2—C2—C3	128.8 (2)	C7—C6—H6A	109.5
O1—C2—C3	110.0 (2)	C5—C6—H6B	109.5
C2—C3—C3A	103.38 (18)	C7—C6—H6B	109.5
C2—C3—H3A	111.1	H6A—C6—H6B	108.1
C3A—C3—H3A	111.1	C7A—C7—C6	114.0 (2)
C2—C3—H3B	111.1	C7A—C7—H7A	108.7
C3A—C3—H3B	111.1	C6—C7—H7A	108.7
H3A—C3—H3B	109.1	C7A—C7—H7B	108.7
C3—C3A—C7A	101.50 (17)	C6—C7—H7B	108.7
C3—C3A—C4	112.80 (19)	H7A—C7—H7B	107.6
C7A—C3A—C4	112.33 (17)	O3—C7A—O1	107.39 (18)
C3—C3A—H3	110.0	O3—C7A—C7	113.4 (2)
C7A—C3A—H3	110.0	O1—C7A—C7	110.14 (18)
C4—C3A—H3	110.0	O3—C7A—C3A	103.91 (17)
C41—C4—C42	107.3 (2)	O1—C7A—C3A	102.82 (18)
C41—C4—C5	110.1 (2)	C7—C7A—C3A	118.23 (19)
C42—C4—C5	108.9 (2)	O3—C8—H8A	109.5
C41—C4—C3A	108.48 (18)	O3—C8—H8B	109.5
C42—C4—C3A	111.02 (19)	H8A—C8—H8B	109.5
C5—C4—C3A	110.99 (19)	O3—C8—H8C	109.5
C6—C5—C4	110.0 (2)	H8A—C8—H8C	109.5
C6—C5—H5A	109.7	H8B—C8—H8C	109.5
C4—C5—H5A	109.7	N42—C42—C4	179.5 (3)
C6—C5—H5B	109.7	N41—C41—C4	178.6 (3)
C4—C5—H5B	109.7		
C7A—O1—C2—O2	−166.8 (2)	C8—O3—C7A—C3A	173.04 (19)
C7A—O1—C2—C3	13.8 (3)	C2—O1—C7A—O3	77.4 (2)
O2—C2—C3—C3A	−169.1 (3)	C2—O1—C7A—C7	−158.74 (19)

O1—C2—C3—C3A	10.3 (3)	C2—O1—C7A—C3A	-31.9 (2)
C2—C3—C3A—C7A	-28.3 (2)	C6—C7—C7A—O3	-160.9 (2)
C2—C3—C3A—C4	92.2 (2)	C6—C7—C7A—O1	78.7 (2)
C3—C3A—C4—C41	78.0 (2)	C6—C7—C7A—C3A	-39.0 (3)
C7A—C3A—C4—C41	-168.0 (2)	C3—C3A—C7A—O3	-75.5 (2)
C3—C3A—C4—C42	-39.6 (2)	C4—C3A—C7A—O3	163.78 (18)
C7A—C3A—C4—C42	74.3 (2)	C3—C3A—C7A—O1	36.4 (2)
C3—C3A—C4—C5	-160.97 (18)	C4—C3A—C7A—O1	-84.4 (2)
C7A—C3A—C4—C5	-47.0 (3)	C3—C3A—C7A—C7	157.9 (2)
C41—C4—C5—C6	-179.2 (2)	C4—C3A—C7A—C7	37.2 (3)
C42—C4—C5—C6	-61.8 (3)	C41—C4—C42—N42	-121 (39)
C3A—C4—C5—C6	60.7 (3)	C5—C4—C42—N42	120 (39)
C4—C5—C6—C7	-61.9 (3)	C3A—C4—C42—N42	-3 (40)
C5—C6—C7—C7A	50.4 (3)	C42—C4—C41—N41	-160 (11)
C8—O3—C7A—O1	64.6 (2)	C5—C4—C41—N41	-42 (11)
C8—O3—C7A—C7	-57.3 (3)	C3A—C4—C41—N41	80 (11)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3A $\cdots$ O3 <sup>i</sup>	0.97	2.59	3.290 (3)	129
C8—H8C $\cdots$ O2 <sup>ii</sup>	0.96	2.46	3.219 (4)	136

Symmetry codes: (i)  $-x+1, -y+2, -z+1$ ; (ii)  $x, y+1, z$ .