

RESEARCH ARTICLE

Open Access

# No evidence that frailty modifies the positive impact of antihypertensive treatment in very elderly people: an investigation of the impact of frailty upon treatment effect in the HYpertension in the Very Elderly Trial (HYVET) study, a double-blind, placebo-controlled study of antihypertensives in people with hypertension aged 80 and over

Jane Warwick<sup>1\*</sup>, Emanuela Falaschetti<sup>2</sup>, Kenneth Rockwood<sup>3</sup>, Arnold Mitnitski<sup>4</sup>, Lutgarde Thijs<sup>5</sup>, Nigel Beckett<sup>6</sup>, Christopher Bulpitt<sup>6</sup> and Ruth Peters<sup>2</sup>

## Abstract

**Background:** Treatment for hypertension with antihypertensive medication has been shown to reduce stroke, cardiovascular events, and mortality in older adults, but there is concern that such treatment may not be appropriate in frailer older adults. To investigate whether there is an interaction between effect of treatment for hypertension and frailty in older adults, we calculated the frailty index (FI) for all available participants from the HYpertension in the Very Elderly Trial (HYVET) study, a double-blind, placebo-controlled study of antihypertensives in people with hypertension aged 80 and over, and obtained frailty adjusted estimates of the effect of treatment with antihypertensive medication on risk of stroke, cardiovascular events, and mortality.

(Continued on next page)

\* Correspondence: j.warwick@warwick.ac.uk

<sup>1</sup>Warwick Clinical Trials Unit, Division of Health Sciences, Warwick Medical School, University of Warwick, Gibbet Hill Campus, Coventry CV4 7AL, UK  
Full list of author information is available at the end of the article

(Continued from previous page)

**Methods:** Participants in HYVET were randomised 1:1 to active treatment with indapamide sustained release 1.5 mg ± perindopril 2 to 4 mg or to matching placebo. Data relating to blood pressure, comorbidities, cognitive function, depression, and quality of life were collected at entry into the study and at subsequent follow-up visits. The FI was calculated at entry, based on 60 potential deficits. The distribution of FI was similar to that seen in population studies of adults aged 80 years and above (median FI, 0.17; IQR, 0.11–0.24). Cox regression was used to assess the impact of FI at entry to the study on subsequent risk of stroke, total mortality, and cardiovascular events. Models were stratified by region of recruitment and adjusted for sex and age at entry. Extending these models to include a term for a possible interaction between treatment for hypertension and FI provided a formula for the treatment effect as a function of FI. For all three models, the point estimates of the hazard ratios for the treatment effect decreased as FI increased, although to varying degrees and with varying certainty.

**Results:** We found no evidence of an interaction between effect of treatment for hypertension and frailty as measured by the FI. Both the frailer and the fitter older adults with hypertension appeared to gain from treatment.

**Conclusions:** Further work to examine whether antihypertensive treatment modifies frailty as measured by the FI should be explored.

**Trial registration:** ClinicalTrials.gov NCT00122811 (July 2005)

**Keywords:** Ageing, Antihypertensives, Frailty, Hypertension

## Background

The global population is ageing and with ageing comes an increased prevalence of both frailty and hypertension [1–3]. Treatment for hypertension with antihypertensive medication has been shown to reduce stroke, cardiovascular events, and mortality in older adults [4,5]. Nevertheless, the potential benefit associated with providing treatment for hypertension must be weighed against the potential risk of overtreatment (excessive blood pressure lowering), polypharmacy, and the impact of side effects. There is concern that the treatment may not be beneficial in all older adults, particularly the frailest [6].

Frailty, a clinical state in which there is an increase in an individual's vulnerability for developing increased dependency and/or mortality when exposed to a stressor [7], has been assessed in a number of ways, varying from evaluation of specific impairments [8] to a holistic operational definition, the Frailty Index (FI), which is quantified by counting the number of diseases, symptoms, or similar health 'deficits' [3,9]. The FI theoretically ranges from 0 to 1.0, with a higher FI indicating a higher level of frailty, but in practice has a maximum observed value of around 0.7 and, perhaps because it is more comprehensive, has been shown to be a better predictor of new disability and mortality than the Cardiovascular Health Study or Study of Osteoporotic Fractures scales [10]. FI may be calculated from any data set with sufficient information related to participant deficits (the recommended minimum is 30 deficits that are not saturated within the data set) and in a number of global population datasets has shown consistent relationships between

ageing and mortality, with higher frailty scores at older ages and higher frailty associated with an increased risk of death [11–16]. The deficits included in the FI do not need to be independent of each other. Population studies of very old adults report a mean FI of around 0.16 to 0.18 in those in their early 80s rising to approximately 0.20 over 90 years [13,17]. The FI is applicable at any age, but of particular use in very elderly people, in whom frailty levels are more widely distributed than in the general population, i.e., ranging from the very frail to the robust fitter older adult [17].

The Hypertension in the Very Elderly Trial (HYVET) was a double-blind, placebo-controlled study of antihypertensives in people with hypertension aged 80 and over, which found that treatment with antihypertensives would lead to a reduction in risk of stroke, cardiovascular events, and total mortality [4,5]. However, participants recruited to HYVET, in common with many clinical trials and other studies, are likely to have been healthier than the general very elderly hypertensive population [18]. In consequence, the applicability of the results to the wider elderly population has been questioned, so that uncertainty remains as to whether treatment benefits also extend to the frailer elderly people [6]. To investigate further, we calculated the FI for all available HYVET study participants and obtained frailty-adjusted estimates of the effect of antihypertensive treatment in very elderly people.

## Methods

The HYVET trial randomised participants 1:1 to active treatment with indapamide sustained release 1.5 mg ±

perindopril 2 to 4 mg or to matching placebo. The full inclusion criteria for entry into HYVET have been reported in full elsewhere [4]; briefly, participants were required to be hypertensive (average sitting systolic blood pressure  $\geq 160$  mmHg), to have no condition likely to limit life to less than a year, no diagnosis of dementia, and no need of 24-hour nursing care. Participants gave written informed consent and were assessed at baseline and at 3-monthly intervals for the first year and 6-monthly intervals thereafter until they either died, withdrew from the study, or the trial ended. Data relating to blood pressure, comorbidities, cognitive function, depression, and quality of life were collected at baseline and at subsequent follow-up visits. Key relevant endpoints were selected prior to the start of the trial and these included incident stroke (the primary endpoint), total mortality, and incident cardiovascular events. Data relating to trial endpoints were collected as these occurred and the endpoints validated by an independent blinded committee. All appropriate ethical and regulatory permissions were obtained. Trial registration number is NCT00122811.

The FI was calculated at entry to the study and based on 60 deficits as detailed in Additional file 1. Each deficit was coded as either present or absent in accordance with previous published methodology [19]. To be eligible for inclusion in the FI, the risk of developing a deficit must increase in the general population with increasing age, be associated with varied organ systems, have negative health associations, and be present in at least roughly 1% of the population under study but not to be saturated in the study data [19]. The number of deficits present was counted for each study participant and divided by 60, the maximum number of deficits possible in our data, to give the FI. Information on all 60 deficits was not available for all participants, partly because some had opted out of the quality of life sub-study and partly because of the usual missing data issues. Where missing data meant that the FI calculation for a particular participant would be based on fewer than the intended 60 possible deficits, the FI was calculated as the number of deficits divided by the number available for that participant. Where missing data meant that the FI calculation for a particular participant would be based on fewer than the recommended minimum of 30 possible deficits, the FI was not calculated but set to missing and the subject excluded from the analysis.

Cox regression [20] was used to assess the impact of FI at entry to the study on subsequent risk of fatal and non-fatal stroke, total mortality, and cardiovascular events (fatal and non-fatal stroke, myocardial infarction and heart failure). Models were stratified by region of recruitment (Western Europe, Eastern Europe, China) to allow each to have a different baseline hazard function (calibrate FI) and adjusted for sex and age at entry to the study. There was

no adjustment for baseline cardiovascular disease as this forms part of the FI. The validity of the proportional hazards assumption was assessed using diagnostic plots and Grambsch and Therneau tests [21], and the overall model fit was assessed graphically by plotting the Nelson-Aalen cumulative hazard function versus the Cox-Snell residuals and comparing to a 45° reference line.

We fitted Cox regression models (stratified by region) with terms for baseline age, sex, FI, treatment, and an interaction between treatment and FI to obtain a formula for the treatment effect as a function of FI [22] to illustrate the impact of FI upon the estimate of treatment effect obtained from our models. The estimates of treatment effect from this model, and associated 95% confidence intervals (CIs), were then presented graphically.

The relationship between baseline FI and subsequent drop out was investigated to establish whether participants for whom the FI could not be calculated, owing to incomplete data, differed substantially from those for whom the FI could be calculated.

## Results

### Characteristics of participants

The HYVET trial randomised 3,845 participants, of whom only the 2,656 who consented to complete an additional quality of life questionnaire provided sufficient data to allow the calculation of the FI. The baseline characteristics of these participants are given in Table 1. There did not appear to be any imbalance between the treatment groups.

We also compared those for whom the FI was calculable with those for whom it was not. There was no difference between them in terms of age, sex, previous cardiovascular disease, or baseline sitting systolic blood pressure but there was a difference in cognitive test score; the median Mini-Mental State Exam (MMSE) score at entry to the study in those in whom FI was calculable was 26.0 (Interquartile range (IQR), 22–28) and in those in whom FI was not calculable was 27.0 (IQR, 24–29) ( $P < 0.001$ , Mann–Witney test).

### Frailty index (FI)

The distribution of the constituents used to calculate the FI was similar in the two treatment groups (Additional file 2: Table S1). The distribution of FI at entry to the study was skewed with mean and median values 0.19 (standard deviation (SD), 0.10) and 0.17 (IQR, 0.11–0.24), respectively, and range 0.01 to 0.63 (Figure 1). Median FI was higher ( $P < 0.01$ , Mann–Whitney rank-sum test) for women (0.18; IQR, 0.12–0.26) compared to men (0.15; IQR, 0.10–0.21). On average, FI scores increased slowly with age (0.003 per year of age (95% CI, 0.002–0.004)). Greater FI at entry to the study was associated with an increased risk of death (HR, 1.24 per 0.05

**Table 1 Baseline characteristics of the 2,656 participants of HYVET for whom the frailty index was calculable, by treatment group**


	Placebo	Active
n	1,324	1,332
Age (yrs), mean (SD)	83.4 (3.0)	83.6 (3.2)
Male	520 (39.3%)	526 (39.5%)
Female	804 (60.7%)	806 (60.5%)
Body mass index*		
Underweight	39 (3%)	58 (4%)
Normal weight	587 (44%)	605 (46%)
Overweight	566 (43%)	530 (40%)
Obese	132 (10%)	138 (10%)
Sitting SBP, mean (SD)	173.1 (8.9)	173.3 (8.8)
Sitting DBP, mean (SD)	90.0 (8.9)	89.9 (8.8)
Standing SBP, mean (SD)	168.0 (11.8)	168.2 (11.9)
Standing DBP, mean (SD)	87.9 (9.9)	88.1 (9.8)
Cardiovascular disease, n (%)	177 (13.4%)	159 (11.9%)
Antihypertensive treatment prior to entry into the trial, n (%)	830 (62.7%)	828 (62.2%)
Mini Mental State Examination, median (IQR)	26.0 (22–28)	26.0 (22–28)
Frailty Index, median (IQR)	0.17 (0.11–0.24)	0.16 (0.11–0.24)

\*Adjusted for region of recruitment.  
 DBP, Diastolic blood pressure; IQR, Interquartile range; SBP, Systolic blood pressure; SD, Standard deviation.

increase in FI; 95% CI, 1.18–1.30), cardiovascular events (HR, 1.23 per 0.05 increase in FI; 95% CI, 1.16–1.30), and stroke (HR, 1.26 per 0.05 increase in FI; 95% CI, 1.15–1.37). Adjustment for treatment group, age, and sex, and stratification by region of recruitment did not alter these findings.

**Effect of frailty on estimates of treatment effect from HYVET**

Median follow-up was similar in the two treatment groups. For time to death, median follow-up was 22 months (IQR, 13–34) in the placebo group compared to 23 months (IQR, 13–35) in the active treatment group; for time to cardiovascular events, median follow-up was 21 months in the placebo group (IQR, 12–33) and 23 months (IQR, 13–35) in the active treatment group; and for time to stroke, median follow-up was 21 months (IQR, 12–34) in the placebo group and 23 months (IQR, 13–34) in the active treatment group. The proportionality assumption was not violated (*P* values for the global Grambsch and Therneau tests of the proportional hazards assumption for the models for time to death, time to cardiovascular events, and time to stroke were 0.78, 0.14, and 0.39, respectively) and overall model fit was adequate. The estimate of treatment effect obtained from the Cox regression models (Table 2) did not change with adjustment for baseline FI. Treatment with an anti-hypertensive was associated with a 36% reduction in risk of fatal and non-fatal stroke (HR, 0.64; 95% CI, 0.42–0.96; *P* = 0.03) after adjustment for baseline FI, sex, and age, and stratification for region of recruitment. Similarly, active treatment within HYVET significantly reduced the risk of fatal and non-fatal cardiovascular events (HR, 0.59; 95% CI,



**Figure 1** Histogram showing the distribution of frailty index (FI) among 2,656 participants of HYVET at entry to the study.


**Table 2 Hazard ratios and associated 95% confidence intervals from Cox regression models showing the effect of adjusting for frailty on the estimate of treatment effect in those for whom a frailty index was calculated n = 2,656**

Variables included in the model	Stroke (95 events)	Cardiovascular events (231 events)	Total mortality (294 events)
Treatment group	0.65 (0.43–0.98)	0.59 (0.45–0.77)	0.83 (0.66–1.05)
Treatment group, sex, and age	0.65 (0.43–0.98)	0.59 (0.45–0.77)	0.83 (0.66–1.05)
Treatment group, sex, age, and FI at entry to the study	0.64 (0.42–0.96)	0.59 (0.45–0.77)	0.83 (0.66–1.04)

All models stratified by region of recruitment.

0.45–0.77) but there was no significant difference in total mortality between the placebo and active treatment groups (HR, 0.83; 95% CI, 0.66–1.04). The adjusted hazard ratios (HR) for all three endpoints were similar to those seen in the main trial results (which were HR, 0.68; 95% CI, 0.47–0.98 for stroke; HR, 0.66; 95% CI, 0.53–0.82 for cardiovascular events, and HR, 0.79; 95% CI, 0.65–0.95 for mortality [23]). Repeating these analyses with previous cardiovascular disease excluded from the calculation of FI did not materially affect the results.

There was no significant interaction between treatment effect and frailty for any of the three endpoints (*P* values for the interaction term in the interaction models for stroke, cardiovascular events, and total mortality were 0.52, 0.73, and 0.61, respectively). Estimates of the log HR for the treatment effect obtained from these interaction models are plotted versus FI, with the associated CIs (Figure 2) and HRs for selected values of FI (FI = 0.1, 0.2, 0.3, 0.4, 0.5, and 0.6) are reported in Table 3. For all three models the point estimate of the HR for the treatment



**Table 3 Estimated hazard ratios for treatment effect (active treatment versus placebo) and associated 95% confidence intervals, by frailty index**

Frailty index	Stroke		Cardiovascular events		Total mortality	
	HR	95% CI	HR	95% CI	HR	95% CI
0.1	0.75	0.40–1.38	0.62	0.42–0.92	0.89	0.63–1.25
0.2	0.66	0.43–1.01	0.60	0.45–0.78	0.84	0.66–1.07
0.3	0.59	0.36–0.96	0.57	0.42–0.79	0.80	0.61–1.04
0.4	0.52	0.25–1.09	0.55	0.34–0.89	0.76	0.50–1.14
0.5	0.47	0.16–1.33	0.53	0.26–1.06	0.72	0.40–1.29
0.6	0.41	0.10–1.65	0.50	0.20–1.27	0.68	0.32–1.48

All models adjusted by age, sex, and interaction between treatment and frailty index, and stratified by region of recruitment.

effect decreases as FI increases, although to varying degrees and with varying certainty.

#### Association between frailty and premature withdrawal

There were 347 patients for whom baseline FI was calculable but who withdrew from the study and were therefore censored in the analysis at the date of withdrawal. The distribution of baseline FI was not the same for those who withdrew and those who did not withdraw from the study (Mann–Whitney test,  $P < 0.0001$ ). The median FI was 0.19 (IQR, 0.13–0.27) in those who withdrew from the study and 0.16 (IQR, 0.11–0.24) in those who did not withdraw. The withdrawal rate in the least frail category (FI  $\leq 0.10$ ) was 9% compared to 17% in the frailest category (FI  $\geq 0.35$ ) and there was no difference in withdrawal rate between the treatment groups (14.38% placebo group vs. 14.90% active treatment group). Within the most frail category (FI  $\geq 0.35$ ), the withdrawal rate was 21% in the active treatment group ( $n = 99$ ) compared to 14% ( $n = 94$ ) in the placebo group, but this was not significant ( $P = 0.18$ ,  $\chi^2$  test).

#### Discussion

Frailty as measured by the FI was a strong predictor of stroke, total mortality, and cardiovascular events in the HYVET trial, which is in agreement with multiple analyses from observational datasets [8–13]. We found no evidence of an interaction between baseline FI and treatment with antihypertensives on risk of stroke, death from all causes, or cardiovascular events in very elderly people. Furthermore, the burden of frailty amongst HYVET participants at baseline was similar to that seen in population studies [13–17].

Overall this suggests both that the HYVET population is more representative in terms of frailty than may have been supposed, and that benefits associated with blood pressure lowering treatment are accrued in both frailer and fitter older adults. These results would imply that frailty alone should not be used as a criteria for determining whether or not the treatment of an individual aged 80 and over with an antihypertensive to lower blood pressure

to a goal of  $<150/80$  mmHg is justified. Nevertheless, further work is required to fully characterize the benefit risk balance in this age group, with particular attention paid to the impact of any treatment adverse effects and/or a diagnosis of dementia, especially as there is epidemiological evidence to suggest that high blood pressure may not be harmful, and may even be beneficial, in the very elderly who are frail or have a functional disability [24–27]. In general, as the degree of frailty increases, so does the chance of functional impairment [28,29] or mobility impairment [30]. Furthermore, as the associations between risk factors and adverse outcomes may differ at extreme age and in frailty subgroups, and be dependent on risk factor change over time, a more nuanced approach to the interpretation of results may be required.

However, our results must be interpreted with caution. The number of items (at least 30) required to complete an FI meant that, of the 3,845 participants randomized, only the 2,656 (69%) who consented to complete an additional quality of life questionnaire had sufficient data. Nevertheless, it is reassuring that those for whom an FI was not calculable differed chiefly only in relation to cognition, scoring on average one point higher on the MMSE at entry, statistically but not clinically different. It is also possible that treatment with an anti-hypertensive may have affected participant withdrawals differentially, leading to bias in our estimate of the treatment effect. However, we found no difference in withdrawal rate between the treatment groups (14.38% placebo group vs. 14.90% active treatment group) and even amongst the frailest participants (FI  $\geq 0.35$ ) the withdrawal rate was only 17% overall. Although only 2,656 participants were available from the HYVET trial for these analyses these data still represent a significant number of older adults.

One must also be careful not to over interpret the results from the interaction models presented in Table 3 and Figure 2. Although the estimate of the log HR decreases as FI increases for all three endpoints, the interaction term in the respective models were not significant (overall the relationship between frailty and treatment effect was not strong). Furthermore, the wider CIs in the

extremes are to be expected, given that very little of our data lies in the region  $FI < 0.1$  and  $FI > 0.4$  (Figure 2).

This analysis has some strengths. In particular, the use of data from a double-blind, placebo-controlled clinical trial also allows exploration of this by randomised group providing a more robust finding that would be possible from observational data alone. The similarity of the FI at baseline with large observational datasets supports the potential applicability of the results. As far as the authors know, this is the first time the FI has been used in analysing the results of a clinical trial, particularly a hypertensive very elderly group. Although only 69% of the randomized patients were included, this still constitutes a significant number within the literature base for this age group.

## Conclusions

Our analyses show that in the HYVET study participants there was no evidence of an interaction between treatment effect and frailty. Both the frailer and the fitter older adults with hypertension appeared to gain from treatment. Further work in is needed to confirm these findings in other similar datasets and to explore whether antihypertensive treatment modifies frailty as measured by the FI. Additionally, examining this more holistically, looking at impact of treatment over time, would be of additional benefit and is motivating further research by our group.

## Ethical approval

All appropriate ethical and regulatory permissions were obtained. This included approval in all participating countries by national drug or healthcare agencies, national ethics committees, and where required local ethical committees. The documentation is therefore large and complex. Full copies are archived by Imperial College London. HYVET was approved in the UK by the Multi-Centre Research Ethics Committee for Wales (MREC 98/9/16).

## Additional files

**Additional file 1: Constituents of the Frailty Index.**

**Additional file 2: Table S1.** Variables included in the calculation of the Frailty Index (percent present in each treatment group).

## Abbreviations

CI: Confidence interval; FI: Frailty Index; HR: Hazard ratio; HYVET: Hypertension in the Very Elderly Trial; IQR: Interquartile range; MMSE: Mini-Mental State Exam; SD: Standard deviation.

## Competing interests

NB has received honoraria from Servier International. JW, EF, KR, AM, LT, CB, and RP have declared that no competing interests exist.

## Authors' contributions

JW designed the study, analysed the data, interpreted the results, and drafted the paper. EF and LT analysed the data, interpreted the results, and reviewed the paper. KR and AM advised on the construction and analysis of the frailty index, interpreted the results, and reviewed the paper.

NB designed the study, interpreted the results, and drafted the paper. CB designed the study, interpreted the results, and reviewed the paper. RP designed the study, analysed the data, interpreted the results, and drafted the paper. All authors read and approved the final manuscript.

## Acknowledgements

Financial disclosure: This work was supported by The Dunhill Medical Trust [grant number: R287/0213] and benefited from facilities funded through Birmingham Science City Translational Medicine Clinical Research and Infrastructure Trials Platform, with support from Advantage West Midlands. The data used for these analyses comes from the HYVET study, which was funded by grants from the British Heart Foundation and the Institute de Recherches Internationales Servier, and sponsored by Imperial College London. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. All persons mentioned in the acknowledgements have given written consent. The committee members and investigators for HYVET were as follows: Co-ordinating Centre: CJ Bulpitt (lead investigator), AE Fletcher (co-investigator), NS Beckett (trial coordinator), R Peters (deputy trial coordinator), HYVET coordinating team at Imperial College London (1999 to 2008). HYVET Committees: Steering Committee: T McCormack, J Potter, BG Extremera, P Sever, F Forette, D Dumitrascu, C Swift, J Tuomilehto, J Coope (retired in 2001), C Nachev (deceased); Data Monitoring Committee: J Staessen, L Thijs, R Clarke, K Narkiewicz; End Points Committee: C Davidson (retired in 2003), J Duggan, G Leonetti, N Gainsborough, MC De Vernejoul, J Wang, V Stoyanovsky; Dementia Validation Committee: J Tuomilehto, R Clarke, A Waldman, I Walton, C Ritchie; Ethics Committee: R Fagard, J Grimley Evans, B Williams; Investigators: (\*National Co-ordinators) Australia — R Warne\* and I Puddey\*, M Woodward, R Penhall, C Inderjeeth, S Roger, R Scholes, C Johnson; Belgium — H Celis\*, G Adriaens, W Onsea, K Cornelli, D Vantroyen, P Cleen, P de Voogt; Bulgaria — C Nachev\* (deceased) (national coordinator from 1998 to 2005), V Stoyanovsky\* (national coordinator after 2005), P Solakov, R Prokopova, E Mantova, D Smilkova, S Mantov, K Yankulova, R Kermova, D Popov, V Sirakova, V Gergova, D Kamenova, F Grigorov, T Vassileva, R Alahverdian, M Tzekova; A Postadjian, M Geneva, V Mincheva, T Petrusheva, A Toncheva, I Gruev, V Tzanova; China — L Liu\*, H Ge, S Wang, J Wang, W Zhang, S Jin, L Ge, YF Lu, S Ma, L Shen, J Guo, Z Lv (deceased), R Huang, X Li, B Guo, GE Yuan, T Zhang, L Zhang, J Feng, Z He, J Wang, L Deng, L Liu, Q Yuan, F Zhang, H Li, D Wang, K Yang, M Sun, H Liu, X Yan, F Ren, J Tang, M Zhao, X Luo, H Zhou, H Sang, J Wang, L Yan, Z Wang, J Zhang, C Wang; Finland — R Antikainen\*, T Strandberg, T Konttila, A Hynninen, M Jääskivi, J Airas, T Jääskeläinen, J Tuomilehto, H Litmanen, T Karhi, H Yliharsila; France — F Forette\*, J Doucet, J Belmin, A Benetos, G Berrut, T Boge, M Bonnefoy, A Carre, N Charas, J Covillard, T Dantoine, M Escande, Y Frances, R Joire, C Jeandel, S Legrain, A Lion, M Maillat-Vioud, JP Escaillas, S Meaume, P Pfitzenmeyer, F Puisieux, E Quercy, O Rodat, J Soubeyrand, B de Wazieres, H Hindennach, L Lugassy, J Rossi, M Martel, J-M Paladel, C Ravier, A Visconti, JP Gallet, D Zygouritsas, D Charles, F Flamand, G Grandmottet, M Grandmottetegermann, C Gevrey, PL Mesnier, G Robert, C Besset-Prat, A Brousse, P Lafont, J Morelli, P Vernede, A Volkmann, X Bodin, B Destrube, R Eoche, A Boye, F Seropian, P Gernigon, D Meker, J Thomere, Y Thual, F Volny, E Grassart, M Herent, D Lejay, J-P Lopez, B Mannessier, G Pruvost, J-C Urbina; Ireland — J Duggan\*; New Zealand — C Anderson\*, S Lillis, J Gommans, H Senior; Poland — T Grodzicki\*, Z Chodorowski, Z Gacjong; Romania — D Dumitrascu\*, M Comsa, V Sandru, G Prada, M Dunca-Moisin, D Jianu, D Jinga-Lazar, V Enachescu, C Zaharia; Russia — Y Nikitin\*, A Kirichenko, L Olbinskaya, A Martynov, V Zadionchenko, V Moiseev, G Storozhakov, S Nedogoda, RS Karpov, O Barbarash, G Efremushkin, V Kostenko, M Boyarkin, S Churina, T Tyurina, M Ballyuzek, L Ermoshkina, A Timofeev, S Yakusheva, N Shilkina, V Barbarich, L Latunceva, S Burakova, T Ripp, S Pekarsky, V Mordovini; Tunisia — A Belhani\*, E Boughzela, S Soraya, B Youssef-Zouari, AB Khalfallah, MH Houman, AK Abida; United Kingdom — C Rajkumar\*, M Wilkins, ND Pandita-Gunawardena, J Potter, E Ekpo, M Price, N de Kare-Silver, A Starczewski, S Chandran, N Nasar, M Datta-Chaudhuri, T McCormack, N Majmudar, A Gordon, L Brawn, T Solanki, F Dockery, R Schiff. We wish to acknowledge the work of Professor C Nachev (Steering committee member, National Co-ordinator of Bulgaria and HYVET investigator from 1998 until his death in 2005).

## Author details

<sup>1</sup>Warwick Clinical Trials Unit, Division of Health Sciences, Warwick Medical School, University of Warwick, Gibbet Hill Campus, Coventry CV4 7AL, UK.

<sup>2</sup>Imperial Clinical Trials Unit, School of Public Health, Faculty of Medicine, Imperial College London, St Mary's Campus, Norfolk Place, London W2 1PG, UK. <sup>3</sup>Geriatric Medicine, Dalhousie University, Queen Elizabeth II Health Sciences Centre, 1421-5955 Veterans' Memorial Lane, Halifax B3H 1C6, Nova Scotia, Canada. <sup>4</sup>Department of Medicine, Dalhousie University, Centre for Health Care of the Elderly, 1421-5955 Veterans' Memorial Lane, Halifax B3H 2E1, Nova Scotia, Canada. <sup>5</sup>Studies Coordinating Centre, Research Unit Hypertension and Cardiovascular Epidemiology, Department of Cardiovascular Sciences, University of Leuven, Campus Sint Rafaël, Kapucijnenvoer 35, Box 7001, BE-3000 Leuven, Belgium. <sup>6</sup>Department of Medicine, Imperial College London, Hammersmith Campus, London W12 0NN, UK.

Received: 12 November 2014 Accepted: 17 March 2015

Published online: 09 April 2015

## References

- United Nations Population Division DESA. World Population Ageing 1950–2050. 2002. <http://www.un.org/esa/population/publications/worldageing19502050/>.
- Health Survey for England 2009, Trend tables: adult trend tables. 2009. <http://www.hscic.gov.uk/pubs/hse09trends>
- Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. *Sci World J*. 2001;1:323–36.
- Beckett NS, Peters R, Fletcher AE, Staessen JA, Liu LS, et al. on behalf of the HYVET Study Group. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med*. 2008;358:1887–98.
- Beckett N, Peters R, Tuomilehto J, Swift C, Sever P, Potter J, et al. Immediate and late benefits of treating very elderly people with hypertension: results from the active treatment extension to Hypertension in the very elderly randomised controlled trial. *BMJ*. 2011;344:d7541.
- Abramov D, Cheng H. Controversy in treating the oldest old with hypertension: is the hypertension in the very elderly trial the final answer? *J Am Geriatr Soc*. 2009;57:570–1.
- Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R, et al. Frailty consensus: A call to action. *J Am Med Dir Assoc*. 2013;14:392–7.
- de Vries NM, Staal JB, van Ravensberg CD, Hobbelen JSM, Olde Rikkert MG, der Sanden MW N-v. Outcome instruments to measure frailty: a systematic review. *Ageing Res Rev*. 2011;10:104–14.
- Rockwood K, Fox RA, Stolee P, Robertson D, Beattie BL. Frailty in elderly people: an evolving concept. *CMAJ*. 1994;150:489–95.
- Malmstrom TK, Miller DK, Morley JE. A comparison of four frailty models. *J Am Geriatr Soc*. 2014;67:721–6.
- Gu D, Dupre M, Sautter J, Zhu H, Liu Y, Yi Z. Frailty and mortality among Chinese at advanced ages. *J Gerontol B Psychol Sci Soc Sci*. 2009;64:279–89.
- Shi J, Song X, Yu P, Tang Z, Mitnitski A, Fang X, et al. Analysis of frailty and survival from late middle age in the Beijing longitudinal study of aging. *BMC Geriatr*. 2011;11:17.
- Mitnitski A, Song X, Skoog I, Broe GA, Cox JL, Grunfeld E, et al. Relative fitness and frailty of elderly men and women in developed countries and their relationship with mortality. *J Am Geriatr Soc*. 2005;53:2184–9.
- Mitnitski A, Bao L, Rockwood K. Going from bad to worse: a stochastic model of transitions in deficit accumulation, in relation to mortality. *Mech Ageing Dev*. 2006;127:490–3.
- Mitnitski A, Bao L, Skoog I, Rockwood K. A cross-national study of transitions in deficit counts in two birth cohorts: Implications for modelling ageing. *Exp Gerontol*. 2007;42:241–6.
- Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet*. 2013;381:752–62.
- Collerton J, Martin-Ruiz C, Davies K, Hilken CM, Isaacs J, Kolenda C, et al. Frailty and the role of inflammation, immunosenescence and cellular ageing in the very old: Cross-sectional findings from the Newcastle 85+ Study. *Mech Ageing Dev*. 2012;133:456–66.
- Strandberg TE, Salomaa VV, Vanhanen HT, Naukkarinen VA, Sarna SJ, Miettinen TA. Mortality in participants and non-participants of a multifactorial prevention study of cardiovascular diseases: a 28 year follow up of the Helsinki Businessmen Study. *Br Heart J*. 1995;74:449–54.
- Searle S, Mitnitski A, Gahbauer E, Gill T, Rockwood K. A standard procedure for creating a frailty index. *BMC Geriatr*. 2008;8:24.
- Cox D. Regression models and life-tables. *J R Statist Soc B*. 1972;34:187–220.
- Grambsch PM, Therneau TM. Proportional hazards tests and diagnostics based on weighted residuals. *Biometrika*. 1994;81:515–26.
- Hosmer DW, Lemeshow S, May S. Applied survival analysis: regression modelling of time-to-event data. New York: John Wiley & Sons, Inc.; 2008.
- Peters R, Beckett N, McCormack T, Fagard R, Fletcher A, Bulpitt C. Treating hypertension in the very elderly—benefits, risks, and future directions, a focus on the hypertension in the very elderly trial. *Eur Heart J*. 2014;35:1712–8.
- Sabayan B, van Vliet P, de Ruijter W, Gusselkoo J, de Craen AJM, Westendorp RGJ. High blood pressure, physical and cognitive function, and risk of stroke in the oldest old: the Leiden 85-plus Study. *Stroke*. 2012;44:15–20.
- Muller M, Smulders YM, de Leuw PW, Stehnowwer CDA. Treatment of hypertension in the oldest old: a critical role for frailty? *Hypertension*. 2014;63:433–41.
- Post Hespers G, Smulders YM, Maier AB, Deeg DJ, Muller M. Relation between blood pressure and mortality risk in an older population: role of chronological and biological age. *J Int Med*. 2014. ahead of print.
- Odden MC, Peralta CA, Haan MN, Covinsky KE. Rethinking the association of high blood pressure with mortality in elderly adults: the impact of frailty. *Arch Intern Med*. 2012;172:1162–8.
- Theou O, Rockwood MR, Mitnitski A, Rockwood K. Disability and co-morbidity in relation to frailty: how much do they overlap? *Arch Gerontol Geriatr*. 2012;55:e1–8.
- Woo J, Leung J, Morley JE. Comparison of frailty indicators based on clinical phenotype and the multiple deficit approach in predicting mortality and physical limitation. *J Am Geriatr Soc*. 2012;60:1478–86.
- Davis DH, Rockwood MR, Mitnitski AB, Rockwood K. Impairments in mobility and balance in relation to frailty. *Arch Gerontol Geriatr*. 2011;53:79–83.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

