

# Molecular heterogeneity of large-conductance calcium-activated potassium channels in canine intracardiac ganglia

Elisabet Selga,<sup>1</sup> Alexandra Pérez-Serra,<sup>1</sup> Alba Moreno-Asso,<sup>1,†</sup> Seth Anderson,<sup>2,‡</sup> Kristen Thomas,<sup>2,§</sup> Mayurika Desai,<sup>2</sup> Ramon Brugada,<sup>1</sup> Guillermo J Pérez<sup>1,2,¶</sup> and Fabiana S Scornik<sup>1,2,¶,\*</sup>

<sup>1</sup>Cardiovascular Genetics Center; Institut d'Investigació Biomèdica de Girona (IdIBGi); Department of Medical Sciences; School of Medicine; University of Girona (UdG); Girona, Spain; <sup>2</sup>Masonic Medical Research Laboratory; Utica, NY USA

<sup>†</sup>Current affiliations: Diabetes and Obesity Research Laboratory; Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS); Centre Esther Koplowitz; Barcelona, Spain; Centro de Investigación Biomédica en Red de Diabetes y Enfermedades Metabólicas Asociadas (CIBERDEM); Spain; <sup>‡</sup>University of Rochester; Rochester, NY USA; <sup>§</sup>Daemen College Amherst; NY South Buffalo Mercy Hospital; Buffalo, NY USA

<sup>¶</sup>These authors contributed equally to this work.

**Keywords:** BK channels, alternative splicing,  $\beta$  subunits, intracardiac neurons, autonomic

**Abbreviations:** BK channel, large conductance calcium-activated potassium channel; ICG, intracardiac ganglia;  $I_{BK}$ , BK current; SS, splice site

Submitted: 05/31/13

Revised: 06/20/13

Accepted: 06/20/13

<http://dx.doi.org/10.4161/chan.25485>

\*Correspondence to: Fabiana S. Scornik;  
Email: [fabianasilvia.scornik@udg.edu](mailto:fabianasilvia.scornik@udg.edu)

Addendum to: Pérez GJ, Desai M, Anderson S, Scornik FS. Large-conductance calcium-activated potassium current modulates excitability in isolated canine intracardiac neurons. *Am J Physiol Cell Physiol* 2013; 304:C280-6; PMID:23195072; <http://dx.doi.org/10.1152/ajpcell.00148.2012>.

**L**arge conductance calcium-activated potassium (BK) channels are widely expressed in the nervous system. We have recently shown that principal neurons from canine intracardiac ganglia (ICG) express a paxilline- and TEA-sensitive BK current, which increases neuronal excitability. In the present work, we further explore the molecular constituents of the BK current in canine ICG. We found that the  $\beta 1$  and  $\beta 4$  regulatory subunits are expressed in ICG. Single channel voltage-dependence at different calcium concentrations suggested that association of the BK $\alpha$  with a particular  $\beta$  subunit was not enough to explain the channel activity in this tissue. Indeed, we detected the presence of several splice variants of the BK $\alpha$  subunit. In conclusion, BK channels in canine ICG may result from the arrangement of different BK $\alpha$  splice variants, plus accessory  $\beta$  subunits. The particular combinations expressed in canine IC neurons likely rule the excitatory role of BK current in this tissue.

## Introduction

Large conductance calcium-activated potassium (BK) channels are expressed in a wide variety of cells within the nervous system, including intracardiac (IC) neurons of different species,<sup>1-3</sup> where they contribute to numerous cellular functions,

including action potential repolarization and afterhyperpolarization. Diverse properties have been reported for  $I_{BK}$  in different tissues, in different cells in a given tissue, in different regions of a single cell, and in the same cell under different stimuli.<sup>4-7</sup> This behavior is accepted to be associated to heterogeneity in the molecular composition of BK channels, which can be achieved, among other mechanisms, by differential expression of BK $\beta$  subunits and BK $\alpha$  splice variants. BK channels are usually composed by four BK $\alpha$  pore-forming subunits and auxiliary  $\beta$  subunits. To date, four different BK $\beta$  subunits have been ascertained (encoded by KCNMB1 to 4), which display tissue-specific distribution and differentially modulate  $I_{BK}$ .<sup>8-13</sup> On the contrary, BK $\alpha$  subunits are encoded by a single gene, KCNMA1, from which a wide diversity of variants can be originated through alternative splicing.<sup>14</sup> At least 13 splice sites have been described for KCNMA1.<sup>6,15</sup> Some of them have been identified in many species and tissues, and thus are considered to be ubiquitous and conserved across species (Table 1). However, the particular molecular composition of BK channels and how this identity modulates  $I_{BK}$  in different tissues still remains greatly unknown.

We have recently shown that principal neurons from canine intracardiac ganglia (ICG) express a conspicuous BK current

**Table 1.** Description of the analyzed splice sites

Splice site	Variant (aa)	Named
5'	+66	MANG
	+36	MSS
	0	MDALI
SS1	+4	+4
	+58/61	STREX
SS2	+29	e22
	+3	+3
SS3	+8	+8
SS4	+27	Slo27
3'	+8	VYR
	+61	DEC
	+8	ERL

The table shows the more common variants described for each of the splice sites (SS) analyzed, the number of aminoacids (aa) added to originate each variant and the name by which they are usually known. The Zero variant, which corresponds to the absence of insertion, has been described for SS1 to SS4. The VYR and ERL splice variants have the same number of aa but different sequences.

( $I_{BK}$ ), which plays a role in their excitability. We demonstrated that these channels do not present intrinsic inactivation and that the decline of the BK whole cell current was most likely dependent on the intracellular calcium concentration. In the present work, we aimed to further explore the molecular constituents of BK channels in canine IC neurons. Our results suggest that the presence of several splice variants of the BK $\alpha$  subunit, modulated by  $\beta 4$  and/or  $\beta 1$  subunits, may explain the voltage- and calcium-dependence of the BK channel observed in this native preparation.

## Results

**BK channel  $\beta 1$  and  $\beta 4$  regulatory subunits are expressed in canine ICG.** It is well known that BK channel  $\beta$  subunits modulate channel's biophysical properties, including activation voltage-dependence, activation and deactivation kinetics, and inactivation.<sup>16</sup> In particular, the  $\beta 2$  and  $\beta 3$  regulatory subunits are known to be responsible for  $I_{BK}$  inactivation.<sup>13</sup> Our recent studies showed that the inactivation of  $I_{BK}$  from canine ICG is not due to intrinsic inactivation of


BK channels. To provide molecular evidence to this conclusion, we sought to determine which of the four described BK channel  $\beta$  subunits are expressed in IC neurons and would thus confer  $I_{BK}$  its particular characteristics. RT-PCR studies evidenced that  $\beta 1$  and  $\beta 4$  subunits were clearly expressed in canine ICG, while a faint expression was observed for  $\beta 2$  and  $\beta 3$  subunits in this tissue (Fig. 1A). To further confirm our results, we performed immunofluorescence staining of partially dissociated ICG. IC neurons expressed the  $\beta 4$  subunit (Fig. 1Ba), and all those cells that expressed this subunit also expressed the BK channel  $\alpha$  subunit (Fig. 1Bb). On the contrary, staining for  $\beta 2$  and  $\beta 3$  subunits was only observed in a low percentage of neurons (Fig. 1Bc and 1Bd, respectively). These data are in agreement with the lack of single channel inactivation observed in our previous article. In addition, they strongly suggest that  $\beta 4$  subunit (and  $\beta 1$  to a lesser extent) is a likely candidate for modulating  $I_{BK}$  in these cells.

**Calcium- and voltage-dependence of single BK channels in IC neurons.** Our previous whole cell current studies could not provide detailed information on BK channel calcium- and voltage-dependence. To cover this aspect, we studied the single channel voltage-dependence at different calcium concentrations in excised patch experiments from our native preparation. Figure 2 depicts examples of single channel recordings at increasing calcium concentrations and the open probability ( $P_o$ ) vs. voltage relationship for the calcium concentrations assayed. The analysis of voltage- and calcium-dependence led to  $V_{1/2}$  values of  $-35.7 \pm 11.1$  mV,  $29.8 \pm 2.2$  mV and  $70.0 \pm 15.1$  mV; at 10, 3 and 0.3  $\mu$ M  $Ca^{2+}$ , respectively. These results deviate from similar analysis using heterologous expression of BK $\alpha$  and either  $\beta 1$  or  $\beta 4$  subunits.<sup>8,9,17,18</sup> These deviations could be due to the possibility that in our native preparation, more than one  $\beta$  subunit modulates the BK $\alpha$  subunit. Alternatively, or in addition, a BK $\alpha$  isoform different from that used in the heterologous expression studies could be expressed in canine IC neurons.

**A combination of BK channel  $\alpha$ -subunit variants is expressed in canine**

ICG. Multiple splice sites (SS) have been described for the BK channel  $\alpha$  subunit.<sup>6</sup> Some of the resulting splice variants have been reported to modulate BK channel electrophysiological properties. In the present study, we have focused on six particular SS (Table 1) to determine which of the associated splice variants were expressed in canine ICG. A summary of the results obtained is shown in Table 2. Of note, PCR amplification of the SS1 region originated two amplicons (Fig. 3A, lane 2). Their molecular weights were compatible with the SS1-insertless variant (lower band) and the variant bearing an insertion of 4aa at the SS1 site (upper band). Sequencing of both PCR products confirmed their anticipated identities. When the SS2 region was amplified in ICG, we observed a band compatible with the shorter (58aa) STREX splice variant (Fig. 3A, lane 3, upper band). Additionally, two bands with lower mobility were also observed in this tissue. Their molecular weight was compatible with the Zero and +4 variants. Their identities were confirmed by sequencing. PCR analysis of the SS3 region and sequencing of the obtained band evidenced the expression of the SS3-Zero variant. Regarding the SS4 region, a band compatible with the slo27 variant was detected (Fig. 3A, lane 5, middle band). Two additional bands were also observed. Sequencing of these 3 amplicons confirmed the expression of the slo27 variant and the SS4-Zero variant (lower band). The upper band corresponded to a heteroduplex of Zero and slo27, a phenomenon previously reported by Lai and McCobb respect to SS2.<sup>19</sup>

We also determined the splicing variants at the N- and C-terminus of the BK channel  $\alpha$  subunit in canine ICG. Sequencing of the PCR products obtained upon amplification of the N-terminal region evidenced the presence of transcripts bearing the MSS translational start site. As this start site is in frame with the 3' start site beginning MDALI, we cannot ensure the presence of transcripts starting with the latter start site in ICG. Regarding the C-terminus, we were able to detect the presence of the 3 most common variants at this site, ending VYR, DEC and ERL. A schematic




**Figure 1.** BK channel  $\beta 1$  and  $\beta 4$  regulatory subunits are expressed in canine ICG. **(A)** Picture depicts the results of RT-PCRs performed as described in Methods using specific primers for BK channel  $\beta$  regulatory subunits. PCR products were resolved in 2% agarose gels. The bands correspond to the expected amplicons (KCNMB1, 278 bp; KCNMB2, 137 bp; KCNMB3, 201 bp; KCNMB4, 216 bp). A molecular marker (MK) was run side by side with the samples, and the corresponding base pair (bp) sizes of two of its bands are indicated on the left.  $n = 1-3$ . **(B)** Pictures show representative images of individual confocal sections obtained from partially dissociated ganglia stained with the standard immunohistochemical technique described in Methods. The antibody used is indicated at the top right corner of each image, and the scale bar was 40  $\mu\text{m}$ . Percentage of positive cells with respect to PGP 9.5 was 47.6 for  $\beta 2$  ( $n = 63$  and 30 cells, PGP 9.5 and  $\beta 2$ , respectively), 6.3 for  $\beta 3$  ( $n = 32$  and 2 cells, PGP 9.5 and  $\beta 3$ , respectively) and 129.2 for  $\beta 4$  ( $n = 24$  and 31 cells, PGP 9.5 and  $\beta 2$ , respectively). Percentage of  $\beta 4$  positive cells with respect to BK $\alpha$  was 116 ( $n = 263$  and 305 cells, BK $\alpha$  and  $\beta 4$ , respectively).

representation of the genomic organization leading to these 3' ends is presented in **Figure 3B**. Note that only one of the canine previously described assemblies (**Fig. 3B, d**) was present in ICG. The other two assemblies identified (**Fig. 3B, b** and **c**) had not been formerly reported in canine tissue.

## Discussion

In the present work, we have explored the molecular identity of BK channels in canine ICG. Our results point at two different levels of the BK channel macromolecular complexity: (1) The principal  $\beta$  subunit expressed in ICG is  $\beta 4$ . This

subunit has been reported to be mainly expressed in brain and has complex  $\text{Ca}^{2+}$ -dependent effects.<sup>9</sup> However, the  $V_{1/2}$  values obtained at different  $\text{Ca}^{2+}$  concentrations in canine IC neurons do not completely match those previously reported in similar conditions in heterologous expression systems. (2) Multiple BK $\alpha$  splice



**Figure 2.** Calcium- and voltage-dependence of single BK channels in IC neurons. *Left*, representative single channel traces elicited by voltage steps to +40 mV at the indicated calcium concentrations. Arrows indicate the closed state of the channel. *Right*, plot of the open probability ( $P_o$ ) as a function of voltage at increasing calcium concentrations. Data are displayed as the mean  $\pm$  SE ( $n = 1-10$ ).  $V_{1/2}$  are  $-35.7 \pm 11.1$  mV,  $29.8 \pm 2.2$  mV and  $70.0 \pm 15.1$  mV (10, 3 and  $0.3 \mu\text{M Ca}^{2+}$ , respectively).

variants are expressed in canine ICG. We have characterized, for the first time, the BK $\alpha$  splice variants expressed in this native preparation. However, we cannot address to which extent each of them will effectively contribute to BK current in IC neurons. It is notable that the specific splice variants' combination identified in canine ICG is somewhat different from that detected in canine brain (Table 2). This observation supports the accepted idea of the occurrence of BK $\alpha$  tissue-specific hallmarks given by a particular set of splice variants, which confer unique tissue-specific identity to  $I_{BK}$ . Interestingly, we have identified the presence of STREX at SS2 and of slo27 at SS4. These insertions result in the inclusion of 58 and 27 extra aminoacids, respectively, and thus are suggested to cause bigger effects on  $I_{BK}$  than the shorter insertions that we identified in canine tissues. In fact, several studies have been published that show major effects of the independent inclusion of either variant on  $I_{BK}$ .<sup>4,5,18,20</sup> Also, some attempts have been made to study the effects of the simultaneous presence of multiple variants.<sup>21,22</sup> Again, our  $V_{1/2}$  values obtained in canine IC neurons deviate from published results. Our observations

suggest that the MSS N-terminus may be expressed in canine ICG, and that the original transcripts will carry one or more insertions. Although at present we cannot define the exact combination of BK $\alpha$  splice variants in ICG, the anticipated portrait is quite different from the expression vectors used in heterologous transfection experiments. Thus, the mismatching between the results presented here and those published elsewhere may not be unexpected. Moreover, one has to bear in mind that the heterologous expression studies are often performed with a single variant, which excludes the possibility of heterotetramers, a channel arrangement likely to occur in our native preparation. Additionally, up to four BK $\beta$  can be assembled with the BK $\alpha$  tetramer to render functional channels. Both the number of  $\beta$  subunits and its identity are unfixed and influence BK channel properties.<sup>23</sup> Little is known about the interplay among alternatively spliced  $\alpha$  subunit variants and  $\beta$  subunits. Petrik and Brenner performed an elegant characterization of the effects of  $\beta 4$  on STREX BK channels.<sup>18</sup> These studies were performed using a mouse zero BK $\alpha$  subunit, a framework that does not resemble much the BK $\alpha$

in our native preparation. Also, Erxleben and collaborators studied the effects of STREX inclusion and  $\beta 1$  modulation on  $I_{BK}$ .<sup>20</sup> Although their assays were performed in the BK $\alpha$  N-terminus MSS background, our results present differences with their observations in terms of calcium- and voltage-dependence, probably due to the predominant expression of  $\beta 4$  in IC neurons. Comparisons among native and heterologous expression preparations are difficult to interpret. Still, our molecular results suggest that BK channels expressed in canine ICG are probably composed of BK $\alpha$  bearing more than one splice variant plus  $\beta 4$  and/or  $\beta 1$ . In this line, we recently demonstrated that 100 nM Iberitoxin blocked BK channels in IC neurons. However, this block occurred at times between 3 and 14 min (data not shown). This behavior perhaps reflects the heterogeneous subunit arrangements of BK channels in ICG, in agreement with the molecular evidence presented here. This heterogeneous expression anticipated in canine ICG for both  $\alpha$  and  $\beta$  subunits adds further complexity to the biophysical response of the channel, and warrants further investigation. For example, experiments showing inhibition



of BK current by Protein Kinase A, or an apparent sensitivity to inhibition by oxidation would be indicative of the predominance of BK $\alpha$ -STREX channels<sup>20</sup> in IC neurons.

In summary, in the present addendum we provide molecular evidence for the presence, in canine ICG, of BK channel components known to affect channel function. Although our study does not cover all the possible splice variants reported for this type of channel, it is reasonable to postulate that the splice variants we identified collectively contribute to the functional effect of the BK channel on membrane excitability observed in this native tissue.

## Materials and Methods

**Canine IC ganglia and neuron isolation.** Parasympathetic ganglia and neurons from the atrial ganglionated plexi of the dog were dissociated as previously described.<sup>3</sup> Briefly, canine hearts were obtained, and the fat pads on the ventral, lateral and dorsal aspects of the atrium were quickly removed and placed in a normal Krebs (NK) solution in ice. Individual ganglia were removed from the fat pads and cleaned under a dissection scope. The ganglia were either flash-frozen in liquid N<sub>2</sub> for subsequent RNA extraction or dissociated with 0.1% collagenase-elastase and 0.2% trypsin. Individual cells were obtained by triturating the remaining tissue with a Pasteur pipette. Cells were resuspended in Dulbecco's modified eagle media (DMEM), supplemented with 1% fetal bovine serum, 100  $\mu$ g/ml penicillin-streptomycin, 2 mM glutamax, 10  $\mu$ g/ml S7 and 0.11 g/ml piruvic acid; and plated on collagen coated bottom glass Petri dishes (MatTek Corp., Ashland, MA). Cells were placed overnight in a CO<sub>2</sub> incubator at 37°C. To obtain partially dissociated ganglia, the trituration step was minimized.

**Immunohistochemistry.** Partially dissociated ganglia were obtained as described above. Ganglia were mounted on glass slides and stained with a standard immunohistochemical technique. Briefly, after fixation with a 4% paraformaldehyde-0.2% picric acid solution and permeabilization with 0.2% Triton

X-100, tissue was incubated overnight at 4°C with 1:100/200 dilutions of primary rabbit polyclonal antibodies against the BK channel  $\alpha$  subunit or the neuronal marker PGP9.5 and a monoclonal antibody against the BK channel  $\beta$  subunits 2–4 ( $\beta$ 2–4). Immunostaining of the  $\beta$ 1 subunit was not performed due to the lack of availability of a reliable antibody. Tissue was then incubated with a 1:1,000 dilution of anti-rabbit Alexa-594 and/or anti-mouse Alexa-488-conjugated secondary antibodies for 2 h at room temperature. Specificity of polyclonal primary antibodies was assessed by preincubation with a control peptide antigen. Ganglia were mounted using Pro-Long antifade mounting media and visualized under a Fluoview Olympus laser scanning confocal microscope (40 oil immersion objective) equipped with argon and He/Ne lasers. Optical sections were taken through the entire volume of the cell with the XY frame set to 512  $\times$  512 pixels and the Z-axis was changed in 0.5 to 1 micron increments. Sections were scanned sequentially to avoid bleeding artifacts. Images shown are individual confocal sections.

**Electrophysiological recordings.** Current measurements were obtained using the patch-clamp technique in excised patch configuration. Solutions were applied with a gravity flow system (speed 1–3 ml/min) to a 150  $\mu$ l bath chamber. Electrode shanks were coated with dental wax and tips fire polished to a tip diameter of approx. 1  $\mu$ m. Electrode resistances were of 8–10 M $\Omega$ . Data acquisition and analysis: Experiments were controlled with an Axopatch 200A amplifier and pClamp 9.0/Digidata 1440A acquisition system (Molecular Devices). All experiments were acquired online for later analyses with Clampfit (Molecular Devices). Voltage activation curves for single-channel experiments were fitted with a two state Boltzmann equation of following form:  $P_o = P_{o_{max}} + (P_{o_{min}} - P_{o_{max}}) / 1 + \exp[(V - V_{1/2})/dV]$ , where  $P_{o_{max}}$  and  $P_{o_{min}}$  are the maximum and minimum  $P_o$  asymptotes, respectively,  $V$  is the holding potential,  $V_{1/2}$  is the voltage for half maximal activation, and  $dV$  represents the slope factor. Single channel data analysis was performed using Clampfit as previously described in Scornik et al.<sup>24</sup> Briefly,

**Table 2.** Splice variants identified in canine ICG and brain

Splice site	Splice variant	ICG	Brain
5'	MANG	–	–
	MSS	+	+
	MDALI	+ (?)	+ (?)
SS1	ZERO	+	–
	+4	+	+
SS2	ZERO	+	+
	+3	+	+
	STREX	+	–
SS3	e22	–	–
	ZERO	+	+
SS4	+8	–	–
	ZERO	+	+
3'	Slo27	+	+
	VYR	+	+
	DEC	+	+
	ERL	+	+


The table displays the results of sequencing the RT-PCR products for canine ICG and brain, as described in methods. The presence (+) or the absence (–) of each splice variant is indicated. Question marks are used to state that we cannot ensure the presence of transcripts starting with the MDALI start site (the three start sites are in the same reading frame, and cDNA sequencing does not allow discrimination). The STREX variant identified in ICG corresponds to the short isoform published (+58aa).

$P_o$  was determined by either all-point amplitude histogram or event detection with 50% amplitude criteria. All recordings were performed at 20–22°C.

**Solutions.** Recordings were done in symmetrical solutions of the following composition (in mM): 140 KCl, 10 HEPES, 2 MgCl<sub>2</sub>. Calcium concentration was titrated to 0.3, 3 or 10  $\mu$ M with glycol-bis(2-aminoethylether)-N,N,N',N'-tetraacetic acid or N-hydroxyethyl-ethylenediamine-triacetic acid and measured with a calcium electrode (Kwik-Tip, World Precision Instruments).

**Drugs.** Collagenase type I, trypsin and trypsin inhibitor (Worthington), D-MEM, NGF 7S, FBS and glutamax (Invitrogen Corp.). All other drugs were obtained from Sigma.

**Antibodies.** Rabbit polyclonal anti-PGP9.5 was from EMD Chemical, mouse monoclonal anti  $\beta$ 2,  $\beta$ 3 and  $\beta$ 4 were from



**Figure 3.** Multiple BK channel splice variants are expressed in canine ICG. **(A)** Picture depicts the results of RT-PCRs performed as described in Methods using specific primers for some BK channel splice sites (SS). A molecular marker (MK) and the PCR products were resolved in 2% agarose gels. Expected amplicons, in bp: SS1: 107 (without insert, -) and 119 (including insert, +); SS2: 172 (-), 346 (+, STREX-short isoform), 355 (+, STREX-long isoform), 259 (+, e22), and 181 (+, +3); SS3: 79 (-) and 103 (+); SS4: 208 (-) and 289 (+). n = 1–4. **(B)** Schematic representation of the genomic organization that leads to the alternative C-terminus identified in canine ICG. The uppermost scheme (a) illustrates the genomic organization of KCNMA1 C-terminus (not in actual scale). The schemes below represent the assemblies identified in ICG (b, c and d) and those previously reported for canine tissue (d and e). Boxes represent exons and are labeled with roman numbers, and dashed lines represent introns. Arrowheads mark the approximate localization of stop codons. Exons that are included in a given splice variant are connected by solid lines. Exons not included are dashed. In e, the box labeled “del a” indicates a deletion of an adenine at the indicated splice site. The resulting amino acid sequences are shown on the right. The arrows indicate splice sites and asterisks mark stop codons. Note that only one of the assemblies previously described (d) was present in ICG. The other two assemblies identified (b and c) had not been formerly reported in canine tissue.

NeuroMab and secondary Alexa conjugated antibodies were from Invitrogen.

**RT-PCR.** Total RNA was extracted from ICG using the RNeasy Micro Kit (Qiagen) and from brain using the RNeasy Fibrous tissue Mini Kit (Qiagen), following the manufacturer’s recommendations. DNase treatment of the RNA samples was performed using the DNA-free Kit from Ambion. Reverse transcription was performed using High Capacity cDNA Reverse Transcription Kit (Applied Biosystems) as per manufacturer’s directions. Primers used are available upon request. The PCR products were analyzed by electrophoresis on 2% agarose gels.

**Sequencing.** DNA was extracted from bands in the agarose gels using the PCR clean-up Gel extraction kit (Macherey-Nagel GmbH and Co. KG). All DNAs were directly sequenced in both directions (Big Dye Terminator v3.1 cycle sequencing kit and 3130XL Genetic Analyzer, both from Applied Biosystems) and the obtained sequences were compared with their respective reference sequences (KCNMA1 from *Canis lupus familiaris* when available or alternatively from *Mus musculus*; GenBank gene IDs 403984 and

16531, respectively. *Canis lupus familiaris* KCNMB1 to B4 GenBank gene IDs: 403983, 488179, 488085 and 474447, respectively).

#### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

#### Acknowledgments

The authors would like to thank Yuesheng Wu for technical assistance in initial molecular biology experiments. This work was supported in part by a National Institute of Health grant HL073161 to GJP, an American Heart Association grant 0335446T to FSS, a grant from the Obra Social “La Caixa” to RB and a Sara Borrell postdoctoral fellowship CD11/00063 from the Instituto de Salud Carlos III to ES.

#### References

1. Franciolini F, Hogg R, Catacuzzeno L, Petris A, Trequatrini C, Adams DJ. Large-conductance calcium-activated potassium channels in neonatal rat intracardiac ganglion neurons. *Pflugers Arch* 2001; 441:629-38; PMID:11294244; <http://dx.doi.org/10.1007/s004240000471>.
2. Merriam LA, Scornik FS, Parsons RL. Ca(2+)-induced Ca(2+) release activates spontaneous miniature outward currents (SMOCs) in parasympathetic cardiac neurons. *J Neurophysiol* 1999; 82:540-50; PMID:10444654.
3. Pérez GJ, Desai M, Anderson S, Scornik FS. Large-conductance calcium-activated potassium current modulates excitability in isolated canine intracardiac neurons. *Am J Physiol Cell Physiol* 2013; 304:C280-6; PMID:23195072; <http://dx.doi.org/10.1152/ajp-cell.00148.2012>.
4. Chen L, Tian L, MacDonald SH, McClafferty H, Hammond MS, Huibant JM, et al. Functionally diverse complement of large conductance calcium- and voltage-activated potassium channel (BK) alpha-subunits generated from a single site of splicing. *J Biol Chem* 2005; 280:33599-609; PMID:16081418; <http://dx.doi.org/10.1074/jbc.M505383200>.
5. Ha TS, Jeong SY, Cho SW, Jeon Hk, Roh GS, Choi WS, et al. Functional characteristics of two BKCa channel variants differentially expressed in rat brain tissues. *Eur J Biochem* 2000; 267:910-8; PMID:10651830; <http://dx.doi.org/10.1046/j.1432-1327.2000.01076.x>.
6. Poulsen AN, Wulf H, Hay-Schmidt A, Jansen-Olesen I, Olesen J, Klaerke DA. Differential expression of BK channel isoforms and beta-subunits in rat neuro-vascular tissues. *Biochim Biophys Acta* 2009; 1788:380-9; PMID:18992709; <http://dx.doi.org/10.1016/j.bbame.2008.10.001>.
7. Xie J, McCobb DP. Control of alternative splicing of potassium channels by stress hormones. *Science* 1998; 280:443-6; PMID:9545224; <http://dx.doi.org/10.1126/science.280.5362.443>.
8. Behrens R, Nolting A, Reimann F, Schwarz M, Waldschütz R, Pongs O. hKCNMB3 and hKCNMB4, cloning and characterization of two members of the large-conductance calcium-activated potassium channel beta subunit family. *FEBS Lett* 2000; 474:99-106; PMID:10828459; [http://dx.doi.org/10.1016/S0014-5793\(00\)01584-2](http://dx.doi.org/10.1016/S0014-5793(00)01584-2).
9. Brenner R, Jegla TJ, Wickenden A, Liu Y, Aldrich RW. Cloning and functional characterization of novel large conductance calcium-activated potassium channel beta subunits, hKCNMB3 and hKCNMB4. *J Biol Chem* 2000; 275:6453-61; PMID:10692449; <http://dx.doi.org/10.1074/jbc.275.9.6453>.

10. Contreras GF, Neely A, Alvarez O, Gonzalez C, Latorre R. Modulation of BK channel voltage gating by different auxiliary  $\beta$  subunits. *Proc Natl Acad Sci U S A* 2012; 109:18991-6; PMID:23112204; <http://dx.doi.org/10.1073/pnas.1216953109>.
11. Meera P, Wallner M, Toro L. A neuronal beta subunit (KCNMB4) makes the large conductance, voltage- and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel resistant to charybdotoxin and iberiotoxin. *Proc Natl Acad Sci U S A* 2000; 97:5562-7; PMID:10792058; <http://dx.doi.org/10.1073/pnas.100118597>.
12. Orio P, Torres Y, Rojas P, Carvacho I, Garcia ML, Toro L, et al. Structural determinants for functional coupling between the beta and alpha subunits in the  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  (BK) channel. *J Gen Physiol* 2006; 127:191-204; PMID:16446507; <http://dx.doi.org/10.1085/jgp.200509370>.
13. Xia XM, Ding JP, Lingle CJ. Molecular basis for the inactivation of  $\text{Ca}^{2+}$ - and voltage-dependent BK channels in adrenal chromaffin cells and rat insulinoma tumor cells. *J Neurosci* 1999; 19:5255-64; PMID:10377337.
14. Tseng-Crank J, Foster CD, Krause JD, Mertz R, Godinot N, DiChiara TJ, et al. Cloning, expression, and distribution of functionally distinct  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel isoforms from human brain. *Neuron* 1994; 13:1315-30; PMID:7993625; [http://dx.doi.org/10.1016/0896-6273\(94\)90418-9](http://dx.doi.org/10.1016/0896-6273(94)90418-9).
15. Fodor AA, Aldrich RW. Convergent evolution of alternative splices at domain boundaries of the BK channel. *Annu Rev Physiol* 2009; 71:19-36; PMID:18694345; <http://dx.doi.org/10.1146/annurev.physiol.010908.163124>.
16. Orio P, Rojas P, Ferreira G, Latorre R. New disguises for an old channel: MaxiK channel beta-subunits. *News Physiol Sci* 2002; 17:156-61; PMID:12136044.
17. Gruslova A, Semenov I, Wang B. An extracellular domain of the accessory  $\beta 1$  subunit is required for modulating BK channel voltage sensor and gate. *J Gen Physiol* 2012; 139:57-67; PMID:22155735; <http://dx.doi.org/10.1085/jgp.201110698>.
18. Petrik D, Brenner R. Regulation of STREX exon large conductance, calcium-activated potassium channels by the beta4 accessory subunit. *Neuroscience* 2007; 149:789-803; PMID:17945424; <http://dx.doi.org/10.1016/j.neuroscience.2007.07.066>.
19. Lai GJ, McCobb DP. Regulation of alternative splicing of Slo  $\text{K}^+$  channels in adrenal and pituitary during the stress-hyporesponsive period of rat development. *Endocrinology* 2006; 147:3961-7; PMID:16675526; <http://dx.doi.org/10.1210/en.2005-1551>.
20. Erxleben C, Everhart AL, Romeo C, Florance H, Bauer MB, Alcorta DA, et al. Interacting effects of N-terminal variation and strex exon splicing on slo potassium channel regulation by calcium, phosphorylation, and oxidation. *J Biol Chem* 2002; 277:27045-52; PMID:12016222; <http://dx.doi.org/10.1074/jbc.M203087200>.
21. Jones EM, Gray-Keller M, Fettiplace R. The role of  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel spliced variants in the tonotopic organization of the turtle cochlea. *J Physiol* 1999; 518:653-65; PMID:10420004; <http://dx.doi.org/10.1111/j.1469-7793.1999.0653p.x>.
22. Ramanathan K, Michael TH, Fuchs PA. beta subunits modulate alternatively spliced, large conductance, calcium-activated potassium channels of avian hair cells. *J Neurosci* 2000; 20:1675-84; PMID:10684869.
23. Wang YW, Ding JP, Xia XM, Lingle CJ. Consequences of the stoichiometry of Slo1 alpha and auxiliary beta subunits on functional properties of large-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels. *J Neurosci* 2002; 22:1550-61; PMID:11880485.
24. Scornik FS, Bucciero RS, Wu Y, Selga E, Bosch Calero C, Brugada R, et al. DiBAC4(3) hits a "sweet spot" for the activation of arterial large-conductance  $\text{Ca}^{2+}$ -activated potassium channels independently of the  $\beta 1$ -subunit. *Am J Physiol Heart Circ Physiol* 2013; 304:H1471-82; PMID:23542916; <http://dx.doi.org/10.1152/ajpheart.00939.2012>.