

Article

# Application of Ultra-High-Performance Liquid Chromatography Coupled with LTQ-Orbitrap Mass Spectrometry for the Qualitative and Quantitative Analysis of *Polygonum multiflorum* Thumb. and Its Processed Products

Teng-Hua Wang<sup>1,2</sup>, Jing Zhang<sup>1</sup>, Xiao-Hui Qiu<sup>1</sup>, Jun-Qi Bai<sup>1</sup>, You-Heng Gao<sup>2,\*</sup> and Wen Xu<sup>1,\*</sup>

Received: 13 July 2015; Accepted: 3 December 2015; Published: 26 December 2015

Academic Editor: Thomas J. Schmidt

<sup>1</sup> Lab of Chinese Materia Medica Preparation, the Second College of Clinic Medicine, Guangzhou University of Chinese Medicine; Guangdong Province Institute of TCM, Guangzhou 510006, China; wangtenghua88@126.com (T.-H.W.); ginniezj@163.com (J.Z.); gxhqxh@medmail.com.cn (X.-H.Q.); baijunqibai@126.com (J.-Q.B.)

<sup>2</sup> School of Chinese Materia Medica, Guangzhou University of Chinese Medicine, Guangzhou 510006, China

\* Correspondence: gaoyouheng@gzucm.edu.cn (Y.-H.G.); freexuwen@163.com (W.X.); Tel./Fax: +86-20-3931-8571 (W.X.)

**Abstract:** In order to quickly and simultaneously obtain the chemical profiles and control the quality of the root of *Polygonum multiflorum* Thumb. and its processed form, a rapid qualitative and quantitative method, using ultra-high-performance liquid chromatography coupled with electrospray ionization-linear ion trap-Orbitrap hybrid mass spectrometry (UHPLC-LTQ-Orbitrap MS<sup>n</sup>) has been developed. The analysis was performed within 10 min on an AcQuity UPLC™ BEH C<sub>18</sub> column with a gradient elution of 0.1% formic acid-acetonitrile at flow rate of 400 μL/min. According to the fragmentation mechanism and high resolution MS<sup>n</sup> data, a diagnostic ion searching strategy was used for rapid and tentative identification of main phenolic components and 23 compounds were simultaneously identified or tentatively characterized. The difference in chemical profiles between *P. multiflorum* and its processed preparation were observed by comparing the ions abundances of main constituents in the MS spectra and significant changes of eight metabolite biomarkers were detected in the *P. multiflorum* samples and their preparations. In addition, four of the representative phenols, namely gallic acid, *trans*-2,3,5,4'-tetra-hydroxystilbene-2-O-β-D-glucopyranoside, emodin and emodin-8-O-β-D-glucopyranoside were quantified by the validated UHPLC-MS/MS method. These phenols are considered to be major bioactive constituents in *P. multiflorum*, and are generally regarded as the index for quality assessment of this herb. The method was successfully used to quantify 10 batches of *P. multiflorum* and 10 batches of processed *P. multiflorum*. The results demonstrated that the method is simple, rapid, and suitable for the discrimination and quality control of this traditional Chinese herb.

**Keywords:** *Polygonum multiflorum*; LTQ-Orbitrap; qualitative and quantitative

## 1. Introduction

The root of *Polygonum multiflorum* Thumb. (*Fallopia multiflora*), well known as He-shou-wu in China, has been widely used as a tonic and purgative in traditional Chinese medicine for thousands of years [1]. Previous phytochemical studies revealed that the constituents of *P. multiflorum* are mainly two kinds of active constituents—anthraquinones and stilbenes—as well as other compounds such

as flavonoids, tannins and phospholipids [2–4]. Recently, four novel stilbene derivatives, named polygonumosides A–D, were isolated from the processed root of *P. multiflorum* [5].

The anthraquinones, such as emodin (EM) and emodin-8-*O*- $\beta$ -*D*-glucopyranoside (EMG), provide immunomodulating, anti-inflammatory, anticancer, antimutagenic, antibacterial, gastrointestinal smooth muscle prokinetic actions, dose-dependent protection against myocardial ischemia-reperfusion injury and cerebral ischemia-induced infarct volume reduction biological effects [6–9]. Some research supports the notion that the stilbenes in PM possess anti-inflammatory [10–12], antioxidant activity [13–16], anti-hyperlipidaemia [17], anti-melanogenic activity, prophylactic and therapeutic activity against Alzheimer's disease and Parkinson's disease [18–20], free radical scavenging activity [21], and hair growth properties [22–25]. According to Chinese medicine theory, the crude *P. multiflorum* (CPM) should be processed before use, involving steaming the crude roots with or without black soybean extract (namely *Paozhi*), which could reduce the laxative effects and strengthen the tonic effects [26]. Both the crude *P. multiflorum* (CPM) and processed *P. multiflorum* (PPM) have been officially listed in the successive editions of the Chinese Pharmacopoeia.

As a consequence of the potential medicinal value of PM, several methods have been published for identification and/or quantification of its chemical components, such as high performance liquid chromatography coupled with photodiode array (HPLC-PDA) detection [27], capillary zone electrophoresis (CZE) [28], and HPLC-DAD coupled with electrospray ionization tandem mass spectrometry (ESI-MS<sup>n</sup>) [29,30]. The present method of using several ingredients in evaluating PM preparations cannot reflect the overall changes of chemical composition that occur during the course of processing. Any new method should overcome this problem by using multi-target ingredient determination and fingerprinting analysis technologies simultaneously, which could control the preparation process and product quality precisely, and ensure the stability and effectiveness of products.

In our previous work, the fragmentation pathways of typical constituents and chemical profiles of PM have been studied by an on-line UHPLC-ESI-linear ion trap-Orbitrap hybrid mass spectrometry (LTQ-Orbitrap) method [31,32]. UHPLC coupled with the high resolution tandem mass spectrometric techniques have been proven to be a powerful tool for the rapid identification of unknown components in botanical extracts. In this paper, a simple and rapid method for the comprehensive qualitative and quantitative analysis of the major constituents was successfully developed for quality evaluation of CPM and PPM. Up to now, this is the first report on the simultaneous identification and determination of multiple components in botanical herb products by an UHPLC-LTQ-Orbitrap MS<sup>n</sup> technique.


## 2. Results and Discussion

### 2.1. Optimization of the LC-MS Conditions


The chromatographic conditions, such as the chromatographic column and mobile phase, were optimized to achieve the best separation efficiency. Three reversed-phase chromatographic columns, including a Hypersil C<sub>18</sub> (2.1 mm × 100 mm, 5 μm, Thermo Fisher Scientific, Carlsbad, CA, USA), a Kinetex XB C<sub>18</sub> (2.1 mm × 100 mm, 1.7 μm, Phenomenex, Torrance, WA, USA) and an AcQuity UPLC<sup>TM</sup> BEH C<sub>18</sub> column (2.1 mm × 50 mm, 1.7 μm, Waters, Milford, MA, USA), were selected to test the separation ability of the four investigated compounds. It was shown that an UHPLC system with a 1.7 μm small particle size column had more powerful separation ability with a higher peak resolution and the total analysis time was less than 10 min, which was approximately fourfold faster than that for a conventional column packed with 5 μm particles (Figure 1). Thus the AcQuity UPLC<sup>TM</sup> BEH C<sub>18</sub> column (2.1 mm × 50 mm, 1.7 μm) was proved to be the best in this application.

The experimental results showed that an acetonitrile-H<sub>2</sub>O mobile phase separated the target compounds more effectively (Figure 2B) than a methanol-H<sub>2</sub>O system (Figure 2A). Thus acetonitrile was selected as the organic phase, so as to obtain the optimal separation of adjacent peaks and to avoid peak tailing. Formic acid (Figure 2B), acetic acid (Figure 2C), ammonium acetate (Figure 2D) and ammonium formate (Figure 2E) were added to the mobile phase, respectively, to achieve high MS

sensitivity and restrain peak tailing. Formic acid was screened as an effective mobile phase additive with which improvements of peak shape, peak width and decreased ion suppression effects could be directly and efficiently obtained, compared to others.



**Figure 1.** The extracted ion chromatograms (EIC) of gallic acid (GA), *trans*-2,3,5,4'-tetrahydroxystilbene-2-*O*- $\beta$ -D-glucopyranoside (THSG), emodin-8-*O*- $\beta$ -D-glucopyranoside (EMG) and emodin (EM) with three different reversed-phase columns. (A) AcQuity UPLC™ BEH C<sub>18</sub> column (2.1 mm  $\times$  50 mm, 1.7  $\mu$ m); (B) Kinetex XB C<sub>18</sub> (2.1 mm  $\times$  100 mm, 1.7  $\mu$ m) and (C) Hypersil C<sub>18</sub> (2.1 mm  $\times$  100 mm, 5  $\mu$ m). The mobile phases consisted of acetonitrile (a) and water containing 0.1% formic acid (b), with the following elution gradient program: (A) used the optimized mobile phase and gradient (see Section 3.2); (B) 13% a (0 min), 35% a (3.5 min), 90% a (7.5 min), 95% a (8.5 min) and 95% a (12 min); (C). 13% a (0 min), 35% a (12 min), 90% a (32 min), 95% a (35 min) and 95% a (40 min).



**Figure 2.** The extracted ion chromatograms (EIC) of gallic acid (GA), *trans*-2,3,5,4'-tetrahydroxystilbene-2-*O*- $\beta$ -D-glucopyranoside (THSG), emodin-8-*O*- $\beta$ -D-glucopyranoside (EMG) and emodin (EM) under five different chromatographic conditions: (A) methanol/water system containing 0.1% formic acid; acetonitrile/water system; (B) containing 0.1% formic acid; (C) containing 0.1% acetic acid; (D) containing 0.1% ammonium acetate and (E) containing 0.1% ammonium formate.

## 2.2. Sample Extraction Optimization

In order to optimize the sample extraction conditions, the factors affecting the extraction efficiency of the four main compounds, including extraction solvents, solid-liquid ratio and extraction times, were studied. In our research, the extraction efficiency was evaluated by using a single factor method and choosing the extraction yield (extraction yield = the extract weight  $\times$  the content of target analyte/the weight of crude drug  $\times$  100%) as the index. The results suggested that single-solvent system (pure methanol) was superior to a binary solvent system in the extraction of all four compounds (Table 1). The extraction yield was as high as 5.27% (summation of the four test components) as a maximum value with methanol at 100%. Furthermore, the extraction yield increased by 75.9% (summation of the four test components) when the amount of liquid was increased from 10 to 25 mL (*i.e.*, solid to liquid ratio from 1:50 to 1:125). When the liquid amount was 50 mL (solid to liquid ratio 1:250), the average extraction yield and extraction yield of gallic acid showed no obvious increase. Thus the solid to liquid ratio of 1:125 was recommended. It was also found that when the extraction time was within 30 min, the extraction yield increased significantly with extraction time. The total extraction yield was 5.85% and the highest yield was reached at an extraction time of 30 min. After that, the yield decreased slightly with time. According to the above results, the optimum extraction conditions were: solid-liquid ratio 1:125 and extraction with 25 mL of 100% methanol at reflux for 30 min.

**Table 1.** Effects of methanol concentration, solid-liquid ratio and extraction time on extraction yield <sup>a</sup>.

Compound	Methanol Concentration (%)			Solid-Liquid Ratio			Extraction Time (min)		
	50	70	100	1:50	1:125	1:250	15	30	60
Gallic acid	0.46	0.52	0.67	0.69	0.88	0.89	0.26	0.77	0.73
Emodin	1.25	1.42	1.55	1.23	1.75	1.77	0.37	1.88	1.68
<i>trans</i> -2,3,5,4'-Tetrahydroxy-stilbene-2-O- $\beta$ -D-glucopyranoside	1.26	1.32	1.61	1.18	1.93	1.69	0.79	1.78	1.86
Emodin-8-O- $\beta$ -D-glucopyranoside	0.86	1.46	1.44	0.43	1.65	1.44	0.73	1.42	1.22

<sup>a</sup> extraction yield (%) = the extract weight  $\times$  the content of target analyte/the weight of crude drug  $\times$  100%.

## 2.3. Tentative Identification of the Major Compounds

The four reference compounds (gallic acid, EM, THSG and EMG) were initially analyzed to obtain the corresponding retention times and characteristic fragmentation pathway data (Figure 3). The precise quasi-molecular and fragment ions were determined within a reasonable degree of measurement error using the Orbitrap instrument (the mass error is less than 2 ppm in most cases). Other clues such as potential elemental composition, degree of unsaturation and fractional isotope abundance of the compounds were also utilized for structural conformation. In our previous work, the fragmentation rules and diagnostic fragment ions of anthraquinones, stilbenes, tannins and naphthalenes have been investigated, which provided useful information for the elucidation of the chemical structures of the prescription components. As for the untargeted compounds, the characteristic diagnostic ions could be used to filter and classify them into a particular chemical family.

Twenty five compounds, including anthraquinones, stilbenes, tannins and naphthalenes were primarily identified in both of the extracts of CPM and PPM, and three compounds were only detected in PPM, which were probably new components produced during the steaming process (Table 2).


**Table 2.** Identification of the chemical constituents in methanol extracts of CPM and PPM by UHPLC-LTQ-Orbitrap MS.

No.	$t_R$ (min)	Precursor ions [M – H]	Formula	Mass Error (ppm)	MS <sup>n</sup>	Identification
1	0.89	341.1078 377.0843 [M + Cl]	C <sub>12</sub> H <sub>21</sub> O <sub>11</sub>	0.18	MS <sup>2</sup> [377]: 341 (100), 215 (15) MS <sup>3</sup> : 179 (100), 161 (23), 143 (22), 113 (17)	Sucrose *
2	1.12	179.0556 225.0609 [M + HCOO]	C <sub>6</sub> H <sub>11</sub> O <sub>6</sub>	3.38	MS <sup>2</sup> : 161 (100), 143 (90), 119 (44), 113 (40), 89 (40)	Glucose *
3	2.17	169.0136	C <sub>7</sub> H <sub>5</sub> O <sub>5</sub>	2.66	MS <sup>2</sup> : 125 (100) MS <sup>3</sup> : 81	gallic acid *
4	4.24	577.1330	C <sub>30</sub> H <sub>25</sub> O <sub>12</sub>	−1.8	MS <sup>2</sup> : 425 (100), 407 (48), 457 (20), 471 (17), 289 (10) MS <sup>3</sup> : 407 (100)	procyanidin B
5	4.34	289.0708	C <sub>15</sub> H <sub>13</sub> O <sub>6</sub>	0.47	MS <sup>2</sup> : 245 (100), 205 (42), 179 (19) MS <sup>3</sup> : 203 (100), 227 (23), 187 (22), 161 (20)	epicatechin/catechin
6 #	4.48	531.1488	C <sub>26</sub> H <sub>27</sub> O <sub>12</sub>	−2.2	MS <sup>2</sup> : 369 (100), 351 (29), 405 (21), 243 (18) MS <sup>3</sup> : 351 (100)	unknown
7 #	4.63	549.1594	C <sub>26</sub> H <sub>29</sub> O <sub>13</sub>	−0.97	MS <sup>2</sup> : 387 (100), 459 (73), 531 (22), 297 (16)	unknown
8	4.82	577.1330	C <sub>30</sub> H <sub>25</sub> O <sub>12</sub>	−1.53	MS <sup>2</sup> : 425 (100), 407 (48), 457 (20), 471 (17), 289 (10) MS <sup>3</sup> : 407 (100)	procyanidin B
9 #	4.98	421.1123	C <sub>20</sub> H <sub>21</sub> O <sub>10</sub>	−1.38	MS <sup>2</sup> : 259 (100) MS <sup>3</sup> : 259 (100), 331 (50), 128 (20)	6-methoxyl-2-acetyl-3-methyljuglone-8-O-glu
10	5.02	613.1751 [M + HCOO]	C <sub>27</sub> H <sub>33</sub> O <sub>16</sub>	−1.21	MS <sup>2</sup> : 405 (100), 567 (36) MS <sup>3</sup> : 243 (100)	tetrahydroxystilbene-O-di-glu
11	5.37	405.1177 811.2428 [2M – H]	C <sub>20</sub> H <sub>21</sub> O <sub>9</sub>	−0.74	MS <sup>2</sup> : 243 (100) MS <sup>3</sup> : 225 (100), 149 (79), 137 (73), 215 (70), 173 (36)	THSG *
12	5.41	557.1286	C <sub>27</sub> H <sub>25</sub> O <sub>13</sub>	−0.66	MS <sup>2</sup> : 313 (100), 243 (30), 405 (20), 169 (5) MS <sup>3</sup> : 169 (100), 125 (20), 151 (20), 295 (17)	tetrahydroxystilbene-O-(galloyl)-glu
13	5.71	557.1285	C <sub>27</sub> H <sub>25</sub> O <sub>13</sub>	−0.82	MS <sup>2</sup> : 313 (100), 243 (80), 405 (70), 169 (10) MS <sup>3</sup> : 169 (100), 125 (20), 151 (20), 295 (17)	tetrahydroxystilbene-O-(galloyl)-glu
14	5.72	431.0970	C <sub>21</sub> H <sub>19</sub> O <sub>10</sub>	−0.63	MS <sup>2</sup> : 269 (100) MS <sup>3</sup> : 225 (100), 241 (21), 181 (4)	emodin-1-O-glu
15	5.75	567.1488	C <sub>29</sub> H <sub>27</sub> O <sub>12</sub>	−1.59	MS <sup>2</sup> : 243 (100) MS <sup>3</sup> : 225 (100), 215 (72), 149 (67)	tetrahydroxystilbene-O-(caffeoyl)-glu

Table 2. Cont.

No.	$t_R$ (min)	Precursor ions [M – H]	Formula	Mass Error (ppm)	MS <sup>n</sup>	Identification
16	5.90	551.1543	C <sub>29</sub> H <sub>27</sub> O <sub>11</sub>	–0.89	MS <sup>2</sup> : 405 (100), 243 (31)	tetrahydroxystilbene-2-O-(coumaroyl)-glu
17	5.92	447.0919	C <sub>21</sub> H <sub>19</sub> O <sub>11</sub>	–0.64	MS <sup>2</sup> : 303 (100), 285 (100) MS <sup>3</sup> : 285 (100), 177 (11), 125 (8)	citreorosein-O-glu
18	6.02	407.1334	C <sub>20</sub> H <sub>23</sub> O <sub>9</sub>	–0.64	MS <sup>2</sup> : 245 (100) MS <sup>3</sup> : 230 (100)	torachryson-O-glu
19	6.11	431.0973	C <sub>21</sub> H <sub>19</sub> O <sub>10</sub>	0.11	MS <sup>2</sup> : 269 (100) MS <sup>3</sup> : 225 (100), 241 (21), 197 (5)	emodin-8-O-glu *
20	6.26	517.0978	C <sub>24</sub> H <sub>21</sub> O <sub>13</sub>	0.26	MS <sup>2</sup> : 473 (100), 431 (10) MS <sup>3</sup> : 269 (100), 311 (12), 225 (5)	emodin-O-(malonyl)-glu
21	6.38	445.1127 491.1182 [M + HCOO]	C <sub>22</sub> H <sub>21</sub> O <sub>10</sub>	–0.22	MS <sup>2</sup> : 283 (100), 445 (42) MS <sup>3</sup> : 240 (100), 268 (36)	physcion-8-O-glu
22	6.42	313.0345	C <sub>16</sub> H <sub>9</sub> O <sub>7</sub>	0.96	MS <sup>2</sup> : 269 MS <sup>3</sup> : 225, 241, 197	carboxyl emodin
23	7.21	269.0447	C <sub>15</sub> H <sub>9</sub> O <sub>5</sub>	0.33	MS <sup>2</sup> : 225, 241	Emodin *
24	7.63	269.0446	C <sub>15</sub> H <sub>9</sub> O <sub>5</sub>	0.17	MS <sup>2</sup> : 225, 241, 254	aloe-emodin
25	8.30	283.0605	C <sub>16</sub> H <sub>11</sub> O <sub>5</sub>	1.41	MS <sup>2</sup> : 240	physcion

\* Compared with standard compounds; # Detected only in methanol extracts of PPM.



**Figure 3.** The selected reaction monitoring (SRM) chromatogram of four compounds (A) and the total ion chromatogram (TIC) of typical *P. multiflorum* sample (B). (1) gallic acid (GA), (2) *trans*-2,3,5,4'-tetrahydroxystilbene-2-*O*- $\beta$ -D-glucopyranoside (THSG), (3) emodin-8-*O*- $\beta$ -D-glucopyranoside (EMG) and (4) emodin (EM).


### 2.3.1. Identification of the Main Stilbene Glycosides

Most of the stilbene glycosides in PM showed common fragmentation pathways and two diagnostic fragment ions, 405.1177 ( $C_{20}H_{21}O_9$ ) and 243.0661 ( $C_{14}H_{11}O_4$ ), were used for rapidly extracting unknown tetrahydroxystilbene glucosides. For example, peaks 12 and 13 exhibited a molecular ion at  $m/z$  557.1286 ( $C_{27}H_{25}O_{13}$ ) in the negative ion mode. It produced a base peak at 313.0548 ( $C_{13}H_{13}O_9$ ) and two prominent fragment ions at  $m/z$  405.1164 ( $C_{20}H_{21}O_9$ ) and 243.0649 ( $C_{14}H_{11}O_4$ ) in the  $MS^2$  spectra, indicating it was a tetrahydroxystilbene derivative. Then, the  $m/z$  313 ion generated the characteristic ions of gallic acid, such as  $m/z$  169 and 125. Thus, peaks 12 and 13 were tentatively identified as tetrahydroxystilbene-*O*-(galloyl)-glucoside and the proposed fragmentation pathways are shown in Figure 4.

### 2.3.2. Identification of the Main Anthraquinones

Most of the anthraquinones in PM are emodin and physcion derivatives and they generated typical adducts  $[M + HCOO]^-$  and/or  $[M - H]^-$  with high intensity in negative ion mode. Diagnostic ions at  $m/z$  269.0447, 225.0544 and 241.0492 could be used for rapid extraction and identification of emodin derivatives. For physcion derivatives, two odd-electron ions ( $OE^-$ ) at  $m/z$  240 and 268 were produced from the simultaneous loss of free radicals on the side chains, which could be used for the identification of such derivatives.

For example, peak 20 gave a  $[M - H]^-$  ion at  $m/z$  517.0978 ( $C_{24}H_{21}O_{13}$ ) and a prominent fragment ion at  $m/z$  473.1063 ( $C_{23}H_{21}O_{11}$ ), as well as a characteristic ion at  $m/z$  431.0960 ( $C_{21}H_{19}O_{10}$ ). Additional fragmentation yielded diagnostic ions at  $m/z$  269 and 225 in the  $MS^3$  spectra. High resolution MS detection indicated it was malonyl-substituted glycoside derivative, which was tentatively identified as emodin-*O*-(malonyl)-glucoside.




**Figure 4.** The proposed fragmentation pathways of peaks 12 and 13.

#### 2.4. Changes of the Relative Intensity of the Main Chemical Components

Statistical analysis based on the main chemical metabolic components clearly showed differences among 10 batches of CPM and 10 batches of PPM by two-way ANOVA and Bonferroni correction. In examining the relative intensity of the 12 main chemical components in more detail, a histogram of compounds revealed the chemical profiles differ between PM and its processed products. In the present study, obvious differences existed in the compounds sucrose, gallic acid, procyanidin B, catechin, THSG, torachryson-*O*-glu, emodin-8-*O*-glu and emodin-*O*-(malonyl)-glu between the 10 batches of CPM and 10 batches of PPM (Figure 5). The contents of those compounds were destroyed may due to long-time steaming [33,34].

Histogram of relative intensity of 12 biomarkers between CPM and PPM



**Figure 5.** Relative intensity of the extracted ion chromatograms (EIC; mean  $\pm$  SD) of 12 biomarkers between PM and its processed products. The peak names were originated from Table 2. Statistics are based on three independent experiments for each sample. \*  $p < 0.05$ , \*\*  $p < 0.01$  and \*\*\*  $p < 0.001$ .

#### 2.5. Quantification of Four Major Phenolic Components of the UHPLC-MS/MS System

A rapid method was developed to quantify four representative phenolic compounds (gallic acid, EM, THSG and EMG) in 10 batches of CPM and 10 batches of PPM that were collected from different regions in China (Supplementary Data, Table S1). Since the four investigated bioactive compounds



vary greatly in polarity, it is difficult to accomplish a satisfactory separation using a conventional HPLC-UV method. The UHPLC-ESI-MS/MS method has demonstrated higher selectivity, sensitivity and shorter running times, thus it was utilized for rapid determination of these biomarkers. The UHPLC conditions and spectrometric parameters transition were optimized as described in Section 3.2. (Figure 3).

### 2.5.1. Calibration Curves, LOD and LOQ

A methanol stock solution containing all the reference standards was prepared and diluted to an appropriate concentration range for the construction of calibration curves. At least six concentrations of the solution were analyzed in triplicate, and then the calibration curves were constructed from peak areas of the reference standards *vs.* their concentrations. The high correlation coefficient values ( $r^2 > 0.9916$ ) indicated good linearity between their peak areas ( $y$ ) and investigated compound concentration ( $x$ ,  $\mu\text{g/mL}$ ) in relatively wide concentration ranges (Table 3).

The limits of detection (LOD) and the limits of quantification (LOQ) under the chromatographic conditions were determined by injecting a series of standard solutions until the signal-to-noise ratios (S/N) for each compound was 3 for LOD and 10 for LOQ (Table 3).

**Table 3.** Calibration curves, LOD and LDQ of the investigated compounds.

Analytes	Linear Regression Data Regression Equation	Test Range ( $\mu\text{g/mL}$ )	$r^2$	LOD <sup>a</sup> ( $\text{ng/mL}$ )	LOQ <sup>b</sup> ( $\text{ng/mL}$ )
Gallic acid	$y = 3598.5x + 205.82$	0.03–4.00	0.9997	0.03	0.10
Emodin	$y = 25639x + 2283.1$	0.01–1.40	0.9916	0.01	0.02
<i>trans</i> -2,3,5,4'-Tetrahydroxy- stilbene-2- <i>O</i> - $\beta$ -D-gluco-pyranoside	$y = 652.2x - 5013.2$	1.15–105.00	0.9997	0.01	0.02
Emodin-8- <i>O</i> - $\beta$ -D-gluco-pyranoside	$y = 7690.7x + 14652$	0.29–114.00	0.9998	0.01	0.06

<sup>a</sup> LOD refers to the limit of detection,  $s/n = 3$ ; <sup>b</sup> LOQ refers to the limit of quantification,  $s/n = 10$ .

### 2.5.2. Precision

The precision of the developed assay was evaluated by analyzing the mixed standard solution at three different concentration levels (high, middle and low) (Table 4). The experiment was repeated six times within one day to determine intra-day precision, while for inter-day variability test, the solution was examined in triplicate for consecutive 3 day. Relative standard deviation (RSD) for each of the marker compounds was calculated respectively, and it was no more than 3.27%.

**Table 4.** Intra- and inter-day variability for the assay of four components.

Compound	Intra-Day ( $n = 6$ )		Inter-Day ( $n = 3$ )	
	Mean ( $\mu\text{g/mL}$ )	RSD <sup>a</sup> (%)	Mean ( $\mu\text{g/mL}$ )	RSD (%)
Gallic acid	0.01	1.27	0.07	3.27
	0.25	1.15	0.25	1.62
	1.00	0.92	0.97	2.74
Emodin	0.00	1.90	0.04	2.14
	0.34	1.67	0.34	1.87
	2.56	1.48	2.50	2.39
<i>trans</i> -2,3,5,4'-Tetrahydroxystilbene-2- <i>O</i> - $\beta$ -D-gluco-pyranoside	2.17	1.45	2.08	1.68
	10.10	1.23	10.12	2.71
	107.67	0.77	106.33	1.65
Emodin-8- <i>O</i> - $\beta$ -D-gluco-pyranoside	0.37	1.14	0.37	2.45
	1.52	1.49	1.50	1.49
	67.88	1.30	68.13	2.57

<sup>a</sup> RSD (%) =  $(\text{SD}/\text{mean}) \times 100\%$ .

### 2.5.3. Repeatability, Stability, Specificity and Selectivity

The repeatability of this method was determined by analyzing six samples from the same batch (Sample A6) using the same preparation procedure described in Section 3.3 of the paper. The RSD values of the four component contents were all less than 3.81% (Table 5). The stability was tested by analysis of the same sample solution (Sample A6) which was at room temperature at different times for 24 h. The RSD values of the four analytes were less than 2.01%, which suggested that it was feasible to analyze samples within 24 h (Table 5).

**Table 5.** Repeatability and stability of the four analytes, expressed as RSD (%).

Compound	Repeatability ( <i>n</i> = 6)		Stability ( <i>n</i> = 6)	
	Mean (μg/mL)	RSD <sup>a</sup> (%)	Mean (μg/mL)	RSD (%)
Gallic acid	0.33	3.81	0.34	2.01
Emodin	1.04	2.53	1.02	1.17
<i>trans</i> -2,3,5,4'-Tetrahydroxystilbene-2-O-β-D-glucopyranoside	27.34	1.85	27.83	1.49
Emodin-8-O-β-D-glucopyranoside	0.83	2.98	0.80	1.37

<sup>a</sup> RSD (%) = (SD/mean) × 100%.

### 2.5.4. Accuracy

Recovery test was used to evaluate the accuracy of this method. A known amount (low, medium and high) of each standard solution was spiked into known amounts of CPM samples (simple A6), and then extracted according to the section of sample pretreatment. The recoveries were counted by the formula: recovery (%) = concentration found/original concentration × 100%. The recoveries of analytes varied from 97.34% to 100.75% (Table 6). The result indicated the reliability and accuracy for the quantitative determination of the constituents.

**Table 6.** Recoveries of the four determined constituents.

Compound	Amount (μg/mL)		Recovery <sup>a</sup> (%)	RSD <sup>b</sup> (%)
	Spiked	Found		
Gallic acid	0.26	0.26 ± 0.00	98.39	1.19
	0.33	0.32 ± 0.01	98.87	1.81
	0.39	0.39 ± 0.00	98.54	0.04
Emodin	0.90	0.91 ± 0.00	100.75	0.06
	1.12	1.13 ± 0.02	100.53	1.43
	1.35	1.33 ± 0.01	98.73	1.41
<i>trans</i> -2,3,5,4'-Tetrahydroxystilbene-2-O-β-D-glucopyranoside	23.46	22.83 ± 0.14	97.34	0.63
	29.32	29.43 ± 0.37	100.37	1.26
	35.19	34.54 ± 0.40	98.17	1.18
Emodin-8-O-β-D-glucopyranoside	7.35	7.17 ± 0.06	97.56	0.77
	9.19	9.16 ± 0.13	99.75	1.46
	11.03	11.07 ± 0.05	100.40	0.43

<sup>a</sup> Recovery (%) = (detected amount – original amount)/spiked amount × 100%. <sup>b</sup> RSD (%) = (SD/mean) × 100%.

### 2.5.5. Sample Analysis

Twenty batches of *P. multiflorum* acquired from different regions of China were determined using the described method. This work disclosed that the variation between different locations and processing methods of this traditional Chinese medicine was obvious. Such variations may presumably be attributed to differences in the cultivation conditions (Table 7).

Table 7. Contents of four components in 20 samples.

No.	Content (mg/g) ( <i>n</i> = 3)			
	Gallic Acid (GA)	Emodin (EM)	2,3,5,4'-Tetrahydroxystilbene-2-O-β-D-glucoside (THSG)	Emodin-8-O-β-D-glucoside (EMG)
A1	0.63 ± 0.01	1.01 ± 0.02	34.43 ± 0.59	21.11 ± 0.34
A2	0.44 ± 0.05	0.23 ± 0.00	4.53 ± 0.15	1.32 ± 0.05
A3	0.32 ± 0.01	0.18 ± 0.00	24.54 ± 0.93	7.29 ± 0.20
A4	0.30 ± 0.04	0.52 ± 0.06	16.27 ± 0.57	7.98 ± 0.17
A5	0.50 ± 0.01	0.82 ± 0.01	24.01 ± 0.76	14.44 ± 0.65
A6	0.31 ± 0.00	2.46 ± 0.04	26.57 ± 0.46	15.77 ± 0.25
A7	0.58 ± 0.01	0.32 ± 0.00	20.55 ± 0.67	3.85 ± 0.08
A8	0.43 ± 0.01	0.10 ± 0.00	24.97 ± 0.58	7.83 ± 0.25
A9	0.60 ± 0.01	2.38 ± 0.05	19.71 ± 0.23	15.87 ± 0.15
A10	0.23 ± 0.00	3.27 ± 0.04	23.51 ± 0.55	14.59 ± 0.31
B1	0.02 ± 0.00	0.49 ± 0.01	2.54 ± 0.04	2.12 ± 0.03
B2	0.09 ± 0.00	0.36 ± 0.00	1.85 ± 0.03	0.35 ± 0.00
B3	0.28 ± 0.00	0.45 ± 0.01	2.63 ± 0.04	3.11 ± 0.04
B4	0.15 ± 0.01	0.50 ± 0.01	1.97 ± 0.05	2.59 ± 0.04
B5	0.90 ± 0.01	0.85 ± 0.03	10.42 ± 0.37	2.51 ± 0.04
C1	0.31 ± 0.01	0.56 ± 0.02	2.05 ± 0.03	3.49 ± 0.04
C2	0.02 ± 0.01	1.13 ± 0.02	5.94 ± 0.10	3.92 ± 0.04
C3	1.31 ± 0.04	0.32 ± 0.01	2.55 ± 0.06	0.47 ± 0.01
C4	0.47 ± 0.01	1.31 ± 0.04	11.71 ± 0.41	4.92 ± 0.09
C5	0.39 ± 0.01	0.67 ± 0.02	8.21 ± 0.10	2.62 ± 0.09

### 3. Experimental

#### 3.1. Reagents and Chemicals

HPLC-grade acetonitrile, methanol and formic acid were purchased from Sigma Aldrich (St. Louis, MO, USA). Ultrapure Water (18.2 MΩ) was produced by Milli-Q water system (Millipore, Bedford, MA, USA). Twenty samples of roots of CPM and PPM were collected from various habitats in China, and authenticated by Professor Zhihai Huang in our lab. The related information is summarized in the Supplementary Data (Table S1). The standards of gallic acid (GA), emodin (EM) and *trans*-2,3,5,4'-tetrahydroxystilbene-2-O-β-D-glucopyranoside (THSG) were obtained from the National Institutes for Food and Drug Control (Beijing, China). Emodin-8-O-β-D-glucopyranoside (EMG; over 98% purity by HPLC) was isolated in our laboratory.

#### 3.2. Chromatography and MS Conditions

LC analyses were performed on a Thermo Accela UHPLC system (Thermo Fisher Scientific, San Jose, CA, USA) equipped with a quaternary pump, a diode-array detector (DAD), an auto-sampler, and a thermostatically column compartment.

After also running optimization analyses on both Hypersil C<sub>18</sub> (2.1 mm × 100 mm, 5 μm) and Kinetex XB C<sub>18</sub> (2.1 mm × 100 mm, 1.7 μm) columns, we found the best overall resolution on an AcQuity UPLC™ BEH C<sub>18</sub> column (2.1 mm × 50 mm, 1.7 μm) at room temperature (see Section 2.1) and used this column for all subsequent runs. The mobile phase was also optimized by comparing an acetonitrile-H<sub>2</sub>O mobile to a methanol-H<sub>2</sub>O system and with the addition of various modifiers to the acetonitrile-H<sub>2</sub>O mobile, including formic acid, acetic acid, ammonium acetate, and ammonium formate, all at 0.1% (see Section 2.1). The final mobile phase was composed of acetonitrile (A) and water containing 0.1% formic acid (B) using the following gradient program: 13% A (0 min), 35% A (3.5 min), 90% A (7.5 min), 95% A (8.5 min) and 95% A (10 min). A pre-equilibration period of 4 min was used between individual runs. The mobile phase flow rate was 400 μL/min, and the injection volume was 2 μL. The online UV spectra were recorded in the range of 200–400 nm.

Mass spectra were acquired using a Thermo-Fisher LTQ-Orbitrap XL hybrid mass spectrometer, which was connected to LC system via an electrospray ionization (ESI) source as interface. The basic conditions of MS analysis were as follows: the mass spectrometer parameters were negative ion mode, ion spray voltage at 3500 V, capillary voltage at 37 V, capillary temperature at 300 °C, sheath gas flow rate at 40 psi and auxiliary gas flow rate at 4 psi. The scan spectra were from *m/z* 150 to 1200.

For qualitative analysis of CPM and PPM, the Orbitrap resolution of survey scan was set as 30000 and MS<sup>n</sup> scan was set 15000. The data-dependent MS<sup>n</sup> scanning was performed to trigger fragmentation spectra of target ions and to prevent repetition by dynamic exclusion settings. For quantitative determinations of four compounds, the MS detection was operated in linear ion trap (LTQ) with selected reaction monitoring (SRM). The ion trap collision induced dissociation (CID) mode was used for SRM fragmentation and the ions transitions are as follows: GA ( $m/z$  169→125), EM ( $m/z$  269→225), THSG ( $m/z$  405→243) and EMG ( $m/z$  431→269). The selected ion width was  $m/z \pm 1$  and the normalized collision energy was set 35%.

### 3.3. Sample Preparation

The dried roots were powdered to a homogeneous size by a mill, sieved through a No.60 mesh (250  $\mu\text{m}$ ), and further dried at 50 °C in the oven for 6 h to constant weight. The powdered sample accurately weighed 0.2 g was extracted with 25 mL of methanol in a round-bottomed flask and the mixture was heated under reflux for 0.5 h at 70–75 °C, and cooled at room temperature. Methanol was added to compensate for the lost weight. The extracted solution was centrifuged at 12,000 rpm for 10 min, after filtration through a 0.45  $\mu\text{m}$  membrane, an aliquot of 10  $\mu\text{L}$  of the filtrate was injected into the UHPLC-MS system for LC-MS analysis.

### 3.4. Preparation of Standard Solutions

The reference standards were accurately weighed and dissolved in methanol to prepare stock solutions. All standards were completely dissolved in the mixed standard working solution of 8  $\mu\text{g}/\text{mL}$  for gallic acid, 2.75  $\mu\text{g}/\text{mL}$  for EM, 210  $\mu\text{g}/\text{mL}$  for THSG, 228  $\mu\text{g}/\text{mL}$  for EMG, respectively. For the construction of calibration plots, the standard stock solution was further diluted with methanol to make seven different concentrations at 1/2, 1/4, 1/8, 1/16, 1/32, 1/64, and 1/128 of the working solutions. All solutions were stored in a refrigerator at 4 °C for analysis.

### 3.5. Statistical Analysis

The mass data acquired were imported into the Xcalibur software (version 2.1) (Thermo Fisher Scientific, CA, USA) for peak detection and alignment. Prism 5.0 was used to run two-way ANOVAs with Bonferroni corrections on ion peak areas of twelve investigated components (sucrose, gallic acid, procyanidin B, catechin, 6-methoxyl-2-acetyl-3-methyl-juglone-8-O-glu, tetrahydroxy stilbene-O-di-glu, THSG, citreorosein-O-glu, torachryson-O-glu, emodin-8-O-glu, emodin-O-(malonyl)-glu and emodin) to test for differences between PM and its processed products. When the peak areas of investigated compounds were difficult to integrate or not detected in the samples, the values of such data was considered to be zero.

## 4. Conclusions

In this study, a simple, rapid and accurate UHPLC-MS method was established to qualitatively and quantitatively determine the major components of PM and PPM. The method was used to successfully quantify four components in twenty batches of PM and PPM samples. This novel approach is a highly useful technique to identify constituents and control the quality of CPM and PPM, it also offers incredible advantages, including speed, simplicity, and a reduction in solvent consumption. The method would become an important quality control technique for Chinese medicine and could be adopted widely.

**Supplementary Materials:** Supplementary materials can be accessed at: <http://www.mdpi.com/1420-3049/20/12/19813/s1>.

**Acknowledgments:** This research was supported by the National Natural Science Foundations of China (81073052, 81373967), the Funds for Distinguished Young Scholars from Guangzhou University of Chinese Medicine (No.10) and supported by Science and Technology Planning Project of Guangdong Province, China (2013B021800236).

**Author Contributions:** Wen Xu and You-Heng Gao conceived and designed the experiments; Teng-Hua Wang performed the experiments; Wen Xu and Teng-Hua Wang analyzed the data; Jing Zhang, Xiao-Hui Qiu, Jun-Oi Bai and You-Heng Gao contributed reagents/materials/analysis tools; Wen Xu and Teng-Hua Wang wrote the paper.

**Conflicts of Interest:** The authors declare that there are no conflicts of interest.

## References

1. Wang, H.Y.; Song, L.X.; Feng, S.B.; Liu, Y.C.; Zuo, G.; Lai, F.L.; He, G.Y.; Chen, M.J.; Huang, D. Characterization of proanthocyanidins in stems of *Polygonum multiflorum* Thunb. as strong starch hydrolase inhibitors. *Molecules* **2014**, *18*, 2255–2265. [[CrossRef](#)] [[PubMed](#)]
2. Sun, Y.N.; Li, W.; Kim, J.H. Chemical constituents from the root of *Polygonum multiflorum* and their soluble epoxide hydrolase inhibitory activity. *Arch. Pharm. Res.* **2015**, *38*, 998–1004. [[CrossRef](#)] [[PubMed](#)]
3. Li, L.F.; Ni, B.R.; Lin, H.M.; Zhang, M. Traditional usages, botany, phytochemistry and toxicology of *Polygonum multiflorum* Thunb.: A review. *J. Ethnopharmacol.* **2015**, *159*, 158–183.
4. Choi, S.G.; Kim, J.; Sung, N.D.; Son, K.H.; Cheon, H.G.; Kim, K.R.; Kwon, B.M. Anthraquinones, Cdc25B phosphatase inhibitors, isolated from the roots of *Polygonum multiflorum* Thunb. *Nat. Prod. Res.* **2007**, *6*, 487–493. [[CrossRef](#)] [[PubMed](#)]
5. Yan, S.Y.; Su, Y.F.; Chen, L. Polygonumosides A–D, Stilbene Derivatives from Processed Roots of *Polygonum multiflorum*. *J. Nat. Prod.* **2014**, *77*, 397–401. [[CrossRef](#)] [[PubMed](#)]
6. Han, J.W.; Shim, D.W.; Shin, W.Y. Anti-inflammatory effect of emodin via attenuation of NLRP3 inflammasome activation. *Int. J. Mol. Sci.* **2015**, *4*, 8102–8109. [[CrossRef](#)] [[PubMed](#)]
7. Srinivas, G.; Anto, R.J.; Srinivas, P.; Vidhyalakshmi, S. Emodin induces apoptosis of human cervical cancer cells through poly (ADP-ribose) polymerase cleavage and activation of caspase-9. *Eur. J. Pharmacol.* **2003**, *473*, 117–125. [[CrossRef](#)]
8. Thiruvengadam, M.; Praveen, N.; Kim, E.H.; Kim, S.H.; Chung, I.M. Production of anthraquinones, phenolic compounds and biological activities from hairy root cultures of *Polygonum multiflorum* Thunb. *Protoplasma* **2014**, *251*, 555–566. [[CrossRef](#)] [[PubMed](#)]
9. Chan, Y.C.; Wang, M.F.; Chen, Y.C.; Yang, D.Y.; Lee, M.S.; Cheng, F.C. Long-term administration of *Polygonum multiflorum* Thunb. reduces cerebral ischemia-induced infarct volume in gerbils. *Am. J. Chin. Med.* **2003**, *31*, 71–77. [[CrossRef](#)] [[PubMed](#)]
10. Zhang, Y.Z.; Shen, J.F.; Xu, J.Y.; Xiao, J.H.; Wang, J.L. Inhibitory effects of 2,3,5,4'-tetrahydroxystilbene-2-O- $\beta$ -D-glucoside on experimental inflammation and cyclo-oxygenase 2 activity. *J. Asian Nat. Prod. Res.* **2007**, *9*, 355–363. [[CrossRef](#)] [[PubMed](#)]
11. Wang, X.M.; Zhao, L.B.; Han, T.Z.; Wang, J.L. Protective effects of 2,3,5,4'-tetrahydroxystilbene-2-O- $\beta$ -D-glucoside, an active component of *Polygonum multiflorum* Thunb. on experimental colitis in mice. *Eur. J. Pharmacol.* **2008**, *578*, 339–348. [[CrossRef](#)] [[PubMed](#)]
12. Zeng, C.; Xiao, J.H.; Chang, M.J.; Wang, J.L. Beneficial effects of THSG on acetic acid-induced experimental Colitis: Involvement of upregulation of PPAR- $\gamma$  and inhibition of the NF- $\kappa$ B Inflammatory Pathway. *Molecules* **2011**, *16*, 8552–8568. [[CrossRef](#)] [[PubMed](#)]
13. Lv, L.S.; Gu, X.H.; Tang, J.; Ho, C.T.; Tang, J. Stilbene glycoside from the roots of *Polygonum multiflorum* Thunb. and their *in vitro* antioxidant activities. *J. Food Lipids* **2006**, *13*, 131–144.
14. Lv, L.S.; Gu, X.H.; Tang, J.; Ho, C.T. Antioxidant activity of stilbene glycoside from *Polygonum multiflorum* Thunb *in vivo*. *Food Chem.* **2007**, *104*, 1678–1681. [[CrossRef](#)]
15. Liu, Q.L.; Xiao, J.H.; Ma, R.; Ban, Y.; Wang, J.L. Effect of 2,3,5,4'-tetrahydroxystilbene -2-O- $\beta$ -D-glucoside on lipoprotein oxidation and proliferation of coronary arterial smooth cells. *J. Asian Nat. Prod. Res.* **2007**, *9*, 689–697. [[CrossRef](#)] [[PubMed](#)]
16. Chiang, Y.C.; Huang, G.H.; Ho, Y.L.; Hsieh, P.C.; Chung, H.P.; Chou, F.I.; Chang, Y.S. Influence of gamma irradiation on microbial load and antioxidative characteristics of *Polygoni multiflori*. Radix. *Process. Biochem.* **2011**, *46*, 777–782. [[CrossRef](#)]
17. Xie, W.D.; Zhao, Y.N.; Du, L.J. Emerging approaches of traditional Chinese medicine formulas for the treatment of hyperlipidemia. *J. Ethnopharmacol.* **2012**, *140*, 345–367. [[CrossRef](#)] [[PubMed](#)]
18. Cheung, F.W.; Leung, A.W.; Liu, W.K.; Che, C.T. Tyrosinase inhibitory activity of a glucosylated hydroxystilbene in mouse melan—A melanocytes. *J. Nat. Prod.* **2014**, *77*, 1270–1274. [[CrossRef](#)] [[PubMed](#)]

19. Um, M.Y.; Choi, W.H.; Aan, J.Y.; Kim, S.R.; Ha, T.Y. Protective effect of *Polygonum multiflorum* Thunb. on amyloid  $\beta$ -peptide 25–35 induced cognitive deficits in mice. *J. Ethnopharmacol.* **2006**, *104*, 144–148. [[CrossRef](#)]
20. Wang, R.; Tang, Y.; Feng, B.; Ye, C.; Fang, L.; Zhang, L.; Li, L. Changes in hippocampal synapses and learning-memory abilities in age-increasing rats and effects of tetrahydroxystilbene glucoside in aged rats. *Neuroscience* **2007**, *149*, 739–746. [[CrossRef](#)] [[PubMed](#)]
21. Luo, A.X.; Fan, Y.J.; Luo, A.S. *In vitro* free radicals scavenging activities of polysaccharide from *Polygonum multiflorum* Thunb. *J. Med. Plants Res.* **2011**, *5*, 966–972.
22. Guan, S.Y.; Su, W.W.; Wang, N.; Li, P.B.; Wang, Y.G. A potent tyrosinase activator from *Radix Polygoni multiflori* and its melanogenesis stimulatory effect in B16 melanoma cells. *Phytother. Res.* **2008**, *22*, 660–663. [[CrossRef](#)] [[PubMed](#)]
23. Jiang, Z.Q.; Xu, J.M.; Long, M.H.; Tu, Z.M.; Yang, G.X.; He, G.Y. 2,3,5,4'-Tetrahydroxystilbene-2-O- $\beta$ -D-glucoside (THSG) induces melanogenesis in B16 cells by MAP kinase activation and tyrosinase upregulation. *Life Sci.* **2009**, *85*, 345–350. [[CrossRef](#)] [[PubMed](#)]
24. Park, H.J.; Zhang, N.N.; Park, D.K. Topical application of *Polygonum multiflorum* extract induces hair growth of resting hair follicles through upregulating Shh and  $\beta$ -catenin expression in C57BL/6 mice. *J. Ethnopharmacol.* **2011**, *135*, 369–375. [[CrossRef](#)] [[PubMed](#)]
25. Li, S.; Zhao, S.J.; Cui, T.B.; Liu, Z.Y.; Zhao, W. 2,3,5,4'-Tetrahydroxystilbene-2-O- $\beta$ -D-glycoside biosynthesis by suspension cells cultures of *Polygonum multiflorum* Thunb. and production enhancement by methyl jasmonate and salicylic acid. *Molecules* **2012**, *17*, 2240–2247.
26. Han, L.F.; Wu, B.; Pan, G.X.; Wang, Y.F.; Song, X.B.; Gao, X.M. UPLC-PDA analysis for simultaneous quantification of four active compounds in crude and processed rhizome of *Polygonum multiflorum* Thunb. *Chromatographia* **2009**, *70*, 657–659. [[CrossRef](#)]
27. Yan, H.J.; Fang, Z.J.; Fu, J.; Yu, S.X. The correlation between bioactive components of *Fallopia multiflora* root and environmental factors. *Am. J. Chin. Med.* **2010**, *38*, 473–483. [[CrossRef](#)] [[PubMed](#)]
28. Zhang, F.; Chen, W.S.; Sun, L.N. LC-VWD-MS determination of three anthraquinones and one stilbene in the quality control of crude and prepared roots of *Polygonum multiflorum* Thunb. *Chromatographia* **2008**, *67*, 869–874. [[CrossRef](#)]
29. Lin, Z.L.; Liu, Y.Y.; Wang, C.; Guo, N.; Song, Z.Q.; Wang, C.; Xin, L.; Lu, A.P. Comparative analyses of chromatographic fingerprints of the roots of *Polygonum multiflorum* Thunb. and their processed products using RRLC/DAD/ESI-MS<sup>n</sup>. *Planta. Med.* **2011**, *77*, 1855–1860.
30. Yi, T.; Leung, K.S.Y.; Lu, G.H.; Zhang, H.; Chan, K. Identification and determination of the major constituents in traditional Chinese medicinal plant *Polygonum multiflorum* Thunb. by HPLC coupled with PAD and ESI/MS. *Phytochem. Anal.* **2007**, *18*, 181–187. [[CrossRef](#)] [[PubMed](#)]
31. Xu, W.; Zhang, J.; Huang, Z.H.; Qiu, X.H. Identification of new dianthrone glycosides from *Polygonum multiflorum* Thunb. using high-performance liquid chromatography coupled with LTQ-Orbitrap mass spectrometry detection: A strategy for the rapid detection of new low abundant metabolites from traditional Chinese medicines. *Anal. Methods* **2012**, *4*, 1806–1812.
32. Qiu, X.H.; Zhang, J.; Huang, Z.H.; Zhu, D.Y.; Xu, W. Profiling of phenolic constituents in *Polygonum multiflorum* Thunb. by combination of ultra-high-pressure liquid chromatography with linear ion trap-Orbitrap mass spectrometry. *J. Chromatogr. A* **2013**, *1292*, 121–131. [[CrossRef](#)] [[PubMed](#)]
33. Liu, Z.L.; Song, Z.Q.; Wang, C. Content variances of catechin and gallic acid in *Polygonum multiflorum* after steaming. *Chin. Tradit. Pat. Med.* **2009**, *31*, 1392–1394.
34. Liang, Z.T.; Chen, H.B.; Yu, Z.L. Comparison of raw and processed *Radix Polygoni multiflori* (Heshouwu) by HPLC-MS. *Chin. Med.* **2010**, *5*, 29–32. [[CrossRef](#)] [[PubMed](#)]

**Sample Availability:** Samples of the compounds gallic acid (GA), *trans*-2,3,5,4'-tetra-hydroxystilbene-2-O- $\beta$ -D-glucopyranoside (THSG), emodin-8-O- $\beta$ -D-glucopyranoside (EMG), emodin (EM) and 20 batches CPM and PPM (crude *P. multiflorum* and processed *P. multiflorum*) are available from the authors.



© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).