

BMJ Open Mutation screening for thalassaemia in the Jino ethnic minority population of Yunnan Province, Southwest China

Shiyun Wang,¹ Rong Zhang,¹ Guangxin Xiang,² Yang Li,² Xuhong Hou,¹ Fusong Jiang,¹ Feng Jiang,¹ Cheng Hu,¹ Weiping Jia¹

To cite: Wang S, Zhang R, Xiang G, *et al.* Mutation screening for thalassaemia in the Jino ethnic minority population of Yunnan Province, Southwest China. *BMJ Open* 2015;**5**:e010047. doi:10.1136/bmjopen-2015-010047

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2015-010047>).

Received 20 September 2015
Revised 4 November 2015
Accepted 2 December 2015



CrossMark

¹Shanghai Diabetes Institute, Shanghai Key Laboratory of Diabetes Mellitus, Shanghai Clinical Center for Diabetes, Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai, People's Republic of China
²National Engineering Research Center for Beijing Biochip Technology, Beijing, People's Republic of China

Correspondence to Professor Weiping Jia; wjia@sjtu.edu.cn

ABSTRACT

Objectives: This study aimed to detect α - and β -thalassaemia mutations in the Jino ethnic minority population of Yunnan Province, Southwest China.

Design: A total of 1613 Jino adults were continuously recruited from February 2012 to April 2012. Fasting venous blood samples were obtained to determine haematological variables. Haemoglobin analysis was conducted using high-performance liquid chromatography. Participants with hypochromic microcytic anaemia or positive haemoglobin analysis profiles were confirmed by α - and β -globin genetic testing, including DNA microarray analysis, direct sequencing methods and multiplex gap-PCR assays.

Setting: Shanghai Diabetes Institute, Shanghai Key Laboratory of Diabetes Mellitus, Shanghai Jiao Tong University Affiliated Sixth People's Hospital.

Results: We found 363 suspected cases by primary screening of haematological variables and haemoglobin analysis. After further genetic testing, four types of α - and β -thalassaemia mutation were detected in 203 out of 363 individuals. Both α^0 - and α^+ -thalassaemia mutations, $-^{SEA}$ and $-\alpha^{3,7}$, were identified. β -Thalassaemia mutations included CD17 (HBB:c.52A>T) and CD26 (HbE or HBB:c.79G>A). In addition, 13 HbE carriers had coexisting α^0 - or α^+ -thalassaemia deletions. Clinical haematological variables indicated that, in this study, carriers of all thalassaemic genotypes had more severe hypochromic microcytic anaemia than non-thalassaemic individuals.

Conclusions: Our results provide information on the Jino ethnic minority that may be useful for further genetic counselling, prenatal screening and clinical diagnosis of thalassaemia in this region.

INTRODUCTION

As a group of monogenic disorders, thalassaemia is a serious health problem worldwide, especially in Mediterranean areas, Southeast Asia and Southern China.^{1–3} Yunnan Province, which is located along the border areas of China–Myanmar–Laos, is notable for its ethnic diversity. According to a previous study of children under 10 years

Strengths and limitations of this study

- As Jino, the last ethnic minority confirmed in China, was reported to have a high prevalence of thalassaemia according to previous research on children under 10 years of age, this study aimed to detect mutations of α - and β -thalassaemia in Jino adults.
- The α - and β -thalassaemia mutation spectrum shown in this research may help to explain further genotype–phenotype correlations and to establish a thalassaemia-prevention programme in this area.
- The sample size we used in genetic testing was relatively small and may not have the validity to identify the rare thalassaemias in this ethnic group.

of age, several ethnic minorities in this region have a high prevalence of thalassaemia, with the prevalence of α -thalassaemia (α -thal) being highest (22.1%) in Dai from Xishuangbanna and the prevalence of β -thalassaemia (β -thal) being highest in Achang (40.6%).⁴

Jino is the last ethnic minority confirmed in China, and the prevalence of α -thal and β -thal among Jino children are 3.1% and 29.3%, respectively. Thalassaemic children may exhibit various clinical symptoms; some are asymptomatic carriers, whereas others have severe haemolytic anaemia.⁵ Blood transfusion therapy, which is needed for severe carriers, imposes a heavy burden on families and public health management.⁶ Although genetic screening is essential to prevent and control this inherited disease, systematic investigations of thalassaemia mutations in Jino adults are rare.

The Jino population comprises nearly 20 000 individuals, and most (~90%) live around Jino Mountain, which is located in East–Central Yunnan Province.⁷ A large number of thalassaemic mutations have been found in the general population worldwide;^{8–10} however,

little is known about this isolated population. Indeed, the molecular mechanism and genetic variations of thalassaemia in Jino individuals may be different from those in other ethnicities. Our study aimed to detect α -thal and β -thal gene mutations in Jino adults to provide basic information for further prenatal consulting and thalassaemia diagnosis.

MATERIALS AND METHODS

Participants and clinical screening

Ethics approval for the study was granted according to the Declaration of Helsinki (paragraph II) by the institutional review board of Shanghai Jiao Tong University affiliated with the Sixth People's Hospital, Shanghai, China. This cross-sectional study was conducted between February 2012 and April 2012 in eight villages (Luote, Jiama, Balai, Situ, New Situ, Baka, Baya, Dapingzhang) around Jino Mountain in Jinghong, Southern Yunnan Province, China (figure 1). A list of Jino adults from these eight villages was obtained from local villager committee offices. Participants were sampled by a simple computer programme of randomisation from these villages. Staff at the local health centre, who understand both Chinese and Jino languages, contacted the subjects and introduced the purpose of the study. Oral and written informed consent was obtained from all the individuals. Basic demographic information and fasting venous blood samples were collected by researchers.

A total of 1613 Jino adults, including 762 men and 851 women, participated in this survey (haematological and demographic characteristics of the total population included in the study are given in table 1). The following haematological variables were measured: haemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), red blood cell (RBC) and red cell distribution width (RDW). Haemoglobin was analysed by high-performance liquid chromatography (HPLC) using the Variant II haemoglobin analysing system (Bio-Rad Laboratories, Hercules, California, USA). α - and β -globin genetic testing was performed in participants (n=363) with hypochromic microcytic anaemia (MCV <80fL and/or MCH <27 pg) and/or positive HPLC profiles. In order to evaluate the validity of primary screening approaches for detection of thalassaemia carriers, we randomly selected some of the individuals with negative screening results (n=50) from the remaining participants (n=1250) for further genetic testing.

Genetic testing

Genomic DNA was extracted from venous blood leucocytes. Three methods were used to detect thalassaemic mutations.

A CapitalBio Thalassaemia Gene Mutation Detection Kit (CapitalBio, Beijing, China) was used to determine 25 common mutations in globin genes in the Chinese population via DNA microarray. Six α -thal gene




Figure 1 Geographical location of the Jino ethnic minority populations in Yunnan Province, southwest China. The solid black triangle represents Jino Mountain. The solid black circles represent the eight villages in Jinghong where the 1613 subjects were randomly selected.

Table 1 Haematological and demographic characteristics of 1613 Jino ethnic minority adults included in the study

Variable	Total	Male	Female
Samples (n)	1613	762	851
Age (years)	40.43±14.78	40.11±15.21	40.71±14.38
BMI (kg/m ²)	21.79±3.20	22.12±3.17	21.50±3.21
RBC (10 ¹² /L)	4.85±0.56	5.08±0.57	4.65±0.47
RDW (%)	12.76±1.29	12.60±1.10	12.89±1.42
MCV (fL)	83.42±8.08	84.97±7.54	82.03±8.30
MCH (pg)	29.21±3.26	29.91±3.07	28.59±3.30
HCT (%)	40.25±4.36	42.88±3.74	37.90±3.44
Hb (g/dL)	14.09±1.70	15.09±1.49	13.20±1.34

Data are shown as mean±SD.

BMI, body mass index; HCT, haematocrit; Hb, haemoglobin; MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume; RBC, red blood cell; RDW, red cell distribution width.

mutations and 19 β -thal gene mutations were included. Among them, there were three α -thal deletions—that is, the Southeast Asian deletion ($-\text{SEA}$), rightward deletion ($-\alpha^{3.7}$) and leftward deletion ($-\alpha^{4.2}$)—and three non-deletional α -thal mutations—that is, Hb Constant Spring (HBA2:c.427T>C), Hb Quong Sze (HBA2:c.377T>C or HBA1) and Hb Westmead (HBA2:c.369C>G). Nineteen β -thal gene mutations were CD14/15 (HBB:c.45_46insG), CD27/28 (HBB:c.84_85insC), CD41/42 (HBB:c.126_129delTCTT), CD71/72 (HBB:c.216_217insA), -32 (HBB:c.-82T>C), -30 (HBB:c.-80T>C), -29 (HBB:c.-79A>G), -28 (HBB:c.-78A>G), CD17 (HBB:c.52A>T), CD26 (HBB:c.79G>A), CD30 (HBB:c.91A>G), CD37 (HBB:c.113G>A), CD43 (HBB:c.130G>T), IVS1-1 (HBB:c.92+1G>T), IVS1-5 (HBB:c.92+5G>T), IVS2-5 (HBB:c.315+5G>C), IVS2-654 (HBB:c.316-197C>T), Int (HBB:c.2T>G), CAP (HBB:c.-11_-8delAAAC). A BioMixer II Microarray Hybridisation Station (CapitalBio) was used for hybridisation after multiplex PCR amplification. Then, chips were scanned using a LuxScan 10K-B Microarray Scanner (CapitalBio).

To validate β -thal mutations, three fragments of the β -globin gene were amplified. The first fragment was amplified with 5'-CCT AAG CCA GTG CCA GAA GAG C-3' as the forward primer and 5'-TGC CCA GTT TCT ATT GGT CTC C-3' as the reverse primer, the second fragment was amplified with 5'-TAG AAA CTG GGC

ATG TGG AG-3' as the forward primer and 5'-TGT ACC CTG TTA CTT ATC CC-3' as the reverse primer, and the third fragment was amplified with 5'-TCA GGG CAA TAA TGA TAC AA-3' as the forward primer and 5'-TTA GTA GTT GGA CTT AGG GA-3' as the reverse primer. The fragments were sequenced using a 3500 Genetic Analyser (Applied Biosystems, Foster City, California, USA).

We also confirmed three α -thal deletions—that is, the Southeast Asian deletion ($-\text{SEA}$), rightward deletion ($-\alpha^{3.7}$) and leftward deletion ($-\alpha^{4.2}$)—using multiplex gap-PCR assays. Primers and PCR conditions were designed as described in the literature.^{11 12}

Statistical analysis

A statistical analysis was carried out using SAS for Windows (V.9.2). All quantitative traits were tested for normality, and skewed quantitative traits were logarithmically transformed to approximate univariate normality. Data are shown as means±SD. Quantitative traits (RBC, Hb, MCV, MCH, RDW) were compared between two groups using the Wilcoxon test, and analysis of variance was performed to compare the differences in the three subgroups of thalassaemia carriers (α -Thal, β -Thal and $\alpha\beta$ -Thal). Two-tailed statistical significance was considered at $p<0.05$.

RESULTS

Mutations identified in Jino

Owing to mutations in different globin genes, we observed three groups of thalassaemic carriers, including individuals with only α -thal gene deletions or β -thal gene mutations and individuals with combined $\alpha\beta$ -thalassaemia ($\alpha\beta$ -thal) gene mutations. Four different thalassaemia mutations were detected in 203 individuals among 363 suspected cases. No mutations were observed in 50 individuals with negative primary screening results. Table 2 shows the allele frequency of α - and β -thal mutations found in our study.

Mutations in the α -thal gene


None of the three common non-deletional α^+ -thal mutations, Hb Constant Spring (HBA2:c.427T>C), Hb Quong

Table 2 Allele frequency of α - and β -thalassaemia mutations found in our study

Mutation	Phenotype	n	Number of alleles	Allele frequency (ratio)
α -Thalassaemia				
$-\text{SEA}$	α^0/α	42	42	70.00% (42/60)
$-\alpha^{3.7}$	α^+/α	16	16	30.00% (16/60)
$-\alpha^{3.7}/-\alpha^{3.7}$	α^+/α^+	1	2	–
β -Thalassaemia				
CD17	β^0/β^A	20	20	12.35% (20/162)
HbE	β^E/β^A	132	132	87.65% (142/162)
HbE/HbE	β^E/β^E	5	10	–

α , the normal α -globin chain; α^0 , the α -globin chain is totally deleted; α^+ , the α -globin chain is partly deleted; β^A , the normal β -globin chain; β^E , the abnormal β -globin chain of HbE mutation; β^0 , the β -globin chain is totally deleted.

Figure 2 Gel electrophoresis of PCR amplifying results in α -thal deletions. M, marker, 200 bp DNA Ladder; lane 1, rightward deletion (genotype of $-\alpha^{3.7}/-\alpha^{3.7}$); lanes 2 and 3, rightward deletion (genotype of $-\alpha^{3.7}/\alpha\alpha$); lanes 4 and 5, Southeast Asia deletion (genotype of $--^{SEA}/\alpha\alpha$).



Sze (HBA2:c.377T>C (or HBA1)) and Hb Westmead (HBA2:c.369C>G), were found in 203 participants with thalassaemia mutations. Forty-six of 203 participants carried α -thal deletions only; $-\text{SEA}$ and $-\alpha^{3.7}$ were observed, accounting for 16.7% (34/203) and 5.9% (12/203) of the mutations, respectively. Among these individuals, we identified both α^+/α and α^+/α^+ for $-\alpha^{3.7}$ and α^0/α for $-\text{SEA}$ (gel electrophoresis of PCR amplifying results are shown in figure 2). However, no $-\alpha^{4.2}$ deletion was observed.

Mutations in the β -thal gene

We observed mutations in CD17 (HBB:c.52A>T) (figure 3) and CD26 (HbE or HBB:c.79G>A) (figure 4). CD17,

which accounted for 9.9% (20/203) of mutations, was found to be β^0/β^A in this population. Participants with HbE variant only, either β^E/β^A or β^E/β^E , accounted for 61.1% (124/203) of mutations. Furthermore, 13 HbE carriers harboured $-\text{SEA}$ (n=8) or $-\alpha^{3.7}$ (n=5) at a combined frequency of 6.4% (13/203).

Haematological features of different thalassaemia genotypes

The haematological data of different thalassaemia genotypes are summarised in table 3. Compared with normal individuals, thalassaemic carriers had significantly lower Hb, MCV and MCH levels ($p<0.001$, respectively) and higher RBC and RDW levels ($p<0.001$, respectively).

Figure 3 Heterozygous CD17 (A>T) mutation (A) and the corresponding normal sequence of β -globin. Red arrow indicates the position of this point mutation.





Figure 4 Heterozygous CD26 (G>A) mutation (A) and the corresponding normal sequence of β -globin. Red arrow indicates the position of this point mutation.



Furthermore, we compared the differences in these five indexes among the three groups of carriers (α -thal, β -thal and $\alpha\beta$ -thal). Significant differences in MCV ($p=0.0111$), MCH ($p=0.0002$) and RBC ($p=0.0012$) were observed between these groups. MCV and MCH levels in

the α -thal group were significantly lower than those in the β -thal group ($p<0.05$), whereas RBC levels were higher ($p<0.05$). In contrast, no difference was observed between the $\alpha\beta$ -thal and α -thal groups/ β -thal groups. Moreover, there was a tendency towards increased RDW

Table 3 Haematological data of 1613 Jino ethnic minority individuals with different thalassaemia subtypes

Thalassaemia type	n	RBC ($10^{12}/L$)	Hb (g/dL)	MCV (fL)	MCH (pg)	RDW (%)
α -Thalassaemia	46	5.63 \pm 0.78	13.03 \pm 1.61	68.99 \pm 6.50	23.33 \pm 2.45	13.64 \pm 1.38
α^0/α	34	5.88 \pm 0.68	12.98 \pm 1.62	65.69 \pm 2.65	22.08 \pm 0.81	14.01 \pm 1.37
α^+/α	11	4.86 \pm 0.58	13.19 \pm 1.70	78.96 \pm 4.43	27.14 \pm 1.87	12.52 \pm 0.74
α^+/α^+	1	5.34	12.7	71.5	23.8	13.1
β -Thalassaemia	144	5.27 \pm 0.48	13.31 \pm 1.70	72.46 \pm 7.09	25.29 \pm 2.75	13.21 \pm 0.96
β^0/β^A	20	5.41 \pm 0.38	10.68 \pm 0.53	58.70 \pm 2.31	19.77 \pm 0.83	14.23 \pm 0.40
β^E/β^A	120	5.23 \pm 0.48	13.77 \pm 1.41	75.17 \pm 3.88	26.34 \pm 1.49	12.98 \pm 0.85
β^E/β^E	4	5.91 \pm 0.51	12.55 \pm 1.22	60.23 \pm 3.61	21.23 \pm 1.15	15.00 \pm 0.57
$\alpha\beta$ -Thalassaemia	13	5.67 \pm 0.70	13.63 \pm 1.67	69.40 \pm 6.85	24.17 \pm 2.59	13.12 \pm 0.92
β^E/β^A with α^0/α	7	6.02 \pm 0.69	13.57 \pm 1.58	65.94 \pm 2.82	22.57 \pm 1.07	13.66 \pm 0.71
β^E/β^A with α^+/α	5	5.21 \pm 0.53	14.08 \pm 1.88	76.46 \pm 3.86	27.00 \pm 1.33	12.22 \pm 0.36
β^E/β^E with β^0/β^A	1	5.57	13.8	58.3	21.2	11.8
Total thalassaemia	203	5.38 \pm 0.60	13.27 \pm 1.68	71.48 \pm 7.08	24.77 \pm 2.79	13.30 \pm 1.08
Non-thalassaemia	1410	4.78 \pm 0.51	14.21 \pm 1.67	85.14 \pm 6.64	29.85 \pm 2.79	12.68 \pm 1.30
p Value*		<0.001	<0.001	<0.001	<0.001	<0.001
p Value†		0.0012	>0.05	0.0111	0.0002	0.0573

Data are shown as n, mean \pm SD, or raw data when necessary.

*Non-thalassaemic individuals compared with thalassaemia group.

†Compared among three subgroups of thalassaemia (α -Thal, β -Thal and $\alpha\beta$ -Thal).

Hb, haemoglobin; MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume; RBC, red blood cell; RDW, red cell distribution width.

levels in the α -thal group compared with the β -thal group ($p=0.0573$).

DISCUSSION

Thalassaemia is a common monogenic disease with a relatively high prevalence in Southeast Asia. In China, this disease is mainly prevalent in areas near the southern bank of the Yangtze River, such as Guangdong, Guangxi, Fujian and Yunnan Provinces.^{13–15} Prenatal screening and related molecular diagnoses are crucial for preventing and treating thalassaemia. Many thalassaemia studies have been conducted in Yunnan Province.^{16–17} However, data on the Jino population are limited because this population is the last ethnic minority confirmed in China.

We randomly selected 1613 Jino adults from eight villages around Jino Mountain in Jinghong, Southern Yunnan. Among the gene mutations identified, the most prevalent α -thal and β -thal genotypes in this region were $-\text{SEA}$ and HbE, in agreement with previous data from Yunnan Province.^{18–19} According to our results, the overall prevalence of thalassaemia in Jino was nearly 12.6%, which is similar to the prevalence observed in Kunming.²⁰ Prevalence of $\alpha\beta$ -thal (8%) in our population was equal to that in the Li population in Hainan Province (7.99%), where thalassaemia prevalence is high.²¹ Although Yunnan Province has a high prevalence of thalassaemia with diverse genotypes, the spectrum of globin gene mutations among the Jino population is relatively limited.

HbE, a type of haemoglobinopathy, can be observed in most regions of Southeast China.²² Due to a point mutation in the β -globin gene, the balance of various globin products is disrupted, leading to a structural haemoglobin variant. Although HbE carriers may only have slight anaemia, their offspring will exhibit severe clinical symptoms in the presence of other β -thal types.²³ Therefore, potential HbE carriers should undergo genetic testing and prenatal counselling. HPLC is often used as an efficient primary screening method to detect abnormal Hb, as was carried out in this study and previous studies.^{24–25} In our study, 95.6% (131/137) of HbE carriers were identified by HPLC, and 13 of these individuals had concomitant α -thal deletions.

Different genotypes lead to different clinical phenotypes.²⁶ We found that thalassaemic carriers had significantly lower MCV and MCH levels. Regarding those with β -thal mutations, MCV and MCH levels were significantly decreased in CD17 carriers compared with HbE carriers, suggesting that a nonsense mutation in the β -globin gene causes greater erythrocyte impairment. Hypochromic microcytic anaemia was moderate in individuals with β^E/β^A and α^+/α compared with β^E/β^A carriers. This paradox may be explained by the fact that changes in the α - and β -globin chains may balance each other out when both mutations coexist in an individual. Accordingly, rapidly estimating the genetic state of an

illness based on haematological variables is difficult. Therefore, genetic screening of both α - and β -globin gene mutations in potential parents is of utmost importance to prevent births with severe defects.²⁷

However, there are some limitations in this study. First, the sample size we used in the genetic testing was relatively small and may not have the validity to identify the rare thalassaemia variants from this ethnic group. Second, investigations of population and family structure were not performed in this study, although α -thal and β -thal gene mutations were common among the Jino ethnic minority. As a result, further studies on the inbreeding levels and consanguinity structure are warranted to reveal the underlying mechanism of gene flow and then assess the occurrence and persistence of α -thal and β -thal gene mutations, especially coexisting α -thal and β -thal gene mutations within Jino individuals.

In conclusion, this study revealed α - and β -thal mutations in the Jino ethnic minority population in Yunnan Province. Of these mutations, $-\text{SEA}$ and HbE were the most prevalent α -thal and β -thal gene mutation types, respectively. In addition, data based on clinical haematological variable analysis indicated that the severity of hypochromic microcytic anaemia is associated with the genotype of thalassaemia. Our results provide evidence that may be useful for further genetic counselling, prenatal screening and clinical diagnosis of thalassaemia in this region.

Acknowledgements We thank all of the participants for their dedication. We acknowledge the skilful technical support of all nursing and medical staff at the Shanghai Clinical Center for Diabetes.

Contributors WJ and CH conceived and designed the experiments. SW and RZ performed the experiments and analysed the data. GX, YL, XH, FuJ and FeJ contributed materials and analysis tools. SW prepared the article. CH and WJ revised the manuscript. All the authors have read and approved the final version of this manuscript.

Funding This work was financially supported by a grant from the National Natural Science Foundation of China (91331110).

Competing interests None declared.

Patient consent Obtained.

Ethics approval According to the Declaration of Helsinki (paragraph II), ethics approval for the study was granted by the institutional review board of Shanghai Jiao Tong University affiliated with the Sixth People's Hospital, Shanghai, China. Written and Oral informed consent was obtained from all participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Weatherall DJ. Thalassaemia as a global health problem: recent progress toward its control in the developing countries. *Ann N Y Acad Sci* 2010;1202:17–23.

2. Fucharoen S, Winichagoon P. Thalassemia in SouthEast Asia: problems and strategy for prevention and control. *Southeast Asian J Trop Med Public Health* 1992;23:647–55.
3. Li B, Zhang XZ, Yin AH, *et al.* High prevalence of thalassemia in migrant populations in Guangdong Province, China. *BMC Public Health* 2014;14:905.
4. Yao LQ, Zou TB, Yang FB, *et al.* [Epidemiological study of thalassaemia among children in Xishuangbanna, Dehong and Nujiang of Yunnan province]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2011;28:579–82.
5. Sankaran VG, Weiss MJ. Anemia: progress in molecular mechanisms and therapies. *Nat Med* 2015;21:221–30.
6. Goss C, Giardina P, Degtyaryova D, *et al.* Red blood cell transfusions for thalassemia: results of a survey assessing current practice and proposal of evidence-based guidelines. *Transfusion* 2014;54:1773–81.
7. Xu JW, Liao YM, Liu H, *et al.* Use of bed nets and factors that influence bed net use among Jinuo Ethnic Minority in southern China. *PLoS ONE* 2014;9:e103780.
8. Piel FB, Weatherall DJ. The α -thalassemias. *N Engl J Med* 2014;371:1908–16.
9. Rund D, Rachmilewitz E. Beta-thalassemia. *N Engl J Med* 2005;353:1135–46.
10. Saller E, Dutly F, Frischknecht H. Two Novel alpha2 Gene Mutations Causing Altered Amino Acid Sequences Produce a Mild (Hb Kinshasa, HBA2: c.428A > T) and Severe (HBA2: c.342-345insCC) alpha-Thalassemia Phenotype. *Hemoglobin* 2015:1–3.
11. Zhou Y, Zhang Y, Li L, *et al.* [Rapid detection of three common deletion alpha thalassemias in Chinese by single-tube multiplex PCR]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2005;22:180–4.
12. Chong SS, Boehm CD, Higgs DR, *et al.* Single-tube multiplex-PCR screen for common deletion determinants of alpha-thalassemia. *Blood* 2000;95:360–2.
13. Xiong F, Sun M, Zhang X, *et al.* Molecular epidemiological survey of haemoglobinopathies in the Guangxi Zhuang Autonomous Region of southern China. *Clin Genet* 2010;78:139–48.
14. Yin A, Li B, Luo M, *et al.* The prevalence and molecular spectrum of α - and β -globin gene mutations in 14,332 families of Guangdong Province, China. *PLoS ONE* 2014;9:e89855.
15. Huang H, Xu L, Lin N, *et al.* Molecular spectrum of β -thalassemia in Fujian Province, Southeastern China. *Hemoglobin* 2013;37:343–50.
16. Zhu BS, He J, Zhang J, *et al.* [A study on gene mutation spectrums of α - and β -thalassemias in populations of Yunnan Province and the prenatal gene diagnosis]. *Zhonghua Fu Chan Ke Za Zhi* 2012;47:85–9.
17. Zou T, Yao L, Li Q, *et al.* The family-based research and genetic diagnosis of β -thal major in Dai ethnic. *Zhonghua Xue Ye Xue Za Zhi* 2014;35:260–1.
18. Zhang J, He J, Zeng XH, *et al.* Genetic heterogeneity of the β -globin gene in various geographic populations of Yunnan in Southwestern China. *PLoS ONE* 2015;10:e0122956.
19. Zhang J, Zhu BS, He J, *et al.* The spectrum of α - and β -thalassemia mutations in Yunnan Province of Southwestern China. *Hemoglobin* 2012;36:464–73.
20. Wen BP, Fan M, Dai HJ, *et al.* Biochemical screening and genetic diagnosis of thalassemia in children from Kunming. *Zhongguo Dang Dai Er Ke Za Zhi* 2011;13:104–6.
21. Yao H, Chen X, Lin L, *et al.* The spectrum of α - and β -thalassemia mutations of the Li people in Hainan Province of China. *Blood Cells Mol Dis* 2014;53:16–20.
22. Chen W, Zhang X, Shang X, *et al.* The molecular basis of beta-thalassemia intermedia in southern China: genotypic heterogeneity and phenotypic diversity. *BMC Med Genet* 2010;11:31.
23. Li YQ, Huang HP, Qin GF, *et al.* [Phenotype and genotype analysis of hemoglobin E]. *Zhonghua Xue Ye Xue Za Zhi* 2012;33:861–4.
24. Eastman JW, Wong R, Liao CL, *et al.* Automated HPLC screening of newborns for sickle cell anemia and other hemoglobinopathies. *Clin Chem* 1996;42:704–10.
25. Khera R, Singh T, Khuana N, *et al.* HPLC in characterization of hemoglobin profile in thalassemia syndromes and hemoglobinopathies: a clinicohematological correlation. *Indian J Hematol Blood Transfus* 2015;31:110–15.
26. Bozdogan ST, Yuregir OO, Buyukkurt N, *et al.* Alpha-thalassemia mutations in Adana Province, southern Turkey: genotype-phenotype correlation. *Indian J Hematol Blood Transfus* 2015;31:223–8.
27. Cao A, Kan YW. The prevention of thalassemia. *Cold Spring Harb Perspect Med* 2013;3:a011775.